Soil Survey Investigations Report No. 24

Soil Survey

Laboratory Data and

Soil survey investigations reports already published:

SSIR No. 1	Soil Survey Laboratory Methods and Procedures for Collecting Soil Samples
	Soil Survey Laboratory Data and Descriptions for Some Soils of:
SSIR No. 2	North Dakota
SSIR No. 3	Iowa
SSIR No. 4	Kansas
SSIR No. 5	Nebraska
SSIR No. 6	Arkansas, Louisiana, and Missouri
SSIR No. 7	Montana
SSIR No. 8	Wyoming
SSIR No. 9	Minnesota
SSIR No. 10	Colorado
SSIR No. 11	Oklahoma
SSIR No. 12	Puerto Rico and the Virgin Islands
SSIR No. 13	Mississippi
SSIR No. 14	Kentucky
SSIR No. 15	Tennessee
SSIR No. 16	North Carolina, South Carolina, and Georgia
SSIR No. 17	Wisconsin
SSIR No. 18	Indiana
SSIR No. 19	Illinois
SSIR No. 20	New England States
SSIR No. 21	A Toposequence of Soils in Tonalite Grus in the [illegible]
	Soil Survey Laboratory Data and Descriptions for Some Soils of:
SSIR No. 22	Alabama and Florida
SSIR No. 23	Nevada

Soil Survey Investigations Report No. 24

Soil Survey Laboratory Data and Descriptions for Some Soils of...

...CALIFORNIA

June 1973

SOIL CONSERVATION SERVICE • U.S. DEPARTMENT OF AGRICULTURE
In cooperation with
CALIFORNIA AGRICULTURAL EXPERIMENT STATION

METHODS CODE SYMBOLS

1. SAMPLE COLLECTION AND PREPARATION
   A. Field sampling
      1. Site selection
      2. Soil sampling
         a. Stony soils
         b. Marsh and swamp soils
   B. Laboratory preparation

5. ION-EXCHANGE ANALYSES (contd.)
   3. Sum of cations
      a. Acidity by $BaCl_2$-TEA, pH 8.2; bases by NH_4OAc, pH 7.0
      b. Sum of bases plus Al
   6. NH_4OAc, pH 7.0, leaching tube

6. CHEMICAL ANALYSES (contd.)
   M. Nitrate
      1. Saturation extract
         a. PDS acid colorimetry
         b. Diphenylamine
   N. Calcium
      1. Saturation extract

PREFACE

The Soil Survey Investigations Report (SSIR) series was established to preserve and make available technical information resulting from soil survey investigations. SSIR No. 1 "Soil Survey Laboratory Methods and Procedures for Collecting Soil Samples," revised April 1972, describes [illegible]

remaining reports are a collection of pedon descriptions and laboratory data for pedons from individual states and groups of states. SSIR No. 21 is a technical report on a specific study. The reports already published are listed on the inside front cover.

This report contains data from soil survey investigations in California. These investigations have been going on since 1945 but most of these data have been accumulated since 1955. Some of the data and descriptions have appeared in technical journals, in regional bulletins, in USDA technical bulletins, and in the text of published soil surveys. But most have been distributed in unpublished form to those immediately concerned and were not available to all who might use them.

This report contains all the data from the soil survey laboratories that form reasonably complete characterization of soils in California. While these data were being accumulated there were many changes in laboratory methods. Consequently, laboratory data for different soils cannot always be directly compared without allowance for the method. For data collected since 1955, however, these changes have been minor. The methods used are identified by symbol in the column headings of the data tables. The codes are identified briefly on the opposite page.

Ways of describing soils have also changed. The soil descriptions published here were prepared as working documents. The soil scientist who wrote them had no idea they would be published. Editing has been limited for the most part to that necessary for conformance to the "Soil Survey Manual," AD 18. Field estimates of texture have been retained, even though some are at variance with laboratory data, because the field estimates themselves are important data.

Many pedons are no longer representative of the soil series with which they were identified at the time of sampling. Hence, during the winter of 1972-73 all pedons for California were classified in existing series and in families and higher categories. Taxonomic criteria were applied rigorously, and a pedon was considered outside a given series if it

CLASSIFICATION INDEX

ALFISOLS

Page

AQUALFS

ALBAQUALFS

Typic Albaqualfs
Fine, montmorillonitic, mesic
Wright, taxadjunct 3

Mollic Albaqualfs
Fine, montmorillonitic, mesic
Wright, taxadjunct 5

XERALFS

DURIXERALFS

Typic Durixeralfs
Fine-loamy, mixed, thermic
Exeter 7
Monserate 9
Monserate 11
Monserate 13
Monserate, taxadjunct 15
San Joaquin, taxadjunct 17

Fine, montmorillonitic, thermic
Tuscan, taxadjunct 19

Abruptic Durixeralfs
Fine-loamy, mixed, thermic
Fiddyment, taxadjunct 21
San Joaquin, taxadjunct 23

Fine, illitic, thermic
Gloria 25

Fine, mixed, thermic
Redding, taxadjunct 27
Redding, taxadjunct 29

Fine, mixed, thermic, shallow
Fiddyment, taxadjunct 31

Fine, montmorillonitic, thermic
Monserate, taxadjunct 33

Abruptic Haplic Durixeralfs
Fine-loamy, mixed, thermic
Huichica, taxadjunct 35

Haplic Durixeralfs
Coarse-loamy, mixed, thermic
Arlington 37

Natric Durixeralfs
Coarse-loamy, mixed, thermic
Fresno, taxadjunct 39

HAPLOXERALFS

Typic Haploxeralfs
Coarse-loamy, mixed, mesic
Bonta variant 41

Coarse-loamy, mixed, thermic
Greenfield 43
Greenfield 45
Greenfield 47
Ramona, taxadjunct 49

Page

Fine-loamy, mixed, thermic
Fallbrook 51
Greenfield, taxadjunct 53
Huichica variant 55
Montpellier 57
Montpellier 59
Montpellier, taxadjunct 61
Ramona 63
Tehama, taxadjunct 65
Tehama, taxadjunct 67

Fine-silty, mixed, thermic
Unnamed 69

Lithic Haploxeralfs
Loamy-skeletal, mixed, thermic
Pardee 71
Pardee, taxadjunct 73

Loamy, mixed, thermic
Millerton, taxadjunct 75
Millerton, taxadjunct 77

Mollic Haploxeralfs
Loamy-skeletal, mixed, thermic
Livermore, taxadjunct 79

Coarse-loamy, mixed, mesic
Steinbeck, taxadjunct 81

Coarse-loamy, mixed, thermic
Ahwahnee, taxadjunct 83

Fine-loamy, mixed, mesic
Hilt 85
Hilt 87
Steinbeck 89

Fine-loamy, mixed, thermic
Fallbrook, taxadjunct 91
Pachappa, taxadjunct 93
Pleasanton 95
Pleasanton, taxadjunct 97
Rescue 99
Rescue 101
Trabuco, taxadjunct 103

Fine, mixed, mesic
Secca 105

Fine, mixed, thermic
Trabuco 107

Fine, montmorillonitic, thermic
Argonaut 109

Fine, serpentinitic, mesic
Dubakella variant 111

Natric Haploxeralfs
Coarse-loamy, mixed, thermic
Traver 113
Traver 115

Fine-loamy, mixed, thermic
Bonsall, taxadjunct 117
Redding, taxadjunct 119

Fine, mixed, thermic
San Miguel variant 121

Psammentic Haploxeralfs
Coarse-loamy, mixed, thermic
Garey 123
Garey 125

CLASSIFICATION INDEX

Page

ALFISOLS, cont'd.

XERALFS, cont'd.

HAPLOXERALFS, cont'd.

Ultic Haploxeralfs
Coarse-loamy, mixed, frigid
Jabu variant 127
Jabu variant 129
Jabu 131

Coarse-loamy, mixed, mesic
Ahwahnee, taxadjunct 133

Fine-loamy, mixed, frigid
Jabu variant 135

Fine-loamy, mixed, mesic
Boomer 137
Elkhorn, taxadjunct 139
Elkhorn, taxadjunct 141
Holland 143
Holland, taxadjunct 145
Josephine variant 147
Josephine, taxadjunct 149
Josephine, taxadjunct 151
Josephine, taxadjunct 153
Josephine, taxadjunct 155
Josephine, taxadjunct 157
Josephine, taxadjunct 159
Mariposa, taxadjunct 161
Musick 163
Musick 165

Fine-loamy, mixed, thermic
Boomer, taxadjunct 167
Boomer, taxadjunct 169
Ramona, taxadjunct 171
Red Bluff variant 173
Sierra 175
Sierra 177

Fine, kaolinitic, mesic
Aiken, taxadjunct 179
Hoda 181
Hoda 183
Hoda 185

Fine, kaolinitic, thermic
Red Bluff, taxadjunct 187

Fine, montmorillonitic, thermic
Argonaut, taxadjunct 189

Typic Natrixeralfs
Fine-loamy, mixed, thermic
Placentia, taxadjunct 191
Solano 193

Fine, montmorillonitic, thermic
Antioch 195
Placentia 197
San Miguel 199

Aquic Natrixeralfs
Fine-loamy, mixed, thermic
Solano, taxadjunct 201

Fine, montmorillonitic, thermic
Pescadero 203
Pescadero 205
Pescadero 207

Page

PALEXERALFS

Typic Palexeralfs
Fine-loamy, mixed, thermic
Cometa, taxadjunct 211
Corning, taxadjunct 213

Fine, montmorillonitic, thermic
Corning 215
Corning 217
Corning 219
San Ysidro 221

Mollic Palexeralfs
Fine, mixed, mesic
Secca, taxadjunct 223

Fine, montmorillonitic, thermic
Millsap, taxadjunct 225
Positas 227
Positas 229

Natric Palexeralfs
Fine, montmorillonitic, thermic
Bonsall 231

Ultic Palexeralfs
Fine, kaolinitic, mesic
Sites, taxadjunct 233

Fine, kaolinitic, thermic
Red Bluff 235
Red Bluff 237

ARIDISOLS

ARGIDS

DURARGIDS

Abruptic Xerollic Durargids
Fine, montmorillonitic, mesic 239
Lovejoy

Xerollic Durargids
Fine-loamy, mixed, mesic
Lovejoy, taxadjunct 241

HAPLARGIDS

Typic Haplargids
Fine-loamy, mixed, thermic
Mohave, taxadjunct 243

Xerollic Haplargids
Coarse-loamy, mixed, mesic
Beckwourth, taxadjunct 247

NADURARGIDS

Typic Nadurargids
Fine-silty, mixed, mesic
McKay, taxadjunct 249

NATRARGIDS

Typic Natrargids
Fine-loamy, mixed, thermic
Casa Grande, taxadjunct 251
Casa Grande, taxadjunct 253

Aquic Natrargids
Fine-loamy, mixed, mesic

CLASSIFICATION INDEX

Page

ARIDISOLS, cont'd.

ARGIDS, cont'd.

NATRARGIDS, cont'd.

Duric Natrargids
Fine, montmorillonitic, mesic
Churchill, taxadjunct 259

Xerollic Natrargids
Fine-silty, mixed, mesic
Skumpah, taxadjunct 261

PALEARGIDS

Typic Paleargids
Fine-loamy, mixed, mesic
Calneva, taxadjunct 263

ORTHIDS

CALCIORTHIDS

Typic Calciorthids
Fine, montmorillonitic, thermic
Morales 265
Morales, taxadjunct 267

Aquic Calciorthids
Coarse-loamy, mixed, mesic
Balman, taxadjunct 269

Fine-loamy, mixed, mesic
Balman 271

Aquic Duric Calciorthids
Coarse-loamy, mixed, mesic
Kisring, taxadjunct 273

CAMBORTHIDS

Typic Camborthids
Coarse-loamy, mixed, thermic
Laveen, taxadjunct 275

Fine-loamy, mixed, thermic
Milham, taxadjunct 277

SALORTHIDS

Typic Salorthids
Very-fine, montmorillonitic, thermic
Morales, taxadjunct 279

ENTISOLS

FLUVENTS

TORRIFLUVENTS

Typic Torrifluvents
Fine, montmorillonitic, thermic
Rosamond, taxadjunct 281

XEROFLUVENTS

Typic Xerofluvents
Coarse-loamy, mixed, calcareous, thermic
San Emigdio 283
San Emigdio 285
San Emigdio 287

Aquic Xerofluvents
Fine-silty, mixed, nonacid, thermic
Campbell, taxadjunct 289

Fine, montmorillonitic, nonacid, thermic
Oxalis, taxadjunct 291

ORTHENTS

TORRIORTHENTS

Typic Torriorthents
Fine-loamy, mixed, calcareous, thermic
Rosamond 293

XERORTHENTS

Typic Xerorthents
Coarse-loamy, mixed, calcareous, thermic
San Emigdio, taxadjunct 295

Coarse-loamy, mixed, nonacid, thermic
Hanford, taxadjunct 297
Hanford 299
Honcut 301

Fine-loamy, mixed, calcareous, thermic
Shedd, taxadjunct 303

Fine-silty, mixed, calcareous, thermic
Shedd 305

Lithic Xerorthents
Loamy, mixed, nonacid, thermic
Daulton 307

PSAMMENTS

CRYOPSAMMENTS

Typic Cryopsamments
Mixed
Cagwin 309
Cagwin 311
Cagwin variant 313
Cagwin 315

Mixed, shallow
Toem 317
Toem 319
Toem 321

TORRIPSAMMENTS

Xeric Torripsamments
Mixed, mesic
Ormsby variant 323

XEROPSAMMENTS

Typic Xeropsamments
Mixed, mesic
LaPosta, taxadjunct 325

Mixed, shallow, thermic
Cieneba variant 327

CLASSIFICATION INDEX

Page

ENTISOLS, cont'd.

PSAMMENTS, cont'd.

XEROPSAMMENTS, cont'd.

Alfic Xeropsamments
Mixed, frigid
Elmira 329
Elmira, taxadjunct 331
Elmira 333

Mixed, thermic
Oceano 335
Oceano 337

Aquic Durorthidic Xeropsamments
Mixed, mesic
Tangair 339
Tangair 341

INCEPTISOLS

ANDEPTS

CRYANDEPTS

Typic Cryandepts
Cindery
Waca 343
Waca 345

DYSTRANDEPTS

Typic Dystrandepts
Cindery, frigid
Windy 347
Windy 349

VITRANDEPTS

Typic Vitrandepts
Medial-skeletal, mesic
McCarthy 351

AQUEPTS

HAPLAQUEPTS

Aeric Haplaquepts
Sandy, mixed, calcareous, thermic
Piper, taxadjunct 353

Sulfic Haplaquepts
Fine, mixed, acid, thermic
Reyes 355
Reyes 357

Vertic Haplaquepts
Fine, montmorillonitic, calcareous, thermic
Marcuse, taxadjunct 359

HUMAQUEPTS

Histic Humaquepts
Coarse-loamy, mixed, calcareous, thermic
Piper, taxadjunct 361

OCHREPTS

CRYOCHREPTS

Dystric Cryochrepts
Loamy-skeletal, mixed
Umpa, taxadjunct 363

Fine-loamy, mixed
Umpa variant 365

Page

DUROCHREPTS

Typic Durochrepts
Coarse-loamy, mixed, thermic
Arlington, taxadjunct 367
Pentz, taxadjunct 369
Pentz, taxadjunct 371
Trigo variant 373
Trigo variant 375
Trigo variant 377

XEROCHREPTS

Typic Xerochrepts
Coarse-loamy, mixed, mesic
Crouch, taxadjunct 379
Hugo variant 381
Stoner 383
Stoner 385

Coarse-loamy, mixed, thermic
Vista 387
Vista 389
Vista 391
Vista 393
Vista 395
Vista 397
Vista 399

Coarse-loamy, mixed, thermic, shallow
Cieneba, taxadjunct 401

Fine-silty, mixed, thermic
Millsholm variant 403

Calcixerollic Xerochrepts
Coarse-loamy, mixed, thermic
San Emigdio, taxadjunct 405

Dystric Xerochrepts
Coarse-loamy, mixed, frigid
Inville, taxadjunct 407

Coarse-loamy, mixed, mesic
Hugo variant 409

Lithic Xerochrepts
Loamy, mixed, thermic
Temescal 411
Toomes 413
Toomes 415

Ruptic-Lithic Xerochrepts
Loamy, mixed, thermic
Auburn 417
Auburn 419

UMBREPTS

CRYUMBREPTS

Aquentic Cryumbrepts
Sandy-skeletal, mixed
Celio 421

Entic Cryumbrepts
Loamy-skeletal, mixed
Tallac 423

HAPLUMBREPTS

Duric Haplumbrepts
Coarse-loamy, mixed, frigid
Tallac variant 425

CLASSIFICATION INDEX

Page

INCEPTISOLS, cont'd.

UMBREPTS, cont'd.

XERUMBREPTS

Pachic Xerumbrepts
Loamy-skeletal, mixed, mesic
Fiddletown 427
Fiddletown 429

Clayey-skeletal, montmorillonitic, thermic
Santa Lucia, taxadjunct 431

MOLLISOLS

ALBOLLS

ARGIALBOLLS

Xeric Argialbolls
Fine, montmorillonitic, mesic
Pomponio variant 433
Pomponio variant 435

AQUOLLS

ARGIAQUOLLS

Abruptic Argiaquolls
Fine, montmorillonitic, mesic
Watsonville, taxadjunct 437
Watsonville, taxadjunct 439

CALCIAQUOLLS

Typic Calciaquolls
Fine-silty, mixed, mesic
Logan 441

HAPLAQUOLLS

Typic Haplaquolls
Fine-loamy, mixed, noncalcareous, mesic
Ramelli 443

Clayey over loamy, montmorillonitic, noncalcareous, mesic
Ramelli, taxadjunct 445

Vertic Haplaquolls
Very-fine, montmorillonitic, noncalcareous, thermic
Sacramento 447
Sacramento 449

XEROLLS

ARGIXEROLLS

Typic Argixerolls
Loamy-skeletal, mixed, mesic
Secca variant 451

Loamy-skeletal, mixed, thermic
Livermore, taxadjunct 453

Coarse-loamy, mixed, thermic
Chualar, taxadjunct 455

Fine-loamy, mixed, thermic
Carlsbad variant 457
Chualar 459
Chualar, taxadjunct 461

Fine-silty, mixed, mesic
Lobitos, taxadjunct 463

Page

Aquic Argixerolls
Coarse-loamy, mixed, mesic
Beckwourth 465

Aridic Argixerolls
Ashy, mesic, shallow
Casuse, taxadjunct 467

Loamy, mixed, mesic, shallow
Casuse 469

Calcic Argixerolls
Fine-loamy, mixed, thermic
Chualar variant 471

Lithic Argixerolls
Clayey, montmorillonitic, thermic
Vallecitos, taxadjunct 473
Vallecitos, taxadjunct 475

Pachic Argixerolls
Fine-loamy, mixed, isomesic
Denison, taxadjunct 477

Fine-loamy, mixed, thermic
Lockwood 479
Lockwood 481

Fine, montmorillonitic, isomesic
Denison 483

Pachic Ultic Argixerolls
Fine, montmorillonitic, mesic
Lobitos, taxadjunct 485

Ultic Argixerolls
Loamy-skeletal, mixed, mesic
Supan, taxadjunct 487

Loamy-skeletal, mixed, thermic
Supan, taxadjunct 489

Coarse-loamy, mixed, mesic
Kitchen Creek 491
Kitchen Creek 493
Kitchen Creek 495
Kitchen Creek 497

Fine-loamy, mixed, mesic
Felton 499

CALCIXEROLLS

Aquic Calcixerolls
Fine-loamy, mixed, thermic
Chino, taxadjunct 501

Fine-silty, mixed, thermic
Merrill 503

Aridic Calcixerolls
Fine-silty, mixed, mesic
Calita variant 505

Pachic Calcixerolls
Fine-silty, mixed, thermic
Linne, taxadjunct 507

DURIXEROLLS

Orthidic Durixerolls
Fine-loamy, mixed, thermic
Pozo 509

CLASSIFICATION INDEX

Page

MOLLISOLS, cont'd.

XEROLLS, cont'd.

HAPLOXEROLLS

Typic Haploxerolls
Coarse-loamy, mixed, mesic
Crouch, taxadjunct 511

Coarse-loamy, mixed, thermic
San Andreas 513
San Andreas, taxadjunct 515
Sheridan, taxadjunct 517
Sheridan, taxadjunct 519
Sheridan, taxadjunct 521
Sheridan, taxadjunct 523

Fine-loamy, mixed, mesic
Sweeney, taxadjunct 525

Fine-loamy, mixed, thermic
San Andreas, taxadjunct 527

Fine, montmorillonitic, thermic
Rincon, taxadjunct 529

Aquic Haploxerolls
Fine, montmorillonitic, thermic
Landlow 531
Landlow, taxadjunct 533

Aridic Haploxerolls
Coarse-loamy, mixed, mesic
Calpine 535
Calpine 537

Cumulic Haploxerolls
Fine-loamy, mixed, mesic
Soquel, taxadjunct 539

Pachic Haploxerolls

Coarse-loamy, mixed, thermic
Sheridan 541
Sheridan 543
Sheridan 545

Coarse-silty, mixed, thermic
Salinas, taxadjunct 547

Fine-loamy, mixed, mesic
Linne 549
Sweeney, taxadjunct 551

Fine-loamy, mixed, thermic
Salinas 553

Fine, montmorillonitic, thermic
Rincon, taxadjunct 555

Pachic Ultic Haploxerolls
Clayey-skeletal, montmorillonitic, thermic
Santa Lucia, taxadjunct 557

Fine-loamy, mixed, frigid
Tournquist 559

Ultic Haploxerolls
Sandy, mixed, mesic
Crouch, taxadjunct 561

Loamy, mixed, mesic, shallow
Sheephead 563

Page

Ultic Haploxerolls, cont'd.
Coarse-loamy, mixed, mesic
Crouch 565
Crouch 567
Crouch 569
Stoner, taxadjunct 571

Coarse-loamy, mixed, thermic
San Andreas, taxadjunct 573
San Andreas, taxadjunct 575

Vertic Haploxerolls
Fine, montmorillonitic, thermic
Zaca 577

NATRIXEROLLS

Typic Natrixerolls
Fine-loamy, mixed, thermic
Waukena variant 579

ULTISOLS

HUMULTS

HAPLOHUMULTS

Xeric Haplohumults
Clayey, kaolinitic, mesic
Aiken 581
Sites 583
Sites 585

XERULTS

HAPLOXERULTS

Typic Haploxerults
Fine-loamy, mixed, mesic
Goldridge 587
Goldridge 589
Josephine 591
Mariposa, taxadjunct 593

Clayey, mixed, thermic
Red Bluff, taxadjunct 595

VERTISOLS

XERERTS

CHROMOXERERTS

Typic Chromoxererts
Fine, montmorillonitic, mesic
Montague 597

Fine, montmorillonitic, thermic
Altamont 599
Altamont 601
Capay 603
Capay 605
Capay 607
Capay, taxadjunct 609

PELLOXERERTS

Typic Pelloxererts
Fine, montmorillonitic, thermic
Clear Lake 611
Clear Lake 613
Diablo, taxadjunct 615
Willows 617
Willows, taxadjunct 619

Page

VERTISOLS, cont'd.

XERERTS, cont'd.

PELLOXERERTS, cont'd.

Chromic Pelloxererts

Fine, montmorillonitic, mesic
Pit, taxadjunct 621

Fine, montmorillonitic, thermic
Bosanko 623
Bosanko 625
Cropley 627
Diablo 629
Diablo 631
Diablo 633
Diablo 635
Diablo 637

GEOGRAPHICAL INDEX

Classification	Soil Series	Page
ALAMEDA COUNTY		
Alfisols		
Xeralfs		
Haploxeralfs	Livermore, taxadjunct	79
	Pleasanton	95
	Pleasanton, taxadjunct	97
	Unnamed	69
Palexeralfs	Positas	227
	Positas	229
Inceptisols		
Ochrepts		
Xerochrepts	Millsholm variant	403
Mollisols		
Xerolls		
Argixerolls	Livermore, taxadjunct	453
	Vallecitos, taxadjunct	473
	Vallecitos, taxadjunct	475
Vertisols		
Xererts		
Chromoxererts	Altamont	599
	Altamont	601
Pelloxererts	Diablo	629
	Diablo	631
AMADOR COUNTY		
Alfisols		
Xeralfs		
Haploxeralfs	Aiken, taxadjunct	179
	Argonaut	109
	Argonaut, taxadjunct	189
	Josephine variant	147
	Josephine, taxadjunct	149
	Mariposa, taxadjunct	161
	Musick	163
	Musick	165
	Pardee	71
	Pardee, taxadjunct	73
	Red Bluff variant	173
	Sierra	175
	Sierra	177
Palexeralfs	Sites, taxadjunct	233
Inceptisols		
Andepts		
Dystrandepts	Windy	351
	Windy	353
Vitrandepts	McCarthy	355
Ochrepts		
Durochrepts	Pentz, taxadjunct	369
	Pentz, taxadjunct	371
Xerochrepts	Auburn	411
	Auburn	413
Umbrepts		
Xerumbrepts	Fiddletown	427
	Fiddletown	429
Mollisols		
Xerolls		
Argixerolls	Supan, taxadjunct	487
	Supan, taxadjunct	489
Haploxerolls	San Andreas, taxadjunct	573
	San Andreas, taxadjunct	575
Ultisols		
Humults		
Haplohumults	Aiken	581
Xerults		
Haploxerults	Josephine	591
	Mariposa, taxadjunct	593
	Red Bluff, taxadjunct	595
CONTRA COSTA COUNTY		
Entisols		
Fluvents		
Xerofluvents	Campbell, taxadjunct	289
	Oxalis, taxadjunct	291
Inceptisols		
Aquepts		
Haplaquepts	Piper, taxadjunct	353
Humaquepts	Piper, taxadjunct	361
EL DORADO COUNTY		
Alfisols		
Xeralfs		
Haploxeralfs	Ahwahnee, taxadjunct	133
	Boomer, taxadjunct	167
	Boomer, taxadjunct	169
	Holland	143
	Jabu variant	127
	Jabu variant	129
	Jabu	131
	Jabu variant	135
	Rescue	99
	Rescue	101
Entisols		
Psamments		
Cryopsamments	Cagwin	313
	Cagwin	315
	Cagwin variant	317
	Cagwin	319
	Toem	321
	Toem	323
	Toem	325
Xeropsamments	Elmira	333
	Elmira, taxadjunct	335
	Elmira	337
Inceptisols		
Ochrepts		
Xerochrepts	Inville, taxadjunct	407
Umbrepts		
Cryumbrepts	Celio	421
	Tallac	423
Haplumbrepts	Tallac variant	425
FRESNO COUNTY		
Alfisols		
Xeralfs		
Durixeralfs	Fresno, taxadjunct	39
GLENN COUNTY		
Alfisols		
Xeralfs		
Palexeralfs	Corning, taxadjunct	213
	Corning	217

GEOGRAPHICAL INDEX

Classification	Soil Series	Page
LASSEN COUNTY		
Aridisols		
Argids		
Nadurargids	McKay, taxadjunct	249
Natrargids	Churchill, taxadjunct	259
	Skumpah, taxadjunct	261
Paleargids	Calneva, taxadjunct	263
Orthids		
Calciorthids	Kisring, taxadjunct	273
Mollisols		
Aquolls		
Calciaquolls	Logan	441
Xerolls		
Calcixerolls	Calita variant	505
Vertisols		
Xererts		
Palloxererts	Pit, taxadjunct	621
MADERA COUNTY		
Mollisols		
Xerolls		
Durixerolls	Pozo	509
MERCED COUNTY		
Mollisols		
Xerolls		
Calcixerolls	Chino, taxadjunct	501
Haploxerolls	Landlow	531
Natrixerolls	Waukena variant	579
MODOC COUNTY		
Mollisols		
Xerolls		
Argixerolls	Casuse, taxadjunct	467
	Casuse	469
Haploxerolls	Tournquist	559
MONTEREY COUNTY		
Alfisols		
Xeralfs		
Durixeralfs	Gloria	25
Natrixeralfs	Placentia, taxadjunct	191
	Placentia	197
Entisols		
Orthents		
Xerorthents	Hanford	301
	Shedd, taxadjunct	305
	Shedd	307
Inceptisols		
Ochrepts		
Xerochrepts	Stoner	383
	Vista	395
	Vista	397
	Vista	399
Mollisols		
Xerolls		
Argixerolls	Chualar, taxadjunct	455
	Chualar	459
	Lockwood	479
	Lockwood	481
Calcixerolls	Linne, taxadjunct	507
Haploxerolls	Salinas, taxadjunct	547
	Salinas	553
	San Andreas	513
	San Andreas, taxadjunct	527
	Sheridan, taxadjunct	517
	Sheridan, taxadjunct	519
	Sheridan, taxadjunct	521
	Sheridan, taxadjunct	523
	Sheridan	541
	Sheridan	543
	Sheridan	545
	Zaca	577
Vertisols		
Xererts		
Palloxererts	Diablo	637
NEVADA COUNTY		
Alfisols		
Xeralfs		
Haploxeralfs	Dubakella variant	111
	Hoda	183
	Hoda	185
	Secca	105
	Trabuco, taxadjunct	103
	Trabuco	107
Palexeralfs	Secca, taxadjunct	223
Mollisols		
Xerolls		
Argixerolls	Secca variant	451
Ultisols		
Humults		
Haplohumults	Sites	583
	Sites	585
PLACER COUNTY		
Alfisols		
Xeralfs		
Durixeralfs	Fiddyment, taxadjunct	21
	Fiddyment, taxadjunct	31
	Monserate, taxadjunct	15
	San Joaquin, taxadjunct	17
	San Joaquin, taxadjunct	23
Palexeralfs	Cometa, taxadjunct	211
Inceptisols		
Andepts		
Cryandepts	Waca	347
	Waca	349
Ochrepts		
Cryochrepts	Umpa, taxadjunct	363
	Umpa variant	365
Durochrepts	Trigo variant	373
	Trigo variant	375
	Trigo variant	377
Xerochrepts	Cieneba, taxadjunct	401
	Toomes	417
	Toomes	419

Classification	Soil Series	Page
PLACER COUNTY, cont'd.		
Mollisols		
Xerolls		
Haploxerolls	San Andreas, taxadjunct	515
PLUMAS COUNTY		
Aridisols		
Argids		
Durargids	Lovejoy	239
Haplargids	Beckwourth, taxadjunct	247
Natrargids	Loyalton	255
	Loyalton	257
Orthids		
Calciorthids	Balman	271
Entisols		
Psamments		
Torripsamments	Ormsby variant	327
Mollisols		
Aquolls		
Haplaquolls	Ramelli	443
	Ramelli, taxadjunct	445
Xerolls		
Argixerolls	Beckwourth	465
Haploxerolls	Calpine	535
RIVERSIDE COUNTY		
Alfisols		
Xeralfs		
Durixeralfs	Arlington	37
	Exeter	7
	Monserate	11
	Monserate, taxadjunct	33
Haploxeralfs	Greenfield	43
	Greenfield	45
	Greenfield	47
	Millerton, taxadjunct	75
	Millerton, taxadjunct	77
	Montpellier	59
	Montpellier, taxadjunct	61
	Ramona, taxadjunct	49
	Ramona	63
	Ramona, taxadjunct	171
	Traver	113
	Traver	115
Entisols		
Fluvents		
Xerofluvents	San Emigdio	285
	San Emigdio	287
	San Emigdio	289
Orthents		
Xerorthents	Daulton	311
	Honcut	303
	San Emigdio, taxadjunct	297
Psamments		
Xeropsamments	Cieneba variant	331
Inceptisols		
Aquepts		
Haplaquepts	Marcuse, taxadjunct	361
Ochrepts		
Xerochrepts	San Emigdio, taxadjunct	405
	Temescal	411
	Vista	387
	Vista	389
Mollisols		
Xerolls		
Haploxerolls	Landlow, taxadjunct	533
Vertisols		
Xererts		
Chromoxererts	Capay	607
	Capay, taxadjunct	609
SAN BERNARDINO COUNTY		
Xeralfs		
Haploxeralfs	Bonta variant	41
	Pachappa, taxadjunct	93
Aridisols		
Argids		
Haplargids	Mohave, taxadjunct	243
	Mohave	245
Natrargids	Casa Grande, taxadjunct	251
	Casa Grande, taxadjunct	253
Orthids		
Calciorthids	Morales	265
	Morales, taxadjunct	267
Camborthids	Laveen, taxadjunct	275
	Milham, taxadjunct	277
Salorthids	Morales, taxadjunct	279
Entisols		
Fluvents		
Torrifluvents	Rosamond, taxadjunct	283
Orthents		
Torriorthents	Rosamond	295
Mollisols		
Xerolls		
Argixerolls	Chualar, taxadjunct	461
	Chualar variant	471
Calcixerolls	Merrill	503
Vertisols		
Xererts		
Pelloxererts	Cropley	627
SAN DIEGO COUNTY		
Alfisols		
Xeralfs		
Durixeralfs	Monserate	13
	Redding, taxadjunct	29
	Tuscan, taxadjunct	19
Haploxeralfs	Bonsall, taxadjunct	117
	Boomer	137
	Fallbrook	51
	Fallbrook, taxadjunct	91
	Hoda	181
	Holland, taxadjunct	145
	Redding, taxadjunct	119
	San Miguel variant	121
Natrixeralfs	San Miguel	199
Palexeralfs	Bonsall	231

GEOGRAPHICAL INDEX

Classification	Soil Series	Page
SAN DIEGO COUNTY, cont'd.		
Entisols		
Psamments		
Xeropsamments	LaPosta, taxadjunct	329
Inceptisols		
Ochrepts		
Durochrepts	Arlington, taxadjunct	367
Xerochrepts	Crouch, taxadjunct	379
	Vista	391
	Vista	393
Mollisols		
Xerolls		
Argixerolls	Carlsbad variant	457
	Felton	499
	Kitchen Creek	491
	Kitchen Creek	493
	Kitchen Creek	495
	Kitchen Creek	497
Haploxerolls	Crouch, taxadjunct	511
	Crouch, taxadjunct	561
	Crouch	565
	Crouch	567
	Crouch	569
	Sheephead	563
Vertisols		
Xererts		
Pelloxererts	Bosanko	623
	Bosanko	625
SAN LUIS OBISPO COUNTY		
Alfisols		
Xeralfs		
Haploxeralfs	Ahwahnee, taxadjunct	83
Palexeralfs	Millsap, taxadjunct	225
Mollisols		
Xerolls		
Haploxerolls	Soquel, taxadjunct	539
Vertisols		
Xererts		
Pelloxererts	Diablo, taxadjunct	615
SAN MATEO COUNTY		
Alfisols		
Xeralfs		
Haploxeralfs	Elkhorn, taxadjunct	139
	Elkhorn, taxadjunct	141
Inceptisols		
Ochrepts		
Xerochrepts	Hugo variant	381
	Hugo variant	409
Mollisols		
Albolls		
Argialbolls	Pomponio variant	433
	Pomponio variant	435
Aquolls		
Argiaquolls	Watsonville, taxadjunct	437
	Watsonville, taxadjunct	439
Xerolls		
Argixerolls	Denison, taxadjunct	477
	Denison	483
	Lobitos, taxadjunct	463
	Lobitos, taxadjunct	485
Haploxerolls	Sweeney, taxadjunct	525
	Sweeney, taxadjunct	551
SANTA BARBARA COUNTY		
Alfisols		
Xeralfs		
Haploxeralfs	Garey	123
	Garey	125
Entisols		
Psamments		
Xeropsamments	Oceano	339
	Oceano	341
	Tangair	343
	Tangair	345
Inceptisols		
Umbrepts		
Xerumbrepts	Santa Lucia, taxadjunct	431
Mollisols		
Xerolls		
Haploxerolls	Linne	549
	Santa Lucia, taxadjunct	557
SHASTA COUNTY		
Alfisols		
Xeralfs		
Palexeralfs	Red Bluff	235
SIERRA COUNTY		
Aridisols		
Argids		
Durargids	Lovejoy, taxadjunct	241
Orthids		
Calciorthids	Balman, taxadjunct	269
Mollisols		
Xerolls		
Haploxerolls	Calpine	537
SISKIYOU COUNTY		
Alfisols		
Xeralfs		
Haploxeralfs	Hilt	85
	Hilt	87
	Josephine, taxadjunct	151
	Josephine, taxadjunct	153
	Josephine, taxadjunct	155
	Josephine, taxadjunct	157
	Josephine, taxadjunct	159
Inceptisols		
Ochrepts		
Xerochrepts	Stoner	385
Mollisols		
Xerolls		
Haploxerolls	Stoner, taxadjunct	571
Vertisols		
Xererts		
Chromoxererts	Montague	597

GEOGRAPHICAL INDEX

Classification	Soil Series	Page
SOLANO COUNTY		
Alfisols		
Xeralfs		
Natrixeralfs	Antioch	195
	Pescadero	203
	Pescadero	205
	Pescadero	207
	Solano	193
	Solano, taxadjunct	201
Palexeralfs	San Ysidro	221
Vertisols		
Xererts		
Chromoxererts	Capay	603
SONOMA COUNTY		
Alfisols		
Aqualfs		
Albaqualfs	Wright, taxadjunct	3
	Wright, taxadjunct	5
Xeralfs		
Durixeralfs	Huichica, taxadjunct	35
Haploxeralfs	Huichica variant	55
	Steinbeck, taxadjunct	81
	Steinbeck	89
Inceptisols		
Aquepts		
Haplaquepts	Reyes	357
	Reyes	359
Ultisols		
Xerults		
Haploxerults	Goldridge	587
	Goldridge	589
Vertisols		
Xererts		
Pelloxererts	Clear Lake	611
	Clear Lake	613
	Diablo	633
	Diablo	635
TEHAMA COUNTY		
Alfisols		
Xeralfs		
Durixeralfs	Monserate	9
	Redding, taxadjunct	27
Haploxeralfs	Greenfield, taxadjunct	53
	Montpellier	57
	Red Bluff, taxadjunct	187
Palexeralfs	Corning	215
	Corning	219
	Red Bluff	237
Entisols		
Orthents		
Xerorthents	Hanford, taxadjunct	299
YOLO COUNTY		
Alfisols		
Xeralfs		
Haploxeralfs	Tehama, taxadjunct	65
	Tehama, taxadjunct	67
Natrixeralfs	Pescadero	209
Mollisols		
Aquolls		
Haplaquolls	Sacramento	447
	Sacramento	449
Xerolls		
Haploxerolls	Rincon, taxadjunct	529
	Rincon, taxadjunct	555
Vertisols		
Xererts		
Chromoxererts	Capay	605
Pelloxererts	Willows	617
	Willows, taxadjunct	619

SOIL SERIES INDEX

Series	Soil Survey No. 1/	Classification	Page
Ahwahnee, taxadjunct	S55Calif-40-1	Haploxeralf	83
Ahwahnee, taxadjunct	S61Calif-9-6	Haploxeralf	133
Aiken, taxadjunct	S59Calif-3-11	Haploxeralf	179
Aiken	S59Calif-3-12	Haplohumult	581
Altamont	S58Calif-1-3	Chromoxerert	599
Altamont	S58Calif-1-4	Chromoxerert	601
Antioch	S64Calif-48-8	Natrixeralf	195
Argonaut, taxadjunct	S59Calif-3-5	Haploxeralf	189
Argonaut	S59Calif-3-6	Haploxeralf	109
Arlington	S57Calif-33-10	Durixeralf	37
Arlington, taxadjunct	S66Calif-37-3	Durochrept	367
Auburn	S59Calif-3-7	Xerochrept	411
Auburn	S59Calif-3-8	Xerochrept	413
Balman	S61Calif-32-12	Calciorthid	271
Balman, taxadjunct	S61Calif-46-4	Calciorthid	269
Beckwourth	S61Calif-32-15	Argixeroll	465
Beckwourth, taxadjunct	S61Calif-32-16	Haplargid	247
Bonsall	S64Calif-37-2	Palexeralf	231
Bonsall, taxadjunct	S64Calif-37-5	Haploxeralf	117
Bonta variant	S69Calif-33-1	Haploxeralf	41
Boomer	S60Calif-37-9	Haploxeralf	137
Boomer, taxadjunct	S61Calif-9-3	Haploxeralf	167
Boomer, taxadjunct	S61Calif-9-4	Haploxeralf	169
Bosanko	S64Calif-37-8	Pelloxerert	623
Bosanko	S64Calif-37-9	Pelloxerert	625
Cagwin	S68Calif-9-21	Cryopsamment	309
Cagwin variant	S68Calif-9-27	Cryopsamment	313
Cagwin	S68Calif-9-32	Cryopsamment	315
Calita variant	S54Calif-18-1	Calcixeroll	505
Calneva, taxadjunct	S55Calif-18-3	Paleargid	263
Calpine	S61Calif-32-19	Haploxeroll	535
Calpine	S61Calif-46-5	Haploxeroll	537
Campbell, taxadjunct	S57Calif-7- 2	Xerofluvent	289
Capay	S64Calif-48-2	Chromoxerert	603
Capay	S64Calif-57-13	Chromoxerert	605
Capay	S67Calif-33-1	Chromoxerert	607
Capay, taxadjunct	S67Calif-33-2	Chromoxerert	609
Carlsbad variant	S56Calif-37-1	Argixeroll	457
Casa Grande, taxadjunct	S55Calif-36-7	Natrargid	251
Casa Grande,	S55Calif-36-8	Natrargid	253

Series	Soil Survey No. 1/	Classification	Page
Corning	S4557-3	Palexeralf	215
Corning, taxadjunct	S4557-12	Palexeralf	213
Corning	S4557-13	Palexeralf	217
Corning	S4557-14	Palexeralf	219
Cropley	S69Calif-36-26	Pelloxerert	627
Crouch, taxadjunct	S60Calif-37-5	Haploxeroll	561
Crouch, taxadjunct	S60Calif-37-6	Xerochrept	379
Crouch	S61Calif-37-3	Haploxeroll	565
Crouch, taxadjunct	S61Calif-37-5	Haploxeroll	511
Crouch	S61Calif-37-6	Haploxeroll	567
Crouch	S61Calif-37-7	Haploxeroll	569
Daulton	S57Calif-33-4	Xerorthent	307
Denison	S55Calif-41-1	Argixeroll	483
Denison, taxadjunct	S55Calif-41-2	Argixeroll	477
Diablo, taxadjunct	S55Calif-40-2	Pelloxerert	615
Diablo	S58Calif-1-1	Pelloxerert	629
Diablo	S58Calif-1-2	Pelloxerert	631
Diablo	S61Calif-49-11	Pelloxerert	633
Diablo	S61Calif-49-12	Pelloxerert	635
Diablo	S65Calif-27-10	Pelloxerert	637
Dubakella variant	S68Calif-29-36	Haploxeralf	111
Elkhorn, taxadjunct	S55Calif-41-5	Haploxeralf	139
Elkhorn, taxadjunct	S55Calif-41-6	Haploxeralf	141
Elmira	S68Calif-9-22	Xeropsamment	329
Elmira, taxadjunct	S68Calif-9-26	Xeropsamment	331
Elmira	S68Calif-9-28	Xeropsamment	333
Fallbrook	S64Calif-37-2	Haploxeralf	51
Fallbrook, taxadjunct	S64Calif-37-6	Haploxeralf	91
Felton	S61Calif-37-1	Argixeroll	499
Fiddletown	S60Calif-3-27	Xerumbrept	427
Fiddletown	S60Calif-3-28	Xerumbrept	429
Fiddyment, taxadjunct	S69Calif-31-8	Durixeralf	21
Fiddyment, taxadjunct	S69Calif-31-13	Durixeralf	31
Fresno, taxadjunct	S58Calif-10-1	Durixeralf	39
Garey	S62Calif-42-3	Haploxeralf	123
Garey	S62Calif-42-4	Haploxeralf	125
Gloria	S65Calif-27-16	Durixeralf	25

SOIL SERIES INDEX

Series	Soil Survey No. 1/	Classification	Page
Huichica, taxadjunct	S61Calif-49-7	Durixeralf	35
Huichica variant	S61Calif-49-8	Haploxeralf	55
Inville, taxadjunct	S68Calif-9-17	Xerochrept	407
Jabu variant	S68Calif-9-29	Haploxeralf	127
Jabu variant	S68Calif-9-30	Haploxeralf	129
Jabu variant	S68Calif-9-33	Haploxeralf	135
Jabu	S68Calif-9-34	Haploxeralf	131
Josephine variant	S59Calif-3-4	Haploxeralf	147
Josephine	S60Calif-3-24	Haploxerult	591
Josephine, taxadjunct	S60Calif-3-29	Haploxeralf	149
Josephine, taxadjunct	S66Calif-47-1	Haploxeralf	151
Josephine, taxadjunct	S66Calif-47-2	Haploxeralf	153
Josephine, taxadjunct	S66Calif-47-3	Haploxeralf	155
Josephine, taxadjunct	S66Calif-47-4	Haploxeralf	157
Josephine, taxadjunct	S66Calif-47-5	Haploxeralf	159
Kisring, taxadjunct	S56Calif-18-4	Calciorthid	273
Kitchen Creek, taxadjunct	S60Calif-37-1	Argixeroll	491
Kitchen Creek, taxadjunct	S60Calif-37-2	Argixeroll	493
Kitchen Creek	S60Calif-37-4	Argixeroll	495
Kitchen Creek	S61Calif-37-4	Argixeroll	497
Landlow	S55Calif-24-1	Haploxeroll	531
Landlow, taxadjunct	S57Calif-33-6	Haploxeroll	533
LaPosta, taxadjunct	S60Calif-37-2	Xeropsamment	325
Laveen, taxadjunct	S55Calif-36-9	Camborthid	275
Linne	S62Calif-42-11	Haploxeroll	549
Linne, taxadjunct	S65Calif-27-4	Calcixeroll	507
Livermore, taxadjunct	S59Calif-01-3	Haploxeralf	79
Livermore, taxadjunct	S59Calif-01-4	Argixeroll	453
Lobitos, taxadjunct	S55Calif-41-13	Argixeroll	463
Lobitos, taxadjunct	S55Calif-41-14	Argixeroll	485
Lockwood	S65Calif-27-1	Argixeroll	479
Lockwood	S65Calif-27-2	Argixeroll	481
Logan	S55Calif-18-5	Calciaquoll	441
Lovejoy	S61Calif-32-21	Durargid	239
Lovejoy, taxadjunct	S61Calif-46-6	Durargid	241
Loyalton	S61Calif-32-10	Natrargid	255
Loyalton	S61Calif-32-11	Natrargid	257
Marcuse, taxadjunct	S57Calif-33-7	Haplaquept	359
Mariposa, taxadjunct	S60Calif-3-25	Haploxerult	593
Mariposa, taxadjunct	S60Calif-3-26	Haploxeralf	161
McCarthy	S62Calif-3-1	Vitrandept	351
McKay, taxadjunct	S55Calif-18-2	Nadurargid	249
Merrill	S69Calif-36-23	Calcixeroll	503
Milham, taxadjunct	S55Calif-36-10	Camborthid	277
Millerton, taxadjunct	S57Calif-33-3	Haploxeralf	75
Millerton, taxadjunct	S57Calif-33-5	Haploxeralf	77
Millsap, taxadjunct	S55Calif-40-4	Palexeralf	225
Millsholm variant	S59Calif-1-5	Xerochrept	403
Mohave, taxadjunct	S55Calif-36-5	Haplargid	243
Mohave	S59Calif-36-1	Haplargid	245
Monserate	S4557-9	Durixeralf	9
Monserate, taxadjunct	S57Calif-33-12	Durixeralf	33
Monserate	S57Calif-33-13	Durixeralf	11
Monserate	S66Calif-37-1	Durixeralf	13
Monserate, taxadjunct	S69Calif-31-10	Durixeralf	15
Montague	S66Calif-47-7	Chromoxerert	597
Montpellier	S4557-11	Haploxeralf	57
Montpellier	S55Calif-33-14	Haploxeralf	59
Montpellier, taxadjunct	S55Calif-33-15	Haploxeralf	61
Morales	S55Calif-36-2	Calciorthid	265
Morales, taxadjunct	S55Calif-36-3	Salorthid	279
Morales, taxadjunct	S55Calif-36-4	Calciorthid	267
Musick	S59Calif-3-13	Haploxeralf	163
Musick	S59Calif-3-14	Haploxeralf	165
Oceano	S62Calif-42-1	Xeropsamment	335
Oceano	S62Calif-42-2	Xeropsamment	337
Ormsby variant	S61Calif-32-17	Torripsamment	323
Oxalis, taxadjunct	S57Calif-7-1	Xerofluvent	291
Pachappa, taxadjunct	S69Calif-36-24	Haploxeralf	93
Pardee	S60Calif-3-22	Haploxeralf	71
Pardee, taxadjunct	S60Calif-3-23	Haploxeralf	73
Pentz, taxadjunct	S60Calif-3-20	Durochrept	369
Pentz, taxadjunct	S60Calif-3-21	Durochrept	371
Pescadero	S64Calif-48-3	Natrixeralf	203
Pescadero	S64Calif-48-5	Natrixeralf	205
Pescadero	S64Calif-48-6	Natrixeralf	207
Pescadero	S64Calif-57-12	Natrixeralf	209
Piper, taxadjunct	S57Calif-7-3	Haplaquept	353
Piper, taxadjunct	S57Calif-7-4	Humaquept	361
Pit, taxadjunct	S54Calif-18-3	Pelloxerert	621
Placentia	S65Calif-27-14	Natrixeralf	197
Placentia, taxadjunct	S65Calif-27-15	Natrixeralf	191
Pleasanton	S60Calif-01-1	Haploxeralf	95
Pleasanton, taxadjunct	S60Calif-01-2	Haploxeralf	97
Pomponio variant	S55Calif-41-3	Argialboll	433
Pomponio variant	S55Calif-41-4	Argialboll	435
Positas	S59Calif-01-1	Palexeralf	227
Positas	S59Calif-01-2	Palexeralf	229
Pozo	S58Calif-20-1	Durixeroll	509
Ramelli	S61Calif-32-13	Haplaquoll	443
Ramelli, taxadjunct	S61Calif-32-14	Haplaquoll	445
Ramona	S57Calif-33-8	Haploxeralf	63
Ramona, taxadjunct	S59Calif-33-1	Haploxeralf	171
Ramona, taxadjunct	S59Calif-33-2	Haploxeralf	49
Red Bluff	S4557-6	Palexeralf	235
Red Bluff	S4557-8	Palexeralf	237
Red Bluff, taxadjunct	S4557-10	Haploxeralf	187
Red Bluff, taxadjunct	S59Calif-3-9	Haploxerult	595
Red Bluff variant	S59Calif-3-10	Haploxeralf	173
Redding, taxadjunct	S4557-5	Durixeralf	27

SOIL SERIES INDEX

Series	Soil Survey No. 1/	Classification	Page
Redding, taxadjunct	S66Calif-37-2	Durixeralf	29
Redding, taxadjunct	S66Calif-37-4	Haploxeralf	119
Rescue	S61Calif-9-1	Haploxeralf	99
Rescue	S61Calif-9-2	Haploxeralf	101
Reyes	S61Calif-49-13	Haplaquept	355
Reyes	S61Calif-49-14	Haplaquept	357
Rincon, taxadjunct	S64Calif-57-16	Haploxeroll	555
Rincon, taxadjunct	S64Calif-57-17	Haploxeroll	529
Rosamond, taxadjunct	S55Calif-36-1	Torrifluvent	281
Rosamond	S55Calif-36-6	Torriorthent	293
Sacramento	S64Calif-57-10	Haplaquoll	447
Sacramento	S64Calif-57-11	Haplaquoll	449
Salinas, taxadjunct	S65Calif-27-3	Haploxeroll	547
Salinas	S65Calif-27-11	Haploxeroll	553
San Andreas, taxadjunct	S59Calif-3-17	Haploxeroll	573
San Andreas, taxadjunct	S59Calif-3-18	Haploxeroll	575
San Andreas	S65Calif-27-6	Haploxeroll	513
San Andreas, taxadjunct	S65Calif-27-28	Haploxeroll	527
San Andreas, taxadjunct	S69Calif-31-22	Haploxeroll	515
San Emigdio, taxadjunct	S55Calif-33-10	Xerochrept	405
San Emigdio	S55Calif-33-11	Xerofluvent	283
San Emigdio, taxadjunct	S55Calif-33-13	Xerorthent	295
San Emigdio	S57Calif-33-1	Xerofluvent	285
San Emigdio	S57Calif-33-2	Xerofluvent	287
San Joaquin, taxadjunct	S69Calif-31-20	Durixeralf	17
San Joaquin, taxadjunct	S69Calif-31-21	Durixeralf	23
San Miguel	S64Calif-37-18	Natrixeralf	199
San Miguel variant	S66Calif-37-5	Haploxeralf	121
Santa Lucia, taxadjunct	S62Calif-42-9	Xerumbrept	431
Santa Lucia, taxadjunct	S64Calif-42-10	Haploxeroll	557
San Ysidro	S64Calif-48-1	Palexeralf	221
Secca, taxadjunct	S68Calif-29-5	Palexeralf	223
Secca	S68Calif-29-7	Haploxeralf	105
Secca variant	S68Calif-29-35	Argixeroll	451
Shedd	S65Calif-27-8	Xerorthent	305
Shedd	S65Calif-27-9	Xerorthent	303
Sheephead	S60Calif-37-7	Haploxeroll	563
Sheridan, taxadjunct	S65Calif-27-20	Haploxeroll	517
Sheridan	S65Calif-27-21	Haploxeroll	541
Sheridan	S65Calif-27-22	Haploxeroll	543
Sheridan, taxadjunct	S65Calif-27-23	Haploxeroll	519
Sheridan	S65Calif-27-24	Haploxeroll	545
Sheridan, taxadjunct	S65Calif-27-25	Haploxeroll	521
Sheridan, taxadjunct	S65Calif-27-26	Haploxeroll	523
Sierra	S59Calif-3-15	Haploxeralf	175
Sierra	S59Calif-3-16	Haploxeralf	177
Sites, taxadjunct	S59Calif-3-3	Palexeralf	233
Sites	S68Calif-29-31	Haplohumult	583
Sites	S68Calif-29-32	Haplohumult	585
Skumpah, taxadjunct	S54Calif-18-2	Natrargid	261
Solano	S64Calif-48-4	Natrixeralf	193
Solano, taxadjunct	S64Calif-48-7	Natrixeralf	201
Soquel, taxadjunct	S55Calif-40-3	Haploxeroll	539
Steinbeck, taxadjunct	S61Calif-49-3	Haploxeralf	81
Steinbeck	S61Calif-49-4	Haploxeralf	89
Stoner	S65Calif-27-19	Xerochrept	383
Stoner, taxadjunct	S66Calif-47-9	Haploxeroll	571
Stoner	S66Calif-47-10	Xerochrept	385
Supan, taxadjunct	S59Calif-3-1	Argixeroll	487
Supan, taxadjunct	S59Calif-3-2	Argixeroll	489
Sweeney, taxadjunct	S55Calif-41-9	Haploxeroll	551
Sweeney, taxadjunct	S55Calif-41-10	Haploxeroll	525
Tallac variant	S68Calif-9-15	Haplumbrept	425
Tallac	S68Calif-9-16	Cryumbrept	423
Tangair	S62Calif-42-7	Xeropsamment	339
Tangair	S62Calif-42-8	Xeropsamment	341
Tehama, taxadjunct	S64Calif-57-18	Haploxeralf	65
Tehama, taxadjunct	S64Calif-57-19	Haploxeralf	67
Temescal	S55Calif-33-9	Xerochrept	411
Toem	S68Calif-9-20	Cryopsamment	317
Toem	S68Calif-9-23	Cryopsamment	319
Toem	S68Calif-9-31	Cryopsamment	321
Toomes	S61Calif-31-1	Xerochrept	413
Toomes	S61Calif-31-2	Xerochrept	415
Tournquist	S66Calif-25-1	Haploxeroll	559
Trabuco	S68Calif-29-29	Haploxeralf	107
Trabuco, taxadjunct	S68Calif-29-30	Haploxeralf	103
Traver	S57Calif-33-14	Haploxeralf	113
Traver	S57Calif-33-15	Haploxeralf	115
Trigo variant	S69Calif-31-3	Durochrept	373
Trigo variant	S69Calif-31-11	Durochrept	375
Trigo variant	S69Calif-31-12	Durochrept	377
Tuscan, taxadjunct	S4557-15	Durixeralf	19
Umpa, taxadjunct	S68Calif-31-7	Cryochrept	363
Umpa variant	S68Calif-31-10	Cryochrept	365
Unnamed	S59Calif-01-6	Haploxeralf	69
Vallecitos, taxadjunct	S59Calif-01-8	Argixeroll	473
Vallecitos, taxadjunct	S59Calif-01-9	Argixeroll	475
Vista	S55Calif-33-6	Xerochrept	387
Vista	S55Calif-33-7	Xerochrept	389
Vista	S64Calif-37-1	Xerochrept	391
Vista	S64Calif-37-4	Xerochrept	393
Vista	S65Calif-27-7	Xerochrept	395
Vista	S65Calif-27-29	Xerochrept	397
Vista	S65Calif-27-30	Xerochrept	399
Waca	S68Calif-31-1	Cryandept	343
Waca	S68Calif-31-2	Cryandept	345
Watsonville, taxadjunct	S55Calif-41-11	Argiaquoll	437
Watsonville, taxadjunct	S55Calif-41-12	Argiaquoll	439
Waukena variant	S55Calif-24-3	Natrixeroll	579
Willows	S64Calif-57-14	Pelloxerert	617
Willows, taxadjunct	S64Calif-57-15	Pelloxerert	619
Windy	S61Calif-3-30	Dystrandept	347
Windy	S61Calif-3-31	Dystrandept	349
Wright, taxadjunct	S61Calif-49-1	Albaqualf	3
Wright, taxadjunct	S61Calif-49-2	Albaqualf	5
Zaca	S65Calif-27-5	Haploxeroll	577

SOIL SERIES INDEX

1/ County numbers (the number following "Calif" in the Soil Survey No.) are as follows:

1. Alameda
3. Amador
7. Contra Costa
9. El Dorado
10. Fresno
11. Glenn
18. Lassen
20. Madera
24. Merced
25. Modoc
27. Monterey
29. Nevada
31. Placer
32. Plumas
33. Riverside
36. San Bernardino
37. San Diego
40. San Luis Obispo
41. San Mateo
42. Santa Barbara
45. Shasta
46. Sierra
47. Siskiyou
48. Solano
49. Sonoma
52. Tehama
57. Yolo

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Wright loam (taxadjunct) **LOCATION** Sonoma County, California

SOIL FAMILY: Typic Albaqualf, fine, montmorillonitic, mesic

SOIL NOS. S61Calif-49-1 **LAB. NOS.** 61377 - 61386

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2	
0- 7	A1	0.9	2.2	4.8	19.1	15.8	45.8	11.4	52.3	22.0	4	1
7-14	A21	1.1	2.5	4.7	18.0	15.9	45.6	12.2	52.0	21.6	2	1
14-22	A23	0.5	3.1	4.7	18.9	15.7	45.3	11.8	51.1	22.7	3	1

WRIGHT LOAM (taxadjunct) 1/
S61Calif-49-1

Location: Sonoma County, California. Gold Ridge SCD - Laguna Addition 1000 feet north, 1700 feet west of junction of Todd and Llano Roads. SW1/4, SW1/4, section 5, projected, T. 6 N., R. 8 W. (Neles Ranch). Date of Sampling: June 19, 1961.
Collectors: Reuben Nelson, Leo C. Klameth, V. C. Miller, G. Lambert, J. H. Kashiwagi, F. Graham.
Classification: **Typic Albaqualf, fine, montmorillonitic, mesic.**
Vegetation: Native: Annual grasses and scattered oak. Cult: Oat hay; dryland improved pasture and some irrigated pasture. Use: Dryland pasture. Climate: Annual precipitation 29.4 inches, average temperature 57.6° F., average January temperature 46.5°F., average July temperature 66.7°F.
Parent Material: Old alluvium - mainly from sedimentary and volcanic rocks. Elevation: 83 feet - taken from Sebastopol Quad. Relief: Slightly hummocky (hog wallow) - 1 to 2 percent slope. Drainage: Imperfect - surface water ponding in pockets. Biological Activity: Some worm and rodent activity in A horizon.

HORIZON	DESCRIPTION
A1 * RSL No. [illegible]	0 to 7 inches, light brownish gray (10YR 6/2) very fine sandy loam, dark grayish brown (10YR 4/2) when moist; medium subangular blocky structure; slightly hard when dry; friable when moist [illegible] and fine interstitial and tubular pores; strongly acid, pH 5.2; clear, smooth boundary.
A21 RSL No. 61378	7 to 14 inches, light brownish gray (10YR 6/2) loam, very dark grayish brown (10YR 3/2) when moist, massive; hard when dry, very friable when moist, slightly sticky, slightly plastic when wet; abundant very fine and fine roots; many very fine and fine tubular and interstitial pores; strongly acid, pH 5.5; clear, wavy boundary.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Wright loam (Taxadjunct) **LOCATION** Sonoma County, California

SOIL FAMILY: Mollic Albaqualf, fine, montmorillonitic, mesic

SOIL NOS. S61Calif-49-2 **LAB. NOS.** 61387 - 61397

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2	
0-7	A1	0.6	1.8	4.5	20.9	14.1	46.4	11.7	48.4	25.8	2	1
7-15	A21	0.6	1.9	4.4	20.6	14.6	46.1	11.8	48.6	25.8	0	1
15-25	A22g	0.3	1.8	4.4	20.2	14.8	44.8	13.7	49.1	24.1	1	1
25-32	B21tg	0.4	1.4	3.5	13.2	8.5	31.1	41.9	29.3	18.7	0	c
32-42	B22tg	0.3	1.6	3.0	11.9	9.2	33.3	40.7	30.1	20.2	0	c-cl
42-47	B23tg	0.1	0.8	3.3	13.8	12.7	39.8	29.5	40.0	22.1	0	cl
47-62	B24tg	0.1	0.6	2.8	26.8	17.8	30.4	21.5	53.5	14.7	0	1

WRIGHT LOAM (taxadjunct) 1/
S61Calif-49-2

Location: Sonoma County, California. Gold Ridge SCD, Laguna Addition. 100 feet west and 1200 feet south of junction of Sebastopol and Llano Roads. Northwest corner of NW1/4, SW1/4, section 31, projected, T. 7 N., R. 8 W. (Buehler Ranch). Date of Sampling: June 20, 1961.
Collectors: Reuben Nelson, Leo C. Klameth, V. C. Miller, G. Lambert, J. H. Kashiwagi, F. Graham.
Classification: **Mollic Albaqualf, fine, montmorillonitic, mesic.**
Vegetation: Native: Annual grasses and scattered oak. Cult: Oat hay; some irrigation pasture and some improved dryland pasture. Use: Dryland pasture. Climate: Annual precipitation 29.4 inches, average temperature 57.6°F., average January temperature 46.5°F., average July temperature 66.7°F. Parent Material: Old alluvium - mainly from sedimentary and volcanic rock sources. Elevation: 82 feet - from Sebastopol Quad. Relief: Slightly hummocky (hog wallow) to gently undulating - 1 to 2 percent slope. Drainage: Imperfect - surface water ponding in concave areas. Biological Activity: Some worm and rodent activity to about 2 feet.

HORIZON	DESCRIPTION
A1 * RSL No. 61387	0 to 7 inches, light brownish gray (10YR 6/2) very fine sandy loam, very dark grayish brown (10YR 3/2) when moist; massive; hard when dry, friable when moist, nonsticky and nonplastic when wet; abundant fine and very fine roots; many fine and medium tubular and interstitial pores; medium acid, pH 5.7; gradual, smooth boundary.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Exeter coarse sandy loam **LOCATION** Riverside County, California

SOIL FAMILY: Typic Durixeralf, fine-loamy, mixed, thermic.

SOIL NOS. S57Calif-33-11 **LAB. NOS.** 57527 - 57533

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-10	Ap	8.4	17.8	11.0	19.1	12.7	22.2	8.8	37.8	7.8	3	cosl
10-16	A3	8.3	15.3	8.9	14.5	10.8	26.4	15.8	33.9	11.8	2	sl
16-25	B21	9.2	12.9	7.2	12.3	10.1	29.1	19.2	33.2	13.3	2	l
25-30	B22	4.8	11.5	6.6	12.9	11.6	33.4	19.2	37.8	14.9	2	l
30-37	B3cam	7.8	14.6	9.3	17.1	13.1	31.2	6.9	40.1	14.5	5	sl
37-50	Cm1	11.9	14.2	7.4	13.2	10.1	27.4	15.8	32.1	13.2	5	cosl
50-60+	Cm2	18.1	18.4	8.6	14.2	8.2	18.5	14.0	26.9	8.0	14	cosl

8C1b SATURATED PASTE 1:1	pH 8C1a 1:10	4A3a Bulk Density 1/3 atm.	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	CO_3 6I1a Meq/l	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM 25°C.	6E1b $CaCO_3$ equivalent %	HCO_3 6J1a Meq/l	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %

EXETER COARSE SANDY LOAM
S57Calif-33-11

Location: 1200 feet west and 400 feet south of north 1/4 corner of Sec. 16, T. 4 S., R. 3 W. On the Hughes Ranch, about 2 miles northeast of Perris, Riverside County, California. Date of Sampling: June 5, 1957.
Description by: Fred Eggers. Collectors: George Borst, R. E. Nelson, Cliff Henry, and Arnold Knecht.
Classification: Typic Durixeralf, fine-loamy, mixed, thermic.
Vegetation: Fallow. This field has been in irrigated rotation many years.
Precipitation: 12 inches. Parent Material: Granitic alluvial. Physiography: Lower end of alluvial fan. Probably occasionally flooded. Elevation: 1440 feet.
Drainage: Well drained.

HORIZON	DESCRIPTION
S57Calif-33-11-1 Ap 0-10"	Brown (10YR 5/3 dry, dark brown 3/3 moist) coarse sandy loam; moderately defined medium subangular blocky breaking to weakly defined fine granular; hard when dry, friable when moist; nonsticky and nonplastic when wet; moderately rapid permeability; many roots; many fine pores; pH 5.5; noncalcareous; abrupt smooth lower boundary.
S57Calif-33-11-2 A3 10-16"	Dark brown (10YR 4/3 dry, dark brown 7.5YR 3/4 moist) sandy loam; moderately defined medium angular blocky breaking to weakly defined fine granular; hard when dry, friable when moist; nonsticky and slightly plastic when wet; moderately rapid permeability; many plant roots; many fine pores; pH 6.5; noncalcareous; gradual smooth lower boundary.
S57Calif-33-11-3 B21 16-25"	Brown to dark brown (7.5YR 5/4-4/4 dry, dark brown 3/2 moist) sandy clay loam; strongly defined medium angular blocky; very hard when dry, firm when moist; slightly sticky and plastic when wet; moderate permeability; occasional roots; many pores; pH 7.5; noncalcareous; weak discontinuous clay skins; gradual smooth lower boundary.
S57Calif-33-11-4 B22 25-30"	Brown (7.5YR 5/4 dry, dark brown 3/2 moist) sandy clay loam; strongly defined medium angular blocky; very hard when dry, firm when moist; sticky and plastic when wet; moderately slow permeability; occasional roots; many pores; pH 7.5; noncalcareous; weak discontinuous clay skins; manganese stainings; slight mottling just above contact; abrupt wavy lower boundary.
S57Calif-33-11-5 B3cam 30-37"	Brown (7.5YR 5/4 dry, dark brown 3/2 moist) sandy clay loam; massive, strongly cemented with lime in seams; extremely hard when dry, extremely firm when moist; slightly sticky and slightly plastic when wet; few roots; many pores; pH 8.2; strong effervescence in seams; moderately strong discontinuous clay skins; gradual wavy lower boundary.
S57Calif-33-11-6 Cm1 37-50"	Brown (7.5YR 5/4 dry, dark brown 3/2 moist) sandy clay loam; massive; weakly cemented; very hard when dry, very firm when moist; slightly sticky and plastic when wet; no roots; many pores; pH 7.5; slight effervescence in seams; diffuse irregular lower boundary.
S57Calif-33-11-7 Cm2 50-60"+	Brown (7.5 YR 5/4 dry, dark brown 3/4 moist) coarse sandy clay loam; massive, weakly cemented; very hard when dry, very firm when moist; slightly sticky and plastic when wet; no roots; many pores; pH 7.5; weak effervescence in seams.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Durixeralf, fine-loamy, mixed, thermic

SOIL Monserate loam SOIL Nos. S4557-9 LOCATION Tehama County, California

SOIL SURVEY LABORATORY _____ LAB. Nos. _____

Depth (in.)	Horizon	1B1b Total: Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Sand: Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt: 0.05–0.02	Int. III (0.02–0.002)	3A1 Int. II (0.2–0.02)	(2–0.1)			Coarse fragments: ≥ 2 Pct.	2–19	19–76
		Pct. of < 2 mm															Pct. of < 76mm	
0-6½	Ap	45.1	45.7	9.2	7.5	7.4	4.5	9.2	16.5	23.9	21.8		28.6					
6½-17	A11	43.6	45.6	10.8	6.8	6.6	4.3	8.9	17.0	24.2	21.4		26.6					
17-25	A12	41.6	46.9	11.5	7.0	6.2	4.0	8.3	16.1	24.0	22.9		25.5					
25-31½	B21t	38.3	42.1	19.6	8.5	6.0	3.4	6.8	13.6	20.0	22.1		24.7					
31½-37	B22t	53.4	24.2	22.4	19.5	14.9	5.5	7.4	6.1	11.1	13.1		47.3					
37-38	C1m																	
38-40	C2m																	

Depth (in.)	6A1a Organic carbon	6B1a Nitrogen	C/N		Carbonate as $CaCO_3$		Bulk density				Water content					pH	8C1a (1:1)
	Pct	Pct.			Pct.		g/cc	g/cc	g/cc		Pct.	Pct.	Pct.				
0-6½	0.38	0.045															5.5
6½-17	0.12	0.026															5.5
17-25	0.09	0.024															5.9
25-31½	0.05	0.022															6.6
31½-37	0.04	0.018															
37-38																	
38-40																	

Depth (in.)	Extractable bases: Ca 6N2b	Mg 6O2b	Na 1/	K 2/ (5B1a)	Sum of bases	6H2a Ext. Acidity	5A3a CEC Sum	Base saturation: 5C3 Sum	
	meq/100 g							Pct.	Pct.
0-6½	1.6	0.6	0.03	0.32	2.5	2.7	5.2	49	
6½-17	1.8	1.1	0.02	0.16	3.1	2.3	5.4	57	
17-25	2.0	2.3	0.03	0.10	4.4	2.1	6.5	68	
25-31½	2.9	5.3	0.27	0.05	8.5	2.2	10.7	80	
31½-37	3.6	7.2	0.59	0.18	11.6	3.2	14.8	78	
37-38									
38-40									

Depth (in.)

1/ Gravimetrically as sodium uranyl magnesium acetate

2/ Precipitated as potassium sodium cobaltinitrite and titrated as potassium permanganate

MONSERATE LOAM
S4557-9

Location: Tehama County, California, 0.13 mile south and 0.03 mile west of bench mark 307 in NE¼ NW¼ Sec. 29, T25N, R3W MDB&M. Date of Sampling: April 4, 1945.

Description by: R. C. Roberts and R. A. Gardner. Collectors: R. C. Roberts, F. O. Youngs, R. A. Gardner, L. T. Alexander, H. Jenny and R. E. Storie.

Classification: Typic Durixeralf, fine-loamy, mixed, thermic.

Vegetation: Used as sheep pasture. Short annual grasses and associated herbaceous plants. Climate: Average annual precipitation is 20 inches. Average annual temperature is 62°F., average January temperature is 45°F., and average July temperature is 82°F. Parent Material: Mixed. Topography: Nearly level to undulating [illegible] terraces of mixed valley fill materials on a general slope of about 1 percent.

Elevation: 310 feet. Drainage: Well drained, very slow permeability, slow runoff. Soil Moisture: Moist to 38 inches, dry below.

Remarks: This soil profile appears to be very similar to the profile of Redding gravelly loam. It has less gravel in the upper part of the profile, less clay above the hardpan, but a more indurated hardpan than Redding gravelly loam. The hardpan layer was extremely hard to excavate even with the use of a heavy pick.

HORIZON	DESCRIPTION
Ap	0 to 6.5 inches, yellowish red (5YR 5/6) light loam containing a slight amount of gravel, dark red (2.5YR 3/6) moist; weak granular structure; slightly hard, friable, slightly plastic; abundant fine roots; strongly acid (pH 5.5); distinct smooth boundary.
A11	6.5 to 17 inches, yellowish red (5YR 5/6) loam containing a slight amount of gravel, red (2.5YR 4/6) moist; weak blocky structure; consistence similar to layer above; plentiful fine roots; strongly acid (pH 5.5); distinct smooth boundary.
A12	17 to 25 inches, yellowish red (5YR 5/7) loam, red (2.5YR 4/6) moist; weak blocky structure; hard, friable, plastic; few fine roots; medium acid (pH 5.9); abrupt smooth boundary.
B21t	25 to 31.5 inches, red (2.5YR 5/6) loam containing a slight amount of gravel, red (2.5YR 4/7) moist; medium to coarse blocky structure with a tendency towards prismatic structure; compact; hard, firm, plastic; very few fine roots; neutral (pH 6.6); diffuse boundary.
B22t	31.5 to 37 inches, red gravelly sandy clay loam, color dry and moist and structure and consistence similar to layer above except weakly cemented; very few roots; medium acid (pH 6.0); about ¼ to ½ inch of lighter colored or more yellowish sandy clay loam of pH about 6.0 occurring in this boundary was not sampled; abrupt wavy boundary.
C1m	37 to 38 inches, red (10R 4/5) gravelly indurated hardpan, dusky red (10R 3/4) moist; massive; extremely hard; very few roots; medium acid (pH 6.0); diffuse boundary.
C2m	38 to 50 inches, red (10R 4/5) gravelly indurated hardpan; very similar to layer above; very few roots; slightly acid (pH 6.5).

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Monserate sandy loam **LOCATION** Riverside County, California

SOIL FAMILY: Typic Durixeralf, fine-loamy, mixed, thermic

SOIL NOS. S57Calif-33-13 LAB. NOS. 57542 - 57548

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	3A1 0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-5	A1	9.9	13.9	9.0	19.0	12.8	27.7	7.7	41.8	10.2	10	sl
5-10	A3/B1	18.6	15.5	6.7	12.4	8.5	21.4	16.9	27.8	9.5	21	cosl
10-20	B21	9.2	13.1	6.7	14.3	7.8	18.1	30.8	25.4	8.9	22	scl
20-28	B22	10.8	14.2	7.4	17.2	9.2	19.6	21.6	30.0	9.2	24	scl
28-45	Cm1	20.8	22.3	10.9	17.9	7.1	14.9	6.1	25.1	6.6	25	lcos*
45-57	Cm2	22.7	25.7	10.5	19.2	7.2	9.6	5.1	23.8	3.5	29	lcos*
57-70+	C3	77.0	4.9	1.8	2.4	0.6	8.3	5.0	6.6	3.5	29	lcos*

8C1b pH SATURATED PASTE 1:1	pH 1:5	8C1a pH 1:10	6A1a ORGANIC MATTER ORGANIC CARBON %	6B1a NITROGEN %	C/N	6I1a CO_3 meq/l	8A1a Elec. Conduc. mmhos/cm at 25°C	6E1b $CaCO_3$ equivalent %	6J1a HCO_3 meq/l	MOISTURE TENSIONS 4B1a 1/10 ATMOS. %	4B1a 1/3 ATMOS. %	4B2 15 ATMOS. %
6.7		6.7	0.57	0.063	9.0	0.0	0.8	0	3.5	20.3	12.1	3.8
6.7		6.4	0.29	0.032	9.1	0.0	0.4	0	1.5			5.8
6.6		6.7	0.24	0.026	9.2	0.0	0.2	0	1.6			10.2
6.9		6.8	0.12	0.015		0.0	0.5	0	1.6			7.9
7.9		7.4	0.03			0.0	0.4	Tr	2.3			4.6
7.9		7.8	0.01			0.0	0.3	Tr	2.4			3.8
7.9		7.7	0.02			0.0	0.4	Tr	2.2			3.8

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a Ca	6O1a Mg	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
7.3	4.3	2.2	1.8	0.3	0.7	81	0.7	1.6	3.9	1.5	1.1	26.9
9.1	5.6	2.6	2.3	0.3	0.4	79	0.6	0.3	1.5	0.9	1.0	24.2
17.8	10.9	4.8	2.6	0.5	0.3	86	1.0	0.1	1.2	0.9	0.5	33.9
18.0	11.7	4.6	1.7	0.7	0.2	91	1.9	0.1	1.7	1.2	1.5	32.1
11.0	7.2	2.5	-	0.8	0.2	100	2.2	0.1	1.1	0.6	1.0	27.5
8.9	6.4	2.1	-	0.5	0.2	100+	2.2	0.1	1.0	0.6	0.5	28.7
8.3	6.4	2.4	-	0.6	0.2	100+	2.3	0.1	1.0	0.7	0.7	27.3

1/ Percent of the whole soil

* Incompletely dispersed

MONSERATE SANDY LOAM
S57Calif-33-13

Location: 1200 feet east and 150 north of the southwest corner of Sec. 33, T. 2 S., R. 3 W. On Maxwell-Taft Ranch, about 3 miles ENE of Sunnymead, Riverside County, California. Date of Sampling: June 7, 1957.
Description by: Fred Eggers. Collectors: George Borst, R. E. Nelson, Cliff Henry, Arnold Knecht, and Fred Eggers.
Classification: Typic Durixeralf, fine-loamy, mixed, thermic.
Vegetation: California sage; wild buckwheat; salvia; turkey mullein and annual grasses. Precipitation: 12 inches. Parent Material: Granitic alluvium. Physiography: Old terrace, strongly dissected site on top of ridge about 300 feet wide. Elevation: 1870 feet. Drainage: External drainage rapid; internal very slow.

HORIZON	DESCRIPTION
S57Calif-33-13-1 A1 0-5"	Brown (7.5YR 5/4 dry, dark reddish brown 5YR 3/4 moist) sandy loam; moderately defined medium granular; slightly hard when dry, friable when moist; nonsticky and nonplastic when wet; moderately rapid permeability; many plant roots; many fine pores; pH 6.0; noncalcareous; clear smooth lower boundary.
S57Calif-33-13-2 A3 or B1 5-10"	Yellowish red (5YR 4/6 dry, dark reddish brown 2.5YR moist) loam; moderately defined medium subangular blocky with some indication of prismatic structure; hard when dry, firm when moist; slightly sticky and slightly plastic when wet; moderate permeability; occasional plant roots; many fine pores; pH 6.5; noncalcareous; clear wavy lower boundary.
S57Calif-33-13-3 B21 10-20"	Reddish brown (5YR 4/4 dry, dark reddish brown 2.5YR 3/4 moist) coarse sandy clay; strongly defined medium prismatic; extremely hard when dry, extremely firm when moist; very sticky and very plastic when wet; very slow permeability; occasional plant roots; occasional pores; pH 6.7; gradual wavy lower boundary.
S57Calif-33-13-4 B22 20-28"	Reddish brown (5YR 5/4 dry, dark reddish brown 3/4 moist) coarse sandy clay; strongly defined medium angular blocky with some indication of prismatic structure; extremely hard when dry, extremely firm when moist; very sticky and very plastic when wet; very slow permeability; occasional roots; occasional pores; pH 7.2; noncalcareous; abrupt wavy lower boundary.
S57Calif-33-13-5 Cm1 28-45"	Dark brown (10YR 4/3 dry, dark reddish brown 5 YR 3/4 moist) loamy coarse sand; massive; strongly cemented; extremely hard when dry; after crushing, loose when moist; slightly sticky and nonplastic when wet; a few roots in fissures; pH 7.7; gradual wavy lower boundary.
S57Calif-33-13-6 Cm2 45-57"	Yellowish brown (10YR 5/4 dry, dark reddish brown 5YR 3/4 moist) coarse sand; massive; weakly cemented; very hard when dry; after crushing, loose when moist; nonsticky and nonplastic when wet; few roots in fissures; pH 7.6; noncalcareous; clear wavy lower boundary.
S57Calif-33-13-7 C3 57-70"+	Yellowish brown (10YR 5/4 dry, dark reddish brown 5YR 3/4 moist) coarse sand; single grain with weakly cemented lenses; loose when dry, loose when moist; nonsticky and nonplastic when wet; very few roots; pH 7.5; noncalcareous.

Remarks: Larger than 2mm subangular blocky rock fragments throughout profile. The C horizons contain many quartz fragments which give a pepper and salt appearance.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Durixeralf, fine-loamy, mixed, thermic

SOIL Monserate loam SOIL Nos. S66Calif-37-1 LOCATION San Diego County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66233 - 66235

Depth (In)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: (0.05-0.02)	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)			1/ Coarse fragments 1A2a > 2 Pct.	2-19	19-76
		Pct. of < 2 mm															Pct. of < 76mm	
0- 2	A11	49.6	35.4	15.0	2.3	13.7	16.5	12.2	4.9	16.7	18.7	26.3	44.7	0	15	13	9	4 2/
2-15	A12	46.8	34.9	18.3	1.8	11.5	15.4	12.7	5.4	15.5	19.4	26.5	41.4	0	18	12	9	3 2/
15-33	B1	46.1	32.9	21.0	2.4	11.8	15.0	11.7	5.2	14.4	18.5	24.7	40.9	0	21	12	8	4 2/

Depth (In.)	6A1a Organic carbon Pct	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1
0- 2	2.01					1.66	1.73		0.91		13.3	6.4	0.013	0.011	5.6
2-15	0.85					1.51	1.58		0.93		12.4	6.5	0.016	0.015	5.3
15-33	0.56					1.60	1.67		0.92		13.5	7.6	0.015	0.014	4.4

MONSERATE LOAM
S66Calif-37-1

Location: San Diego County, California; Miramar Transect No. 1, station 23.8; 950 feet west and 750 feet north of the southeast corner of section 24, T. 15 S., R. 3 W. Date of sampling: June 10, 1966.
Description by: George Borst, David Estrada, Gerald Kester, Reuben Nelson, and Dennis Nettleton. Samples collected by: Klaus Flach, Benny Brasher, and Dennis Nettleton.
Classification: Typic Durixeralf; fine loamy, mixed, thermic family.
Vegetation: Sparse chamise and flattop buckwheat. Climate: Weather record for San Diego, near the sampling site, is mean winter air temperature 55° F., mean summer air temperature 68° F., mean annual precipitation 10 inches. Parent Material: Stream deposit over Pleistocene marine terrace. Topography: Flat marine terrace of less than 1 percent slope. Elevation: About 425 to 450 feet. Drainage: Moderately well drained; runoff is medium; permeability is very slow. Soil Moisture: Dry.
Remarks: 15 percent rounded gravel throughout the profile.

HORIZON	DESCRIPTION
A11 RSL No. 66233	0 to 2 inches, reddish brown (5YR 5/4) loam, reddish brown (5YR 4/4) when moist; massive; slightly hard, friable, slightly sticky and plastic; abundant very fine roots; few very fine tubular pores; medium acid (pH 6.0); abrupt smooth boundary.
A12 RSL No. 66234	2 to 15 inches, reddish brown (5YR 5/4) loam, dark reddish brown (2.5YR 3/4) when moist; massive; hard, firm, sticky, very plastic; few fine to medium roots; many fine tubular pores; strongly acid (pH 5.5); many animal burrows and krotovinas, some approximately 3 inches in diameter are probably pocket gopher burrows, other approximately 3/8 inch in diameter are probably insect burrows; gradual wavy boundary.
B1 RSL No. 66235	15 to 33 inches, reddish brown (5YR 5/4) clay loam, dark reddish brown (2.5YR 3/4) when moist; massive; hard, firm, very sticky, very plastic; few fine roots mostly in krotovinas; many fine tubular pores; very strongly acid (pH 4.5); common krotovinas filled with weakly granular material resembling the surface horizon; clear wavy boundary.
B2t	33 to 35 inches, reddish brown (2.5YR 5/4) heavy clay loam, dark reddish brown (2.5YR 3/4) when moist; weak medium angular blocky structure; hard, firm, very sticky, very plastic; few fine roots; many fine tubular pores; common medium clay films line tubular pores; very strongly acid (pH 4.5); very abrupt wavy boundary.
B3sim	35 inches +, grayish brown (10YR 5/2) duripan, brown to dark brown (10YR 4/2) when moist; dusky red (10R 3/3) coatings of iron oxide on fracture planes over thin, noneffervescent white coatings, probably silica.

CONSECUTIVE PEDON NUMBER C 69 18 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES MONSERATE TAXADJUNCT SOIL NO. S69 CALI. 31-10 SAMPLE NO. 9 84-9 90 PLACER COUNTY AREA

SOIL FAMILY TYPIC DURIXERALF FINE-LOAMY,MIXED,THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1-.5	MS .5-.25	FS .25-.10	VFS .1-.05	SILTS CSI .05-.02	FSI .02-.002	CLAYS CO. 2-.2U	FINE LT .2U	CARBO NATE 3A1A	1A2A INT. II .2-.02	COARSE FRAGMENTS PCT. OF WHOLE SOIL I 75-20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75		
9 84	0- 24	AP	65.0	26.4	8.6	11.1	14.0	9.3	19.7	10.9	13.2	13.2				35.0	--A	--	--	--
9 85	24- 46	A12	59.7	28.9	11.4	4.8	10.9	8.7	21.4	13.9	14.9	14.0				41.5	--A	--	--	--
9 86	46- 60	A3	60.9	28.4	10.7	7.7	13.3	8.9	19.8	11.2	15.1	13.3				37.6	--A	--	--	--
9 87	60- 80	B2 T	53.0	21.5	25.5	6.6	10.1	8.0	18.9	9.4	10.2	11.3				30.4	--A	--	--	--
9 88	80- 92	B3 T	57.8	15.1	27.1	8.7	13.1	10.2	18.6	7.2	6.7	8.4				22.8	--A	--	--	--
9 89	92-105	C1 SIM	72.5	16.8	10.7	9.5	30.1	14.1	12.9	5.9	7.3	9.5				19.0	--G	--	--	--
9 90	105-135	2C 2SIM	36.6	44.6	18.8	3.4	4.6	3.8	14.0	10.8	16.6	28.0				36.2	--G	--	--	--

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	19-2 6E1B	EXT. IRON FE 6C2B	ATTERBERG LIMIT EST LL 4F2	PL 4F1	BULK DENSITY EST 1/3 BAR 4A1F	4A1F	DRY 4AH	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTENSBLTY LEF 4D1	LE 4D1	PH H2O 1/1 8C1A	H2O 1/10 8C1A	CACL2 8C1E
		-PERCENT--		-PERCENT-		PCT.	PCT.	NO.40	GRAMS PER CC				----PERCENT---			-PERCENT--				
9 84	0- 24	0.47	0.042						1.60			1.00			3.6			5.9		5.3
9 85	24- 46	0.20	0.021							1.69	1.76	1.00		11.7	4.2	1.4	1.4	6.0		5.4
9 86	46- 60	0.13							1.68			1.00			4.6			6.0		5.3
9 87	60- 80	0.16								1.66	1.96	1.00		19.7	11.4	5.7	5.7	6.5		5.9
9 88	80- 92								1.67			1.00			11.2			7.0		6.4
9 89	92-105				1				1.35			1.00			9.6			7.7		7.1
9 90	105-135								1.35			1.00			13.5			7.5		7.1

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6Q2D	NA 5B4A 6P2A	K 6Q2A	SUM	BASE SAT.	CEC NA- OAC 5A2A	NH4- OAC 5A6A	EC 8A1A	WATER EXTRACT FROM SATURATED PASTE 8A1 CA 6N1B	MG 6Q1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	B 6T1A	PH SAT. PASTE 8C1B
		MEQ/100G					PCT.	-MEQ/100G		MMHO	MEQ/100D ML								PPM	
9 84	0- 24	4.0	1.7	0.1	0.2	6.0	95		6.3											
9 85	24- 46	4.4	2.6	0.1	0.1	7.2	100+		7.1											
9 86	46- 60	4.2	2.2	0.2	0.1	6.7	96		7.0											
9 87	60- 80	9.8	6.4	0.7	0.2	17.1	98		17.5											
9 88	80- 92	10.9	8.3	0.9	0.2	20.3	100+		18.4											
9 89	92-105	16.5	7.6	1.1	0.2	25.4	100+		19.3	0.86										
9 90	105-135	16.3	9.9	1.3	0.3	27.8	100+		26.4											

SAMPLE NO.	DEPTH CM.	WATER AT SAT. 8A PCT.	EXCHANGBLE NA 5B4B MEQ.	NA 5D2 PCT.	SAR 5E	GYPSUM AMT. REQ. 6F1B PERCENT	6P3A	CLAY MINERALOGY LESS THAN 0.002 MM X RAY I 7A2E	II 7A2E	III 7A2E	IV 7A2E	DTA 7A3 PCT.	L.T. .0002 MM X RAY I 7A2E	II 7A2E	III 7A2E
9 84	0- 24		0.1	2				MI 3	KK 3						
9 85	24- 46		0.1	1				MT 3	KK 3	MI 2					
9 86	46- 60		0.2	3											
9 87	60- 80		0.7	4				MI 3	KK 3	MT 2					
9 88	80- 92		0.9	5											
9 89	92-105		1.1	6											
9 90	105-135		1.3	5											

MINERAL CODE: MI MICA, KK KAOLINIT, MT MONTMORT

RELATIVE AMOUNT: 1 TRACE, 2 SMALL, 3 MODERATE, 4 ABUNDANT, 5 DOMINANT, 6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
G=GROUND WHOLE. HORIZON INDURATED OR GRAVEL TOO SOFT TO BE SEPARATED. ALL DATA FOR WHOLE SOIL.

MONSERATE LOAM (TAXADJUNCT) 1/
S69Calif-31-10

Location: Placer County, California. Approximately 3.5 miles southwest of Lincoln, 300 yards west of California 65 and 150 feet north of Pleasant Grove Road; 1,500 feet west and 150 feet north of the SE corner of sec. 33, T. 12 N., R. 6 E. Pit located approximately 100 feet from Trigo site (S69Calif-31-11). Date of Sampling: May 7, 1969.
Description by: G. M. Kennedy, C. B. Goudey. Collectors: K. W. Flach, W. D. Nettleton, G. M. Kennedy, L. A. Brittan, T. M. Ryan.
Classification: Typic Durixeralf, fine-loamy, mixed, thermic. *
Vegetation: Rose clover, soft chess, ripgut brome, and filaree. Climate: Average annual rainfall 17 to 32 inches. Mean annual air temperature 61°F. Parent Material: Surficial sediments deposited over Laguna Formation. Topography: Undulating terrace slope of 4 percent on southeast aspect. Elevation: 120 feet. Drainage: Moderately well drained; slow to medium runoff; very slow permeability. Soil Moisture: Dry on surface and moist below 12 inches.
Remarks: Texture of A12 horizon feels slightly finer than Ap horizon above. Roots matting out on top of C1sim horizon. Top of IIC1sim has a thin (1/8 to 1/4") silica cementation.
*Varies slightly from range of series as now defined.

HORIZON	DESCRIPTION
Ap RSL 6984	(0-24 cm.) 0 to 9½ inches, grayish brown (10YR 5/2) loam, dark brown (10YR 3/3) moist; massive; hard, friable, slightly sticky, slightly plastic; common very fine vertical roots; common very fine tubular pores; medium acid; (pH 6.0); clear wavy boundary.
A12 RSL 6985	(24-46 cm.) 9½ to 18 inches, brown (10YR 5/3) loam, dark brown (10YR 3/3) moist; massive; hard, friable, slightly sticky, slightly plastic; common very fine vertical roots; common very fine and fine tubular pores; medium acid (pH 6.0); clear wavy boundary.
A3 RSL 6986	(46-60 cm.) 18 to 23½ inches, light brownish gray (10YR 6/2) sandy clay loam, dark brown (10YR 4/3) moist; massive; hard, friable, slightly sticky, slightly plastic; few very fine random roots; common very fine tubular and interstitial pores; medium acid (pH 6.0); abrupt wavy boundary.
B2t RSL 6987	(60-80 cm.) 23½ to 31½ inches, **brown (10YR 5/3)** sandy clay; very dark grayish brown (10YR 3/2) moist; weak coarse prismatic parting to moderate coarse angular blocky structure; very hard, firm, sticky, plastic; very few very fine random roots; few very fine interstitial pores; common thin clay films on faces of peds and bridging mineral grains; slightly acid (pH 6.2); clear wavy boundary.
B3t RSL 6988	(80-92 cm.) 31½ to 36 inches, **brown (10YR 5/3)** sandy clay loam, dark grayish brown (10YR 4/2) moist; weak coarse angular blocky structure; very hard, firm, sticky, slightly plastic; very few very fine random roots; few very fine interstitial pores; common thin clay films on faces of peds and bridging mineral grains; slightly acid (pH 6.5); abrupt smooth boundary.
C1sim RSL 6989	(92-105 cm.) 36 to 42 inches, light yellowish brown (10YR 6/4) strongly cemented silty material, brown (10YR 5/3) moist; common medium distinct yellowish brown (10YR 5/6) mottles; massive; extremely hard, very firm, brittle; abrupt smooth boundary.
IIC2sim RSL 6990	(105-135 cm.) 42 to 53 inches, siltstone.

1/ The Monserate series is in a fine-loamy, mixed, thermic family of Typic Durixeralfs. This pedon is in the same family but has more silt than is typical of the series.

CONSECUTIVE PEDON NUMBER C 69 25 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES SAN JOAQUIN TAXADJUNCT SOIL NO. S69 CALI. 31-20 SAMPLE NO. 9 115-9 122 PLACER COUNTY AREA

SOIL FAMILY TYPIC DURIXERALF FINE-LOAMY,MIXED,THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1-.05	MS .5-.25	FS .25-.10	VFS .1-.05	SILTS CSI .05-.02	FSI .02-.002	CLAYS CO. 2-.2U	FINE LT .2U	CARBONATE 3A1A.02	INT. II .2-.02 WHOLE 20	1A2A COARSE FRAGMENTS PCT. OF SOIL	I 75- PCT. OF LT 75	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM																	
9 115	0- 28	AP	52.4	38.2	9.4	6.3	10.8	6.8	15.6	12.9	19.9	18.3	6.5	2.9		42.2	--A	--	--	--
9 116	28- 38	B11T	48.9	37.4	13.7	6.1	9.8	6.5	14.5	12.0	19.8	17.6	8.9	4.8		40.5	--A	--	--	--
9 117	38- 51	B12T	45.6	37.3	17.1	4.5	8.4	5.9	14.5	12.3	19.0	18.3	8.1	9.0		40.1	--A	--	--	--
9 118	51- 79	2B2 T	24.7	32.3	43.0	0.7	2.7	2.5	9.8	9.0	12.8	19.5	17.2	25.8		28.2	--A	--	--	--
9 119	79- 94	2B31TBIM	56.2	26.1	7.7	10.9	16.3	9.6	19.1	10.3	12.1	14.0				33.3	--G	--	--	--
9 120	94-115	2B32TBIM	49.0	40.8	10.2	2.6	9.6	7.9	17.6	11.3	17.0	23.8				38.4	--G	--	--	--
9 121	115-155	2B33TBIM	31.2	51.6	17.2	1.9	4.9	4.4	10.7	9.3	16.5	35.1				31.9	--G	--	--	--
9 122	155-190	2C SI	31.9	50.4	17.7	2.0	5.2	4.7	10.9	9.1	16.5	33.9				32.0	--A	--	--	--

SAMPLE NO.	DEPTH CM.	ORG. MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 19-2	EXT. IRON FE 6C2B	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST 1/3 BAR 4A1F	DRY 4A1F	4AH	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTENSBLTY LEF 4D1	LE 4D1	PH H2O 1/1 8C1A	H2O 1/10 8C1A	CACL2 8C1E
		PERCENT		PERCENT	PCT.	PCT.	NO.40	GRAMS PER CC				PERCENT			PERCENT				
9 115	0- 28	0.30	0.032		1.0			1.79	1.81	1.00		12.7	4.4	0.4	0.4	5.6		5.0
9 116	28- 38	0.16			1.4			1.70	1.75	1.00		13.3	5.5	1.0	1.0	5.8		5.2
9 117	38- 51	0.13			1.6			1.73	1.81	1.00		14.3	6.9	1.5	1.5	6.1		5.5
9 118	51- 79				2.0			1.62	1.97	1.00		20.4	17.4	6.7	6.7	6.7		6.3
9 119	79- 94				1.4			1.90	1.97	1.00		13.4	12.1	1.2	1.2	7.3		6.6
9 120	94-115				1.0			1.90	1.99	1.00		14.0	11.0	1.6	1.6	7.6		6.6
9 121	115-155				0.9			1.80	1.91	1.00		15.9	12.6	2.0	2.0	7.7		7.3
9 122	155-190				1.0			1.80	1.91	1.00		15.8	14.0	2.0	2.0	7.8		7.2

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT.	C E C NA-OAC 5A2A	NH4-OAC 5A6A	EC 8A1A	WATER EXTRACT FROM SATURATED PASTE CA 6N1B	MG 6O1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	B 6T1A	PH SAT. PASTE 8C1B
		MEQ/100G					PCT.	MEQ/100G		MMHO	MEQ/1000 ML								PPM	
9 115	0- 28	2.9	1.1	0.1	0.1	4.2	74		5.7											
9 116	28- 38	4.0	2.0	0.1	0.1	6.2	91		6.8											
9 117	38- 51	4.6	3.1	0.1	0.1	7.9	92		8.6											
9 118	51- 79	16.5	13.4	0.5	0.3	30.7	99		31.1											
9 119	79- 94	17.9	14.4	0.6	0.2	33.1	100+		31.3											
9 120	94-115	18.1	13.0	0.5	0.2	31.8	100+		27.6											
9 121	115-155	19.3	14.3	0.5	0.3	34.4	100+		30.0											
9 122	155-190	21.2	14.8	0.6	0.3	36.9	100+		29.9											

SAMPLE NO.	DEPTH CM.	WATER AT SAT. 8A PCT.	EXCHANGBLE NA 5B4B MEQ.	NA 5D2 PCT.	SAR 5E	GYPSUM AMT. 6F1B PERCENT	REQ. 6P3A	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 PCT.	L.T. .0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
9 115	0- 28		0.1	2													1 TRACE
[illegible]	[illegible]		[illegible]	[illegible]												MI MICA	2 SMALL

SAN JOAQUIN SANDY LOAM (TAXADJUNCT) 1/
S69Calif-31-20

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Durixeralf, fine, montmorillonitic, thermic

SOIL Tusean gravelly sandy loam (taxadjunct) SOIL Nos. 84557-15 LOCATION San Diego County, California

SOIL SURVEY LABORATORY _____ LAB. Nos. _____

Depth (In.)	Horizon	1B1b Total Sand (2–0.05)	Total Silt (0.05–0.002)	Total Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)			Coarse fragments > 2 Pct.	2 – 19	19 – 76
		Pct. of < 2 mm															Pct. of < 76mm	
0-3/4	A11	55.7	30.9	13.4	2.8	12.4	16.6	16.9	7.0	13.0	17.9		48.7					
3/4-4-1/4	A12	59.1	27.6	13.3	2.0	12.4	18.5	18.9	7.3	11.8	15.8		51.8					
4-1/4-7½	A13	56.4	25.8	17.8	1.9	11.0	17.7	18.5	7.3	10.5	15.3		49.1					
7½-8-3/4	B1t	54.5	24.2	21.3	0.8	9.9	17.5	19.2	7.1	9.9	14.3		47.4					
8-3/4-13-3/4	B21t	47.5	19.3	33.2	0.7	9.2	15.9	16.1	5.6	6.3	13.0		41.9					
13-3/4-18-3/4	B22t	33.5	15.7	50.8	1.2	7.2	10.5	10.4	4.2	4.7	11.0		29.3					
18-3/4-21	B23t	30.7	14.4	54.9	1.6	6.6	10.3	8.5	3.7	3.6	10.8		27.0					
21-23-3/4	B24t	36.6	10.8	52.6	1.9	9.4	14.0	8.1	3.2	3.2	7.6		33.4					
23-3/4-30	Cm																	

Depth (In.)	6A2a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	Pct.			pH		8C1a (1:1)
0-3/4	8.00	0.406																6.1
3/4-4-1/4	1.29	0.085																6.4
4-1/4-7½	0.66	0.055																6.4
7½-8-3/4	0.52	0.047																6.3
8-3/4-13-3/4	0.45	0.046																6.8
13-3/4-18-3/4	0.48	0.048																5.4
18-3/4-21	0.52	0.052																4.8
21-23-3/4	0.48	0.043																4.5
23-3/4-30	0.07	0.007																5.3

Depth (In.)	Extractable bases: Ca 6N2b	Mg 6O2b	Na 1/	K 2/ (5B1a)	Sum of bases	6H2a Ext. Acidity	5A3a CEC Sum									Base saturation 5C3	
	meq/100 g															Pct.	Pct.
0-3/4	24.6	4.6	0.16	0.64	30.0	8.9	38.9									77	
3/4-4-1/4	7.4	2.3	0.05	0.38	10.1	2.7	12.8									79	
4-1/4-7½	5.1	2.7	0.07	0.41	8.3	2.8	11.1									75	
7½-8-3/4	4.3	3.1	0.07	0.32	7.8	2.9	10.7									73	
8-3/4-13-3/4	3.9	4.6	1.24	0.55	10.3	3.0	13.3									77	
13-3/4-18-3/4	4.1	6.7	0.74	0.20	11.7	5.8	17.5									67	
18-3/4-21	6.5	13.4	1.74	0.27	21.9	7.5	29.4									75	
21-23-3/4	7.0	14.6	2.03	0.30	23.9	7.5	31.4									76	
23-3/4-30	3.9	6.7	2.12	0.08	12.8	4.5	17.3									74	

Depth (In.)				

1/ Gravimetrically as sodium uranyl magnesium acetate

2/ Precipitated as potassium sodium cobaltinitrite and titrated as potassium permanganate

TUSCAN GRAVELLY SANDY LOAM (taxadjunct) 1/
S4557-15

Location: San Diego County, California, 1/3 mile south and 1/10 mile west of bench mark 417 located about 2½ miles south of Camp Kearney. Date of Sampling: April 15, 1945.

Description by: R. C. Roberts and R. A. Gardner. Collectors: R. C. Roberts, F. O. Youngs, R. A. Gardner, L. T. Alexander, H. Jenny and R. E. Storie.

Classification: Typic Durixeralf, fine, montmorillonitic, thermic.

Vegetation: Chamise brush. Climate: Average annual precipitation is about 11 inches. Average annual temperature is about 62°F., January 54°F., July 70°F. Parent Material: Mixed older alluvium. Topography: Nearly level higher terraces. Elevation: 415 feet. Drainage: Well drained, very slow permeability, slow run-off. Soil Moisture: Dry.

Remarks: This soil profile is typical of a large area of soils in southern California. The color appears to be about as red as any sample collected for this study; the boundary between the surface and subsoil is as abrupt as in any other profile sampled; the claypan is just as compact; and the layer below the claypan is just as hard or harder than any of the other soils that have indurated layers. There does not appear to be so much manganese in this soil as in samples of the other soils, but the acidity is as strong or stronger, especially in the lower part of the profile. In some areas there is a thin but distinct gray layer immediately above the claypan.

HORIZON	DESCRIPTION
A11	0 to 0.75 inch, dark reddish brown (5YR 3/3) sandy loam containing considerable fine fragments of what appears to be burned organic matter that affects color of layer, dark reddish brown (5YR 2/2) moist; strong fine granular structure; soft, friable, slightly plastic; abundant roots and root remains; slightly acid (pH 6.1).
A12	0.75 to 4.25 inches, dark red (2.5YR 3/6) gravelly sandy loam, dark reddish brown (2.5YR 3/4) moist; very fine granular and fine blocky structure; slightly hard, friable, slightly plastic; slightly acid (pH 6.4).
A13	4.25 to 7.5 inches, dark red (2.5YR 3/6) gravelly heavy sandy loam, color dry and moist, structure and consistence similar to layer above; slightly acid (pH 6.4).
B1t	7.5 to 8.75 inches, dark red (2.5YR 3/6) gravelly light sandy clay loam, color dry and moist similar to layer above; granular structure; compact; hard, firm, plastic; slightly acid (pH 6.3).
B21t	8.75 to 13.75 inches, red (10R 4/6) gravelly sandy clay loam, dark red (10R 3/6) moist; blocky structure; compact; hard, firm, plastic; neutral (pH 6.8).
B22t	13.75 to 18.75 inches, red (10R 4/6) gravelly clay, color dry and moist and structure similar to layer above; very compact; very hard, very firm, very plastic; strongly acid (pH 5.4).
B23t	18.75 to 21 inches, yellowish red (5YR 4/6) gravelly clay, red (2.5YR 4/6) moist; coarse blocky to columnar structure; very compact; extremely hard, extremely firm, very plastic; very strongly acid (pH 4.8).
B24t	21 to 23.75 inches, yellowish red (5YR 4/6) gravelly clay, color dry and moist, structure and consistence similar to layer above; very strongly acid (pH 4.5).
Cm	23.75 to 30 inches, red (2.5YR 5/6) gravelly indurated hardpan, red (2.5YR 4/6) moist; extremely hard and cemented; strongly acid (pH 5.3).

1/ The Tuscan series is in a fine, montmorillonitic, thermic family of Typic Durixeralfs. This pedon has redder hues and lower pH than is typical of the Tuscan series.

CONSECUTIVE PEDON NUMBER C 69 17 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES FIDDYMENT TAXADJUNCT SOIL NO. S69 CALI. 31- 8 SAMPLE NO. 9 74-9 83 PLACER COUNTY AREA

SOIL FAMILY ABRUPTIC DURIXERALF FINE-LOAMY,MIXED,THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL SAND 2-.05	FRACTION SILT .05-.002	CLAY LT .002	SANDS VCS 2-1	CS 1-.5	MS .5-.25	FS .25-.10	VFS .1-.05	SILTS CSI .05-.02	FSI .02-.002	CLAYS CO. 2-.2U	FINE LT .2U	CARBONATE 3A1A	INT. COARSE FRAGMENTS II .2-.02	1A2A PCT. OF WHOLE SOIL I 75-20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75		
9 74	0- 23	AP	42.1	49.6	8.3	0.3	1.7	3.7	19.5	16.9	23.4	26.2				53.3	--A --	--	--
9 75	23- 30	A3	37.3	51.5	11.2	0.2	1.8	3.4	17.1	14.8	21.4	30.1				47.3	--A --	--	--
9 76	30- 48	B21T	31.5	41.9	26.6	0.2	1.5	2.8	14.8	12.2	17.2	24.7				39.5	--A --	--	--
9 77	48- 63	B22T	26.5	40.0	33.5	0.1	0.9	2.1	11.5	11.9	13.3	26.7				33.0	--A --	--	--
9 78	63- 71	B3	19.8	46.6	33.6	0.5	1.8	2.1	7.8	7.6	9.7	36.9				22.2	--A --	--	--
9 79	71- 89	2C1 SIM	46.8	39.2	14.0	6.9	14.2	8.0	12.2	5.5	6.0	33.2				18.0	--G --	--	--
9 80	89-105	2C2 SIM	35.0	51.8	13.2	0.8	7.6	7.5	11.6	7.5	10.6	41.2				24.6	--G --	--	--
9 81	105-120	2C3 SI	41.8	49.2	9.0	0.9	6.1	7.0	12.1	15.7	20.5	28.7				43.5	--A --	--	--
9 82	120-150	3C4	68.9	18.3	12.8	1.3	11.1	18.7	27.0	10.8	12.1	6.2				34.6	--A --	--	--
9 83	0- 0	AP															--A --	--	--

SAMPLE NO.	DEPTH CM.	ORG. CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	19-2 6E1B	EXT. IRON FE 6C2B	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4AH	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTENSIBILITY LEF 4D1	LE 4D1	PH H2O 1/1 8C1A	H2O 1/10 8C1A	CACL2 8C1E
		-PERCENT--		-PERCENT-		PCT.	PCT.	NO.40	GRAMS PER CC				----PERCENT----			-PERCENT--				
9 74	0- 23	0.50	0.042							1.62	1.65	1.00		16.2	4.1	0.6	0.6	5.4		4.6
9 75	23- 30	0.24	0.028							1.71	1.77	1.00		15.3	5.0	1.2	1.2	5.6		4.9
9 76	30- 48	0.24	0.034							1.62	1.88	1.00		20.4	11.4	5.1	5.1	6.0		5.3
9 77	48- 63	0.24	0.032							1.58	1.91	1.00		22.6	14.4	6.5	6.5	6.4		5.8
9 78	63- 71	0.25								1.33	1.66	1.00		32.9	18.0	7.7	7.7	6.9	7.9	6.3
9 79	71- 89									1.32	1.38	1.00		37.3	19.6	1.5	1.5	7.1	8.0	6.8
9 80	89-105									1.50	1.55	1.00		24.2	11.9	1.1	1.1	7.2	7.9	6.8
9 81	105-120									1.59	1.66	1.00		19.3	8.1	1.4	1.4	7.2	7.5	6.9
9 82	120-150									1.48	1.59	1.00		19.9	8.1	2.4	2.4	7.0	7.7	6.8
9 83	0- 0								1.48			1.00			4.3			5.5		4.8

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 602D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT.	CEC NA-OAC 5A2A	NH4-OAC 5A6A	EC 8A1A	WATER EXTRACT FROM SATURATED PASTE CA 6N1B	MG 6O1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	B 6T1A	PH SAT. PASTE 8C1B
		--------MEQ/100G--------					PCT.	-MEQ/100G		MMHO	----------MEQ/1000 ML----------								PPM	
9 74	0- 23	3.6	1.9	0.1	0.1	5.7	80		7.1											
9 75	23- 30	4.6	2.3	0.3	0.1	7.3	86		8.5											
9 76	30- 48	9.6	5.6	1.0	0.2	16.4	93		17.7											
9 77	48- 63	12.5	6.4	1.8	0.2	20.9	89		23.5	0.23										
9 78	63- 71	19.5	10.9	3.0	0.2	33.6	100+		32.3	0.90	1.1	0.9	6.7	0.1		4.3	1.5			6.4
9 79	71- 89	22.7	12.1	4.6	0.1	39.5	100+		36.1	1.50	2.3	1.9	12.3	TR		2.3	8.6			7.1
9 80	89-105	13.3	7.3	2.7	0.2	23.5	100+		22.1	2.36	5.7	4.8	16.8	0.1		2.3	18.1			7.2
9 81	105-120	9.8	5.9	2.0	0.2	17.9	100+		16.2	2.36	5.2	4.3	15.2	0.1		1.7	17.9			7.1
9 82	120-150	9.4	5.2	1.7	0.2	16.5	100+		14.6	2.26	5.7	4.5	15.0	0.1		2.4	17.1			7.0
9 83	0- 0																			

SAMPLE NO.	DEPTH CM.	WATER AT SAT. 8A PCT.	EXCHANGEABLE NA 5B4B MEQ.	NA 5D2 PCT.	SAR 5E	GYPSUM AMT. 6F1B PERCENT	REQ. 6P3A	CLAY MINERALOGY LESS THAN 0.002 MM X RAY I 7A2E	II 7A2E	III 7A2E	IV 7A2E	DTA PCT. 7A3	LT. .0002 MM X RAY I 7A2E	II 7A2E	III 7A2E	MINERAL CODE	RELATIVE AMOUNT
9 74	0- 23		0.1	1				MI 4	KK 3							MI MICA	1 TRACE
9 75	23- 30		0.3	4												KK KAOLINIT	2 SMALL
9 76	30- 48		1.0	6				MI 4	KK 3	MT 1							3 MODERATE
9 77	48- 63		1.8	8												MT MONTMORT	4 ABUNDANT
9 78	63- 71	48.5	2.7	8	7												5 DOMINANT
9 79	71- 89	61.7	3.8	11	8												6 INDETER.
9 80	89-105	48.1	1.9	9	7												
9 81	105-120	42.3	1.4	8	7												
9 82	120-150	36.7	1.1	8	7			MT 3	MI 3	KK 3							
9 83	0- 0																

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
G=GROUND WHOLE. HORIZON INDURATED OR GRAVEL TOO SOFT TO BE SEPARATED. ALL DATA FOR WHOLE SOIL.

FIDDYMENT LOAM (TAXADJUNCT)* 1/
S69Calif-31-8

Location: Placer County, California. Approximately 3.7 miles northwest of Roseville, California. John Fiddyment ranch, 55 yards north and 185 yards east of southeast corner of sec. 30, T.10N., R.6E. Date of Sampling: October 12, 1970.
Description by: C. B. Goudey, T. M. Ryan, L. A. Brittan, G. M. Kennedy. Collectors: L. A. Brittan, C. B. Goudey, T. M. Ryan, G. M. Kennedy, W. D. Nettleton.
Classification: Abruptic Durixeralf, fine-loamy, mixed, thermic.
Vegetation: Tarweed, soft chess, wild oats, and other annual grasses (poor cover). Climate: Average annual precipitation 17 to 22 inches. Mean annual air temperature 61°F. Parent Material: Surficial sediments deposited over Laguna Formation. Topography: Undulating terrace on a southeasterly slope of 5 percent. Elevation: 107 feet. Drainage: Moderately well drained, slow to medium runoff and very slow permeability. Soil Moisture: Dry to 12 inches, slightly moist below.
Remarks: Field has been cropped to annual grain. Poor cover of annuals. Fertility appears to be very low. Roots were dominantly matted on top of B21t horizon. Top of IIC1sim horizon has 1 mm thick discontinuous coating with matted roots above.
*The argillic horizon has slightly less clay than required of the series.

HORIZON	DESCRIPTION
Ap RSL 6974	(0-23 cm.) 0 to 9 inches, light yellowish brown (10YR 6/4) loam, dark brown (7.5YR 4/4) moist; massive; hard, friable, slightly sticky, slightly plastic; many very fine and fine random roots; many very fine tubular and interstitial pores; strongly acid (pH 5.5); clear wavy boundary.
A3 RSL 6975	(23-30 cm.) 9 to 12 inches, light yellowish brown (10YR 6/4) loam, dark brown (7.5YR 4/4) moist; common fine distinct yellowish brown (10YR 5/6) mottles; massive; hard, friable, sticky, slightly plastic; common very fine random roots; common very fine tubular and interstitial pores; medium acid (pH 5.6); abrupt smooth boundary.
B21t RSL 6976	(30-48 cm.) 12 to 19 inches, brown (10YR 5/3) clay loam, dark brown (7.5YR 4/4) moist; moderate medium and coarse prismatic structure; very hard, firm, sticky, plastic; few very fine random roots; many very fine tubular pores; few thin clay films line pores and bridge mineral grains; slightly acid (pH 6.3); gradual smooth boundary.
B22t RSL 6977	(48-63 cm.) 19 to 25 inches, yellowish brown (10YR 5/4) clay, dark brown (7.5YR 4/4) moist; moderate medium and coarse prismatic structure; very hard, firm, sticky, plastic; very few very fine random roots; common very fine tubular pores; common thin clay films line pores and bridge mineral grains and many moderately thick clay films on faces of peds; slightly acid (pH 6.5); abrupt wavy boundary.
B3 RSL 6978	(63-71 cm.) 25 to 28 inches, yellowish brown (10YR 5/4) clay loam, dark yellowish brown (10YR 3/4) moist; moderate coarse prismatic structure; very hard, firm, sticky, plastic; very few very fine random roots; few very fine tubular pores; many moderately thick clay films line pores, bridge mineral grains and on faces of peds; neutral (pH 6.8); abrupt wavy boundary.
IIC1sim RSL 6979	(71-89 cm.) 28 to 35 inches, light yellowish brown (10YR 6/4) and very pale brown (10YR 7/3) with brown (10YR 5/3) coatings on horizontal fracture plates of silty material, dark yellowish brown (10YR 3/4) and yellowish brown (10YR 5/4) moist; massive; extremely hard, extremely firm, brittle, slightly sticky, nonplastic; very few very fine tubular pores; few moderately thick clay films bridge mineral grains and on

CONSECUTIVE PEDON NUMBER C 69 26 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES SAN JOAQUIN TAXADJUNCT SOIL NO. S69 CALI. 31-21 SAMPLE NO. 9 123-9 130 PLACER COUNTY AREA

SOIL FAMILY ABRUPTIC DURIXERALF FINE-LOAMY,MIXED,THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	SIZE, CLASS AND PARTICLE DIAMETER (MM) 181B,3A1 TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SANDS VCS 2-1	CS 1-.5	MS .5-.25	FS .25-.10	VFS .1-.05	SILTS CSI .05-.02	FSI .02-.002	CLAYS CO. 2-.20	FINE LT .20	CARBONATE 3A1A.02	INT. 1A2A II .2-.02	COARSE FRAGMENTS PCT. OF WHOLE SOIL I 75-20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75		
9 123	0- 23	AP	49.5	39.0	11.5	1.2	5.9	11.2	21.2	10.0	19.9	19.1	6.6	4.9		40.0	--A --	--	--
9 124	23- 33	A3	48.8	38.2	13.0	1.7	6.0	11.5	19.8	9.8	20.3	17.9				39.8	--A --	--	--
9 125	33- 46	B21T	43.3	20.4	36.3	0.4	5.6	16.5	16.9	3.9	8.4	12.0	11.9	24.4		18.4	--A --	--	--
9 126	46- 61	B22T	50.8	20.7	28.5	0.3	5.4	20.8	22.0	2.3	8.0	12.7				17.3	--A --	--	--
9 127	61- 79	B31SIM	62.5	21.6	15.9	0.3	7.3	29.3	21.8	3.8	7.4	14.2				17.6	--G --	--	--
9 128	79-100	B32SIM	71.7	19.4	8.9	1.6	15.4	31.5	17.6	5.6	8.2	11.2				20.1	--G --	--	--
9 129	100-120	C 1SIM	74.6	16.4	9.0	17.1	32.7	11.6	8.7	4.5	7.5	8.9				16.3	--G --	--	--
9 130	120-170	C 2SIM	44.4	48.7	6.9	6.5	9.1	3.7	9.4	15.7	24.4	24.3				46.2	--G --	--	--

SAMPLE NO.	DEPTH CM.	ORG. MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	19-2 6E1B	EXT. IRON FE 6C2B	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST 1/3 BAR 4A1A	1/3 BAR 4AH	DRY 3B2	CM WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTENSBLTY LEF 4D1	LE 4D1	PH H2O 1/1 8C1A	H2O 1/10 8C1A	CACL2 8C1E
		PERCENT		PERCENT		PCT.	PCT. NO.40		GRAMS PER CC			PERCENT			PERCENT				
9 123	0- 23	0.45	0.040			1.4			1.52	1.56	1.00		15.2	5.3	0.9	0.9	5.5		4.9
9 124	23- 33	0.30	0.033			1.5			1.62	1.69	1.00		15.5	6.3	1.4	1.4	5.6		4.9
9 125	33- 46	0.23				2.3			1.44	1.87	1.00		26.6	18.1	9.1	9.1	6.0		5.5
9 126	46- 61	0.10				2.0			1.49	1.76	1.00		24.7	15.2	5.7	5.7	6.3		5.8
9 127	61- 79					1.7			1.64	1.70	1.00		17.2	13.3	1.2	1.2	6.9		6.4
9 128	79-100					1.6			1.67	1.70	1.00		16.9	12.7	0.6	0.6	7.5		7.0
9 129	100-120					1.1			1.62	1.64	1.00		16.7	12.7	0.4	0.4	7.6		6.6
9 130	120-170					1.5			1.37	1.43	1.00		28.4	12.5	1.4	1.4	6.9		6.3

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6Q2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT.	C E C NA-OAC 5A2A	NH4-OAC 5A6A	WATER EXTRACT FROM SATURATED PASTE 8A1 EC 8A1A	CA 6N1B	MG 6O1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	B 6T1A	PH SAT. PASTE 8C1B
		MEQ/100G					PCT.	MEQ/100G		MMHO	MEQ/1000 ML								PPM	
9 123	0- 23	4.4	2.7	0.1	0.2	7.4	80		9.2											
9 124	23- 33	4.9	3.3	0.1	0.1	8.4	79		10.7											
9 125	33- 46	15.5	13.7	0.5	0.4	30.1	100+		29.8											
9 126	46- 61	15.0	11.8	0.5	0.3	27.6	98		28.2											
9 127	61- 79	15.0	11.8	0.6	0.2	27.6	100+		27.2											
9 128	79-100	16.7	12.2	0.6	0.2	29.7	100+		26.7											
9 129	100-120	16.5	11.3	0.8	0.1	28.7	100+		26.6											
9 130	120-170	16.6	11.0	0.7	0.2	28.5	100+		27.5											

SAMPLE NO.	DEPTH CM.	WATER AT SAT. 8A PCT.	EXCHANGBLE NA 5B4B MEQ.	NA 5D2 PCT.	SAR 5E	GYPSUM AMT. 6F1B PERCENT	REQ. 6P3A	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 PCT.	L.T. .0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
9 123	0- 23		0.1	1													1 TRACE
9 124	23- 33		0.1	1				MI 3	KK 3							MI MICA	2 SMALL
9 125	33- 46		0.5	2				MI 3	KK 3	MT 1						KK KAOLINIT	3 MODERATE
9 126	46- 61		0.5	2												MT MONTMORT	4 ABUNDANT
9 127	61- 79		0.6	2													5 DOMINANT
9 128	79-100		0.6	2													6 INDETER.
9 129	100-120		0.8	3													
9 130	120-170		0.7	3													

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
G=GROUND WHOLE. HORIZON INDURATED OR GRAVEL TOO SOFT TO BE SEPARATED. ALL DATA FOR WHOLE SOIL.

SAN JOAQUIN LOAM (TAXADJUNCT)* 1/

S69Calif-31-21

Location: Placer County, California. West side of South Sutter Canal, Thorton ranch. 1,000 feet north of the center of sec. 23, T. 12 N., R. 5 E., 1 mile southwest of Sheridan. Date of Sampling: May 9, 1969.

Description by: G. M. Kennedy, T. M. Ryan. Collectors: K. W. Flach, W. D. Nettleton, G. M. Kennedy, L. A. Brittan, T. M. Ryan.

Classification: Abruptic Durixeralf, fine-loamy, mixed, thermic.

Vegetation: Fallowed grain field. Climate: Average annual precipitation 25 inches. Mean annual air temperature 61°F. Parent Material: Mixed sediments over the Laguna Formation. Topography: Undulating terrace on a southwesterly slope of 3 percent. Elevation: 110 feet. Drainage: Well drained; slow runoff; very slow permeability. Soil Moisture: Profile moist below 9 inches.

Remarks: Ap and A3 horizons have about 5 percent quartzitic gravel. B32sim and C2sim horizons have thin continuous white silica sheets.

* Average clay content of the argillic horizon is somewhat lower than characteristic of series.

HORIZON	DESCRIPTION
	(0-23 cm.)
Ap RSL 69123	0 to 9 inches, reddish brown (5YR 4/4) loam, dark reddish brown (5YR 3/4) moist; massive; hard, friable, slightly sticky, slightly plastic; many very fine roots; common very fine, few fine tubular pores; strongly acid (pH 5.3); abrupt smooth boundary.
	(23-33 cm.)
A3 RSL 69124	9 to 13 inches, yellowish red (5YR 4/6) loam, dark reddish brown (5YR 3/4) moist; moderate coarse angular blocky structure; hard, friable, sticky, plastic; common very fine roots; common very fine and fine tubular pores; few thin clay films line pores; strongly acid (pH 5.5); abrupt smooth boundary.
	(33-46 cm.)
B21t RSL 69125	13 to 18 inches, yellowish red (5YR 4/6) clay, reddish brown (5YR 4/4) moist; strong medium and coarse prismatic structure; very hard, firm, sticky, very plastic; few very fine roots; common very fine tubular pores; continuous moderately thick clay films on faces of peds and lining pores; medium acid (pH 6.0); gradual smooth boundary.
	(46-61 cm.)
B22t RSL 69126	18 to 24 inches, variegated yellowish red (5YR 4/8) and dark reddish brown (2.5YR 3/4) sandy clay loam, dark red (2.5YR 3/6) moist; moderate medium and coarse prismatic structure; very hard, firm, sticky, very plastic; very few very fine roots; few very fine tubular pores; many moderately thick clay films on faces of peds, lining pores, and bridging mineral grains; slightly acid (pH 6.5); abrupt wavy boundary.
	(61-79 cm.)
B31sim RSL 69127	24 to 31 inches, yellowish red (5YR 5/6) sandy loam, reddish brown (5YR 4/4) moist; common distinct light brown and brown (7.5YR 6/4, 5/4) mottles; many medium distinct Mn stains; weak coarse prismatic structure; hard, firm, slightly sticky, slightly plastic; no roots; few very fine and fine interstitial pores; continuous thin clay films lining pores and bridging mineral grains, colloidal stains on mineral grains; neutral (pH 6.8); gradual wavy boundary.
	(79-100 cm.)
B32sim RSL 69128	31 to 40 inches, yellowish red (5YR 5/6) sandy loam, yellowish red (5YR 4/6) moist; common medium distinct dark reddish brown (5YR 3/3, 3/2) Mn stains, many medium distinct black Mn stains moist; massive; very hard, very firm, nonsticky, slightly plastic; no roots; few very fine interstitial pores; many thin clay films lining pores and bridging mineral grains; neutral (pH 7.0); abrupt wavy boundary.
	(100-120 cm.)
C1sim RSL 69129	40 to 48 inches, reddish yellow (7.5YR 6/6) sandy loam, yellowish red (5YR 4/6) moist; many medium distinct dark reddish brown (5YR 3/3, 3/2) Mn stains, black moist; indurated; extremely firm, nonsticky, nonplastic; no roots; no pores; common moderately thick clay films bridging mineral grains; mildly alkaline (pH 7.5); abrupt wavy boundary.
	(120-170 cm.)
C2sim RSL 69130	48 to 67 inches, variegated light gray (2.5Y 7/2) and pale brown (10YR 6/3) sandy loam, variegated yellowish brown (10YR 5/4) and dark yellowish brown (10YR 4/4) moist; indurated; extremely hard, extremely firm, nonsticky, nonplastic; no roots; no pores; few thin clay films bridging mineral grains; mildly alkaline (pH 7.6).

1/ The San Joaquin series is in a fine, kaolinitic family of Abruptic Durixeralfs. This pedon has less than 35 percent clay in its control section.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Abruptic Durixeralf, fine, illitic, thermic
SOIL Gloria loam SOIL Nos. S65Calif-27-16 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65456 - 65467

Depth (in.)	Horizon	1B1b Total: Sand (2–0.05)	Total: Silt (0.05–0.002)	Total: Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	* Coarse fragments 1A2a > 2 Pct.	2 – 19 (Pct. of < 76 mm)	19 – 76 (Pct. of < 76 mm)	
		Size class and particle diameter (mm) 3A1																	
		Pct. of < 2 mm																	
0-8	Ap1	54.4	31.6	14.0	7.8	14.6	8.9	14.4	8.7	17.6	14.0	33.6	45.7	0	14	10	10	0	**
8-15	A12	52.9	29.4	17.7	8.1	14.6	8.3	13.1	8.8	15.3	14.1	31.0	44.1	0	18	12	12	0	**
15-16	A2	52.9	31.8	15.3	10.3	13.8	7.6	12.5	8.7	16.9	14.9	32.2	44.2	0	15	12	12	0	**
16-23	B2t	34.6	18.5	46.9	7.6	9.4	5.2	7.6	4.8	9.9	8.6	18.8	29.8	0	47	15	15	0	**
23-32	B31sim-1 1/	78.1	15.6	6.3	15.1	24.7	13.1	16.7	8.5	8.7	6.9	25.8	69.6	0	6				***
32-41	B31sim-2 1/	72.4	20.0	7.6	21.9	20.1	9.6	13.5	7.3	10.0	10.0	24.4	65.1	0	8				***
41-57	B32sim-1 1/	68.6	21.5	9.9	17.9	19.5	8.5	14.6	8.1	9.7	11.8	25.8	60.5	0	10				***
57-68	B32sim-2 1/	65.9	21.3	12.8	16.7	18.7	9.1	13.7	7.7	11.4	9.9	26.5	58.2	0	13				***
68-69+	B32sim-3 1/																		***
0-14	B31sim 2/																		***
14-27	B32sim 2/																		***
27-80	C1 2/	72.1	13.4	14.5	19.7	15.3	9.8	19.2	8.1	6.4	7.0	24.8	64.0	0	15	33	33	0	**
	6A1a	6B1a		6C2a	6B1b		Bulk density			3B2	Water content			4D1	4D1		pH		

GLORIA SANDY LOAM

S65Calif-27-16

Location: Monterey County, California, near center SE¼ NW¼ SE¼ Sec. 25, T16S, R5E, about 4 miles east of Gonzales on Gloria Road. Date of Sampling: May 22, 1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C.J.Weisel, R. D. Waggoner, F. J. Pecarich; site selection by T. D. Cook.
Classification: Abruptic Durixeralf, fine, illitic, thermic. Noncalcic Brown (maximal).
Vegetation: Grain land with volunteer ripgut brome and barley, pastured. Climate: Precipitation is 12 inches; mean annual temperature 58°F., mean January 50°F., and mean July 65°F. Parent Material: Granitic alluvium. Topography: Smooth, slightly concave, gently sloping south slope of 5 percent. Elevation: 550 feet. Drainage: Well drained, very slowly permeable, medium runoff. Soil Moisture: Dry, wilting point or below.
Remarks: No paired profile secured. Soil units mapped as loam, but contain areas of sandy loam with varying amounts of very fine gravel.

HORIZON	DESCRIPTION
Ap RSL 65456	0 to 8 inches, brown (10YR 5/3) sandy loam, dark yellowish brown (10YR 3/4) moist; weak medium subangular blocky structure; hard, very friable, slightly sticky and slightly plastic; plentiful very fine roots; many very fine interstitial pores; medium acid (pH 5.8); 15 percent of horizon consists of gopher holes filled with strong granular structure; gradual smooth boundary.
A1 RSL 65457	8 to 15 inches, brown (7.5YR 5/4) sandy loam, reddish brown (5YR 4/4) moist; massive; hard, very friable, sticky and plastic; plentiful very fine roots; many very fine interstitial and common very fine and fine tubular pores; slightly acid (pH 6.5); small thin blotches of A2 occur in lower part; clear smooth boundary.
A2 RSL 65458	15 to 16 inches, very pale brown (10YR 7/3) sandy loam, brown (7.5YR 5/4) moist; weak medium platy structure; hard, very friable, sticky and plastic; plentiful very fine roots; few very fine interstitial, common very fine and fine tubular pores; neutral (pH 7.0); very abrupt smooth boundary.
B2t RSL 65459	16 to 23 inches, reddish brown (5YR 4/4) moist and dry, light clay, with dark brown (7.5YR 3/2) moist and dry coatings on top and upper sides of peds, some veins and blotches of olive hues (2.5Y) in middle and lower part plus various shades of brown to reddish brown; strong coarse prismatic structure; extremely hard, very firm, very sticky and plastic; plentiful very fine roots on ped exteriors; common very fine tubular pores; continuous thin clay films and few moderately thick films on ped faces; mildly alkaline (pH 7.7); abrupt smooth boundary.
C1sim RSL 65460	23 to 32 inches, reddish brown (5YR 5/4) coarse sandy loam crushed; reddish brown (2.5YR 4/4) moist; massive; indurated, does not soften or slake upon wetting and can be chipped with hand tools with extreme difficulty; no roots; common very fine tubular pores; few moderately thick clay films in pores and continuous thin films as bridges; moderately alkaline (pH 7.9); matrix noncalcareous, very slightly calcareous with lime as filaments in some veins, in parts of the pit 1/8" lime seams occur on top of indurated pan but not in area sampled; horizon has light yellowish brown (2.5Y 6/2d, 4/2m) usually vertical veins, 1/4 to 3/4 inch thick, makes up about 5 to 10 percent of the soil mass, and appears to be silica and clay; this horizon becomes more yellow with depth; boundary indeterminate.
C2sim RSL - 65461	32 to 41 inches, same as the C1sim horizon.
C3sim RSL - 65462	41 to 57 inches, strong brown (7.5YR 5/6) coarse sandy loam crushed, crushed colors are light yellowish brown (10YR 6/4d); massive; strongly cemented, does not slake or soften upon wetting; common very fine tubular pores; this horizon examined by auger with great difficulty, contains few black Mn specks 1 to 2 mm; boundary indeterminate.
C4sim RSL - 65463	57 to 68 inches, same as the C3sim horizon.
C5sim RSL - 65464	68 to 69 inches, same as the C3sim horizon.
- RSL 65465	0 to 14 inches*, yellowish red (5YR 5/8) sandy clay loam; massive; indurated, does not soften or slake upon wetting and can be chipped with hand tools with extreme difficulty; no roots; many very fine and fine and common medium tubular pores; common moderately thick clay skins in tubular pores; gradual boundary.
- RSL 65466	14 to 27 inches, yellowish red (5YR 5/8) fine gravelly sandy loam with few fine distinct red (2.5YR 4/6) streaks; massive; indurated; no roots; common very fine tubular pores; few thin to common thick clay skins in pores; common thick, mostly horizontal seams of silica; gradual boundary.
- RSL 65467	27 to 80 inches, light yellowish brown (10YR 6/4) light gravelly sandy loam; massive; friable; few micro and few fine interstitial pores; few thin clay skins in pores and as bridges between sand grains; few horizontal and vertical seams of silica.

*The description and sampling of Gloria was continued in and below the duripan in a road cut 30 yards east of the profile pit where the lower part of the profile could be observed. Depths are measured from the top of the duripan at the road cut.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Abruptic Durixeralf, fine, mixed, thermic

SOIL Redding gravelly loam (taxadjunct) SOIL Nos. S4557-5 LOCATION Tehama County, California

SOIL SURVEY LABORATORY _____ LAB. Nos. _____

Depth (In.)	Horizon	1B1b Total: Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Size class and particle diameter (mm) 3A1 Sand: Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt: 0.05–0.02	Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)			Coarse fragments: > 2 Pct	2 – 19	19 – 76
		← Pct. of < 2 mm →															← Pct. of < 76mm →	
0-8	A11	50.2	40.2	9.6	9.1	8.6	4.3	10.0	18.2	20.4	19.8		32.0					
8-14½	A12	50.0	39.1	10.9	8.2	7.9	4.5	10.3	19.1	19.2	19.9		30.9					
14½-19½	A13	49.1	38.7	12.2	10.1	7.5	4.3	9.7	17.5	19.2	19.5		31.6					
19½-22	B2t	25.0	19.5	55.5	6.1	4.9	2.2	4.4	7.4	8.4	11.1		17.6					
22-24½	IIC1m	60.1	13.9	26.0	28.5	15.5	4.6	6.5	5.0	3.9	10.0		60.1					
24½-35	IIC2m	54.4	19.6	26.0	25.1	13.9	4.4	5.3	5.7	4.7	14.9		48.7					

Depth (In.)	6A2a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	Pct.			pH		8C1a (1:1)
0-8	0.36	0.040																5.7
8-14½	0.18	0.029																5.6
14½-19½	0.14	0.026																5.5
19½-22	0.35	0.057																5.3
22-24½	0.18	0.044																5.5
24½-35	0.05	0.021																6.8

Depth (In.)	Extractable bases: Ca 6N2b	Mg 6O2b	Na 2/	K 5B1a 1/	Sum of bases	6H2a Ext. Acidity	5A3a CEC Sum									Base saturation: 5C3 Sum Pct.	Pct.
	← meq/100 g →																
0-8	3.2	1.3	tr	0.38	4.9	2.6	7.5									65	
8-14½	3.2	1.9	<0.01	0.25	5.3	2.6	8.0									67	
14½-19½	3.2	2.6	0.03	0.17	6.0	2.6	8.6									70	
19½-22	12.6	13.4	0.29	0.29	26.6	6.2	32.8									81	
22-24½	13.5	13.5	0.37	0.28	27.6	5.4	33.0									84	
24½-35	12.6	12.2	0.20	0.21	25.2	2.9	28.1									90	

Depth (In.)				

1/ Precipitated as potassium sodium cobaltinitrite and titrated as potassium permanganate

2/ Gravimetrically as sodium uranyl magnesium acetate

REDDING GRAVELLY LOAM (taxadjunct) 1/
84557-5

Location: Tehama County, California, 0.42 mile north of bench mark 311 located about 5 miles south of Red Bluff. Date of Sampling: April 2, 1945.
Description by: R. C. Roberts and R. A. Gardner. Collectors: R. C. Roberts, F. O. Youngs, R. A. Gardner, L. T. Alexander, H. Jenny and R. E. Storie.
Classification: Abruptic Durixeralf, fine, mixed, thermic.
Vegetation: Sparse vegetation consists of short annual grasses and associated herbaceous plants dominated by alfilaria. Climate: Average annual precipitation is 19 inches, average annual temperature is 62°F., January 45°F., and July 82°F. Parent Material: Mixed materials. Topography: Nearly level to undulating old terraces of mixed valley-fill materials on a general slope of about 1 percent. Many of the fields have numerous mounds or "hog-wallow" microrelief. Pit was dug about halfway between mound and intermound areas. Elevation: 315 feet. Drainage: Well drained, very slow permeability, slow runoff. Soil Moisture: Moist to 24 inches, dry below.
Remarks: The soil at this site is representative of many of the soils of California that occupy very old terraces with deeply entrenched drainageways. The principal characteristics of these soils are (1) Weakly developed surface-soil structure and very low organic matter content; (2) a yellowish red or reddish brown surface soil that rests abruptly on (3) a distinct claypan, which is generally redder than the layers above or below; (4) an indurated layer below the claypan; and (5) in general a decrease in acidity with depth. This profile appears to represent an advanced stage of soil development, but it appears to be normal for its relief, effective soil climate, vegetation, parent material, and age or length of time without overflow or significant increment of material.

The pit where the samples were collected, like many of the other pits examined in this study, exposed an unconformity, possibly an old rodent burrow in one corner. At the time the samples were collected, the soil material of this unconformity was saturated, whereas the adjacent soil was nearly dry or only slightly moist. The rodent burrow or unconformity was very irregular and appeared as small pits in the hardpan. Some penetrated deeper than others, and one penetrated the hardpan into the gravelly layer below.

HORIZON	DESCRIPTION
A11	0 to 8 inches, yellowish red (5YR 5/6) gravelly light loam, dark reddish brown (2.5YR 3/4) moist; weak granular structure; slightly hard, very friable, slightly plastic; plentiful fine roots; medium acid (pH 5.7); gradual boundary.
A12	8 to 14.5 inches, yellowish red (5YR 5/6) gravelly light loam, red (2.5YR 4/6) moist; weak blocky structure; slightly hard, friable, slightly plastic; plentiful fine roots but somewhat fewer than in layer above; medium acid (pH 5.6); gradual boundary.
A13	14.5 to 19.5 inches, yellowish red (5YR 5/6) gravelly light loam, color dry and moist, structure and consistence similar to layer above; plentiful fine roots; strongly acid (pH 5.5); abrupt wavy boundary.
B24	19.5 to 29 inches, yellowish red (5YR 4/6) clay containing a slight amount of

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Abruptic Durixeralf, fine, mixed, thermic

SOIL Redding loam (taxadjunct) SOIL Nos. S66Calif-37-2 LOCATION San Diego, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66236 - 66242*

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Silt: Int. II (0.2-0.02)	Silt: (2-0.1)			1/ Coarse fragments 1A2a > 2 Pct.	2-19 (Pct. of < 76mm)	19-76 (Pct. of < 76mm)	
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm																	
0- 4	A1	55.3	33.5	11.2	2.9	15.7	18.0	13.7	5.0	14.7	18.8	25.3	50.3	0	11	22	8	14	2/
4-15	A2	52.8	35.9	11.3	3.1	12.9	17.1	13.8	5.9	15.2	20.7	27.0	46.9	0	11	14	9	5	2/
15-18	B1	35.3	27.3	37.4	1.6	8.3	11.2	10.0	4.2	11.1	16.2	19.4	31.1	0	37	8	7	1	2/
18-30	B2t	24.2	15.6	60.2	1.0	6.0	7.6	6.7	2.9	6.9	8.7	12.6	21.3	0	60	6	1	2	3/
30-36	B31sim	65.3	7.1	27.6	17.4	21.8	11.9	10.9	3.3	3.8	3.3	11.7	62.0	0	28	14	3	4	3/
36-45+	B32sim	85.1	6.0	8.9	20.8	28.3	16.5	15.1	4.4	3.6	2.4	14.7	80.7	0	9	20	2	5	3/
27-41	Bm32sim	87.1	7.1	5.8	22.1	28.6	16.1	15.0	5.3	4.5	2.6	16.6	81.8	0	6	19	4	3	3/

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1		
0- 4	2.51						1.27	1.30		0.88		9.6	5.5	0.009	0.008	5.6		
4-15	0.63						1.65	1.68		0.91		9.8	3.7	0.006	0.006	5.5		
15-18	0.41						1.59	1.78		0.95		20.3	12.1	0.039	0.037	4.6		
18-30	0.38						1.47	1.78		0.96		26.8	20.6	0.067	0.065	4.3		
30-36	0.35						1.69	1.77		0.91		17.0	15.8	0.015	0.013	4.3		
36-45+	0.17						1.79	1.81		0.86		14.1	11.2	0.005	0.004	4.4		
27-41							1.92	1.94		0.85		11.2	8.0	0.004	0.003	4.4		

Depth (In.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1a NH_4OAc	5A3a Sum	6G1d KCl Ext. Al^{+++}	Water extract from saturated paste 8A1: 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K	6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1b SO_4	8A1a Electrical conductivity
	meq/100 g									meq/liter								mmho/cm
0- 4	5.2	2.1	0.4	0.6	8.3	5.9	11.1	14.2	0.2	2.5	2.1	1.9	1.0	-	2.8	2.4		0.70
4-15	4.3	2.4	0.4	0.3	7.4	3.5	6.8	10.9	0.3	0.5	0.5	4.2	tr.	-	1.0	0.6		0.25
15-18	3.3	5.5	1.8	0.2	10.8	9.8	16.6	20.6	2.9	0.2	0.3	3.4	tr.	-	0.3	1.5		0.41
18-30	5.5	10.4	4.1	0.2	20.2	15.6	28.6	35.8	5.1	0.4	0.3	7.2	tr.	-	0.2	4.5		0.85
30-36	5.1	9.5	4.9	0.2	19.7	14.3	29.4	34.0	5.2	0.6	0.8	13.8	tr.	-	0.3	9.7		2.12
36-45+	4.7	8.3	4.9	0.2	18.1	11.7	26.6	29.8	4.3	0.6	0.9	15.9	tr.	-	0.2	12.6		1.92
27-41	3.2	5.1	4.2	0.2	12.7	9.8	19.8	22.5	3.7	1.4	2.2	25.5	tr.	-	0.1	17.8	9.5	3.00

Depth (In.)	8A Water at Saturation	5D2 Exchangeable Na	Clay mineralogy (<0.002 mm.) 7A: Mont.	Verm.	Mica	Kaolin.	Base saturation: 5C3 Sum	5C1 NH_4OAc

* RSL 66242 was collected at station 39.8, or 6.8 feet from S66Calif-37-2.

1/ From characterization sample (5 kg)(except as noted).

REDDING LOAM (taxadjunct) 1/
S66Calif-37-2

Location: San Diego County, California; Miramar Transect No. 1, station 33.0; 950 feet west and 750 feet north of the southeast corner of section 24, T. 15 S., R. 3 W. Date of sampling: June 10, 1966.
Description by: George Borst, David Estrada, Gerald Kester, Reuben Nelson, and Dennis Nettleton.
Samples collected by: Klaus Flach, Benny Brasher, and Dennis Nettleton.
Classification: Abruptic Durixeralf, fine, mixed, thermic family.
Vegetation: Sparse chamise and flattop buckwheat. Climate: Weather record for San Diego, near the sampling site, is mean winter air temperature 55° F., mean summer air temperature 68° F., mean annual precipitation 10 inches. Parent Material: Stream deposit over Pleistocene marine terrace.
Topography: Flat marine terrace of less than 1 percent slope. Elevation: About 425 to 450 feet.
Drainage: Moderately well drained; runoff is medium; permeability is very slow. Soil Moisture: Dry.
Remarks: Less than 15 percent rounded gravel throughout.

HORIZON	DESCRIPTION
A1 RSL No. 66236	0 to 4 inches, yellowish brown (10YR 5/6) loam, dark yellowish brown (10YR 4/6) when moist; weak fine granular structure; slightly hard, very friable, nonsticky, nonplastic; abundant very fine roots; many very fine tubular pores; medium acid (pH 5.8); clear smooth boundary.
A2 RSL No. 66237	4 to 15 inches, light brown (7.5YR 6/4) loam, brown (7.5YR 5/4) when moist; massive; hard, friable, slightly sticky, nonplastic; plentiful very fine to medium roots; many very fine tubular pores; strongly acid (pH 5.5); clear smooth boundary.
B1 RSL No. 66238	15 to 18 inches, yellowish red (5YR 5/6) clay, yellowish red (5YR 4/6) when moist; moderate medium angular blocky structure; very hard, firm, sticky, plastic; plentiful very fine to fine roots; common very fine tubular pores; common moderately thick clay skins * on ped faces; very strongly acid (pH 4.5); very abrupt smooth boundary.
B2t RSL No. 66239	18 to 30 inches, red (2.5YR 4/6) clay, dark red (2.5YR 3/6) when moist; strong medium angular blocky structure; very hard, firm, sticky, plastic; plentiful very fine to fine roots; common very fine tubular pores; many thick clay skins * on ped faces; very strongly acid (pH 4.5); abrupt smooth boundary.
B31sim RSL No. 66240	30 to 36 inches, yellowish red (5YR 5/6) duripan, yellowish red (5YR 4/6) when moist; strong fine angular blocky structure; indurated; few fine roots on ped exteriors; few fine tubular pores; many moderately thick clay skins on ped faces; extremely acid (pH 4.2); abrupt smooth boundary.
B32sim RSL No. 66241	36 to 45 inches +, light brownish gray (2.5Y 6/2) duripan, grayish brown (2.5Y 5/2) when moist; massive; indurated; extremely acid (pH 4.0).

* Probably mostly pressure faces --- some are slickensides.

1/ The Redding series is in a fine, kaolinitic, thermic family of Abruptic Durixeralfs. This pedon is in the same subgroup, but has about equal amounts of montmorillonite and kaolinite.

CONSECUTIVE PEDON NUMBER C 69 21 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES FIDDYMENT TAXADJUNCT SOIL NO. S69 CALI. 31-13 SAMPLE NO. 9 98-9 99 PLACER COUNTY AREA

SOIL FAMILY ABRUPTIC DURIXERALF FINE,MIXED,THERMIC,SHALLOW RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A	1A2A INT. II .2- .02	-COARSE FRAGMENTS- PCT. OF WHOLE SOIL I 75- 20	II 20-5 PCT. OF	III 5-2 LT 75
			PERCENT OF LESS THAN 2MM																
9 98	8- 25	A3	48.2	39.4	12.4	2.2	5.2	8.6	19.4	12.8	18.3	21.1				42.3	--A	--	--	--
9 99	26- 43	B2 T	32.3	28.6	39.1	0.9	3.7	7.1	12.3	8.3	9.0	19.6				24.2	--A	--	--	--

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR-BON 6A1A	NITRO GEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	19-2 6E1B	EXT. IRON FE 6C2B	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST 1/3 4A1F	DRY 1/3 BAR 4A1F	CM 4AM	3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTENSBLTY LEF 4D1	LE 4D1	PH H2O 1/1 8C1A	H2O 1/10 8C1A	CACL2 8C1E
		-PERCENT--		-PERCENT-		PCT.	PCT.	NO.40	GRAMS PER CC				----PERCENT---			-PERCENT--				
9 98	8- 25	0.27	0.032						1.68			1.00			5.1			5.6		4.9
9 99	26- 43	0.25							1.66			1.00			17.4			6.2		5.8

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT.	C E C NA- OAC 5A2A	NH4- OAC 5A6A	EC 8A1A	WATER EXTRACT FROM SATURATED PASTE CA 6N1B	MG 6O1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	B 6T1A	PH SAT. PASTE 8C1B
		MEQ/100G					PCT.	-MEQ/100G		MMHO	MEQ/1000 ML								PPM	
9 98	8- 25	5.2	2.7	0.2	0.1	8.2	81		10.1											
9 99	26- 43	18.2	9.6	1.0	0.3	29.1	96		30.3											

SAMPLE NO.	DEPTH CM.	WATER AT SAT. 8A PCT.	EXCHANGBLE NA 5B4B MEQ.	NA 5D2 PCT.	SAR	GYPSUM AMT. REQ. 6F1B PERCENT	CLAY MINERALOGY LESS THAN 0.002 MM X RAY I 7A2E	II 7A2E	III 7A2E	IV 7A2E	DTA PCT. 7A3	L.T. .0002 MM X RAY I 7A2E	II 7A2E	III 7A2E	MINERAL CODE	RELATIVE AMOUNT
9 98	8- 25		0.2	2			MI 4	KK 3	MT 1						MI MICA	1 TRACE
9 99	26- 43		1.0	3			MI 4	MT 3	KK 3						KK KAOLINIT	2 SMALL
															MT MONTMORT	3 MODERATE
																4 ABUNDANT
																5 DOMINANT
																6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.

FIDDYMENT LOAM (taxadjunct) 1/
S69Calif-31-13

Location: Placer County, California. Approximately 4.7 miles south of Lincoln on Highway 65 along embankment of PG&E substation; 15 feet west from Trigo pedon (S69Calif-31-11); 750 feet west and 500 feet south of NE corner of sec. 9, T. 11 N., R. 6 E. Date of Sampling: May 7, 1970.

Description by: L. A. Brittan, T. M. Ryan. Collectors: K. W. Flach, W. D. Nettleton, G. M. Kennedy, L. A. Brittan, T. M. Ryan.

Classification: Abruptic Durixeralf, fine, mixed, thermic, shallow.

Vegetation: Filaree, rose clover, soft chess, lupine, and purple vetch. Climate: Average annual rainfall 17 to 32 inches. Mean annual air temperature 61°F. Parent Material: Surficial sediments deposited over Laguna Formation. Topography: Undulating terrace slope of 2 to 9 percent on a south aspect. Elevation: 125 feet. Drainage: Moderately well drained, slow to medium runoff and very slow permeability. Soil Moisture: Slightly moist.

Remarks: Roots in B2t horizon essentially follow vertical cracks between peds. Tubular pores in IIClsim horizon appear to be root channels. Top of IIClsim has thin (2mm) coating of clay; also has clay bedding in planes of siltstone. Below siltstone has stratified layers of sand.

HORIZON	DESCRIPTION
Ap	(0-8 cm.) 0 to 3 inches, brown (7.5YR 5/4) loam, dark brown (7.5YR 4/4) moist; massive; slightly hard, friable, slightly sticky, slightly plastic; common very fine vertical roots; many very fine tubular and interstitial pores; medium acid (pH 6.0); clear smooth boundary.
A3 RSL 6998	(8-25 cm.) 3 to 10 inches, brown (7.5YR 5/4) loam, dark brown (7.5YR 4/4) moist; massive; slightly hard, friable, slightly sticky, slightly plastic; few very fine vertical roots; many very fine tubular and interstitial pores; medium acid (pH 5.8); abrupt wavy boundary.
B2t RSL 6999	(25-43 cm.) 10 to 17 inches, brown (10YR 5/3) clay, dark reddish brown (5YR 3/4) moist; weak medium and coarse prismatic structure; very hard, very firm, very sticky, very plastic; few very fine vertical roots; common very fine tubular pores; many moderately thick clay films line pores; neutral (pH 7.0); abrupt wavy boundary.
IIClsim	(43-66 cm.) 17 to 26 inches, light gray (2.5Y 7/2) silty clay loam matrix with strong brown (7.5YR 5/6) stains, light olive brown (2.5Y 5/4) moist; massive; extremely hard, extremely firm, brittle, sticky, plastic; common very fine tubular pores; many moderately thick clay films in bedding planes; mildly alkaline (pH 7.8).

1/ The Fiddyment series is in a fine, montmorillonitic, thermic family of Abruptic Durixeralfs. This pedon is in a mixed family of the same subgroup.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Abruptic Durixeralf, fine, montmorillonitic, thermic

SOIL Monserate sandy loam (taxadjunct) SOIL Nos. S57Calif-33-12 LOCATION Riverside County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 57534 - 57541

Depth (in)	Horizon	1B1b Total: Sand (2–0.05)	Total: Silt (0.05–0.002)	Total: Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Non carbonate clay	1/ Coarse fragments 1A2a: > 2	2 – 19	19 – 76
		Pct of < 2 mm														Pct.	Pct. of < 76 mm	
0-6	Ap	51.8	39.0	9.2	8.8	11.0	6.3	12.3	13.4	25.7	13.3	46.7	38.4			18		
6-9	A1	52.5	37.3	10.2	10.7	12.2	5.9	11.6	12.1	24.2	13.1	43.3	40.4			19		
9-12	A3	49.9	31.4	18.7	11.9	11.5	5.8	10.8	9.9	18.6	12.8	35.1	40.0			28		
12-18	B21	37.0	24.1	38.9	10.6	9.1	4.0	6.8	6.5	14.1	10.0	24.6	30.5			20		
18-25	B22	36.1	25.5	38.4	9.3	8.8	4.0	7.0	7.0	14.5	11.0	25.6	29.1			17		
25-37	Cm1	62.9	30.6	6.5	11.9	18.0	8.6	13.7	10.7	17.0	13.6	35.7	52.2			19		
37-58	Cm2	56.1	32.8	11.1	15.3	13.4	5.6	11.4	10.4	17.6	15.2	35.4	45.7			21		
58-64+	Cm3	55.3	29.5	15.2	14.9	12.9	5.8	12.6	9.1	16.0	13.5	32.9	46.2			21		

Depth (in)	6A1a Organic carbon Pct	6B1a Nitrogen Pct	C/N	Ext. Iron as Fe Pct.	6E1b Carbonate as CaCO3 Pct		Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	4A3a Field moist g/cc	3B2 Cm	Water content: 4B1a 1/10 bar Pct.	4B1a 1/3 bar Pct.	4B2 15 bar Pct	4D1 Extensibility LEf Pct	4D1 Extensibility LE Pct.	pH: 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-6	0.25	0.036	7		-						19.7	11.9	4.1			6.1	6.3	
6-9	0.23	0.033			-				1.54		19.1	11.8	4.1			6.1	6.2	
9-12	0.15	0.025			-								6.1			6.8	6.7	
12-18	0.27	0.037	7		-				1.85				12.1			7.1	6.7	
18-25	0.24				-								12.7			7.1	7.0	
25-37	0.01				-								7.6			7.7	7.8	
37-58	0.02				tr.								7.4			7.7	7.5	
58-64+	0.02				-								7.7			7.4	7.1	

Depth (in)	Extractable bases: 6N2b Ca 5B1a	6O2c Mg 5B1a	Na	K	Sum of bases	6H2a Ext. Acidity	Cation Exch Capacity: 5A2a NaOAc	5A6a NH_4OAc	Water extract from saturated paste 8A1: 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1b SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-6	3.4	2.3				2.8	7.9		3.2	1.5	1.1	0.7		-	0.9	0.6		0.7
6-9	3.7	2.2				2.2	7.9		2.2	1.4	1.3	0.3		-	0.6	1.0		0.6
9-12	5.1	3.2				2.4	10.8		3.5	2.8	3.3	0.2		-	1.3	2.7		1.1
12-18	10.3	8.7				3.4	21.9		4.5	4.1	5.9	0.1		-	1.0	7.4		1.6
18-25	12.2	10.4				2.6	24.5		3.3	2.9	5.7	0.1		-	0.9	8.9		1.4
25-37	9.9	7.9				0.6	19.7		3.6	3.1	9.3	0.1		-	2.3	10.5		1.8
37-58	8.8	7.8				0.7	16.9		5.5	5.1	13.3	0.1		-	1.6	18.5		2.3
58-64+	8.9	10.5				1.4	18.0		6.5	6.0	15.2	0.1		-	1.1	22.3		3.0

Depth (in)	8A Water at Saturation Pct	Exch. bases: 5B3b Ca	5B1b 6Q2a K	5B1b 6P2a Na	5D2 Exchangeable Na Pct.	5C3 Base Sat. Pct.	Gypsum Pct.	Clay mineralogy (<0.002 mm) 3A1: Mont.	Verm.	Mica 7A2 X-ray	Kaolin.	
		meq./100 g.										
0-6	20.6		0.5	0.6		71		tr.		xxx	x	
6-9	22.5		0.5	0.3		75						
9-12	22.6		0.3	0.3		79						
12-18	42.8		0.3	1.5		86		xxx		tr.	xx	
18-25	48.5		0.4	1.2		90						
25-37	30.5		0.2	1.5		97						
37-58	28.7		0.2	2.0		96						
58-64+	31.1		0.3	1.9		94						

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

MONSERATE SANDY LOAM (taxadjunct) 1/
S57Calif-33-12

Location: 1400 feet east and 400 feet north of the southwest corner of Sec. 36, T. 2 S., R. 4 W. On the Langdon Maxwell Ranch, about 1 mile north of Sunnymead, Riverside County, California. Date of Sampling: June 6, 1957.
Description by: Fred Eggers. Collectors: George Borst, R. E. Nelson, Cliff Henry, Arnold Knecht, and Fred Eggers.
Classification: Abruptic Durixeralf, fine, montmorillonitic, thermic.
Vegetation: Fallow. In dry farmed grain fallow rotation. Precipitation: 12 inches.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Huichica loam (taxadjunct) **LOCATION** Sonoma County, California

SOIL FAMILY: Abruptic Haplic Durixeralf, fine-loamy, mixed, thermic.

SOIL NOS. S61Calif-49-7 **LAB. NOS.** 61431 - 61437

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0- 7	A11	1.8	5.7	6.1	20.7	14.7	41.7	9.3	48.4	21.2	8	1
7-14	A12	1.6	5.5	5.9	19.3	14.6	42.3	10.8	47.7	21.8	5	1
14-23	A21	2.4	5.3	6.1	18.9	15.8	40.1	11.4	46.8	21.4	6	1
23-30	B2t	6.5	8.0	7.0	15.5	7.7	26.1	29.2	26.8	15.7	31	cl-scl
30-36	C1m	2.1	15.0	12.3	20.7	5.2	15.0	29.7	22.0	8.4	11	scl
36-47	IIC2m	7.6	24.6	18.4	17.9	5.4	13.5	12.6	18.8	8.1	3	cosl
47-57+	IIIC3m	3.7	15.9	11.9	20.9	10.6	24.4	12.6	32.9	14.1	0	sl

pH 8C1a H_2O 1:1	8C1c KCl 1:1	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	Bulk Density 4A3a g/cc	Ext. Iron % Fe 6C1b	4B2 S 11 water 15 atmos. Pct.
5.4	4.2	1.10	0.096	11	1.53	0.4	5.0
5.5	4.0	0.26	0.047		1.50	0.4	5.1
6.1	4.3	0.17	0.040			0.4	5.7
6.7	5.0	0.14				0.4	13.6
7.7	5.9	0.04				0.3	17.5
7.6	5.8	0.02				0.3	15.3
7.3	5.6	0.01				0.4	21.4

5A2a CATION EXCHANGE CAPACITY ← Na →	EXTRACTABLE CATIONS 6N2b Ca	6O2c Mg	6H2a H	6P2a Na	6Q2a K	5B1a BASE SAT. % 5C3	Sum of Cations 5A3a meq/100g.	MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →							
10.0	2.5	0.8	5.7	0.1	0.3	39	9.4	
8.8	2.4	1.5	4.4	0.2	0.3	50	8.8	
9.1	2.6	2.3	3.2	0.3	0.2	63	8.6	
22.1	7.6	9.8	3.2	1.2	0.2	85	22.0	
34.4	12.3	16.4	1.8	2.1	0.2	95	32.8	
32.6	12.0	15.2	2.0	2.1	0.2	94	31.5	
42.6	16.2	20.3	3.0	2.7	0.2	93	42.4	

HUICHICA LOAM(**taxadjunct**) 1/
S61Calif-49-7

Location: Sonoma County, California. Loamite Company. SE1/4, SW1/4, section 19, T. 8 N., R. 8 W. 480 feet east and 220 feet north of the southwest corner of section 19. In field 930 feet southwest of railroad track, site spotted on photo No. CSH-2BB-183.
Date of Sampling: June 23, 1961.
Collectors: Reuben Nelson, Leo C. Klameth, V. C. Miller, G. Lambert, J. H. Kashiwagi, F. Graham.
Classification: **Abruptic Haplic Durixeralf, fine-loamy, mixed, thermic.**

Vegetation: Annual grasses, a few perennial grasses, and a few scattered oaks. Use: Dryland pasture. Climate: Average rainfall 29.4 inches, average temperature 57.6° F., average January temperature 46.5° F., average July temperature 66.7° F. Parent Material: Mixed valley alluvium. Elevation: About 112 feet. Relief: Hummocky on 1 percent slope. Drainage: Imperfectly drained, slow runoff, moderately slow to very slow permeability. Biological Activity: Worm casts to 23 inches.

HORIZON	DESCRIPTION
A11 RSL No. 61431	0 to 7 inches, light brownish gray (10YR 6/2) loam with common medium distinct mottles of yellowish brown (10YR 5/4) when dry, dark brown (10YR 4/3) when moist; massive; slightly hard when dry, friable when moist, slightly sticky and slightly plastic when wet; abundant very fine and fine roots; many very fine and fine tubular pores; strongly acid, pH 5.5; gradual, wavy boundary.
A12 RSL No. 61432	7 to 14 inches, pale brown and brown (10YR 6/3 and 10YR 5/3) loam, dark brown (10YR 4/3) when moist; massive; hard when dry, friable when moist, slightly sticky and slightly plastic when wet; abundant very fine and fine roots; many very fine and fine interstitial and tubular pores; strongly acid, pH 5.5; light gray stains on sand grains; clear, wavy boundary.
A21 RSL No. 61433	14 to 23 inches, light gray (10YR 7/2) sandy clay loam with many fine faint mottles of brown (10YR 5/3) when dry, brown (10YR 5/3) when moist; massive; hard when dry, friable when moist, slightly sticky and slightly plastic when wet; plentiful very fine and fine roots; many very fine and fine tubular and interstitial pores; strongly acid, pH 5.3; abrupt, wavy boundary.
A22 Not sampled	Ashy cap, light gray (10YR 7/1) when dry, grayish brown (10YR 5/2) when moist; 2-3 mm. thick; abrupt, wavy boundary.
B2t RSL No. 61434	23 to 30 inches, light brownish gray (10YR 6/2) clay, very dark grayish brown (10YR 3/2) when moist; strong coarse columnar structure; top and sides of columns coated with ashy gray film; extremely hard when dry, very firm when moist, very sticky and very plastic when wet; abundant very fine roots along ped faces; many very fine and fine tubular and interstitial pores; continuous thick clay films on ped faces, in bridges, and in pores; medium acid, pH 6.0; abrupt, irregular boundary.
C1m RSL No. 61435	30 to 36 inches, white (2.5Y 8/2) sandy loam, dark brown (10YR 4/3) with variegated mottles of light olive brown (2.5Y 5/4) when moist; massive; very hard when dry; firm when moist; slightly sticky and nonplastic when wet; few fine roots; common very fine interstitial and fine tubular pores; continuous, moderately thick to thick clay films in pores and on bridges; moderately alkaline, pH 8.0; firm to weakly cemented; gradual, irregular boundary.
IIC2m RSL No. 61436	36 to 47 inches, white (2.5Y 8/2) loamy sand with common fine distinct mottles of strong brown (7.5YR 5/6) when dry, dark brown (10YR 4/3) when moist; massive breaking to moderate subangular blocks; very hard when dry, firm to extremely firm when moist; nonsticky and nonplastic when wet; common fine and very fine interstitial pores; continuous, thick clay films on fracture planes; moderately alkaline, pH 8.0; varied weak to strong indurations of soil material; (manganese) stains concentrated in fissures; clear, wavy boundary.
IIIC3m RSL No. 61437	47 to 57 inches +, white (10YR 8/2) sandy clay loam with common fine distinct mottles of strong brown (7.5YR 5/6) when dry; grayish brown (10YR 5/2) with common fine faint mottles of yellowish brown (10YR 5/4) when moist; clay films are strong brown when moist; massive breaking to weak subangular blocky structure; very hard when dry, firm when moist, slightly sticky and slightly plastic when wet; continuous, thick clay films on fracture planes; mildly alkaline, pH 7.5.

1/ The Huichica series is in a fine, montmorillonitic, thermic family of Abruptic Haplic Durixeralfs. This pedon has less than 35 percent clay in the argillic horizon.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Arlington sandy loam **LOCATION** Riverside County, California

SOIL FAMILY: Haplic Durixeralf, coarse-loamy, mixed, thermic.

SOIL NOS. S57Calif-33-10 **LAB. NOS.** 57520 - 57526

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-6	Ap	5.9	17.0	12.6	23.0	12.4	20.8	8.3	37.4	9.1	3	sl
6-12	A1	6.1	14.2	12.3	24.5	13.8	22.0	7.1	40.5	9.7	2	sl
12-24	A3	3.7	12.2	11.9	26.7	15.5	22.0	8.0	45.5	8.3	2	fsl
24-35	B21	7.6	14.5	8.7	16.4	11.3	27.5	14.0	36.8	11.6	4	sl
35-38	B22	7.8	13.6	8.8	16.3	11.6	26.4	15.5	36.4	11.1	3	sl
38-55	Cm1	8.3	12.6	7.6	15.0	11.3	26.0	19.2	37.7	8.6	4	fsl
55-60+	Cm2	7.8	12.8	8.3	18.6	10.8	22.8	18.9	34.7	10.0	8	fsl

8C1b SATURATED PASTE 1:1	pH 8C1a 1:10	4A3a Bulk Density g/cc	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	CO_3 6I1a Meq/1	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	HCO_3 6J1a Meq/1	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
6.2	6.7	–	0.46	0.055	8.4	0.0	0.9	0	5.1	16.9	10.4	3.6
5.9	6.6	1.50	0.37	0.046	8.0	0.0	0.8	0	4.9	16.7	10.8	3.5
6.9	7.1	1.77	0.22	0.021		0.0	1.6	0	2.4	16.4	8.2	3.6
7.0	7.4	1.73	0.27	0.016	16.9	0.0	1.6	0	2.4			5.8
7.0	7.4	1.80	0.24			0.0	1.0	0	1.9			5.9
7.3	7.4	1.76	0.24			0.0	1.2	Tr	1.8			7.6
7.5	8.4		0.01			0.0	1.5	<1	2.8			7.4

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	5B Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a Ca	6O1a Mg	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
7.3	4.4	1.4	1.4	0.5	0.3	83	3.4	0.2	4.1	1.8	2.1	23.5
7.3	4.2	1.4	1.5	0.7	0.2	81	3.4	0.1	3.2	1.7	2.4	22.4
7.3	4.0	1.4	0.8	0.4	0.2	88	5.7	0.1	7.7	3.3	6.2	20.3
11.8	7.8	2.6	1.0	0.7	0.2	94	6.8	0.1	7.6	3.2	6.7	23.4
11.7	8.3	2.9	1.3	0.8	0.2	90	5.3	0.1	3.0	1.3	1.6	22.5
15.4	9.8	3.6	0.9	0.9	0.3	94	5.6	0.1	4.2	2.2	2.8	28.5
13.8	15.0	4.0		0.8	0.3		5.4	0.1	6.0	3.0	3.4	28.0

1/ Percent of whole soil

ARLINGTON SANDY LOAM
S57Calif-33-10

Location: 1200 feet east and 100 feet north of the south 1/4 corner of Sec. 21, T. 4 S., R. 3 W. On the Roy Smith Ranch, about 2 miles northeast of Perris, Riverside County, California. Date of Sampling: June 5, 1957.
Description by: Fred Eggers. Collectors: George Borst, R. E. Nelson, Cliff Henry, and Arnold Knecht.
Vegetation: Fallow land. Precipitation: 12 inches. Parent Material: Granitic alluvium. Physiography: Alluvial fan. Elevation: 1425 feet. Drainage: Well drained.
Classification: Haplic Durixeralf, coarse-loamy, mixed, thermic.

HORIZON	DESCRIPTION
S57Calif-33-10-1 Ap 0-6"	Brown (10YR 5/3 dry, dark yellowish brown 3/4 moist) sandy loam; moderately defined medium subangular blocky breaking to weakly defined fine granular; hard when dry, friable when moist; nonsticky and slightly plastic when wet; moderately rapid permeability; many roots; many fine pores; pH 6.0; noncalcareous; clear smooth lower boundary.
S57Calif-33-10-2 A1 6-12"	Brown (10YR 5/3 dry, dark yellowish brown 3/4 moist) sandy loam; moderately defined medium subangular blocky breaking to weakly defined fine granular; hard when dry, firm when moist; nonsticky and slightly plastic when wet; moderately rapid permeability; many roots; many fine pores; pH 6.2; noncalcareous; clear smooth lower boundary.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Fresno loam (taxadjunct) **LOCATION** Fresno County, California

SOIL FAMILY: Natric Durixeralf, coarse-loamy, mixed, thermic.

SOIL NOS. S58Calif-10-1 **LAB. NOS.** 58245 - 58250

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1	2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	
0-6	A1	0.3	2.9	3.7	13.2	17.4	53.4	9.1	57.5	22.5	0	sil
6-12	B2	0.2	2.0	2.4	9.5	12.2	48.3	25.4	41.2	26.3	0	l
12-21	B3t	7.6	23.5	3.2	6.9	2.3	45.1	11.4	27.8	24.5	0	l
21-28	C1sicam	9.7	13.6	6.3	12.1	14.8	37.3	6.2	41.6	18.1	4	2/
28-39	C2sica	8.2	10.8	5.7	15.2	15.2	39.0	5.9	46.6	17.3	1	2/
39-63	C3sica	2.1	3.0	2.3	9.9	18.1	60.4	4.2	55.9	30.1	1	2/

FRESNO LOAM (taxadjunct) 1/
S58Calif-10-1

Location: Fresno County, California. 2,390 feet west and 1,640 feet south of northeast corner, NW1/4 Sec. 18, T. 15 S., R. 18 E. Date of Sampling: October 31, 1958.
Description by: R. Ulrich. Collectors: R. E. Nelson, Jess Glouser, Gene Novak, and R. Ulrich.
Classification: Natric Durixeralf, coarse-loamy, mixed, thermic.
Vegetation: Saltgrass, spikeweed, pickleweed, soft chess, jackass clover. Range pasture land use.
Climate: About 9 inches rainfall. Mean annual temperature about 63° F. Parent Material: Largely water transported, sorted, and deposited sediments from granitic sources in Sierra Nevada to eastward. The position is basin or basin rim by University of California physiographic terminology.
Slope: Essentially level plain, less than 1 percent slope, with minor microrelief including some low mounds. Elevation: 220 feet as shown in Kearney Park USGS Topographic Quadrangle. Drainage: Imperfect natural drainage, very slow internal drainage due to hardpan, very slow runoff.

DESCRIPTION

A1 RSL No. 58245 — 0-6 inches, light gray (2.5Y 7/2) light loam (this texture is questionable; it is being mapped fine sandy loam), grayish brown (2.5Y 5/2) moist, very weak very fine granular moist drying essentially massive with some coarse prismatic cracks appearing late in the dry season; hard, friable, very slightly plastic, nonsticky; few fine roots in soil mass but numerous and concentrated along the prismatic cracks, with mat-like root concentrations occuring in places; many very fine interstitial pores, common extremely fine and very fine tubular pores; slightly calcareous; reaction not determined in field, see laboratory data; abrupt, slightly wavy lower boundary.

B2t RSL No. 58246 — 6-12 inches, grayish brown (2.5Y 5/2) light clay loam, very dark grayish brown (2.5Y 4/2) with darker (2.5Y 3/2 and 3/1) organic films on many ped faces, moist; weak coarse prismatic and moderate medium angular blocky; very hard, very firm, plastic, slightly sticky; few fine roots in soil mass, roots more numerous and somewhat flattened along structure planes; common extremely fine and very fine tubular pores; slightly calcareous, the lime disseminated; thin to moderately thick patchy clay films; clear slightly wavy lower boundary.

B3t RSL No. 58247 — 12-21 inches, light brownish gray (2.5Y 6/2) loam, grayish brown (2.5Y 5/2) with darker (2.5Y 4/2 and 3/2) organic films on some ped faces, moist; weak medium subangular blocky; very hard, very firm, very slightly plastic, nonsticky; few fine roots in soil mass; common extremely fine and very fine tubular pores; slightly calcareous, the lime disseminated; thin patchy clay films; abrupt, slightly wavy boundary.

C1sicam RSL No. 58248 — 21-28 inches, variable light brownish gray and light gray (2.5Y 6/2 and 7/2) indurated lime-silica hardpan, grayish brown (2.5Y 5/3) moist, with a very thin, very dense, dark grayish brown (2.5Y 4/2) upper crust; this extremely cemented crusty cementing material appears elsewhere in the hardpan but it is thinner, more lenselike and less continuous than on the upper surface of the hardpan; very thin light colored lime segregations and crusts may also appear on the surface of the hardpan and at various places within the hardpan horizon; the lower portion of the hardpan becomes more platy, lenselike, and fragmental and softer materials are also present; roots are very few, fine and in places flattened and matted - they appear to follow cracks and joint planes and plates in the hardpan; calcareous (amount uncertain) with little if any softening upon treatment with dilute acid; abrupt slight wavy lower boundary.

C2sica RSL No. 58249 — 28-39 inches, light gray (2.5Y 7/2) stratified fine sandy loam and loam, light brownish gray (2.5Y 6/2) moist, with some thin silt lenses; few faint fine yellowish brown (10YR 5/6 and 5/8) moist mottles; massive; hard, firm to friable, very slightly to nonplastic, nonsticky; no roots; common extremely fine and very fine interstitial pores and few extremely fine and very fine tubular pores; moderately calcareous, most of the lime being softly segregated in a reticulate pattern or in very thin slightly hardened masses and lenses; clear smooth lower boundary.

C3sica RSL No. 58250 — 39-63 inches, light gray (2.5Y 7/1) micaceous, stratified, finely to coarsely laminated fine sands and silts, light brownish gray (2.5Y 6/2) moist, variable few to numerous fine faint to distinct mottles of 10YR 5/6 and 5/8 and 7.5YR 5/6 and 5/8; variable consistency but mostly hard, firm to friable, very slightly plastic to nonplastic, and nonsticky; no roots; common extremely fine and few very fine interstitial pores; variable slightly to strongly calcareous in very thin lenses and seams of softly to somewhat hardened lime segregations, probably will average moderately calcareous; clear smooth lower boundary.

C4 RSL No. 58251 — 63 inches +, similar to material above but very slightly calcareous to noncalcareous in most of mass.

1/ The Fresno series is in a fine-loamy, mixed, thermic family of Natric Durixeralfs. This pedon has an argillic horizon with less than 18 percent clay.

CONSECUTIVE PEDON NUMBER C 69 59 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES BONTA DEEP VARIANT SOIL NO. S69 CALI. 33- 1 SAMPLE NO. 9 309-9 315 SAN BERNARDINO AREA

SOIL FAMILY TYPIC HAPLOXERALF COARSE-LOAMY,MIXED,MESIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A	INT. II .2- .02	1A2A COARSE FRAGMENTS PCT. OF WHOLE SOIL I 75- 20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75		
9 309	0- 5	AP1	82.1	11.7	6.2	13.4	22.0	14.3	22.7	9.7	5.7	6.0	0.2	6.0		27.1	--	--	--	--
9 310	5- 20	AP2	81.0	12.9	6.1	12.9	21.1	13.9	23.3	9.8	6.7	6.2	TR	6.0		28.9	--	--	--	--
9 311	20- 30	AP3	78.2	15.2	6.6	12.2	19.6	12.6	23.3	10.5	7.6	7.6		7.0		30.8	--	--	--	--
9 312	30- 58	A14	75.0	15.4	9.6	11.4	19.7	11.6	22.4	9.9	6.4	9.0		10.0		28.3	--	--	--	--
9 313	58- 79	A15	75.6	16.5	7.9	9.3	18.9	12.5	24.7	10.2	4.7	11.8		8.0		28.1	--	--	--	--
9 314	79-135	B21T	74.9	15.5	9.6	7.8	15.7	12.3	27.1	12.0	6.7	8.8		10.0		33.6	--	--	--	--
9 315	135-165	2B22T	78.7	13.6	7.7	10.5	21.4	12.9	24.0	9.9	5.4	8.2		8.0		27.0	--	--	--	--

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	19-2 6E1B	EXT. IRON FE 6C2B	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4AH	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTENSBLTY LEF 4D1	LE 4D1	PH H2O 1/1 8C1A	H2O 1/10 8C1A	CACL2 8C1E
		-PERCENT--		-PERCENT-		PCT.	PCT.	NO.40	GRAMS PER CC				----PERCENT---			-PERCENT--				
9 309	0- 5	2.11													4.6			6.1		5.7
9 310	5- 20	1.11	0.064												3.8			7.0		6.5
9 311	20- 30	0.50	0.035												3.7			6.8		6.1
9 312	30- 58	0.13													4.7			6.2		5.6
9 313	58- 79	0.20													4.1			6.3		6.1
9 314	79-135	0.13													4.8			5.8		5.3
9 315	135-165	0.13													4.1			6.0		5.5

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT.	C E C NA- OAC 5A2A	NH4- OAC 5A6A	EC 8A1A	WATER EXTRACT FROM SATURATED PASTE CA 6N1B	MG 6O1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	B 6T1A	PH SAT. PASTE 8C1B
		--------MEQ/100G--------					PCT.	-MEQ/100G		MMHO	----------MEQ/1000 ML----------								PPM	
9 309	0- 5																			
9 310	5- 20																			
9 311	20- 30																			
9 312	30- 58																			
9 313	58- 79																			
9 314	79-135																			
9 315	135-165																			

SAMPLE NO.	DEPTH CM.	WATER EXCHANGBLE AT 8A PCT.	NA 5B4B MEQ.	NA 5D2 PCT.	SAR 5E	GYPSUM AMT. 6F1B PERCENT	REQ. 6P3A	CLAY MINERALOGY LESS THAN 0.002 MM 7A2E I	7A2E II	7A2E III	7A2E IV	L.T. .0002 MM 7A3 PCT.	7A2E I	7A2E II	7A2E III	CODE	AMOUNT
9 309	0- 5																1 TRACE
9 310	5- 20																2 SMALL
9 311	20- 30																3 MODERATE
9 312	30- 58																4 ABUNDANT
9 313	58- 79																5 DOMINANT
9 314	79-135																6 INDETER.
9 315	135-165																

SAMPLE PREPARATION CODES

BONTA SANDY LOAM, DEEP VARIANT 1/
S69Calif-33-1

Location: Riverside County, California; the U. S. Forest Service Nixon Plots nursery area, on North Mountain, on the northwest flanks of the San Jacinto Mountain mass, south of Beaumont, California, SE¼ SE¼ sec. 8, T.4S., R1E.,SBBM. Date of Sampling: 12/2/66**

Description by: F. F. Peterson and C. S. Holzhey. Collectors: F. F. Peterson and C. S. Holzhey.**Resampled 6/27/69 by E. C. Nord.

Classification: Typic Haploxeralf, coarse-loamy, mixed, mesic.

and forbs. Climate: Mediterranean. Estimated total annual precipitation 23 inches, but varies from 13 to 43 inches that comes mostly between October and April.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Greenfield sandy loam **LOCATION** Riverside County, California

SOIL FAMILY: Typic Haploxeralf, coarse-loamy, mixed, thermic

SOIL NOS. S57Calif-33-9 **LAB. NOS.** 57513 - 57519

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	3A1 2A2 > 2 1/	TEXTURAL CLASS
0-5	A1	8.9	15.3	9.8	18.9	10.8	25.6	10.7	36.2	11.2	6	sl
5-12	B1	6.0	14.4	9.4	18.1	11.3	25.8	15.0	36.0	11.9	6	fsl
12-24	B21	9.7	16.7	8.5	15.0	8.1	22.2	19.8	28.8	9.9	6	cosl
24-32	B22	10.6	19.7	11.3	17.2	8.2	19.4	13.6	29.8	7.2	3	cosl
32-44	C1	6.2	14.5	9.3	18.4	11.4	26.9	13.3	38.1	10.8	3	sl
44-52	C2	8.8	16.4	10.2	17.3	10.0	23.8	13.5	33.7	9.9	6	cosl
52-60+	B2b	7.3	12.0	7.3	13.3	7.3	16.9	35.9	25.6	6.1	4	sc

pH 8C1b Saturated Paste 1:1	8C1a 1:10	4A3a Bulk Density g/cc	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	6I1a CO_3 meq/l	8A1a Elec. Conduc. mmhos/cm. at 25° C.	6E1b $CaCO_3$ equivalent %	6J1a HCO_3 meq/l	MOISTURE TENSIONS 4B1a 1/10 ATMOS. %	4B1a 1/3 ATMOS. %	4B2 15 ATMOS. %
5.9	7.6		0.48	0.075	6.4	0.0	14.0	0	2.8	19.4	13.4	4.1
6.8	7.0	1.90	0.28	0.044	6.4	0.6	2.8	0	5.5			5.4
6.8	7.0	1.84	0.10	0.027		0.0	2.3	0	4.3			6.5
7.2	7.5	1.82	0.25	0.014		0.0	2.9	0	4.7			5.0
7.4	8.1	1.83	0.24	0.011		0.6	4.0	Tr	4.3			5.2
7.4	8.3	1.74	0.03			0.0	4.4	Tr	3.6			5.1
7.1	7.5	1.77	0.01			0.0	4.8	Tr	3.0			11.3

5A2a CATION EXCHANGE CAPACITY (+ Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	5D1 Exch. Na %	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a Ca	6O1a Mg	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
7.5	7.6	2.7	1.4	0.5	0.5	7	34.0	3.6	77.1	28.7	31.4	25.9
9.3	6.8	1.9	1.2	1.1	0.3	12	11.0	0.3	11.9	3.3	15.3	25.3
12.6	8.7	2.7	1.2	1.5	0.2	12	11.1	0.2	6.0	3.5	13.2	28.2
13.0	8.7	2.5	0.8	1.4	0.2	11	14.3	0.1	9.5	3.8	17.8	25.4
11.4	10.8	2.6	0.4	0.8	0.2	7	17.0	0.2	15.1	6.1	27.8	26.3
11.8	12.9	2.8	1.0	2.1	0.1	18	16.4	0.2	18.0	7.4	34.6	23.6
24.5	17.3	6.7	1.4	0.6	0.3	2	17.2	0.2	20.7	7.8	39.4	49.5

1/ Percent of the whole soil

GREENFIELD SANDY LOAM
S57Calif-33-9

Location: Riverside County, California. 150 feet north and 150 feet east of the south 1/4 corner of Sec. 3, T. 5 S., R. 3 W. On the Clyde Smith Ranch, about 2 miles southeast of Perris. Date of Sampling: June 5, 1957.
Description by: Fred Eggers. Collectors: George Borst, R. E. Nelson, Cliff Henry, and Arnold Knecht.
Classification: Typic Haploxeralf, coarse-loamy, mixed, thermic.
Vegetation: Fallow land. It was in irrigated alfalfa many years, potatoes two years, and is now being leveled. Precipitation: 12 inches. Parent Material: Granitic alluvium. Physiography: Alluvial fan. Elevation: 1426 feet.
Drainage: Well drained.

HORIZON	DESCRIPTION
S57Calif-33-9-1 A1 0-5"	Grayish brown to brown (10YR 5/2-5/3 dry, 3/3 moist) sandy loam; moderately defined fine granular; slightly hard when dry, friable when moist; nonsticky and slightly plastic when wet; moderately rapid permeability; occasional plant roots; many fine pores; pH 6.0; noncalcareous; clear smooth lower boundary.
S57Calif-33-9-2 B1 5-12"	Brown to yellowish brown (10YR 5/3-5/4 dry, dark yellowish brown 3/4 moist) loam; strongly defined medium granular; hard when dry, firm when moist; slightly sticky and plastic when wet; moderately slow permeability; few roots; many fine pores; pH 6.7; gradual smooth lower boundary.
S57Calif-33-9-3 B21 12-24"	Brown (7.5YR 5/4 dry, dark brown 3/4 moist) sandy clay loam; massive to strongly defined medium angular blocky; very hard when dry; firm when moist; sticky and plastic when wet; few roots; many fine pores; pH 7.7; noncalcareous; weak discontinuous clay skins; gradual smooth lower boundary.
S57Calif-33-9-4 B22 24-32"	Brown (7.5YR 5/4 dry, dark brown 3/4 moist) gritty sandy clay loam; massive to strongly defined medium angular blocky; very hard when dry, firm when moist; sticky and plastic when wet; moderately slow permeability; no plant roots; many fine pores; pH 8.0; noncalcareous; moderately strong clay skins. Clear smooth lower boundary.
S57Calif-33-9-5 C1 32-44"	Brown (7.5YR 5/4 dry, dark brown 3/4 moist) sandy clay loam; massive; breaking to moderately defined medium granular; slightly hard when dry, friable when moist; slightly sticky and slightly plastic when wet; moderate permeability; no plant roots; occasional fine pores; pH 8.0; slight effervescence; fragmentary clay skins; gradual smooth lower boundary.
S57Calif-33-9-6 C2 44-52"	Brown (10YR 5/3 dry, dark brown 3/4 moist) gritty sandy clay loam; massive; breaking to strongly defined medium angular blocky; very hard when dry, firm when moist; sticky and plastic when wet; moderate permeability; no plant roots; occasional fine pores; pH 8.0; slight effervescence; fragmentary clay skins; abrupt smooth lower boundary.
S57Calif-33-9-7 B2b 52-60"+	Brown (7.5YR 5/4 dry, dark brown 4/4 moist) sandy clay; massive; extremely hard when dry, very firm when moist; sticky and very plastic when wet; very slow permeability; no roots; occasional fine pores; pH 8.0; slight effervescence with soft lime segregated in seams; heavy clay skins on all peds.

Remarks: Fertilizers used on potatoes in 1955 totaled approximately 200 pounds of P per acre. Slight mottling observed in B1 horizon.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Greenfield coarse sandy loam **LOCATION** Riverside County, California

SOIL FAMILY: Typic Haploxeralf, coarse-loamy, mixed, thermic

SOIL NOS. S57Calif-33-16 **LAB. NOS.** 57564 - 57570

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1	2A2	TEXTURAL CLASS
		VERY COARSE	COARSE	MEDIUM	FINE	VERY FINE						

GREENFIELD COARSE SANDY LOAM
S57Calif-33-16

Location: Riverside County, California. 1500 feet south and 1100 feet west of the intersection of San Diego aqueduct and Long Valley road, 1/2 mile north and 1¼ miles east of Temecula School, Temecula. Date of Sampling: June 10, 1957.
Description by: Fred Eggers. Collectors: George Borst, R. E. Nelson, Cliff Henry, Arnold Knecht, and Fred Eggers.
Classification: Typic Haploxeralf, coarse-loamy, mixed, thermic.
Vegetation: Wild oats, bur clover, dove weed, lupine, soft chess, Turkey mullein and alfileree. Precipitation: 14 inches. Parent Material: Granitic material.
Physiography: A very old non-marine terrace thoroughly dissected.
Elevation: 1260 feet. Drainage: Well drained.
Remarks: All horizons very gritty and micaceous. Evidence of much animal activity in All horizon. Comparable soil profiles are found only on ridges and other tops of terrace remnants. Prominent discontinuous clay skins in B horizons.

HORIZON	DESCRIPTION
S57Calif-33-16-1 A_{p1} 0- 1"	Grayish brown (10YR 5/2 dry, dark brown 3/3 moist) coarse sandy loam; weakly defined fine granular; soft when dry, very friable w hen moist; nonsticky and nonplastic when wet; moderately rapid permeability; many roots; many pores; pH 6.5; noncalcareous; clear smooth lower boundary.
S57Calif-33-16-2 A_{p2} 1- 8"	Grayish brown (10YR 5/2 dry, dark brown 3/3 moist) coarse sandy loam; moderately defined medium subangular blocky breaking to weakly defined fine granular; slightly hard when dry, very friable when moist; nonsticky and nonplastic when wet; moderately rapid permeability; many roots; many pores; pH 6.0; gradual wavy lower boundary.
S57Calif-33-16-3 A_{11} 8-18"	Grayish brown (10YR 5/2 dry, dark brown 3/3 moist) coarse sandy loam; moderately defined medium subangular blocky breaking to weakly defined fine granular; hard when dry, very friable when moist; nonsticky and nonplastic when wet; moderately rapid permeability; many roots; many pores; pH 5.5; noncalcareous; gradual wavy lower boundary.
S57Calif-33-16-4 A_{12} 18-32"	Grayish brown (10YR 5/2-6/2 dry, dark brown 3/3 moist) sandy loam; moderately defined medium subangular blocky breaking to weakly defined fine granular; slightly hard when dry, very friable when moist; nonsticky and nonplastic when wet; moderately rapid permeability; many roots; many pores; pH 6.0; noncalcareous; abrupt wavy lower boundary.
S57Calif-33-16-5 B_{21m} 32-41"	Light brownish gray (10YR 6/2 dry, dark brown 4/3 moist) sandy clay loam; massive; extremely hard when dry, firm when moist; slightly sticky and slightly plastic when wet; moderately slow permeability; few roots; many pores; pH 6.0; noncalcareous; gradual wavy lower boundary.
S57Calif-33-16-6 B_{22m} 41-47"	Light reddish brown (5YR 6/3 dry, reddish brown 4/3 moist) sandy clay loam; massive; very hard when dry, firm when moist; slightly sticky and slightly plastic when wet; moderately slow permeability; no roots; occasional pores; pH 5.5; noncalcareous; clear wavy lower boundary.
S57Calif-33-16-7 C 47-64"+	Light reddish brown (5YR 6/3 dry, reddish brown 4/3 moist) loamy coarse sand containing lenses and layers of clay material; single grain; very hard when dry, very friable when moist; nonsticky and nonplastic when wet; no roots; occasional pores; pH 6.0.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Greenfield coarse sandy loam **LOCATION** Riverside County, California

SOIL FAMILY: Typic Haploxeralf, coarse-loamy, mixed, thermic.

SOIL NOS. S57Calif-33-17 **LAB. NOS.** 57571 - 57577

DEPTH INCHES	HORIZON	1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	3A1 2A2 > 2 1/	TEXTURAL CLASS
		PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)										
0-10	Ap	12.5	17.5	10.0	18.6	9.4	22.2	9.8	31.0	11.1	10	cosl
10-18	A11	11.4	17.0	10.4	20.0	10.9	22.9	7.4	31.4	13.8	10	cosl
18-25	A12	10.0	20.2	10.8	17.6	9.2	21.2	11.0	28.5	11.6	11	cosl
25-32	B21	9.0	17.1	10.4	19.2	9.8	22.6	11.9	32.1	11.4	14	cosl
32-43	B22	13.9	19.6	9.8	17.0	8.3	18.2	13.2	26.2	9.9	10	cosl
43-55	B3	14.1	23.0	11.6	17.4	7.2	16.2	10.5	23.8	9.2	10	cosl
55-72	C	18.7	31.3	12.4	14.3	4.7	9.8	8.8	15.8	6.1	20	lcos

	pH		ORGANIC MATTER		6I1a	8A1a	6E1b	6J1a	MOISTURE TENSIONS		
8C1b SATURATED PASTE		8C1a	6A1a ORGANIC CARBON	6B1a NITROGEN	CO_3	Elec. Conduc. mmhos/	$CaCO_3$ equiv.	HCO_3	4B1a 1/10	4B1a 1/3	4B2 15

GREENFIELD COARSE SANDY LOAM
S57Calif-33-17

Location: Riverside County, California. North of Long Valley Road, 500 feet southwest of Buck Messa benchmark. On the Vail Company Ranch, Pauba Grant, 2½ miles north and 6½ miles east of Temecula School, Temecula. Date of Sampling: June 10, 1957.
Description by: Fred Eggers. Collectors: George Borst, R. E. Nelson, Cliff Henry, Arnold Knecht, and Fred Eggers.
Classification: Typic Haploxeralf, coarse-loamy, mixed, thermic.
Vegetation: Barley. Climate: 14 inches rainfall. Parent Material: Granitic alluvium; ancient nonmarine terrace.
Elevation: 1,448 feet. Drainage: Well drained.
Remarks: All horizons very gritty and micaceous.

HORIZON	DESCRIPTION
S57Calif-33-17-1	Grayish brown (10YR 5/2 dry, dark brown 3/3 moist) [redacted] sticky and nonplastic when wet; moderately rapid permeability; many roots; many pores; pH 6.0; noncalcareous; clear smooth lower boundary.
S57Calif-33-17-2	Grayish brown (10YR 5/2 dry, dark brown 3/3 moist)

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Ramona fine sandy loam (taxadjunct) **LOCATION** Riverside County, California

SOIL FAMILY: Typic Haploxeralf, coarse-loamy, mixed, thermic

SOIL NOS. S59Calif-33-2 **LAB. NOS.** 59556 - 59566

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	3A1 0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-9	Ap	9.2	17.1	11.9	20.9	12.2	24.4	4.3	40.8	7.8	13	cosl
9-13	A12	7.8	16.8	11.4	20.1	11.9	25.2	6.8	40.0	8.9	15	sl
13-17	A13	11.7	18.3	10.8	17.5	10.3	22.4	9.0	34.4	8.2	20	cosl
17-24	B1t	11.2	17.3	9.9	16.4	9.8	21.4	14.0	32.5	7.9	24	cosl
24-31	B21t	16.1	18.1	8.2	14.7	8.5	19.0	15.4	28.3	7.5	15	cosl
31-45	B22t	11.3	14.5	8.6	17.9	9.6	19.2	18.9	30.9	8.4	13	cosl
45-51	B31t	11.2	16.5	10.6	19.8	9.8	17.0	15.1	29.8	8.3	15	cosl
51-58	B32t	12.2	17.4	11.5	23.3	9.7	14.6	11.3	30.6	6.7	20	cosl
58-67	C1	23.4	24.5	10.9	17.9	5.6	11.6	6.1	20.7	5.5	24	lcos
67-87	C2	17.7	22.7	13.2	20.3	7.3	12.8	6.0	25.3	5.5	29	lcos
87-96+	C3	12.3	19.8	11.8	24.7	10.6	13.8	7.0	33.0	5.5	23	lcos

RAMONA FINE SANDY LOAM (taxadjunct) 1/
S59Calif-33-2

Location: Riverside County, California. About 3 miles northeast of the town of Sunnymead, approximately 1370 feet north, 1220 feet west of southeast corner of Sec. 34, T. 2 S., R. 3 W. Date of Sampling: September 11, 1959.
Description by: Arnold Knecht and George Borst. Collectors: Arnold Knecht, Cliff Henry, Reuben Nelson and Lothair Grant.
Classification: Typic Haploxeralf, coarse-loamy, mixed, thermic.
Vegetation: Grain stubble. Climate: Average annual precipitation about 11 inches; average January temperature about 50° F., average July temperature about 70° F., and average annual temperature about 62° F.
Parent Material: Moderately coarse textured alluvium, predominantly from granitic sources. Topography: Gently sloping alluvial terrace, with about 3 percent southeasterly slope. Elevation: 1,880 feet. Drainage: Well drained. Soil Moisture: None.
Remarks: This profile is characterized by an ochric epipedon and an argillic horizon, according to the revised system of soil classification (7th Approximation). A weak duripan, with coarse polygonal structure, was observed in the road cut about 100 feet west of sampling site.

HORIZON	DESCRIPTION
Ap RSL No. 59556	0-9 inches (0-23 cm.), brown (10YR 5/3) coarse sandy loam, dark yellowish brown (10YR 3/4) when moist; massive; hard, very friable, nonsticky and nonplastic, few fine roots; common very fine tubular pores; slightly acid (pH 6.2); abrupt, smooth boundary.
A12 RSL No. 59557	9-13 inches (23-33 cm.), brown (10YR 5/3) coarse sandy loam, dark yellowish brown (10YR 3/4) when moist; massive; very hard, very friable; nonsticky and nonplastic; plentiful fine roots; common very fine tubular pores; slightly acid (pH 6.2); abrupt, smooth boundary. This horizon may be a tillage pan.
A13 RSL No. 59558	13-17 inches (33-43 cm.), brown (10YR 5/3) coarse sandy loam, dark yellowish brown (10YR 3/4) when moist; massive; hard, very friable, nonsticky and nonplastic; plentiful fine roots; many fine and very fine tubular pores; slightly acid (pH 6.5); clear, smooth boundary.
B1t RSL No. 59559	17-24 inches (43-61 cm.), brown (10YR 5/3) coarse sandy loam, dark yellowish brown (10YR 3/4) when moist; weak coarse prismatic structure; very hard, friable, slightly sticky and slightly plastic; few fine roots; many fine to very fine tubular pores; few very thin discontinuous clay films on ped faces; neutral (pH 6.8); clear, smooth boundary.
B21t RSL No. 59560	24-31 inches (61-79 cm.), reddish brown (5YR 4/3) coarse sandy loam near sandy clay loam; dark reddish brown (5YR 3/3) when moist; moderate coarse prismatic structure; very hard, firm, sticky and plastic; few fine roots; many fine to very fine tubular pores; few moderately thick clay films on ped faces; neutral (pH 7.2); gradual smooth boundary.
B22t RSL No. 59561	31-45 inches (79-115 cm.), reddish brown (5YR 4/3) coarse sandy loam near sandy clay loam, dark reddish brown (5YR 3/3) when moist; moderate coarse prismatic structure; very hard, firm, sticky and plastic; common fine to very fine tubular pores; few moderately thick clay films on ped faces; neutral (pH 7.2); gradual, smooth boundary.

SCS-420
10-64

U S DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haploxeralf, fine-loamy, mixed, thermic

SOIL Fallbrook sandy loam SOIL Nos. S64Calif-37-2 LOCATION San Diego County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 6414 - 6422

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Silt: Int. II (0.2-0.02)	(2-0.1)	* Clay <.002	** Clay <.002	Coarse fragments 1A2a: > 2	2 - 19	19 - 76
		Pct. of < 2 mm														Pct.	Pct. of < 76 mm	
0- 2	A11	72.9	19.5	7.6	13.3	19.2	10.2	19.5	10.7	10.1	9.4	31.3	62.2	7.8	7.2	8	8	0
2- 6	A12	72.4	19.4	8.2	13.1	18.2	10.1	19.7	11.3	9.6	9.8	31.5	61.1	8.6	7.8	9	9	0
6-12	A3	66.4	21.0	12.6	11.8	16.4	9.4	18.3	10.5	9.4	11.6	29.3	55.9	12.5	11.6	3	3	0
12-20	B21t	54.4	19.5	26.1	11.6	12.1	7.6	14.1	9.0	8.4	11.1	25.1	45.4	28.0	27.0	4	4	0
20-28	B22t	51.8	17.0	31.2	12.1	11.6	7.4	13.4	7.3	7.0	10.0	21.3	44.5	31.4	30.4	3	3	0
28-47	B3	72.4	12.8	14.8	17.0	21.6	10.5	16.7	6.6	5.1	7.7	19.7	65.8	13.6	12.1	11	11	0
47-68	C1	79.6	11.2	9.2	23.8	25.8	9.6	14.9	5.5	4.1	7.1	16.3	74.1	8.3	6.7	19	19	0
68-85	C2	80.7	13.9	5.4	21.7	25.0	10.4	16.9	6.7	6.7	7.2	21.8	74.0	4.5	3.6	21	21	0
85-90+	C3	80.2	14.7	5.1	27.1	23.4	9.1	14.8	5.8	7.6	7.1	20.3	74.4	8.1	5.2	36	36	0

Depth (In.)	6A1a * Organic carbon	6B1a* Nitrogen	C/N	*6C2a Ext. Iron as Fe	6E1b Carbonate as $CaCO_3$		Bulk density: 4A1f 1/3 bar	Bulk density: 4A1h Oven dry		3B2 C'	Water content	4B1c 1/3 bar	4B2* 15 bar	4D1 Extensibility COLEF	4D1 Extensibility COLE	pH: 8C1b Saturated Paste	pH: 8C1a H_2O 1:1	pH: 8C1a H_2O 1:10
	Pct.	Pct.		Pct.	Pct.		g/cc	g/cc	g/cc		Pct.	Pct	Pct.	in./in.	in./in.			
0- 2	0.73	0.070	10	0.8			1.42	1.44		0.96		10.2	2.9	0.005	0.004	6.4	6.6	
2- 6	0.50	0.047	10	0.7			1.60	1.62		0.94		10.5	3.6	0.003	0.003	6.3	6.5	
6-12	0.30			0.7			1.78	1.85		0.98		11.0	5.0	0.013	0.013	6.5	6.8	
12-20	0.27			1.4			1.78	1.92		0.97		15.0	10.1	0.026	0.025	6.6	6.8	
20-28	0.24			1.4			1.73	1.89		0.98		16.5	11.3	0.029	0.029	6.7	7.0	
28-47	0.07			0.9			1.80	1.89		0.92		13.7	7.8	0.016	0.015	7.0	7.4	
47-68	0.02			0.7			1.81	1.90		0.86		10.9	5.9	0.016	0.014	7.4	7.6	
68-85	0.01			0.6									3.7			7.9	8.2	
85-90+	0.02			0.8									4.2			7.9	8.1	

Depth (In.)	Extractable bases 6N2a * Ca	6O2a* Mg	6P2a* Na	5B1a 6Q2a* K	Sum of bases	6H2a Ext.* Acidity	Cation Exch. Capacity 5A2a* NaOAc	5A3a * Sum	Clay mineralogy (<0.0002 mm) 7A2e Verm.		Illite		Kaol.	Water extract from saturated paste** 8A1 6P1a Na	6Q1a K	6L1b SO_4	8A1a Electrical conductivity
	meq/100 g								X-ray					meq/liter			mmho/cm
0- 2	4.6	1.3	0.3	0.4		1.5	9.6	8.1									
2- 6	5.2	1.3	0.3	0.1		1.6	9.2	8.5	tr.				xx				
6-12	6.5	2.0	0.4	0.1		1.9	10.8	10.9	x				xx				
12-20	10.0	4.7	0.4	0.1		2.9	18.3	18.1	x				xxx				
20-28	10.0	5.0	0.5	0.1		2.9	19.7	18.5	tr.				xxx				
28-47	11.3	5.0	0.9	0.1		1.7	19.3	19.0	tr.		tr.		xx				
47-68	10.3	4.3	0.9	0.1		1.2	16.7	16.8						1.1	tr.		0.18
68-85	8.5	3.9	0.8	tr.		0.8	13.7	14.0									
85-90+	9.4	3.7	0.9	0.2		0.8	15.1	15.0						0.8	tr.		0.07

Depth (In.)	8A Water at Saturation	5D2 Exchangeable Na	5B1b Exch. Na meq / 100g	Base Sat. Cations	Base Sat. NaOAc CEC	Clay mineralogy (<0.002 mm) 7A Relative abundance based on height of diagnostic X-ray peak (7A2): Mont.	Verm.	Hydrobiotite	Illite	Kaol.	% Kaol. (DTA) 7A3	% Kaol. in ground whole soil (DTA)
	Pct.	Pct.		Pct.	Pct.	X-ray						
0- 2		3		81	69	-	-	t	xxxx	xx	20	tr.
2- 6		4		82	75	-	-	-	xxxx	xx		
6-12		4		82	81	-	-	xxxx	x	xx		
12-20		2		84	83	-	x	xxx	x	xxx		
20-28		3		84	80	-	t	t	xx	xxxx	33	17
28-47		5		91	90	-	t	t	xx	xxxx	29	12
47-68	38.2	5		93	93	-	x	t	xx	xxx	12	tr.
68-85		6		94	96							
85-90+	44.1	6		95	94	x	xx	t	xx	xx	0	1

1/ From characterization sample; determined by shaking overnight in a sodium hexametaphosphate solution.

* Analysis of ground whole soil; results expressed on <2-mm basis.

** Analysis of ground whole soil; results expressed on whole-soil basis.

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

FALLBROOK SANDY LOAM
S64Calif-37-2

Location: San Diego County, California. A soil profile pit located about 2-1/4 miles south and 3/4 mile east of Fallbrook in the NE1/4 of the SW1/4 of section 31, T. 9 S., R. 3 W. Date of Sampling: February 17, 1964.
Description By: Roy H. Bowman and Gerald Kester. Collectors: Klaus Flach, Gerald Kester, George Borst, Benny Brasher and R. E. Nelson.
Classification: Typic Haploxeralf, fine-loamy, mixed, thermic.
Vegetation: Wild oats, annual weeds and grasses. Climate: Annual precipitation is 15 inches and mean annual temperature is 60° F. Parent Material: Tonalite grus. Topography: Southwest-facing, smooth 12 percent slopes in middle backslope position on rolling upland. Elevation: 555 feet. Drainage: Well drained; medium runoff; moderately slow permeability. Soil Moisture: Dry when sampled.
Remarks: The B3 horizon appears to be the most dense of the profile, and the B3 horizon the lowest in permeability. This soil occurs on a slope 320 feet SE of the Vista site S64Calif-37-1.

HORIZON	DESCRIPTION
A11 RSL No. [illegible]	0 to 2 inches, brown (10YR 5/3) sandy loam, dark brown (10YR 3/3) moist; weak fine crumb structure; slightly hard, very friable, nonsticky and nonplastic; abundant [illegible]
A12	2 to 6 inches, brown (10YR 5/3) sandy loam, dark brown (10YR 3/3) moist; massive; [illegible]

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haploxeralf, fine-loamy, mixed, thermic
SOIL Greenfield gravelly loam (taxadjunct) SOIL Nos. S4557-4 LOCATION Tehama County, California

SOIL SURVEY LABORATORY _____ LAB. Nos. _____

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)			Coarse fragments > 2 Pct.	2-19	19-76
		Size class and particle diameter (mm) 3A1																
		Pct. of < 2 mm															Pct. of < 76mm	
0-9	Ap	39.1	49.9	11.0	6.6	6.9	3.9	7.5	14.2	22.5	27.4		24.9					
9-21	A1	39.7	45.5	14.8	6.2	7.1	4.4	7.6	14.4	20.8	24.7		25.3					
21-31	B21t	41.9	41.2	16.9	7.5	7.9	4.9	8.5	13.1	18.4	22.8		28.8					
31-43	B22t	46.0	33.7	20.3	9.7	9.1	6.0	10.4	10.8	13.0	20.7		35.2					
43-51	B3t	54.3	27.1	18.6	11.6	11.3	7.2	13.6	10.6	9.7	17.4		43.7					
57-65	C?	65.0	19.4	15.6	15.9	17.4	10.5	13.7	7.5	6.1	13.3		57.5					
84-107	C?	77.6	12.0	10.4	29.9	26.0	9.9	8.0	3.8	3.8	8.2		73.8					

Depth (In.)	6A2a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	Pct.			pH		8C1a (1:1) H_2O
0-9	0.69	0.081																5.5
9-21	0.30	0.051																5.9
21-31	0.14	0.037																6.4
31-43	0.12	0.038																6.6
43-51	0.10	0.034																6.7
57-65	0.08	0.028																6.9
84-107	0.08	0.021																6.8

Depth (In.)	Extractable bases 5B1a: Ca 6N2b	Mg 6O2b	Na 2/	K 1/	Sum of bases	6H2a Ext. Acidity	5A3a CEC Sum									Base saturation 5C3 Sum Pct.	Pct.
	meq/100 g																
0-9	3.4	0.8	<0.01	0.53	4.7	4.4	9.1									52	
9-21	5.0	1.3	tr.	0.25	6.5	2.6	9.2									71	
21-31	5.8	3.8	0.02	0.13	9.8	2.7	12.4									79	
31-43	5.9	5.7	tr.	0.16	11.8	3.2	15.0									78	
43-51	5.3	6.4	0.00	0.14	11.8	1.8	13.6									87	
57-65	5.4	6.5	0.00	0.12	12.0	2.1	14.1									85	
84-107	3.7	5.8	tr.	0.08	9.6	1.5	11.1									86	

Depth (In.)					

1/ Precipitated as potassium sodium cobaltinitrite and titrated as potassium permanganate

2/ Gravimetrically as sodium uranyl magnesium acetate

GREENFIELD GRAVELLY LOAM (taxadjunct) 1/
S4557-4

Location: Tehama County, California. From pit located 55 yards southwest of northeast corner Sec. 6, R24N, R3W, MDB&M. Date of Sampling: April 2, 1945.

Description by: R. C. Roberts and R. A. Gardner. Collectors: R. C. Roberts, F. O. Youngs, R. A. Gardner, L. T. Alexander, H. Jenny and R. E. Storie.

Classification: Typic Haploxeralf, fine-loamy, mixed, thermic.

Vegetation: Used for sheep pasture - formerly cultivated. Cover consists of a fairly dense stand of short annual grasses and associated herbaceous plants. Climate: Average annual precipitation is about 19 inches. Mean annual temperature is 62°F., January temperature is 45°F., and July 82°F. Parent Material: Alluvium from sedimentary rocks. Topography: Nearly level fans adjacent to stream channels. Elevation: 320 feet. Drainage: Well drained, moderate permeability, slow runoff. Soil Moisture: Moist.

Remarks: This soil has a distinct texture and color profile that does not appear to be due to stratification. The structural units of the subsoil are slightly coated with reddish-colored colloidal material.

HORIZON	DESCRIPTION
Ap	0 to 9 inches, yellowish brown (10YR 5/4) gravelly loam (very near gravelly silt loam), brown (10YR 4/3) moist; weak coarse granular structure; slightly hard, very friable or nearly loose, slightly plastic; abundant small roots present; strongly acid (pH 5.5); abrupt smooth boundary.
A1	9 to 21 inches, yellowish brown gravelly loam; weak granular structure; color dry and moist and consistence similar to layer above; abundant small roots; medium acid (pH 5.9); clear wavy boundary.
B21t	21 to 31 inches, strong brown (7.5YR 5/5) gravelly loam, brown (7.5YR 4/4) moist; weak granular structure; slightly compact; slightly hard, friable, slightly plastic; abundant small roots; slightly acid (pH 6.4); gradual boundary.
B22t	31 to 43 inches, strong brown gravelly heavy loam (7.5YR 5/5), brown (7.5YR 4/4) moist; blocky structure; compact; hard, firm, plastic; plentiful small roots; neutral (pH 6.6); gradual boundary.
B3t	43 to 51 inches, strong brown gravelly heavy sandy loam, color dry and moist similar to layer above but very slightly less reddish; blocky structure; slightly compact; hard, friable, slightly to moderately plastic; few roots; neutral (pH 6.7); clear wavy boundary.
C?	57 to 65 inches, light yellowish brown (10YR 6/4) very gravelly coarse sandy loam, color moist similar to layer above; massive; slightly hard, loose, nonplastic; very few roots; neutral (pH 6.9).
C ?	84 to 107 inches, light yellowish brown (10YR 6/4) very gravelly coarse sandy loam (near gravelly loamy coarse sand), yellowish brown (10YR 5/5) moist; structure and consistence similar to layer above; very few roots present; neutral (pH 6.8).

1/ The Greenfield series is in a coarse-loamy, mixed, thermic family of Typic Haploxeralfs. This pedon is in a fine-loamy family of the same subgroup.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Huichica loam variant **LOCATION** Sonoma County, California

SOIL FAMILY: Typic Haploxeralf, fine-loamy, mixed, thermic.

SOIL NOS. S61Calif-49-8 **LAB. NOS.** 61438 - 61445

DEPTH INCHES	HORIZON	1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
		PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1		
0- 8	Ap	1.7	2.9	5.6	19.4	13.7	45.4	11.3	47.0	24.4	5	1
8-19	A12	1.7	3.0	5.3	19.1	12.8	46.3	11.8	46.3	25.0	5	1
19-24	A2	1.2	3.0	5.3	19.1	13.3	46.5	11.6	47.1	25.2	5	1
24-31	B2t	0.9	3.0	5.1	17.6	11.8	38.2	23.4	44.0	17.5	4	1
31-42	B3tg	2.2	5.2	5.8	20.0	11.5	30.7	24.6	37.5	17.5	12	1
42-48	C1mg	9.2	17.6	13.7	18.6	6.7	14.4	19.8	24.3	7.1	21	cosl - scl
48-54	IIC2mg	18.3	23.1	9.4	17.5	8.9	15.5	7.3	26.9	7.2	0	cosl-lcos
54-66	IIC3	4.3	6.9	7.0	22.1	13.6	32.1	14.0	44.3	15.4	12	fsl

8C1a pH H_2O 1:1	8C1c pH KCl 1:1		6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N		Ext. Iron 6C1b % Fe					4B2 Soil water 15 atmos. Pct.
			ORGANIC MATTER									
5.7	4.4		0.82	0.079	10		0.4					5.2
5.6	4.2		0.29	0.045			0.4					5.3
5.9	4.3		0.18	0.035			0.3					5.2
5.8	4.0		0.15				0.4					11.6
5.4	4.0		0.08				0.3					12.1
5.4	3.8		0.03				0.6					10.7
5.5	3.9		0.02				0.5					14.7
6.1	4.3		0.01				0.6					14.1

5A2a CATION EXCHANGE CAPACITY ←(Na)	6N2b Ca	6O2c Mg	6H2a H	6P2a Na	6Q2a K	BASE SAT. % 5C3	Sum of Cations 5A3a meq/100g.					MOISTURE AT SATURATION %
	EXTRACTABLE CATIONS 5B1a											
	milliequivalents per 100g. soil →											
11.5	4.2	1.7	4.9	0.1	0.2	56	11.1					
10.4	3.6	1.9	4.2	0.2	0.2	58	10.1					
9.8	3.0	2.7	3.5	0.2	0.2	64	9.6					
18.5	5.2	7.6	4.0	0.7	0.2	77	17.7					
19.7	5.5	8.5	3.7	1.0	0.2	80	18.9					
19.7	5.4	8.1	3.5	1.2	0.1	81	18.3					
35.4	10.7	15.7	5.5	2.8	0.1	84	34.8					
30.7	10.1	14.7	4.6	2.1	0.2	85	31.7					

HUICHICA LOAM variant 1/
S61 Calif-49-8

Location: Sonoma County, California. Northwest corner of SE1/4, SE1/4, section 5, T. 7 N., R. 8 W. (Matteri Ranch). 165 feet west of fence corner. Date of Sampling: June 26, 1961. Collectors: Reuben Nelson, Leo C. Klameth, V. C. Miller, G. Lambert, J. H. Kashiwagi, F. Graham. Classification: **Typic Haploxeralf, fine-loamy, mixed, thermic.**

Vegetation: Annual grasses, a few perennial grasses and a few scattered oaks. Climate: Annual precipitation 29.4 inches, average temperature 57.6° F., average January temperature 46.5° F., average July temperature 66.7° F. Parent Material: Mixed valley alluvium. Elevation: About 135 feet. Relief: Hummocky on 1 - 2 percent slope. Drainage: Imperfect drainage, slow runoff, moderately slow to very slow permeability. Biological Activity: Abundant worm casts in zone

Ap RSL No. 61438 — 0 to 8 inches, light brownish gray (10YR 6/2) loam, dark grayish brown (10YR 4/2) when moist; massive; slightly hard when dry, very friable when moist, slightly sticky and plastic when wet; abundant very fine and fine roots; many very fine and fine tubular and interstitial pores; strongly acid, pH 5.3; clear, wavy boundary.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haploxeralf, fine-loamy, mixed, thermic

SOIL Montpellier gravelly loam SOIL Nos. S4557-11 LOCATION Tehama County, California

SOIL SURVEY LABORATORY _____ LAB. Nos. _____

Depth (In.)	Horizon	Size class and particle diameter (mm)															
		Total			Sand					Silt					Coarse fragments		
		Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	0.05–0.02	Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)			> 2 Pct.	2 – 19 / 19 – 76 Pct. of < 76mm
		Pct. of < 2 mm															

MONTPELLIER GRAVELLY LOAM
S4557-11

Location: Tehama County, California. Pit located 3/4 mile N. of the County road bend 6 miles west and 1 mile north of Corning. Date of Sampling: April 4, 1945.

Description by: R. C. Roberts and R. A. Gardner. Collectors: R. C. Roberts, F. O. Youngs, R. A. Gardner, L. T. Alexander, H. Jenny and R. E. Storie.

Classification: Typic Haploxeralf, fine-loamy, mixed, thermic.

Vegetation: Used for grazing. Vegetation is short annual grasses and associated herbaceous plants. Climate: Mean annual precipitation is about 19 inches; average annual temperature is about 62°F., January 45°F., July 82°F. Parent Material: Gravelly terrace alluvium. Topography: Gently undulating older terraces. On a slope of about 2 percent. Site has a distinct "hogwallow" microrelief. Elevation: 350 feet. Drainage: Well drained, very slow permeability, slow runoff. Soil Moisture: Moist.

Remarks:
This soil has some of the characteristics of the Redding soils, as indicated by some cementation below layer 3. Deep road cuts near the pit show that the material immediately below the claypan is in various stages of cementation. and within a distance of a few feet will range from indurated to no cementation. The material at lower depths is loose and gravelly.

HORIZON	DESCRIPTION
A11	0 to 7 inches, yellowish red (5YR 5/6) gravelly loam, red (2.5YR 4/6) moist; weak granular structure; hard, friable, slightly plastic; abundant small roots; strongly acid (pH 5.5); clear smooth boundary.
A12	7 to 13.5 inches, yellowish red (5YR 5/5) dry and moist, gravelly loam; weak granular structure; hard, friable, slightly plastic; few small roots; medium acid (pH 5.9); abrupt wavy boundary.
B21t	13.5 to 20 inches, yellowish red (5YR 5/5) gravelly clay loam, yellowish red (5YR 5/5) moist; weak blocky structure; compact; very hard, firm, plastic; very few roots; neutral (pH 6.6); clear boundary.
B22t	20 to 26 inches, yellowish red gravelly clay loam containing a slight amount of gravel; color dry and moist similar to layer above; coarse blocky structure; compact, weakly cemented; very hard, firm, plastic; very few roots; mildly alkaline (pH 7.5); gradual boundary.
B23t	26 to 32 inches, red (2.5YR 4/6) gravelly clay loam, red (10R 4/6) moist; strong coarse blocky structure; compact, weakly cemented; very hard, firm, plastic; very few roots; mildly alkaline (pH 7.5); wavy gradual boundary.
B24t	32 to 41 inches, yellowish red (5YR 4/6) gravelly clay loam (near gravelly sandy clay), red (2.5YR 4/6) moist; reticulate mottling; massive; very compact; extremely hard, very firm, plastic; very few roots; mildly alkaline (pH 7.5); clear boundary.
C	41 to 50 inches, yellowish red, sandy clay loam (near sandy clay) containing a slight amount of gravel. Color dry and moist, consistence, and structure similar to layer above but slightly less compact. Reticulate mottling more pronounced than above; weathered gravel also cause yellowish streaking; very few roots; neutral (pH 7.3).

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Montpellier fine sandy loam **LOCATION** Riverside County, California

SOIL FAMILY: Typic Haploxeralf, fine-loamy, mixed, thermic

SOIL NOS. S55Calif-33-14 **LAB. NOS.** 55593-597

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 >2 1/	TEXTURAL CLASS
0-10	A1	4.1	8.4	6.7	17.3	20.7	34.2	8.6	53.8	12.5	5	sl
10-24	B2	2.4	4.7	4.3	12.4	14.7	30.2	31.3	40.4	12.8	0	cl
24-38	B2ca	2.0	4.4	4.4	15.3	19.1	31.7	23.1	48.0	13.4	1	l
38-60	Cca	5.4	9.4	7.4	17.8	18.4	27.7	13.9	47.3	10.4	1	sl
60+	C	28.0	22.6	8.3	10.8	6.7	14.0	9.6	18.8	7.8	67	cosl

DEPTH INCHES	pH 8C1b SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B1a 1/10 ATMOS. %	4B1a 1/3 ATMOS. %	4B2 15 ATMOS. %
0-10	5.4		6.2	0.40	0.041	9.8		0.9			21.9	11.8	3.7
10-24	7.0		8.1	0.24	0.032			1.0					12.0
24-38	7.6		8.7	0.07				3.6	tr.				9.7
38-60	7.3		7.9	0.02				6.5	tr.				7.1
60+	6.6		7.2	0.01				7.2					4.7

DEPTH INCHES	5A2a CATION	EXTRACTABLE CATIONS 5B 5B1a	5B1a	6H2a	Exch.	Exch.	BASE SAT.	SATURATION EXTRACT SOLUBLE 8A 6P1a	6Q1a	6N1a	6K1a	8A MOISTURE

MONTPELLIER FINE SANDY LOAM
S55Calif-33-14

Collectors: Fred Eggers, Arnold Knecht, Floyd G. Stephens, W. G. Harper, Reuben E. Nelson and F. F. Harradine
Date of Sampling: December 8, 1955.
Description by: W. G. Harper.

Location: On Vail Co. Ranch - 1000' E of N 1/4 corner, Sec. 23, T7S, R2W, about 7 miles NE of Temecula, Riverside County, California.

Elevation: About 1,425 feet.

Precipitation: About 12 inches

Classification: Typic Haploxeralf, fine-loamy, mixed, thermic.

California Group: IV

Vegetation: Fallow, following dry-farmed small grain, probably barley.

Parent Material: Old alluvium; Pleistocene non-marine deposits; chiefly derived from granitic materials.

Physiography: Terrace, nearly level top but severely dissected.

Drainage: Surface drainage is good. Internal drainage is very slow.

HORIZON		DESCRIPTION
A1	0-10"	Brown (10YR 5/3 dry, dark brown 4/3 moist) fine sandy loam; moderately defined medium granular; non-sticky when wet, very friable, slightly hard; moderately permeable; many plant roots; pH on dilution 6.0.
B2	10-24"	Reddish-brown (2.5YR 4/4 dry, dark reddish-brown 3/4 moist) fine sandy clay; strongly defined medium prismatic; sticky and plastic, very firm, extremely hard; very slowly permeable; fair amount of plant roots; thin grayish vesicular layer on tops of prisms; pH on dilution 6.5.
B_{2ca}	24-38"	Brown (7.5YR 5/4 dry, dark brown 7.5YR 4/4 moist) fine sandy clay; strongly defined, medium prismatic; sticky and plastic, very firm, extremely hard; few roots; strongly calcareous; very slowly permeable; pH 8.0.
C_{ca}	38-60"	Pale-brown (10YR 6/3 dry, dark grayish-brown 4/2 moist) sandy loam; massive; slightly sticky, firm, hard; strongly calcareous; weakly cemented; pH 7.5; flakes of manganese present.
C	60"+	Brown (10YR 5/3 dry, dark brown 10YR 4/3 moist) gravelly sandy loam; massive; slightly sticky, firm, hard; pH 7.0.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Montpellier fine sandy loam (taxajunct) **LOCATION** Riverside County, California

SOIL FAMILY: Typic Haploxeralf, fine-loamy, mixed, thermic.

SOIL NOS. S55Calif-33-15 **LAB. NOS.** 55598-603

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-8	A11	2.8	5.6	4.6	19.1	26.4	35.7	5.8	63.2	12.8	2	vfsl
8-13	A12	2.2	5.1	4.4	17.9	25.8	36.6	8.0	62.0	13.4	2	vfsl
13-18	A2	2.8	5.0	4.3	15.9	22.9	38.3	10.8	57.2	15.4	2	l
18-31	B2	1.3	3.7	3.1	12.0	16.7	30.3	32.9	43.3	12.5	3	cl
31-39	B2ca	2.2	3.4	2.5	11.9	19.7	30.7	29.6	46.7	12.7	2	cl
39-57	Cca	4.1	6.1	4.4	15.9	21.0	26.5	22.0	48.5	10.2	4	scl

pH 8C1b SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B1a 1/10 ATMOS. %	4B1a 1/3 ATMOS. %	4B2 15 ATMOS. %
5.4		5.9	0.38	0.038	10.0		0.20			20.4	11.5	3.2
5.1		5.7	0.27	0.034	8.0		0.31			21.8	12.8	3.9
6.1		6.8	0.14				0.29			22.1	14.7	4.8
7.2		8.2	0.27				0.85					12.5
8.0		9.0	0.13				2.0	2				12.1
8.0		8.9	0.08				3.4	Tr.				9.9

5A2a CATION EXCHANGE CAPACITY ←(Na)	EXTRACTABLE CATIONS 5B 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg		6K1a	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
7.0	2.3	0.9	2.7	0.2	0.2	57	1.0	0.1	0.6			23.2
7.9	3.0	3.9	3.0	0.2	0.2	71	1.3	0.1	1.0			23.3

MONTPELLIER FINE SANDY LOAM (taxadjunct) 1/
S55Calif-33-15

Location: On right-of-way west side of U. S. Highway 395, 1000 north of junction with State Highway 71, near northwest corner, Sec. 13, T. 8 S., R. 3 W, about 2 miles north of Temecula, Riverside County, California.
Date of Sampling: December 9, 1955.
Description by: W. G. Harper. Collectors: Fred Eggers, Arnold Knecht, Floyd G. Stephens, W. G. Harper, and Reuben E. Nelson.
Classification: Typic Haploxeralf, fine-loamy, mixed, thermic.
Vegetation: Alfileria and annual grasses. This site was probably dry-farmed to grain prior to the construction of the new highway (about 1945). Precipitation: About 12 inches. Parent Material: Old alluvium. Pleistocene non-marine deposits; chiefly derived from granitic materials. Physiography: Terrace with nearly level top but severly dissected. Elevation: About 1100 feet.
Drainage: Surface drainage is good. Internal drainage is very slow.

HORIZON		DESCRIPTION
A11	0- 8"	Brown (10YR 5/3 dry, dark-brown 4/3 moist) fine sandy loam; weakly defined medium granular; slightly sticky, very friable, loose; moderately permeable; many plant roots; pH 6.0; clear smooth boundary.
A12	8-13"	Brown (10YR 5/3 dry, dark-brown 4/3 moist) fine sandy loam; weakly defined medium subangular blocky; slightly sticky, friable, soft; moderately permeable; many plant roots; pH 6.0; clear smooth boundary.
A2	13-18"	Pale-brown (10YR 6/3 dry, dark-brown 4/3 moist) fine sandy loam; weakly defined medium subangular blocky; somewhat vesicular; slightly sticky, friable, soft; moderately permeable; many plant roots; pH 6.2; abrupt smooth boundary.
B2	18-31"	Dark-brown (7.5YR 4/4 dry, dark-brown 3/2 moist) clay; strongly defined coarse columnar with rounded caps on columns; plastic, extremely firm, extremely hard; very slowly permeable; few plant roots; pH 7.5; clear smooth boundary.
B-2ca	31-39"	Brown (7.5YR 5/4 dry, dark-brown 3/2 moist) clay; strongly defined coarse columnar; plastic, extremely firm, extremely hard; very slowly permeable; very few plant roots; strongly calcareous with both disseminated and segregated lime; pH 8.0; clear smooth boundary.
Cca	39-57"	Brown (10YR 5/3 dry, dark reddish-brown 5YR 3/2 moist) sandy clay; massive; plastic, very firm, hard; very slowly permeable; strongly calcareous; pH 8.0.
D	57"+	A gravelly, cobbly layer of mixed rock material.

1/ The color of the argillic horizon is outside the limits recognized for the Montpellier series.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Ramona loamy coarse sand **LOCATION** Riverside County, California

SOIL FAMILY: Typic Haploxeralf, fine-loamy, mixed, thermic

SOIL NOS. S57Calif-33-8 **LAB. NOS.** 57507 - 57512

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 9	Ap	13.4	23.6	14.0	22.0	10.0	12.1	4.9	30.4	3.7	5	lcos
9-20	A1	11.4	21.4	13.6	22.4	11.2	14.3	5.7	34.2	3.7	5	lcos
20-28	B1	5.9	18.2	14.5	22.5	11.7	19.0	8.2	36.7	6.3	3	sl
28-36	B21	6.4	11.8	7.5	12.3	9.9	29.4	22.7	32.7	13.7	2	l
36-48	B22	5.9	10.9	8.2	14.7	10.6	29.2	20.5	35.6	12.7	2	l
48-60+	C1	3.7	9.4	7.8	17.9	12.8	30.7	17.7	42.3	12.0	2	l

pH 8C1b Saturated Paste 1:1	pH 8C1a 1:10	4A3a Bulk Density g/cc	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	6I1a CO_3 meq/l	8A1a Elec. Conduc. mmhos/cm. at 25° C.	6E1b $CaCO_3$ equivalent %	6J1a HCO_3 meq/l	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
6.2	7.6	1.71	0.15	0.019		0.0	1.3	0	1.5	12.8	5.9	1.9
6.1	6.9	1.79	0.11	0.016		0.0	1.7	0	1.4	14.2	7.0	2.2
6.1	6.8	1.81	0.08			0.0	2.2	0	1.1	15.5	8.6	2.7
6.5	7.0	1.82	0.09			0.0	1.1	0	1.7			7.8
6.7	7.0	1.92	0.08			0.0	0.9	0	2.0			7.7
7.6	8.2	1.84	0.01			0.0	1.0	Tr	3.2			6.4

5A2 EXTRACTABLE CATIONS 5B BASE SATURATION EXTRACT SOLUBLE 8A1 8A

RAMONA LOAMY COARSE SAND
S57Calif-33-8

Location: 500 feet south and 30 feet west of east 1/4 corner of Sec. 9, T. 4 S., R. 3 W. On Floyd Bonge Ranch, about 4 miles northeast of Perris, Riverside County, California. Date of Sampling: June 4, 1957.
Description by: Fred Eggers. Collectors: George Borst, R. E. Nelson, Cliff Henry, and Arnold Knecht.

[illegible]

inches. Parent Material: Granitic alluvium. Physiography: Alluvial fan. Elevation: 1,460 feet. Drainage: Well drained.
Remarks: Few roots in the Ap horizon and none below it. Occasional small pores throughout the profile. Weak discontinuous clay skins in the B1 horizon, moderately prominent clay skins in the B21 and B22 horizons and discontinuous moderately prominent clay skins in the C1 horizon. Potatoes in this field have been heavily fertilized with NH_4SO_4. This is the first irrigated crop after dry farmed grain.

HORIZON	DESCRIPTION
S57Calif-33-8-1	**Brown (10YR 5/3 dry, dark brown 3/3 moist) loamy coarse**

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haploxeralf, fine-loamy, mixed, thermic
SOIL Tehama clay loam (taxadjunct) SOIL Nos. S64Calif-57-18 LOCATION Yolo County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64427 - 64434

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	Coarse fragments 1A2a > 2	2 - 19	19 - 76
		Pct. of < 2 mm														Pct.	Pct. of < 76 mm	
0-10	Ap	35.6	49.5	14.9	2.2	2.6	3.6	12.7	14.5	23.1	26.4	45.7	21.1			1		
10-15	B1	29.3	36.0	34.7	1.7	2.4	3.4	10.4	11.4	17.5	18.5	35.4	17.9			0		
15-29	B21t	25.9	42.4	31.7	0.9	1.6	2.8	9.8	10.8	16.8	25.6	33.7	15.1			0		
29-40	B22t	35.3	35.4	29.3	3.9	3.5	4.4	11.9	11.6	15.7	19.7	33.9	38.8			3		
40-53	B31t	44.2	33.3	22.5	6.7	6.3	6.3	12.5	12.4	16.6	16.7	35.7	31.8			24		
53-63	B32t	50.5	29.2	20.3	5.9	6.8	9.5	18.6	9.7	12.3	16.9	30.6	40.8			5		
63-75	C1	60.0	21.9	18.1	1.2	4.3	13.0	32.7	8.8	8.8	13.1	32.4	51.2			0		
75+	IIC2	73.5	14.1	12.4	1.5	6.6	17.1	41.3	7.0	6.4	7.7	30.2	66.5			0		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	Atterberg limits: Plastic limit Pct. of <0.4 mm	Liquid limit	Plastic index	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLE in./in.	4D1 Extensibility COLE in./in.	pH 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-10	0.52	0.052	10		-	14	16	2			6.1				6.3	6.4
10-15	0.32	0.050	6		-						10.7				6.4	6.9
15-29	0.30	0.049	6		-	19	36	17			14.5				7.1	7.5
29-40	0.17				-						11.3				7.5	7.8
40-53	0.10				-						9.1				8.1	7.7
53-63	0.10				-						8.4				8.2	7.9
63-75	0.06				-	14	20	6			7.5				8.3	8.0
75+	0.06				-						5.7				8.3	8.0

Depth (in.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	5B1a 6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 8A1: 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-10	4.5	4.7	0.2	0.4	9.8	3.2	13.2											0.24
10-15	7.5	9.9	0.2	0.4	18.0	2.8	22.8											0.19
15-29	9.5	15.0	0.4	0.5	25.4		26.9											0.25
29-40	8.1	12.0	0.4	0.4	20.9		22.1											0.26
40-53	6.2	10.0	0.4	0.4	17.0		18.0											0.32
53-63	6.4	8.8	0.6	0.3	16.1		16.2											0.34
63-75	5.8	8.4	0.6	0.3	15.1		15.8											0.40
75+	5.0	5.7	0.6	0.2	11.5		12.4											0.28

Depth (in.)	8A Water at Saturation Pct.	Exch. bases 5B3b Ca (meq./100 g.)	5B3b Mg	5B1b Na	5D2 Exchangeable Na Pct.	5C2 Base Sat. Pct.	Gypsum Pct.	Clay mineralogy (<0.002 mm.) X-ray: Mont.	Verm.	Mica	Kaolin.
0-10	26.8				2	74					
10-15	34.6				1	79					
15-29	45.5				2	94					
29-40	39.8				2	95					
40-53	37.9				2	94					
53-63	36.7				4	99					
63-75	32.0				4	96					
75+	32.7				5	93					

TEHAMA CLAY LOAM (taxadjunct) 1/
S64 Cal-57-18

Location: Yolo County, California. A pit located about .2 miles west of the intersection of Roads 89B and 8 and 50 feet south of Road 8 in the northwest corner of the SE¼ Sec. 27, T. 12N., R.1W. Area located about 2 miles south of Dunnigan. Photo ABB-42T-91. Date of Sampling: September 30, 1964.
Description by: G. J. Staidl. Collectors: R. E. Nelson, B. R. Brasher, G. M. Kennedy, W. F. Andrews, L. A. Bates, G. J. Staidl, C. B. Goudey.
Classification: Typic Haploxeralf, fine-loamy, mixed, thermic.
Vegetation: Cultivated, fallow, but usually barley is grown. Climate: Average annual precipitation is 15.3 inches; mean annual temperature 61°F. Parent Material: Alluvium from sedimentary rocks. Topography: Alluvial fan; plane; less than 1 percent slope. Elevation: 65 feet above sea level. Drainage: Runoff is slow; permeability is moderately slow to slow; drainage class is well drained. Soil Moisture: The profile was moist below 15 inches.
Remarks: (1) Truog soil reaction test used in field pH determination; (2) Paired sample No. 64 Cal. 57-19; (3) Diagnostic horizons-ochric epipedon and argillic horizon with over 35 percent clay; (4) B.P.R. Sample horizon Ap inches 0-10, B.P.R. sample horizon B21t inches 15-29, B.P.R. Sample horizon C1 inches 63-75; (5) The lower portion of the Ap has developed some platy structure; (6) The difference in structure of the B22t as compared to the B21t is probably due to the variation in moisture. The structure in the B22t was not readily observable in place but broke out to moderate coarse angular blocky structure; (7) In the B31 lenses of gravel varying in thickness from 1/4 to 3 inches.

HORIZON	DESCRIPTION
Ap RSL 64427	0 to 10 inches, pale brown (10YR 6/3) loam, brown to dark brown (10YR 4/3) when moist; massive lower portion platy structure; hard, friable, nonsticky and slightly plastic; medium acid (pH 6.0); abrupt smooth boundary.
B1 RSL 64428	10 to 15 inches, brown (10YR 5/3) clay loam, brown to dark brown (10YR 4/3) when moist; moderate coarse prismatic structure; very hard, friable, slightly sticky and plastic; very few very fine roots; many micro and few fine tubular pores; common thin clay films on ped faces and in pores; medium acid (pH 6.0); clear smooth boundary.
B21t RSL 64429	15 to 29 inches, yellowish brown (10YR 5/4) clay, dark yellowish brown (10YR 3/4) when moist; moderate coarse prismatic structure; extremely hard, firm, sticky and very plastic; very few micro roots; many micro tubular pores; many moderately thick clay films on ped faces and in pores; slightly acid (pH 6.5); gradual smooth boundary.
B22t RSL 64430	29 to 40 inches, yellowish brown (10YR 5/4) clay, dark yellowish brown (10YR 3/4) when moist; moderate coarse angular blocky structure; extremely hard, firm, sticky and very plastic; many micro tubular and interstitial pores; many thin clay films on ped faces, in pores and as bridges between sand grains; mildly alkaline (pH 7.5); clear smooth boundary.
B31t RSL 64431	40 to 53 inches, light yellowish brown (10YR 6/4) clay loam, dark yellowish brown (10YR 4/4) when moist; weak medium angular blocky structure; very hard, firm, slightly sticky and plastic; many micro tubular and interstitial pores; many thin clay films

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haploxeralf, fine-loamy, mixed, thermic
SOIL Tehama clay loam (taxadjunct) SOIL Nos. S64Calif-57-19 LOCATION Yolo County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64435 - 64442

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	3A1 Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	Coarse fragments 1A2a > 2 Pct.	2 - 19	19 - 76
		Size class and particle diameter (mm)															Pct. of < 76 mm	
		Pct. of < 2 mm																
0-8	Ap	29.2	52.2	18.6	0.2	0.9	2.5	12.5	13.1	33.7	18.5	54.5	16.1			0		
8-15	A12	28.5	47.3	24.2	0.1	0.6	2.3	12.1	13.4	21.0	26.3	41.6	15.1			0		
15-21	B1	25.4	43.2	31.4	0.1	0.6	2.1	10.8	11.8	19.2	24.0	37.7	13.6			0		
21-32	B21t	27.4	41.2	31.4	0.1	0.5	2.4	12.2	12.2	18.7	22.5	38.7	15.2			0		
32-40	B22t	27.2	43.3	29.5	0.1	0.6	2.7	12.3	11.5	20.2	23.1	39.5	15.7			0		
40-55	B23t	31.7	41.3	27.0	0.7	2.4	5.1	13.7	9.8	17.9	23.4	35.3	21.9			0		
55-67	B3	39.0	34.9	26.1	1.5	4.9	9.2	15.8	7.6	13.5	21.4	28.6	31.4			0		
67-78+	C1	42.7	31.6	25.7	1.6	5.8	10.6	17.0	7.7	10.4	21.2	26.1	35.0			0		

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density: 4A1f 1/3 bar g/cc	Bulk density: 4A1h Oven dry g/cc	Bulk density: g/cc	3B2 Cm	Water content: Pct.	Water content: 4B1c 1/3 bar Pct.	Water content: 4B2 15 bar Pct.	4D1 Extensibility COLE in./in.	4D1 Extensibility COLE in./in.	pH: 8C1b Saturated Paste	pH: 8C1a H_2O 1:1	pH: 8C1a H_2O 1:10
0-8	0.80	0.080	10	1.3	-		1.60	1.65		1.0		17.8	5.9	0.011	0.011		6.2	6.4
8-15	0.40	0.054	7	1.4	-		1.57	1.69		1.0		17.7	8.8	0.025	0.025		6.3	6.6
15-21	0.28	0.049	6	1.8	-		1.59	1.78		1.0		19.2	12.7	0.039	0.039		6.4	6.8
21-32	0.18			1.9	-		1.67	1.89		1.0		19.4	13.7	0.043	0.043		6.6	7.0
32-40	0.15			2.0	-		1.66	1.88		1.0		19.7	12.8	0.042	0.042		6.7	7.0
40-55	0.08			1.8	-		1.69	1.86		1.0		18.2	12.0	0.033	0.033		7.0	7.2
55-67	0.10			2.0	-		1.72	1.87		1.0		16.8	11.4	0.029	0.029		7.0	7.3
67-78+	0.09			2.0	-		1.68	1.84		1.0		16.6	11.5	0.030	0.030		7.1	7.3

Depth (In.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 8A1: 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1a SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-8	6.8	4.0	0.3	0.6	11.7	2.5	14.9											0.29
8-15	8.7	4.9	0.3	0.4	14.3	2.5	18.3											0.12
15-21	12.2	7.8	0.3	0.4	20.7	3.3	24.3											0.10
21-32	13.6	9.4	0.3	0.5	23.8	2.4	29.1											0.13
32-40	12.9	9.2	0.4	0.4	22.9	2.1	26.4											0.13
40-55	12.8	8.7	0.4	0.4	22.3	1.7	25.3											0.13
55-67	11.2	7.5	0.4	0.4	19.5	1.9	23.0											0.23
67-78+	11.6	7.4	0.4	0.4	19.8	1.4	21.7											0.25

Depth (In.)	8A Water at Saturation Pct.	Exch. bases 5B3b Ca	5B3b Mg	5B1b Na	5D2 Exchangeable Na Pct	5C2 Base Sat. Pct.	Gypsum Pct.	Clay mineralogy (<0.002 mm.) 7A1: Mont.	Verm.	Mica	Kaolin.	
		meq./100 g.						X-ray 7A2				
0-8	35.9				2	78						
8-15	35.8				2	78		xxxx	-	-	x	
15-21	42.8				1	85						
21-32	43.5				1	82						
32-40	41.6				2	87		xxxx	-	-	x	
40-55	40.6				2	88						
55-67	39.7				2	85						
67-78+	36.4				2	91		xxxx	-	-	x	

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

TEHAMA CLAY LOAM (taxadjunct) 1/
S64Cal-57-19

Location: Yolo County, California. A pit located 100 feet northwest of the southeast corner of Sec. 29, T. 9N., R.1W., 5 miles north of Winters. Photo ABB-2K-40. **Date of Sampling:** September 28, 1964.

Description by: C. B. Goudey. **Collectors:** R. E. Nelson, B. Brasher, G. M. Kennedy, W. F. Andrews, L. A. Bates, G. J. Staidl, C. B. Goudey.

Classification: Typic Haploxeralf, fine-loamy, mixed, thermic.

Vegetation: Barley Stubble. **Climate:** 19.5 inches average annual precipitation and mean annual temperature of 60°F. **Parent Material:** Alluvium from sedimentary rocks. **Topography:** Alluvial fan; plane; one percent slope (east; near crest). **Elevation:** 193 feet. **Drainage:** Runoff-slow; permeability-moderately slow to slow; well drained. **Soil Moisture:** Profile was moist throughout below 15 inches.

Remarks: (1) Truog field pH determinations; (2) Paired profile 64 Cal. 57-18; (3) Diagnostic horizons-ochric epipedon, argillic horizon, with over 30 percent clay; (4) 25.5° C soil temperature at 72 inches.

HORIZON	DESCRIPTION
Ap RSL 64435	0 to 8 inches, pale brown (10YR 6/3) loam, brown to dark brown (10YR 4/3) when moist; massive; hard, friable, slightly sticky and plastic; plentiful micro to very fine, few fine and few medium vertical roots; common very fine interstitial and common fine tubular pores; medium acid (pH 6.0); clear smooth boundary.
A12 RSL 64436	8 to 15 inches, pale brown (10YR 6/3) loam, brown to dark brown (10YR 4/3) when moist; massive; hard, friable, slightly sticky and plastic; plentiful micro to very fine, few fine random and few medium vertical roots; common very fine to fine tubular pores; few thin clay films in pores; medium acid (pH 6.0); clear smooth boundary.
B1 RSL 64437	15 to 21 inches, pale brown (10YR 6/3) clay loam, brown to dark brown (10YR, 4/3) when moist; moderate coarse prismatic structure; hard, firm, sticky and plastic; plentiful [illegible]
	fine tubular pores; common clay films on ped faces and pores; slightly acid (pH 6.2); clear smooth boundary.
B21t RSL 64438	21 to 32 inches, light yellowish brown (10YR 6/4) clay, dark yellowish brown (10YR 4/4) when moist; moderate coarse prismatic breaking to strong coarse angular blocky structure; very hard, very firm, sticky and plastic; plentiful micro to very fine horizontal and few fine to medium vertical roots; common micro tubular pores; many moderately thick clay films on ped faces and pores; slightly acid (pH 6.5); gradual smooth boundary.
B22t RSL 64439	32 to 40 inches, light yellowish brown (10YR 6/4) clay loam, dark yellowish brown (10YR 4/4) when moist; moderate coarse to medium angular blocky structure; very hard, very firm, sticky and plastic; few micro to very fine random roots; common micro tubular pores; many moderately thick clay films on ped faces and pores; slightly acid (pH 6.5); gradual smooth boundary.
B23t RSL	40 to 55 inches, light yellowish brown (10YR 6/4) clay loam, dark yellowish brown (10YR 4/4) when moist; moderate coarse to medium angular blocky structure; very hard, very firm,

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Unnamed silt loam **LOCATION** Alameda County, California

SOIL FAMILY: Typic Haploxeralf, fine-silty, mixed, thermic.

SOIL NOS. S59Calif-01-6 **LAB. NOS.** 59234 thru 59237

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE	COARSE	MEDIUM	FINE	VERY FINE				3A1	2A2	TEXTURAL CLASS

UNNAMED SILT LOAM
S59Calif-01-6

Collectors: Dick Huff, Larry Welch, Lothair Grant.
Description by: Larry E. Welch.
Date of Sampling: May 6, 1959.

Location: Alameda County, California, in the SW1/4 of the SE1/4 of the NW 1/4 of Sec. 7, T3S, R1W, about 3 miles east of the center of Hayward.

Elevation: 450 feet.

Relief: Hilly uplands on a 25 percent southwest-facing, convex slope.

Drainage: Well drained, moderate runoff, moderate permeability.

Parent Material: Shale.

Climate: Rainfall about 20 inches.

Vegetation: Annual grasses and forbs.

Classification: Typic Haploxeralf, fine-silty, mixed, thermic.

HORIZON		DESCRIPTION
A_{11}	0- 6"	Olive (5Y 5/3) silt loam, olive brown (2.5Y 4/3) when moist; massive; hard when dry, friable when moist, slightly sticky and slightly plastic when wet; abundant very fine roots; many very fine pores; medium acid, pH 6.0; gradual, smooth lower boundary.
A_{12}	6-13"	Light olive brown (2.5Y 6/3) silt loam, olive brown (2.5Y 4/3) when moist; massive; hard when dry, friable when moist; slightly sticky and slightly plastic when wet; plentiful very fine roots; many very fine and few fine pores; few thin nearly continuous clay films in pores; medium acid, pH 6.0; clear, smooth lower boundary.
B_2	13-20"	Light yellowish brown (2.5Y 6/3) silty clay loam near silt loam, olive brown (2.5Y 4/3) when moist; massive; hard when dry, friable when moist, plastic and sticky when wet; plentiful fine roots; many fine and very fine pores; common thin continuous clay films in pores; medium acid, pH 6.0; clear, smooth lower boundary.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Pardee gravelly loam **LOCATION** Amador County, California

SOIL FAMILY: Lithic Haploxeralf, loamy-skeletal, mixed, thermic

SOIL NOS. 60Calif-3-22 **LAB. NOS.** 6096 - 60101

1B1b **PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)** 3A1

PARDEE GRAVELLY LOAM 1/
S60Calif-3-22

Location: Amador County, California. On Strohm's Ranch, 0.3 mile west and 0.1 mile north of the southeast corner of Sec. 25, T. 5 N., R. 9 E., 1/2 mile east of Jackson Valley-Comanche Road, 3.5 miles northeast of Comanche; 1.6 miles southwest of Buena Vista Peaks.
Date of Sampling: May 3, 1960.
Description by: Grant Kennedy. Collectors: Grant Kennedy, Reuben Nelson, Robert Sketchley, Clifford Henry, John Rogers, Paul Nazar, and Jack McClelland.
Classification: Lithic Haploxeralf, loamy-skeletal, mixed, thermic.
Vegetation: Annuals - soft chess, filaree, ripgut, occasional stipa, scattered oaks. Climate: Mean annual rainfall - 22 inches; average temperature at Ione Station: January - 46° F.; July - 88° F.; mean annual temperature - 61° F. Parent material: Mixed coarse alluvium containing a high percentage of quartz and quartzite gravel and cobbles, overlying consolidated andesitic conglomerate. Topography: Gently sloping terrace in the lower foothills on a relatively smooth slope of 4 percent to the west with slightly hummocky micro-relief. Elevation: 550 feet. Drainage: Well drained; moderate runoff; moderate permeability in the upper profile, moderately slow below.
Remarks: The lower IIB23t horizon is variable in thickness, the clay tonguing into the upper IIC1 horizon along cracks. The upper part of this clayey horizon appears to be degraded and has distinct gray coatings. Most of the IIB23t horizon is very thin (less than 1½ inches) but in one pocket in the pit it is 7 inches thick and the upper 2 inches is columnar. The soil was moist when described.

HORIZON	DESCRIPTION
A11 RSL No. 6096	0-2 inches, brown (7.5YR 4.5/3) gravelly loam, dark brown (7.5YR 3/4) when moist; massive; slightly hard, friable, slightly sticky and slightly plastic; abundant very fine roots; many very fine interstitial and tubular pores; slightly acid (pH 6.1); abrupt smooth boundary.
A12 RSL No. 6097	2-9 inches, brown (7.5YR 5/3.5) gravelly loam with about 15 percent cobbles, dark brown (7.5YR 3/4) when moist; massive; hard, friable, slightly sticky and slightly plastic; abundant very fine roots, many very fine pores; slightly acid (pH 6.3); gradual smooth boundary.
B21t RSL No. 6098	9-14 inches, brown (7.5YR 5/4) cobbly gravelly light clay loam with about 35 percent cobbles, reddish brown (5YR 4/4) when moist; massive; hard, friable, slightly sticky and plastic; plentiful very fine roots; many very fine, common fine, few medium, mainly tubular pores; thin, discontinuous clay films in pores; medium acid (pH 6.0); gradual smooth boundary.
B22t RSL No. 6099	14-17 inches, brown (7.5YR 5.5/4) gravelly clay loam with about 60 percent cobbles; reddish brown (5YR 4/4) when moist; massive; hard, friable, slightly sticky and plastic; plentiful very fine roots; many very fine and fine pores; thin discontinuous clay films in pores; medium acid (pH 5.8); abrupt slightly wavy lower boundary.
IIB23t RSL No. 60100	17-18 inches, brown (7.5YR 5/3) clay with 35 percent cobbles and with flecks of light gray (10YR 7/2) weathered sand, brown (7.5YR 4/2) when moist; massive; very hard, very firm, sticky and very plastic; plentiful very fine roots; common very fine tubular pores; thick clay films nearly fill all voids; strongly acid (pH 5.3); abrupt slightly wavy boundary, 1/4 to 2 inches thick, rarely up to 7 inches thick.
IIC1 RSL No. 60101	18-41 inches +, very pale brown, light brownish gray, brownish yellow, and gray (10YR 7/3, 6/2, 6/6, and 5Y 5/1) consolidated andesitic conglomerate, grayish brown, brown, gray, and dark gray (10YR 5/2, 7.5YR 4/4, and 5Y 4/1 and 5/1) when moist; massive; very hard, extremely firm, essentially no roots or pores; medium acid (pH 6.0).

1/ Note that the IIC horizon is treated as a lithic contact. Parts of the argillic horizon have base saturation of less than 75 percent, but no Lithic Ultic Haploxeralf has been provided for.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Pardee gravelly loam (taxadjunct) **LOCATION** Amad r County, California

SOIL FAMILY: Lithic Haploxeralf, loamy-skeletal, mixed, thermic

SOIL NOS. 60Calif-3-23 **LAB. NOS.** 60102 - 60107

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)							3A1		2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	
0- 2	A11	9.6	14.7	8.6	15.8	10.4	32.2	8.7	34.0	17.9	47	sl-cosl
2- 7	A12	6.6	12.9	7.8	17.3	12.3	34.0	9.1	39.8	17.6	27	fsl
7-13	B1	6.4	12.1	7.4	14.9	14.0	33.1	12.1	38.5	17.5	23	fsl
13-17	B2t	6.2	10.3	6.3	12.4	9.2	27.5	28.1	28.2	16.1	40	cl- scl
17-29	IIC1	14.6	24.6	8.5	6.7	3.5	16.5	25.6	12.3	11.2	73	scl
29-38+	IIC2	Not determined										

pH | ORGANIC MATTER | ELECTRI- | 6C1b | MOISTURE TENSIONS

PARDEE GRAVELLY LOAM (taxadjunct) 1/
S60Calif-3-23

Location: Amador County, California. 0.1 mile east of the W 1/4 corner of Sec. 26, T. 5 N., R. 9 E., 1/4 mile northwest of Grapevine Gulch; 300 feet southeast of Rabbit Creek Road (Jackson Valley to Clements Road), 2.7 miles west-southwest of Buena Vista Peaks.
Date of sampling: May 3, 1960.
Description by: Grant Kennedy. Collectors: Grant Kennedy, Reuben Nelson, Robert Sketchley, Clifford Henry, John Rogers, Paul Nazar, and Jack McClelland.
Classification: Lithic Haploxeralf, loamy-skeletal, mixed, thermic.
Vegetation: Annuals - filaree, soft chess, ripgut. Climate: Mean annual rainfall - 20 inches; average temperatures (Ione Station): January 46° F.; July - 88° F.; mean annual - 61° F.
Parent material: Mixed coarse alluvium containing a high percentage of quartz and quartzite gravel and cobbles, overlying consolidated andesitic conglomerate. Topography: Gently sloping terrace in the lower foothills of the Sierra Nevada on a smooth but slightly hummocky slope of 3 percent which faces to the southwest. Elevation: 300 feet. Drainage: Well drained; moderate runoff; moderately permeable through the solum, moderately slow or slow below.
Remarks: This soil is a moderately developed Noncalcic Brown soil with many properties in common with the Perkins series but has a thinner solum and is more acid than the Perkins where modal. The soil was moist when described.

HORIZON	DESCRIPTION
A11 RSL No. 60102	0-2 inches, brown (7.5YR 4.5/3.5) gravelly loam, dark reddish brown (5YR 3/3) when moist; weak very fine granular structure; slightly hard, friable, nonsticky and slightly plastic; abundant very fine roots; many very fine tubular and interstitial pores; thin (1/4 inch) dark mulch covers surface; medium acid (pH 6.0); clear smooth boundary.
A12 RSL No. 60103	2-7 inches, brown (7.5YR 5/4) gravelly loam with 15 percent gravel and cobbles over 1 inch in diameter, dark reddish brown (5YR 3/4) when moist; massive; slightly hard, friable, nonsticky and slightly plastic; abundant very fine roots; many very fine interstitial and tubular pores; strongly acid (pH 5.5); gradual smooth boundary.
B1 RSL No. 60104	7-13 inches, brown (7.5YR 5/5) gravelly loam with 40 percent gravel and cobbles 1 to 3 inches in diameter (mainly quartzites), dark reddish brown (5YR 3/4) when moist; massive; hard, friable, slightly sticky and slightly plastic; abundant very fine roots; many very fine interstitial and tubular pores, few fine tubular pores; few thin discontinuous clay films in pores; strongly acid (pH 5.5); clear smooth boundary.
B2t RSL No. 60105	13-17 inches, reddish brown (5YR 5/5) gravelly clay loam with 50 percent gravel and cobbles 1 to 4 inches in diameter (quartzite and andesite), dark reddish brown (5YR 3/5) when moist; massive; hard, firm, sticky and plastic; plentiful very fine roots; many very fine interstitial, common very fine tubular pores; common thick clay films in some pores; strongly acid (pH 5.5); abrupt slightly wavy boundary.
IIC1 RSL No. 60106	17-29 inches, grayish brown (10YR 5/2) weathered tuffaceous conglomerate containing 50 percent andesitic gravel and cobbles over 1 inch in diameter with brown (7.5YR 5/4) seams, dark grayish brown and dark reddish brown (10YR 4/2 and 5YR 3/4) when moist; massive; very hard, extremely firm; few roots along weathered fractured planes; clay nearly fills some of larger voids between cobbles; strongly acid (pH 5.5); abrupt irregular boundary.
IIC2 RSL No. 60107	29-38 inches +, light brownish gray (10YR 6/2) consolidated tuffaceous conglomerate, dark grayish brown (10YR 4/2) when moist; massive; very hard; slightly acid (pH 6.5) when crushed.

1/ The weathered material starting at 17 inches is not as hard as typical of the series.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Millerton loam (taxadjunct) **LOCATION** Riverside County, California

SOIL FAMILY: Lithic Haploxeralf, loamy, mixed, thermic

SOIL NOS. S57Calif-33-3 **LAB. NOS.** 57483 - 57487

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 2	A11	3.3	6.9	5.6	13.8	11.3	41.8	17.3	43.0	18.5	35	1
2- 8	A12	1.3	5.5	4.6	12.2	11.2	45.3	19.9	41.9	22.3	13	1
8-13	B21	1.5	4.3	4.3	11.6	11.6	46.0	20.7	42.8	22.5	13	1
13-19	B22	2.4	5.4	4.0	10.0	9.8	46.2	22.2	45.8	16.9	24	1
19-25+	C1		Parent Material									

8C1b pH SATURATED PASTE 1:1	pH 1:5	8C1a pH 1:10	6A1a ORGANIC MATTER ORGANIC CARBON %	6B1a ORGANIC MATTER NITROGEN %	C/N	6I1a CO_3 meq/l	8A1a Elec. Conduc. mmhos/cm. at 25° C.	6E1b $CaCO_3$ equivalent %	6J1a HCO_3 meq/l	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.7		7.2	1.84	0.152	12.1	0.0	0.9	0	8.0			6.6
6.8		6.7	0.28	0.032	8.8	0.0	0.4	0	2.3			6.8
6.5		7.4	0.22	0.032		0.0	0.4	0	1.8			7.1
6.7		7.1	0.22			0.0	0.4	0	1.9			8.1
		Parent Material										

5A2a CATION EXCHANGE CAPACITY	EXTRACTABLE CATIONS 5B 5B1a	5B1a	6H2a	Exch. 5B1b	Exch. 5B1b	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a	6Q1a	6N1a	6O1a	6K1a	8A MOISTURE AT SATU-

MILLERTON LOAM (taxadjunct) 1/
S57Calif-33-3

Sampled by: George Borst, R. E. Nelson, Clifford Henry, Arnold Knecht and Fred Eggers. Date: June 4, 1957.

Description by: Fred Eggers.

Location: 400' E and 750' N of the center of Sec. 13, T5S, R4W, about 4 miles SW of Perris, Riverside County, California.

Elevation: 1430'

Precipitation: 10"

Classification: Lithic Haploxeralf, loamy, mixed, thermic.

Vegetation: California sage, wild buckwheat, wild oats, cheat grass, some salt grass, annual grasses and weeds.

Parent Material: Bedford Canyon formation. Metamorphosed fine grain sandstone.

Physiography: Low rolling hills.

Drainage: Well drained.

HORIZON	DESCRIPTION
S57Calif-33-3-1 A11 0-2"	Brown (7.5 YR 5/4 dry, dark brown 3/2 moist) loam; moderately defined medium platy breaking to weakly defined fine crumb; slightly hard when dry, friable when moist; slightly sticky and slightly plastic when wet; moderately permeable; many plant roots; pH 6.5; noncalcareous; few pores; clearsmooth lower boundary.
S57Calif-33-3-2 A12 2-8"	Dark brown (7.5YR 4/4 dry, dark brown 3/2 moist) loam; moderately defined medium angular blocky; hard when dry, friable when moist; slightly sticky and plastic when wet; moderately permeable; many plant roots; pH 6.5; noncalcareous; many fine pores; diffuse wavy lower boundary.
S57Calif-33-3-3 B21 8-13"	Dark brown (7.5 YR 4/4 dry, dark brown 3/2 moist) loam; moderately defined fine angular blocky; hard when dry, friable when moist; slightly sticky and plastic when wet; moderately permeable; many plant roots; pH 6.8; many fine pores; diffuse wavy lower boundary.
S57Calif-33-3-4 B22 13-19"	Dark brown (7.5 YR 4/4 dry, dark yellowish-brown 10YR 3/4 moist) loam; strongly defined fine subangular blocky; hard when dry, friable when moist; slightly sticky and plastic when wet; moderately permeable; many plant roots; pH 6.8; noncalcareous; many fine pores, diffuse irregular lower boundary.
S57Calif-33-3-5 C1 19-25"+	Decomposing; strongly metamorphosed fine grain sandstone. There were a few roots extending down the crevices.

Remarks: Weak discontinuous clay skins throughout B horizons.

1/ The Millerton series has a hue of 5 YR in the argillic horizon. Evidence of an argillic horizon is minimal in this pedon.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Millerton gravelly loam (taxadjunct) **LOCATION** Riverside County, California

SOIL FAMILY: Lithic Haploxeralf, loamy, mixed, thermic

SOIL NOS. S57Calif-33-5 **LAB. NOS.** 57491 - 57494

DEPTH INCHES	HORIZON	1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
		PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)										
0- 5	A1	1.3	6.1	8.2	18.9	13.2	44.2	8.1	47.1	22.0	18	1
5-12	B2	0.6	5.9	8.9	18.1	11.3	40.8	14.4	41.6	21.4	19	1
12-18	C	0.4	5.7	12.3	23.3	10.7	36.2	11.4	41.3	18.5	16	fsl
18-23+	Dr	Parent Material										

8C1b SATURATED PASTE 1:1	pH 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	6I1a CO_3 meq/l	8A1a Elec. Conduc. mmhos/cm. at 25°C.	6E1b $CaCO_3$ equivalent %	6J1a HCO_3 meq/l	MOISTURE TENSIONS 4B1a 1/10 ATMOS. %	4B1a 1/3 ATMOS. %	4B2 15 ATMOS. %
6.6		6.4	0.40	0.041	9.8	0.0	0.5	0	3.1	21.2	13.8	3.8
6.5		6.5	0.29	0.029	10.0	0.0	0.4	0	2.4			5.4
6.8		6.5	0.13			0.0	0.4	0	2.2			5.4
		Parent Material										

5A2a	EXTRACTABLE CATIONS 5B	BASE SAT.	SATURATION EXTRACT SOLUBLE 8A1	8A

MILLERTON GRAVELLY LOAM (taxadjunct) 1/
S57Calif-33-5

Location: Riverside County, California. Near the northwest corner of the SE 1/4 of Sec. 14, T. 5 S., R. 4 W., about 5 miles southwest of Perris. Date of sampling: June 18, 1957.
Description by: Fred Eggers. Collectors: George Borst, R. E. Nelson, Clifford Henry, Arnold Knecht and Fred Eggers.
Classification: Lithic Haploxeralf, loamy, mixed, thermic.
Vegetation: California sage, wild buckwheat, dove weed, alfilaree, cactus and annual grasses. Precipitation: 10 inches. Parent material: Bedford Canyon formation. Metamorphic fine grain sandstone. Topography: Low rolling hills. The site is on a convex slope of a gently rolling ridge. Elevation: 1480 feet. Drainage: Well drained.
Remarks: There are many stone fragments throughout the soil and there are rock outcrops. The A1 horizon contains 5 percent rocks. There are weak discontinuous clay skins in the B2 horizon.

HORIZON	DESCRIPTION
A1 RSL No. 57491	0-5 inches, grayish brown to brown (10YR 5/2-5/3 dry, very dark grayish brown to dark brown 3/2-3/3 moist) gravelly loam; weakly defined very fine granular; slightly hard when dry, very friable when moist; nonsticky and slightly plastic when wet; moderately permeable; many plant roots; pH 6.5; noncalcareous; many fine pores; clear wavy lower boundary.
B2 RSL No. 57492	5-12 inches, brown to yellowish brown (10YR 5/3-5/4 dry, dark brown to dark yellowish brown 3/3-3/4 moist) gravelly loam; weakly defined fine subangular blocky; breaking to weakly defined fine granular; hard when dry, friable when moist; slightly sticky and slightly plastic when wet; moderately permeable; many plant roots; pH 6.5; noncalcareous; many fine pores; clear irregular lower boundary.
C RSL No. 57493	12-18 inches, pale brown (10YR 6/3 dry, dark brown 4/3 moist) gravelly loam; pH 6.8; diffuse irregular lower boundary. This horizon contains 75 percent rock fragmentations. There are occasional clay skins between the rock fragmentations and in crevices. There are also a few roots present.
Dr RSL No. 57494	18-23 inches +, bedrock consisting of strongly metamorphosed fine grain sandstone. Soil only in rock crevices; pH 6.8.

1/ The Millerton series has a hue of 5 YR in the argillic horizon.

SOIL SURVEY LABORATORY Riverside, California

LIVERMORE VERY GRAVELLY SANDY LOAM (taxadjunct) 1/
S59Calif-01-3

Location: Alameda County, California. NW1/4 of the NE1/4 of the SW1/4 of Sec. 23, T. 3 S., R. 2 E., about 3 miles southeast of the center of Livermore.
Date of Sampling: May 5, 1959.
Description by: Larry E. Welch. Collectors: R. C. Huff, L. E. Welch, and L. H. Grant.
Classification: Mollic Haploxeralf, loamy-skeletal, mixed, thermic.
Vegetation: Cultivated, vineyard. Climate: Mean annual rainfall about 15 inches. Parent Material: Alluvium from sedimentary rocks. Relief: Nearly level, convex slope to the northwest of less than 1 percent, on a low terrace. Elevation: 620 feet. Drainage: Well drained; slow runoff; moderate permeability. Soil Moisture: Moist throughout.
Remarks: Few worm holes. Few stones and cobbles, mainly gravelly.

HORIZON		DESCRIPTION
A_{11p}	0-4"	Brown (10YR 5/3) very gravelly sandy loam, dark brown (10YR 3/3) when moist; weak very fine granular structure; loose when dry, friable when moist; slightly acid, pH 6.2; diffuse, smooth lower boundary.
A_{12p}	4-12"	Dark grayish brown (10YR 4/2) very gravelly sandy loam, very dark grayish brown (10YR 3/2) when moist; porous massive; slightly hard when dry, friable when moist, slightly sticky when wet; abundant fine and very fine roots; many fine and very fine pores; slightly acid, pH 6.5; diffuse, smooth lower boundary.
A_{13}	12-21"	Dark grayish brown (10YR 4/2) very gravelly loam, very dark grayish brown (10YR 3/2) when moist; weak fine granular structure; slightly hard when dry, friable when moist, slightly sticky; abundant fine and very fine roots; many fine and very fine pores; slightly acid, pH 6.5; diffuse, smooth lower boundary
B_1	21-28"	Brown (10YR 5/3) heavy very gravelly loam, dark brown (10YR 3/3) when moist; massive; slightly hard when dry, friable when moist, slightly sticky and slightly plastic when wet; few fine roots; many fine and very fine pores, mainly interstitial; few thin discontinuous clay films in pores; slightly acid, pH 6.5; diffuse, smooth lower boundary.
B_2	28-34"	Brown (10YR 5/3) stony, sandy clay loam, dark brown (10YR 4/3) when moist; massive; slightly hard when dry, friable when moist, slightly sticky and slightly plastic when wet; few fine roots; many fine and very fine interstitial pores; very thin, nearly continuous clay films in pores, Most of colloid in bridges; slightly acid, pH 6.5; diffuse, smooth lower boundary.
B_3	34-60"+	Brown (10YR 5/3) gravelly loamy coarse sand, dark brown (10YR 4/3) when moist; massive; slightly hard when dry, friable when moist, nonsticky; few fine roots; many fine and very fine pores; a small amount of colloid occurs in bridges between sand grains and stains the sand and gravel; slightly acid, pH 6.5.

1/ The Livermore series is in a loamy-skeletal, mixed, thermic family of Typic Haploxerolls. This pedon has an argillic horizon and an ochric epipedon.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Steinbeck fine sandy loam (taxadjunct) **LOCATION** Sonoma County, California

SOIL FAMILY: Mollic Haploxeralf, coarse-loamy, mixed, mesic

SOIL NOS. S61Calif-49-3 **LAB. NOS.** 61398 - 61406

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	3A1 0.02-0.002	2A2 > 2	TEXTURAL CLASS
0- 8	Ap	0.4	0.8	0.6	14.5	40.4	26.8	16.5	70.1	10.8	0	vfsl
8-20	A1	0.4	0.7	0.5	14.7	41.3	26.9	15.5	72.7	10.7	0	vfsl
20-34	A21	0.3	0.8	0.7	16.8	46.2	28.7	6.5	80.8	11.2	0	vfsl
34-39	A22	0.4	0.7	0.6	19.2	48.7	26.4	4.0	84.4	9.4	0	vfsl
39-48	B21t	0.2	0.6	0.5	18.5	40.2	22.9	17.1	71.9	9.3	0	vfsl
48-53	B22t	0.1	0.4	0.3	16.3	44.0	22.8	16.1	73.9	8.9	0	vfsl
53-64	B23t	0.4	1.5	0.8	20.1	41.9	20.2	15.1	73.2	8.5	0	vfsl
64-71	C1	0.7	1.7	0.9	22.4	43.3	20.1	10.9	76.5	8.7	1	vfsl
71+	A/C2	0.0	0.2	0.4	19.8	44.1	24.8	10.7	75.6	12.1	0	vfsl

pH 8C1a H_2O 1:1	pH 8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	Bulk Density 4A3a g/cc		Ext. Iron as % Fe 6C1b				4B2 Soil water 15 atmos. Pct.
5.7	4.3		2.11	0.180	12	1.31		0.6				8.0
5.7	4.1		0.98	0.105	9	1.5		0.7				7.3

STEINBECK FINE SANDY LOAM(taxadjunct) 1/
S61Calif-49-3

Location: Sonoma County, California, Northwest corner of SE1/4, NW1/4, section 22, T. 6 N., R. 10 W. (Peruchi Ranch). 41° and 1800 feet from junction of Joy Woods Road and Bodega Highway. Date of Sampling: June 21, 1961.
Collectors: Reuben Nelson, Leo C. Klameth, V. C. Miller, G. Lambert, J. H. Kashiwagi, F. Graham.
Classification: Mollic Haploxeralf, coarse-loamy, mixed, mesic.

Vegetation: Annual and perennial grasses. Use: Improved dry land pasture. Climate: About 24.7 inches annual rainfall, average temperature 57.5° F., average January temperature 46.7° F., average July temperature 66.3° F. Parent Material: Merced formation - soft sedimentary marine formation. Elevation: 18 feet. Relief: Rolling hills on an easterly facing slope of 12

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Ahwahnee fine sand (taxadjunct) **LOCATION** San Luis Obispo Co., California

SOIL FAMILY: Mollic Haploxeralf, coarse-loamy, mixed, thermic.

SOIL NOS. S55Ca1. 40-1 **LAB. NOS.** 55449-454

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0-4	A_{11}	0.3	1.2	14.6	72.1	2.0	4.2	5.6	28.9	2.3	0	fs
4-12	A_{12}	0.2	0.6	12.6	73.8	2.2	4.6	6.0	30.4	2.5	1	fs
12-18	A_3	0.0	0.5	10.2	75.1	2.3	5.7	6.2	33.0	3.4	1	fs
18-30	B_{21}	0.2	0.6	9.3	66.7	2.8	8.1	12.3	31.8	5.1	2	fsl
30-44	B_{22}	0.1	0.8	10.8	65.8	2.3	7.1	13.1	29.5	4.9	1	fsl
44-60	C	0.2	0.6	10.4	64.7	2.1	6.2	15.8	28.3	3.7	1	fsl

pH 8C1b SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
6.6		7.0	1.18	0.092	12.8						6.8	3.7
6.5		7.0	0.52	0.041	12.7						5.5	3.3
6.5		6.9	0.29								6.1	3.5
6.3		7.0	0.22									6.0
6.4		7.1	0.16									6.4
6.4		7.3	0.14									6.8

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	5B1a Na 6P2a	5B1a K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE Na	K				MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
9.2	4.3	2.0	3.1	0.2	1.2	71						
8.1	4.3	1.7	2.0	0.3	1.2	79						
8.0	4.4	2.8	1.6	0.2	0.3	83						
14.0	6.0	6.3	1.9	0.3	0.1	87						
14.2	5.1	6.2	1.9	0.2	0.1	86						
15.1	4.8	7.3	2.1	0.3	0.1	86						

AHWAHNEE FINE SAND (taxadjunct) 1/
S55Calif-40-1

Location: 7/8 mile N.N.E. of farm buildings on the George Mello farm, San Luis Obispo County, California.

Collectors: W. M. Johnson, Rudolph Ulrich, Lewis Leifer, W. G. Harper, L. R. Wohletz, J. M. Williams. Samples taken September 7, 1955. Description by W. M. Johnson.

Elevation: About 110 feet.

Vegetation: Soil was cultivated at one time. Now is pasture in poor condition. Annual fescues, wild oats, ripgut, soft chess, California croton, tarweed, deer vetch, Astragalus sp., and other weeds; scattered live oak; California sage.

Relief: Hilly and steep eroded upland. Residual. Slope about 20 percent, slightly convex, toward the south.

Drainage: Somewhat excessively drained. Both runoff and permeability are rapid.

Parent material: Weathered marine sandstone, not identified as to age.
Classification: Mollic Haploxeralf, coarse-loamy, mixed, thermic.

HORIZON		DESCRIPTION
A_{11}	0-4"	Very dark grayish brown, moist to dark grayish brown, dry (10YR 4/2; 3/2, moist) loamy fine sand. Massive. Slightly hard, very friable, nonplastic, nonsticky. pH 6.5. Clear, smooth lower boundary.
A_{12}	4-12"	Very dark grayish brown (10YR 3/2, moist) loamy fine sand with few fine indistinct mottles (10YR 4/3, dry). Massive. Very hard, very friable, nonplastic, nonsticky. Much worm activity. pH 6.3. Gradual, wavy lower boundary.
A_3	12-18"	Dark brown (10YR 3/3, moist) loamy fine sand. Massive. Slightly hard, very friable, nonplastic, nonsticky. pH 6.3. Clear, wavy lower boundary.
B_{21}	18-30"	Finely mottled dark yellowish brown (10YR 4/4, moist) and dark brown (3/4, moist) fine sandy loam. Weak medium to coarse subangular blocky. Very firm, slightly plastic, nonsticky, numerous distinct patches of clay skins. Slightly porous. pH 6.0. Gradual lower boundary.
B_{22}	30-44"	Dark brown (10YR 3/3, moist) sandy clay loam finely mottled with dark yellowish brown (4/4). Very weak medium subangular blocky. Very firm, plastic, slightly sticky. Compact, hard to dig out. Many patches of clay skins. pH 6.0.
C	44-60+"	Dark brown (10YR 4/3, moist) stratified fine sandy loam and sandy clay loam. Massive. Friable, plastic, slightly sticky. pH 6.0.

1/ The Ahwahnee series is in a coarse-loamy, mixed, thermic family of Mollic Haploxeralfs. This pedon is a taxadjunct to the series because the surface horizon is darker and thicker than 10 inches

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Mollic Haploxeralf, fine-loamy, mixed, mesic

SOIL Hilt loam SOIL Nos. S66Calif-47-6 LOCATION Siskiyou County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66376 - 66383

Depth (In)	Horizon	1B1b Total Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int III (0.02-0.002)	Int II (0.2-0.02)	(2-0.1)	3A1a Non-carbonate clay	1/ Coarse fragments LA2a > 2	2-19	19-76
		Pct. of < 2 mm													Pct.	Pct. of < 76mm	
0-2	A1	50.4	35.2	14.4	0.9	6.0	9.8	20.5	13.2	15.4	19.8	39.9	37.2	14	-	-	-
2-9	A3	49.2	33.8	17.0	0.7	5.7	9.9	19.5	13.4	15.6	18.2	40.0	35.8	17	-	-	-
9-16	B1t	47.4	31.9	20.7	0.5	5.2	9.8	19.1	13.3	14.1	17.8	38.0	34.1	21	-	-	-
16-23	B21t	46.5	30.7	22.8	0.5	5.2	9.3	18.6	12.9	13.8	16.9	37.2	33.6	23	-	-	-
23-37	B22t	45.4	28.9	25.7	0.4	5.3	9.5	18.4	11.8	13.4	15.5	35.2	33.6	26	-	-	-
37-52	B3t	49.1	28.8	22.1	0.8	6.7	10.2	19.5	11.9	13.9	14.9	36.5	37.2	22	-	-	-
52-60+	C	46.2	29.1	24.7	0.8	6.9	9.4	17.6	11.5	13.9	15.2	34.8	34.7	25	-	-	-
0-9	A 2/														-	-	-

Depth (In.)	6A1a Organic carbon Pct	6B1a Nitrogen Pct	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct	Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 C'	Water content: Pct	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1a H_2O 1:1		
0-2	2.54	0.157	16	1.3		1.59	1.62		1.00		15.1	6.7	0.008	0.008	7.0		
2-9	0.88	0.070	13	1.4		1.61	1.65		1.00		13.8	7.0	0.007	0.007	6.7		
9-16	0.38			1.6		1.65	1.72		1.00		13.9	7.6	0.015	0.015	6.4		
16-23	0.29			1.6		1.66	1.77		1.00		14.5	8.6	0.022	0.022	6.0		
23-37	0.25			1.6		1.69	1.80		1.00		15.8	10.0	0.020	0.020	5.5		
37-52	0.18			1.5		1.71	1.82		1.00		14.8	9.2	0.022	0.022	5.7		
52-60+	0.17			1.5		1.69	1.79		1.00		17.9	10.5	0.020	0.020	5.7		
0-9	1.48	0.107	14	1.4								7.5			6.8		

Depth (In)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1a NH_4OAc	5A3a Sum	6G1d KCl Ext. Al^{+++}	Base saturation: 5C1 NH_4OAc	5C3 Sum
	meq/100 g									Pct.	Pct.
0-2	12.4	2.1	0.1	0.8	15.4	2.6	16.9	18.0		91	86
2-9	8.9	1.9	0.1	0.7	11.6	2.6	13.7	14.2		85	82
9-16	7.3	2.8	0.1	0.7	10.9	3.0	13.7	13.9		80	78
16-23	7.1	3.8	0.2	0.6	11.7	4.0	15.6	15.7	0.1	75	75
23-37	9.8	5.7	0.2	0.4	16.1	5.2	20.6	21.3	0.3	78	76
37-52	9.4	6.0	0.4	0.3	16.1	5.2	20.6	21.3	0.4	78	76
52-60+	11.1	7.2	0.6	0.3	19.2	5.0	23.8	24.2	0.4	81	79
0-9	10.6	1.7	0.1	1.0	13.4	3.0	15.4	16.4		87	82

Depth (In.)	Clay mineralogy (<0.002 mm) 7A: Mont.	Verm.	Illite	Kaolin
	X-ray			
0-2	t		xxx	xx
2-9				
9-16				
16-23				
23-37	xx		x	x
37-52				
52-60+	x	xx		t
0-9				

1/ From characterization sample (5 kg).

2/ Composite of 20 subsamples taken in a 15x12' grid adjacent to the master pedon.

- \- = looked for but not found
- t = trace
- x = small
- xx = moderate
- xxx = abundant
- xxxx = dominant

HILT CLAY LOAM
S66Calif-47-6

Location: Siskiyou County, California. A pit located on the Cedar Grove Ranch, about 3 miles north of the California Quarantine Station on Interstate Highway No. 5. Site is located 400 feet west of Interstate Highway No. 5 where the Southern Pacific Railroad and highway intersect, or 2000 feet east and 200 feet north of the SW corner of Sec. 25, T48N, R7W. Date of Sampling: May 24, 1966.
Description by: J. J. Newlun. Collectors: K. W. Flach, S. B. Johnson and L. Mann.

Classification: Mollic Haploxeralf, fine-loamy, mixed, mesic.

Vegetation: In order of dominance - white oak, tall oatgrass, big bluegrass, annual bromes, and other grasses and forbs, (Ceanothus cuneatus), Juniper, and common manzanita. An occasional ponderosa pine occurs in the surrounding area. The site area has been grazed heavily and is heavily trampled. Climate: Average annual precipitation is approximately 16 inches with some in the form of snow. Average January temperature is 32°F., July temperature 72°F., and mean annual temperature 51°F. This site occurs in the upper limits of the precipitation zone in which this soil series occurs in this area. Parent Material: Residium from the sedimentary rock of the Chico Formation. This soil has developed from the sandstone in this formation. The sandstone is fairly fine textured and weathers to heavy sandy clay loam or silty clay loam in most cases. The sandstone has formed from sediments from mixed rock sources that were dominantly basic. The dominant clay mineral is probably montmorillonitic. Topography: The site is located on hilly upland with a single convex slope, a SE aspect and a 17 percent gradient. Site is near midslope. Elevation: 2880 feet. Drainage: Well drained, moderately slow permeability, medium runoff. Soil Moisture: The A horizons approximately 30 percent of field capacity. B horizons about 70 percent of field capacity.

Remarks: Reactions determined by the Truog soil reaction test. Paired profile is S66Calif-47-8. This profile has an ochric epipedon and an argillic horizon. Epipedon not thick enough for mollic. This soil varies mainly in color and depth. Depths to paralithic contact range from 20 to about 60 inches with average depth about 38 inches. Colors range from the 10YR hue to 5YR hue with most having colors in the 7.5YR hue. The A horizons range in texture from sandy loam to light clay loam. The B horizon textures range from clay loam to heavy silty clay loam. The C horizons where present have the structure and fabric of the parent rock but are friable to slightly firm when moist. Some soils in the area may have thicker A horizons. Most of the cultivated soils are eroded and have thinner A horizons. There may be a few quartzite gravels throughout the profile.

HORIZON	DESCRIPTION
A11 RSL 66376	0 to 2 inches, brown (10YR 5/3) light clay loam, dark brown (10YR 3/3) moist; moderate coarse platy structure; slightly hard, friable, slightly sticky and slightly plastic; common fine and very fine roots; few fine and common medium tubular pores; slightly acid (pH 6.5); abrupt smooth boundary.
A3 RSL 66377	2 to 9 inches, brown (7.5YR 5/4) clay loam, dark brown (7.5YR 3/2) moist; weak medium subangular blocky structure to massive; hard, friable, sticky and plastic; common very fine and fine roots; many fine and few medium tubular pores; few thin clay films in pores; slightly acid (pH 6.2); abrupt smooth boundary.
B1 RSL 66378	9 to 16 inches, brown (7.5YR 5/4) heavy clay loam, dark brown (7.5YR 3/4) moist; moderate medium subangular blocky structure; hard, friable, sticky and plastic; many medium and common fine roots; many fine and common medium tubular pores; few moderately thick clay films in pores and many thin films in pores and on ped faces.
B21t RSL 66379	16 to 23 inches, brown (7.5YR 5/4) light silty clay loam, dark reddish brown (5YR 3/3) moist and rubbed, with reddish brown (5YR 4/4) moist ped faces; weak medium prismatic breaking to moderate medium subangular blocky structure; very hard, friable, sticky and plastic; many medium and coarse and few fine roots; many fine and common medium tubular pores; continuous thin clay films in pores and on ped faces with a few moderately thick films in pores; medium acid (pH 5.8); clear wavy boundary.
B22t RSL	23 to 37 inches, strong brown (7.5YR 5/6) silty clay loam, dark brown (7.5YR 4/4) moist; weak medium prismatic breaking to moderate medium subangular blocky structure;

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Mollic Haploxeralf, fine-loamy, mixed, mesic

SOIL Hilt loam SOIL Nos. S66Calif-47-8 LOCATION Siskiyou County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66384 - 66392

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand Very coarse (2-1)	Sand Coarse (1-0.5)	Sand Medium (0.5-0.25)	Sand Fine (0.25-0.1)	Sand Very fine (0.1-0.05)	Silt 0.05-0.02	Silt Int III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Non-carbonate clay	1/ Coarse fragments 1A2a > 2 Pct.	2-19 Pct. of < 76mm	19-76 Pct. of < 76mm
		Pct. of < 2 mm (3A1)															
0-2	A11	64.2	25.4	10.4	3.8	8.3	7.2	29.8	15.1	12.6	12.8	47.4	49.1	10	8	7	1
2-6	A12	60.7	23.0	16.3	2.5	6.2	6.6	28.9	16.5	10.6	12.4	46.9	44.2	16	4	3	1
6-11	A3	57.4	23.8	13.8	2.4	6.7	6.8	26.6	14.9	11.5	12.3	45.0	42.5	19	1	1	-
11-23	B1t	55.6	22.8	21.6	2.0	5.5	6.2	26.5	15.4	10.7	12.1	45.1	40.2	22	2	2	-
23-31	B21t	52.7	21.7	25.6	2.2	6.2	5.8	24.2	14.3	10.1	11.6	41.6	38.4	26	3	2	1
31-38	B22t	46.5	23.7	29.8	2.5	6.5	6.1	20.5	10.9	12.4	11.3	37.2	35.6	30	5	1	4
38-47	B3t	59.2	18.1	22.7	2.5	12.9	10.6	22.6	10.6	9.6	8.5	34.5	48.6	23	1	1	-
47-49+	R																
0-6	A2/														6	5	1
[redacted]	6A1c	6B1c		6C2a	6E1b		Bulk density			3B2	Water content			4D1	4D1	pH	

HILT CLAY LOAM
S66Calif-47-8

Location: Siskiyou County, California. A pit located on the Flying S Ranch about 6 miles N. of Montague off the Yreka-Ager Road. 2600 feet W. and 1750 feet S. of the NE corner of Sec. 26, T46N, R6W. Date of Sampling: May 24, 1966.
Description by: J. J. Newlun. Collectors: K. W. Flach, S. B. Johnson, D. Lucas.
Classification: Mollic Haploxeralf, fine-loamy, mixed, mesic.
Vegetation: Dominantly annual brome grasses, with scattered patches of ceanothus cuneatus, an occasional Juniper and white oak tree, wild buckwheat and other annual grasses and weeds. The site area is being used as grazing land and is overgrazed and well trampled. Climate: Average annual precipitation is approximately 13 inches with some in the form of snow. Average January temperature is 33°F., average July temperature 72°F., mean annual temperature 52°F. Precipitation occurs mainly during the winter and early spring months. Summers are warm and dry. This site occurs in the lower limits of the precipitation zone in which this soil series occurs in this area. Parent Material: Residium from the sedimentary rock of the Chico Formation. This soil has developed in the sandstone of this formation. The sandstone is fairly fine textured and weathers to heavy sandy clay loam or silty clay loam in most cases. The sandstone has formed from mixed rock sources that were dominantly basic. The dominant clay mineral is probably montmorillonitic. Topography: Site is located in hilly upland with a single convex slope, 17 percent gradient, east aspect and near midslope. There was no significant microrelief. Elevation: 2840 feet. Drainage: Well drained, moderately slow permeability, medium runoff. Soil Moisture: The A horizons were nearly air dry. The B horizons were at about 50 percent of field capacity.
Remarks: Reactions determined by Truog soil reaction test. This soil has an ochric epipedon (<1 percent O.M.) and an argillic horizon. Paired profile is S66Calif-47-6. This soil varies mainly in color and depth. Depth to bedrock ranges from 20 to over 60 inches and averages

in the 7.5YR hue. The A horizons range in texture from sandy loam to light clay loam. The B horizon textures range from clay loam to heavy silty clay loam. The C horizons, when present, have the structure and fabric of the parent rock but are friable to slightly firm when moist. The cultivated soils are usually eroded and have thinner A horizons than uncultivated soils. A few quartzite gravels may occur throughout the profile.

HORIZON	DESCRIPTION
A11 RSL 66384	0 to 2 inches, brown (10YR 5/3) light clay loam, very dark grayish brown (10YR 3/2) moist; weak medium platy structure; slightly hard, friable, slightly sticky and slightly plastic; many fine roots; many fine tubular pores; slightly acid (pH 6.5); abrupt smooth boundary.
A12 RSL 66385	2 to 6 inches, brown (10YR 5/3) light clay loam, very dark grayish brown (10YR 3/2) moist; moderate medium platy structure; slightly hard, friable, sticky and slightly plastic; many fine roots; common fine tubular pores; slightly acid (pH 6.5); abrupt smooth boundary.
A3 RSL 66386	6 to 11 inches, dark brown (7.5YR 4/4) sandy clay loam, dark brown (10YR 3/3) moist; moderate medium platy structure; slightly hard, friable, sticky and plastic; many fine roots; many fine tubular pores; few thin clay films in pores; slightly acid (pH 6.5); clear wavy boundary.
B1t RSL 66387	11 to 23 inches, dark brown (7.5YR 4/4) heavy sandy clay loam, dark brown (10YR 3/3) moist and rubbed with dark brown (7.5YR 3/2) ped faces moist; weak medium prismatic structure; hard, friable, sticky and very plastic; common fine roots; common fine and medium tubular pores; common thin clay films on ped faces and in pores; slightly

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Steinbeck loam LOCATION Sonoma County, California

SOIL FAMILY: Mollic Haploxeralf, fine-loamy, mixed, mesic

SOIL NOS. S61Calif-49-4 LAB. NOS. 61407 - 61413

STEINBECK LOAM
S61Calif-49-4

Location: Sonoma County, California. Near center of NE1/4 of section 34, T. 6 N., R. 10 W. (Coronet Ranch) 390 feet south of Valley Ford cutoff road, 5180 feet west of junction with Freestone - Valley Ford Road. Date of Sampling: June 22, 1961.
Collectors: Reuben Nelson, Leo C. Klameth, V. C. Miller, G. Lambert, J. H. Kashiwagi, F. Graham.
Classification: **Mollic Haploxeralf, fine-loamy, mixed, mesic.**

Vegetation: Annual and perennial grasses. Use: Improved dry land pasture. Climate: Average rainfall about 24.7 inches, average temperature 57.5° F., January temperature 46.7° F., July temperature 66.3° F. Parent Material: Merced formation - soft sedimentary marine formation. Elevation: About 160 feet. Relief: Rolling hills on 9 percent slightly concave west facing slope. Drainage: Moderately well drained, medium runoff, moderately rapid over moderately slow permeability. Biological Activity: Worm casts to 28 inches.

HORIZON	DESCRIPTION
Ap RSL No. 61407	0 to 8 inches, dark gray (10YR 4/1) loam (silty), very dark grayish brown (10YR 3/2) when moist; massive breaking to fine and very fine subangular blocky structure; slightly hard when dry, friable when moist, slightly sticky and slightly plastic when wet; abundant very fine and fine roots; common very fine and fine interstitial and tubular pores; strongly acid, pH 5.5; clear, wavy boundary.
A1 RSL No. 61408	8 to 18 inches, dark gray (10YR 4/1) loam (silty), black (10YR 2/1) when moist; massive breaking to very fine subangular blocky structure; slightly hard when dry, friable when moist, slightly sticky and slightly plastic when wet; abundant very fine and fine roots; common fine and very fine tubular and interstitial pores; strongly acid, pH 5.5; gradual, wavy boundary.
A21 RSL No. 61409	18 to 28 inches, gray (10YR 5/1) fine sandy loam, mixed very dark grayish brown and dark grayish brown (10YR 3/2 and 10YR 4/2) when moist; massive; hard when dry, friable when moist, slightly sticky and slightly plastic when wet; abundant very fine and fine roots; many very fine and fine tubular and interstitial pores; strongly acid, pH 5.5; gradual, wavy boundary.
A22 RSL No. 61410	28 to 35 inches, light gray (10YR 7/2) fine sandy loam (loamy fine sand), brown (10YR 5/3) when moist; massive; slightly hard when dry; friable when moist, nonsticky and nonplastic when wet; abundant very fine and fine roots oriented vertically; many very fine and fine tubular and interstitial pores; few moderately thick clay films in pores; strongly acid, pH 5.5; clear, wavy boundary.
B21t RSL No. 61411	35 to 45 inches, light yellowish brown (10YR 6/4) clay with common medium distinct mottles of yellowish brown (10YR 5/8) when dry, yellowish brown (10YR 5/6) with common medium distinct mottles of strong brown (7.5YR 5/8) when moist; massive breaking to weak very coarse prismatic structure; hard when dry, very firm when moist, sticky and plastic when wet; abundant very fine and fine roots oriented vertically; many very fine tubular pores; continuous, thick clay films in pores and on ped faces; medium acid, pH 5.8; scattered shot size (manganese) concretions; gradual, wavy boundary.
B22t RSL No. 61412	45 to 56 inches, light yellowish brown (10YR 6/4) clay with common medium distinct mottles of yellowish brown (10YR 5/8) when dry, yellowish brown (10YR 5/6) with common medium distinct mottles of reddish yellow (7.5YR 6/8) when moist; massive breaking to weak coarse prismatic structure; hard when dry, very firm when moist, very sticky and very plastic when wet; plentiful fine and very fine roots; many very fine tubular pores; continuous, moderately thick and thick clay films on ped faces and in pores; medium acid, pH 6.0; abrupt, wavy boundary in parts of the pit.
B3 RSL No. 61413	56 to 68 inches +, light brownish gray (2.5Y 7/2) sandy loam with many fine distinct mottles of strong brown (10YR 5/8) when dry; yellowish brown (10YR 5/6) with mottles of light brownish gray and strong brown (2.5Y 6/2 and 7.5YR 5/6) when moist; massive; hard when dry, friable when moist, slightly sticky and nonplastic when wet; few discontinuous tubular pores; continuous, thick clay films; slightly acid, pH 6.5; some clam shell imprints are evident; weakly cemented.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Mollic Haploxeralf, fine-loamy, mixed, thermic

SOIL Fallbrook sandy loam (taxadjunct) SOIL Nos. S64Calif-37-6 LOCATION San Diego County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 6423 - 6429

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	* Clay <.002	** Clay <.002	1/ Coarse fragments 1A2a >2	2 - 19	19 -76
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm														Pct.	Pct. of < 76 mm	
0- 3	Ap	64.8	26.4	8.8	10.2	14.7	10.0	17.8	12.1	13.6	12.8	35.5	52.7	8.5	8.1	5	5	0
3-13	A12	61.6	26.2	12.2	7.9	13.9	9.0	17.9	12.9	12.8	13.4	35.0	48.7	9.8	9.4	4	4	0
13-18	B1	58.5	25.2	16.3	8.6	13.7	8.5	16.6	11.1	12.1	13.1	32.4	47.4	18.4	17.3	6	6	0
18-27	B21t	52.8	22.6	24.6	7.7	11.4	7.5	15.7	10.5	10.6	12.0	30.1	42.3	23.3	22.0	5	5	0
27-41	B22t	56.9	23.7	19.4	8.1	12.1	8.0	17.1	11.6	11.3	12.4	32.0	45.3	19.3	17.6	9	9	0
41-62	C1	73.9	14.7	11.4	20.8	19.1	8.4	17.0	8.6	6.7	8.0	24.3	65.3	11.1	7.9	29	29	0
62-70+	C2	76.6	16.4	7.0	23.4	19.4	8.6	16.4	8.8	7.1	9.3	23.9	67.8	6.7	4.6	31	31	0

Depth (In.)	* 6A1a Organic carbon Pct	6B1a * Nitrogen Pct.	C/N	* 6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct		Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 C'	Water content Pct	4B1c 1/3 bar Pct.	4B2* 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0- 3	2.1	0.178	12	0.6									6.6			7.0	7.2	
3-13	0.83	0.042	20	0.8									5.1			6.9	7.3	
13-18	0.30			1.3									7.8			6.9	7.2	
18-27	0.23			1.2									9.5			6.8	7.0	
27-41	0.16			1.1									8.5			6.8	7.0	
41-62	0.03			1.1									7.3			7.0	7.2	
62-70+	0.01			1.0									5.7			7.2	7.3	

Depth (In.)	Extractable bases 6N2a* Ca	6O2a* Mg	6P2a* Na	5B1a 6Q2a* K	* Sum of bases	6H2a* Ext. Acidity	Cation Exch. Capacity 5A2a* NaOAc	5A3a* Sum	Water extract from saturated paste 8A1 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0- 3	8.8	4.2	0.5	0.6	14.1	2.0	16.2	16.1										
3-13	6.6	3.1	0.7	0.1	10.5	1.5	12.4	12.0										
13-18	10.2	4.0	1.3	0.1	15.6	1.9	18.1	17.5										
18-27	11.0	4.6	1.6	0.1	17.3	2.3	20.9	19.6										
27-41	11.3	5.1	1.8	0.1	18.3	2.0	18.7	20.3										
41-62	16.1	6.1	1.6	0.1	23.9	1.4	25.1	25.3										
62-70+	17.3	4.9	2.0	0.1	24.3	1.0	24.8	25.3										

Depth (In.)	8A Water at Saturation Pct.	5D2 Exchangeable Na Pct.	5B1b Exch. Na meq / 100g	Base Sat. Cations Pct.	Base Sat. NaOAc CEC Pct.	Clay mineralogy (<0.002 mm) 7A Mont.	Verm.	Mica	Kaolin.	
						X-ray				
0- 3		3		88	88					
3-13		6		85	85					
13-18		7		89	86					
18-27		8		88	83					
27-41		9		90	98					
41-62		6		94	95					
62-70+		8		96	98					

1/ From characterization sample; determined by shaking overnight in a sodium hexametaphosphate solution.

* Analysis of ground whole soil; results expressed on <2-mm basis.

** Analysis of ground whole soil; results expressed on whole-soil basis.

FALLBROOK SANDY LOAM (taxadjunct) 1/
S64Calif-37-6

Location: San Diego County, California. A soil profile pit located about 1-1/2 miles south and 1-1/2 miles east of Fallbrook in the SW1/4 of the NW1/4 of section 29, T. 9 S., R. 3 W. Date of Sampling: February 20, 1964.
Description By: Roy H. Bowman, Gerald Kester. Collectors: Klaus Flach, Gerald Kester, George Borst, Lothair Grant and Donald Markewich.
Classification: Mollic Haploxeralf, fine-loamy, mixed, thermic.
Vegetation: Wild oats and annual weeds. Climate: Annual precipitation is 15 inches and mean annual temperature is 60° F. Parent Material: Tonalite grus. Topography: East-facing, smooth 9 percent slopes in middle backslope positions on rolling upland. Elevation: 685 feet above sea level. Drainage: Well drained; medium runoff; moderately slowly permeable. Soil Moisture: Moist to 18 inches.

HORIZON	DESCRIPTION
Ap RSL No. 6423	0 to 3 inches, brown (10YR 5/3) sandy loam, dark brown (10YR 3/3) moist; weak fine crumb structure; slightly hard, very friable, slightly sticky and nonplastic; abundant very fine and fine roots; many very fine and fine interstitial pores; neutral (pH 6.6); abrupt smooth boundary.
A12 RSL No. 6424	3 to 13 inches, brown (10YR 5/3) sandy loam, dark brown (10YR 3/3) moist; massive; hard, friable, slightly sticky and nonplastic; abundant very fine and fine roots; many very fine to medium tubular pores; slightly acid (pH 6.5); clear smooth boundary.
B1 RSL No. 6425	13 to 18 inches, reddish brown (5YR 5/3) sandy clay loam, dark reddish brown (5YR 3/3) moist; weak coarse subangular blocky structure; hard, firm, slightly sticky and slightly plastic; plentiful very fine roots; common very fine and fine tubular pores; slightly acid (pH 6.3); clear smooth boundary.
B21t RSL No. [illegible]	18 to 27 inches, reddish brown (5YR 4/3) sandy clay loam, dark reddish brown (5YR 3/3) moist; moderate coarse prismatic structure; very hard, firm, sticky [illegible]

CONSECUTIVE PEDON NUMBER C 69 56 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES PACHAPPA TAXADJUNCT SOIL NO. S69 CALI. 36-24 SAMPLE NO. 9 299-9 301 SAN BERNARDINO AREA

SOIL FAMILY MOLLIC HAPLOXERALF FINE-LOAMY,MIXED,THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	CSI .05- .02	FSI .02- .002	CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A	1A2A INT. II .2- .02	PCT. OF SOIL	-COARSE FRAGMENTS- I 75- 2	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM															PCT. OF LT 75		
9 299	0- 15	AP	40.8	40.1	19.1	3.6	6.0	4.9	10.0	16.3	25.8	14.3				47.8	--A	--	--	--
9 300	15- 28	B1	35.7	33.6	30.7	3.3	6.3	5.1	8.3	12.7	19.1	14.5				36.2	--A	--	--	--
9 301	43- 74	B22T	37.4	32.7	29.9	4.2	6.5	5.4	8.7	12.6	18.7	14.0				36.0	--A	--	--	--

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR- BON 6A1A -PERCENT--	NITRO GEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B -PERCENT-	19-2 6E1B	EXT. IRON FE 6C2B PCT.	ATTERBERG L I M I T LL PCT.NO.40	PL 4F1	BULK DENSITY EST 1/3 BAR 4A1F GRAMS PER CC	1/3 BAR 4A1F	DRY 4AH	CM 3B2	WATER CONTENT 1/10 BAR 4B1C ----PERCENT---	1/3 BAR 4B1C	15 BAR 4B2	EXTENSBLTY LEF 4D1 -PERCENT--	LE 4D1	-----PH------ H2O 1/1 8C1A	H2O 1/10 8C1A	CACL2 8C1E
9 299	0- 15	1.40	0.148							1.46	1.63	1.00		19.6	8.7	3.7	3.7	7.8		7.3
9 300	15- 28	0.46	0.039						1.49			1.00			13.2			7.6		6.9
9 301	43- 74	0.38								1.52	1.93	1.00		24.6	13.2	8.3	8.3	8.1		7.2

SAMPLE NO.	DEPTH CM.	----EXTRACTABLE BASES--- CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT. PCT.	--C E C-- NA- OAC 5A2A -MEQ/100G	NH4- OAC 5A6A	EC 8A1A MMHO	------WATER EXTRACT FROM SATURATED PASTE 8A1 ----- CA 6N1B	MG 6O1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	B 6T1A	PH SAT. PASTE 8C1B
		--------MEQ/100G--------									----------MEQ/1000 ML----------								PPM	
9 299	0- 15	15.0	7.9	0.2	0.6	23.7	100+		19.5	0.91										
9 300	15- 28	18.1	13.3	1.5	0.3	33.2	100+		30.5											
9 301	43- 74	17.1	12.9	2.9	0.3	33.2	100+		30.8	0.58										

SAMPLE NO.	DEPTH CM.	WATER EXCHANGBLE AT SAT. 8A PCT.	NA 5B4B MEQ.	NA 5D2 PCT.	SAR 5E	-GYPSUM- AMT. 6F1B PERCENT	REQ. 6P3A	---------- C L A Y M I N E R A L O G Y ---------- -LESS THAN 0.002 MM--- -L.T. .0002 MM- ------X RAY------ DTA ----X RAY----- I 7A2E	II 7A2E	III 7A2E	IV 7A2E	PCT. 7A3	I 7A2E	II 7A2E	III 7A2E	MINERAL CODE	RELATIVE AMOUNT
9 299	0- 15		0.2	1													1 TRACE
9 300	15- 28		1.5	5												MT MONTMORT	2 SMALL
9 301	43- 74		2.9	9				MT 5	MI 2							MI MICA	3 MODERATE
																	4 ABUNDANT
																	5 DOMINANT
																	6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.

PACHAPPA LOAM (TAXADJUNCT) 1/
S69Calif-36-24

Location: San Bernardino County, California; about 1/2 mile south of Chino-Corona Road and Cucamonga Avenue intersection; about 100 feet east of Cucamonga Avenue; SW¼ NE¼ SE¼ of sec. 5, T. 3 S., R. 7 W. Date of Sampling: June 10, 1969.
Description by: A. A. Knecht, W. D. Nettleton, W. Z. Brock. Collectors: A. A. Knecht, W. D. Nettleton, W. Z. Brock.
Classification: Mollic Haploxeralf, fine-loamy, mixed, thermic.
Vegetation: Small grain stubble, mustard, cheatgrass and sunflower weed. Climate: Average annual precipitation about 16 inches. Mean temperature about 61°F. Parent Material: Acid igneous alluvium. Topography: Alluvial fan, moderately sloping, about 5 percent slopes. Elevation: 550 feet. Drainage: Well drained; medium runoff; moderate permeability. This area is above flood plain and water table is below five feet. Soil Moisture: Profile was dry throughout when sampled.
Remarks: Field reaction was determined by the Truog method. Three samples (0-6", 6-11", 17-29") were analyzed in the Riverside Soil Survey Laboratory. *Color value of the Bt horizon slightly lower than currently defined.

HORIZON	DESCRIPTION
	(0-15 cm.)
Ap RSL 69299	0 to 6 inches, grayish brown (10YR 5/2) loam, very dark grayish brown (10YR 3/2) moist; weak medium subangular blocky structure; slightly hard, friable, slightly sticky, plastic; many very fine roots; many very fine discontinuous tubular pores; moderately alkaline (pH 8.0); clear smooth boundary.
	(15-28 cm.)
B1 RSL 69300	6 to 11 inches, dark grayish brown (10YR 4/2) clay loam, very dark grayish brown (10YR 3/2) moist; weak medium prismatic parting to moderate medium subangular blocky structure; very hard, firm, very sticky, very plastic; many very fine inped roots; common very fine discontinuous tubular pores; moderately alkaline (pH 8.0); clear smooth boundary.
	(28-43 cm.)
B21t	11 to 17 inches, dark grayish brown (10YR 4/2) clay loam, very dark grayish brown (10YR 3/2) moist; moderate medium prismatic parting to moderate medium subangular blocky structure; extremely hard, very firm, very sticky, very plastic; common very fine inped roots; few very fine vertical interstitial pores; very few thin clay films occur on faces of peds; moderately alkaline (pH 8.0); gradual smooth boundary.
	(43-74 cm.)
B22t RSL 69301	17 to 29 inches, dark grayish brown (10YR 4/2) clay loam, very dark brown (10YR 2/2) moist; moderate medium prismatic parting to moderate medium subangular blocky structure; extremely hard, very firm, very sticky, very plastic; few very fine roots; few very fine discontinuous tubular pores; few thin clay films occur on faces of peds; slightly effervescent; strongly alkaline (pH 8.5); clear smooth boundary.
	(74-100 cm.)
B3t	29 to 40 inches, dark grayish brown (10YR 4/2) sandy clay loam, very dark grayish brown (10YR 3/2) moist; weak coarse prismatic parting to moderate medium angular blocky structure; very hard, very firm, sticky, plastic; very few roots; very few discontinuous tubular pores; common moderately thick clay films on faces of peds; strongly effervescent; medium generally rounded segregated lime occurs in soft masses; strongly alkaline (pH 8.5); clear irregular boundary.
C1	40 to 54 inches, reddish yellow (7.5YR 6/6) very coarse loamy sand, strong brown (7.5YR 5/6) moist; massive; hard, friable, nonsticky, nonplastic; strongly effervescent; strongly alkaline (pH 8.5); clear wavy boundary.
	(135-170 cm.)
C2	54 to 67 inches, very pale brown (10YR 7/3) very coarse loamy sand, pale brown (10YR 6/3) moist; massive; hard, firm, nonsticky, nonplastic; moderately alkaline (pH 8.0).

1/ The Pachappa series is in a fine-loamy, mixed, thermic family of Mollic Haploxeralfs. This pedon is in the same family and subgroup but has a Bt horizon with a color value of less than 5.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Pleasanton gravelly loam **LOCATION** Alameda County, Calif rnia

SOIL FAMILY: Mollic Haploxeralf, fine-loamy, mixed, thermic

SOIL NOS. S60Calif-01-1 **LAB. NOS.** 60377 thru 60382

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-9	A_{1p}	6.2	12.7	8.7	20.2	10.1	29.0	13.1	36.1	14.9	55	fsl
9-21	A_{12}	7.0	11.8	8.1	18.7	10.3	29.9	14.2	35.8	15.7	27	fsl
21-32	B_{21}	7.2	12.8	7.9	18.0	10.0	27.2	16.9	24.0	14.0	49	fsl
32-48	B_{22}	6.1	13.5	6.6	17.3	10.3	26.3	19.9	34.1	13.4	40	fsl-scl
48-64	B_3	3.8	7.0	6.4	17.6	14.5	31.6	19.1	40.0	16.7	42	l
64-72	C	3.0	5.4	6.9	14.9	22.0	29.9	17.9	43.6	14.9	30	fsl- l

pH 8C1b Saturated Paste 1:1	pH 8C1a 1:10	Organic Matter 6A1a Organic Carbon Pct.	6B1a Nitrogen Pct.	C/N	6E1b $CaCO_3$ equivalent Pct.	8A1a Elec. conduc. mmhos/cm. at 25°C	Saturation Ext. Soluble 6I1a CO_3	6J1a HCO_3	6L1a SO_4	6M1a NO_3	6C1b Free Fe_2O_3 Pct.	4B2 Soil water 15 atmos. Pct.
						milliequivalents per liter						
6.3	6.6	2.02	0.172	11.7		0.5	0	3.8				3.6
6.8	7.0	0.96	0.087	11.0		0.3	0	2.0				3.5
7.1	7.3	0.45	0.052	8.7		0.3	0	1.5				5.7
7.3	7.7	0.23	0.032			0.2	0	1.0				7.6
7.3	7.6	0.17				0.2	0	1.1				7.4
7.4	8.1	0.12				0.3	0	1.1				4.4

5A2a CATION EXCHANGE CAPACITY	EXTRACTABLE CATIONS 5B 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a Ca	6O1a Mg	6K1a Cl	8A MOISTURE AT SATURATION %
← Na →	milliequivalents per 100g. soil →						← milliequivalents per liter →					
17.3	10.1	2.0	4.6	0.1	0.6	74	0.8	0.3	3.7	0.8	1.3	29.6
14.4	9.3	1.5	2.7	0.1	0.4	81	0.7	0.1	2.2	0.5	0.4	23.8
15.4	9.4	3.6	2.3	0.1	0.3	85	0.9	<0.1	1.6	0.7	0.4	22.7
18.0	8.4	7.1	2.1	0.2	0.3	93	1.0	<0.1	0.9	0.9	0.2	26.1
19.6	8.0	8.9	2.0	0.4	0.3	90	1.1	<0.1	0.6	0.7	0.2	27.8
17.2	7.3	8.0	1.6	0.4	0.3	91	1.6	<0.1	1.0	0.6	0.6	27.6

1/ Percent of the whole soil

PLEASANTON GRAVELLY LOAM
S60Calif-01-1

Location: Alameda County, California. SE1/4 of the NE1/4 of the NE1/4 of Sec. 24, T. 3 S., R. 2 E., about 5 miles southeast of the center of Livermore.
Date of Sampling: August 5, 1960.
Description by: L. E. Welch. Collectors: W. C. Fischer and L. E. Welch.
Classification: Mollic Haploxeralf, fine-loamy, mixed, thermic.
Vegetation: Cultivated, barley. Climate: Mean annual rainfall about 14 inches.
Parent Material: Alluvium from sedimentary rock. Relief: Slightly convex south-facing slope of 1 percent, on low terrace. Elevation: 710 feet. Drainage: Well drained, medium runoff, moderate permeability over moderately slow permeability.
Remarks: Moist below 24 inches. Occasional cobbles throughout. Few worm holes.

HORIZON		DESCRIPTION
A_{1p}	0-9"	**Gray brown (10YR 5/2) gravelly loam, very dark gray brown (10YR 3/2) when moist; massive to very weak granular structure; hard when dry; friable when moist; slightly sticky and slightly plastic when wet; abundant very fine, plentiful fine, plentiful medium, random roots; common very fine, common fine and few medium, random, inped open, interstitial pores; slightly acid, pH 6.5; abrupt, smooth lower boundary.**
A_{12}	9-21"	**Gray brown (10YR 5/2) gravelly loam, very dark gray brown (10YR 3/2) when moist; massive; hard when dry; friable when moist; slightly sticky and slightly plastic when wet; abundant very fine, plentiful fine and few medium random roots; common very fine, common fine and few medium random, inped, open interstitial pores; slightly acid, pH 6.5; clear, smooth lower boundary.**
B_{21}	21-32"	Brown (10YR 4/3) gravelly clay loam, dark brown (10YR 3/3) when moist; moderate, medium, subangular blocky; very hard when dry; friable when moist; sticky and plastic when wet; plentiful very fine, plentiful fine, random roots; many very fine, many fine, and few medium, random, inped, open, tubular pores; common, moderately thick clay films on ped faces and pores; slightly acid, pH 6.5; gradual, wavy lower boundary.
B_{22}	32-43"	Brown (10YR 4/3) gravelly clay loam, dark brown (10YR 3/3) when moist; moderate, medium, subangular blocky structure; very hard when dry; friable when moist; sticky and plastic when wet; plentiful very fine, few fine, random roots; many very fine, common fine, few medium, random, inped, few exped, open, tubular pores; common, moderately thick clay films on ped faces and pores; neutral, pH 7.0; gradual, wavy lower boundary.
B_3	43-64"	Brown (10YR 4/3) gravelly silty clay loam; dark brown (10YR 3/3) when moist; weak, medium subangular structure; very hard when dry; friable when moist; sticky and plastic when wet; very few, very fine roots; common very fine, common fine, random, open, tubular pores; few thick and few thin clay films on ped faces and pores; mildly alkaline, pH 7.5; gradual, wavy lower boundary.
C	64-72"	Yellowish brown (10YR 5/4) gravelly silt loam, dark yellowish brown (10YR 4/4) when moist; massive to weak, angular blocky structure; hard when dry, friable when moist; sticky and slightly plastic when wet; void of roots; common very fine, common fine, and few medium random inped, open, tubular pores; few thin clay films on ped faces and pores; mildly alkaline, pH 7.5.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Pleasanton gravelly loam (taxadjunct) **LOCATION** Alameda County, California

SOIL FAMILY: Mollic Haploxeralf, fine-loamy, mixed, thermic

SOIL NOS. S60Calif-01-2 **LAB. NOS.** 60383 thru 60388

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-6	A_{1p}	8.9	7.9	5.9	13.2	13.1	38.9	12.1	42.4	17.8	38	1
6-17	A_{12}	6.7	8.5	5.6	13.1	13.7	36.8	15.6	42.3	16.4	31	1
17-31	B_{21}	9.1	8.6	5.4	11.5	12.6	37.4	15.4	40.7	16.6	28	1
31-48	B_{22}	8.7	8.8	5.0	8.6	11.9	39.1	17.9	36.3	19.7	23	1
48-61	B_3	9.0	10.6	6.0	11.5	10.9	36.1	15.9	36.6	17.2	32	1
61+	C	7.9	8.0	4.5	9.4	11.5	39.0	19.7	37.5	18.6	21	1

pH 8C1b Saturated Paste 1:1	8C1a 1:10	Organic Matter 6A1a Organic Carbon Pct.	6B1a Nitrogen Pct.	C/N	6E1b $CaCO_3$ Equivalent Pct.	8A1a Elec. Conduc. mmhos/cm. at 25° C	Saturation Ext. Soluble 6I1a CO_3	6J1a HCO_3	6L1a SO_4	6M1a NO_3	6C1b Free Fe_2O_3 Pct.	4B2 Soil water 15 atm s. Pct.
							milliequivalents per liter					
6.4	6.7	1.79	0.140	12.8		0.5	0	3.5				5.9
6.4	6.9	0.85	0.080	10.6		0.3	0	1.9				6.9
6.9	7.1	0.50	0.047	10.6		0.3	0	1.4				6.8
7.2	7.4	0.46	0.046	10.0		0.2	0	1.2				7.6
7.1	7.3	0.28	0.035	8.0		0.2	0	0.8				6.6
6.9	7.1	0.14				0.3	0	0.8				8.5

5A2a CATION EXCHANGE CAPACITY ← (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a Ca	6O1a Mg	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
16.3	8.0	2.8	4.4	0.3	0.7	73	1.0	0.5	3.4	1.2	1.0	31.8
18.6	10.3	4.2	3.2	0.2	0.4	83	0.8	0.1	1.7	0.9	0.3	27.7
18.4	8.1	6.6	2.2	0.2	0.2	87	0.8	<0.1	1.0	1.0	0.4	28.0
19.1	8.7	8.0	2.2	0.2	0.2	87	0.9	<0.1	1.0	0.7	0.3	28.1
16.0	7.0	6.9	1.9	0.3	0.2	88	1.2	<0.1	0.8	0.8	0.2	21.1
19.9	7.7	8.6	2.4	0.5	0.2	88	1.5	<0.1	0.7	0.4	0.6	26.0

1/ Percent of the whole soil

PLEASANTON GRAVELLY LOAM (taxadjunct) 1/
S60Calif-01-2

Location: Alameda County, California, SW1/4 of the NW1/4 of the SW1/4 of Sec. 22, T. 3 S., R. 2 E., about 2 miles southeast of the center of Livermore.
Date of Sampling: August 5, 1960
Description by: L. E. Welch. Collectors: L. E. Welch and W. C. Fischer.
Classification: Mollic Haploxeralf, fine-loamy, mixed, thermic.
Vegetation: Cultivated in past, annual grasses at present. Parent Material: Alluvium from sedimentary rock. Relief: Slightly convex west-facing slope of 1 percent, on low terrace. Elevation: 590 feet. Drainage: Well drained, medium runoff, moderate permeability over moderately slow permeability. Soil Moisture: Moist below 17 inches.
Remarks: Few worm holes. Occasional cobbles throughout.

HORIZON		DESCRIPTION
A_{1p}	0-6"	Gray brown (10YR 5/2) gravelly loam, very dark gray brown (10YR 3/2) when moist; massive; hard when dry; friable when [illegible] very fine, abundant fine, plentiful medium, few coarse, random roots; many very fine, many fine, few medium, inped, open, interstitial pores; slightly acid, pH 6.5; gradual, smooth lower boundary.
A_{12}	6-17"	Gray brown (10YR 5/2) gravelly loam, very dark gray brown (10YR 3/2) when moist; massive; hard when dry, friable when moist, slightly sticky and slightly plastic when wet; abundant very fine, abundant fine, few medium, random roots; many very fine, many fine, common medium, inped, open, interstitial pores; slightly acid, pH 6.5; gradual, smooth lower boundary.
B_{21}	17-31"	Gray brown (10YR 5/2) gravelly clay loam, very dark gray brown (10YR 3/2) when moist; weak medium subangular blocky structure; very hard when dry, friable when moist, sticky and plastic when wet; abundant very fine, abundant fine, few medium, random roots; many very fine, many fine and common medium, inped, open, tubular pores; common thin [illegible]

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Rescue loam **LOCATION** El Dorado County, Calif.

SOIL FAMILY: Mollic Haploxeralf, fine-loamy, mixed, thermic

SOIL NOS. S61Calif-9-1 **LAB. NOS.** 6179 - 6185

DEPTH INCHES	HORIZON	181b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)							3A1		2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	
0- 5	A_1	6.8	16.1	8.1	16.1	9.0	29.8	14.1	29.1	19.3	8	sl
5-10	A_3	7.7	16.2	8.4	15.0	8.5	29.1	15.1	27.3	19.1	5	sl
10-14	B_1 t	9.2	18.6	6.8	12.0	7.4	25.7	20.3	21.8	18.3	4	scl-sl
14-26	B_2 t	9.4	12.1	4.2	13.3	8.6	21.3	31.1	23.6	14.8	4	scl
26-34	B_3 t	10.1	10.6	5.9	21.1	11.7	20.4	20.2	31.6	14.1	6	scl-sl
34-55	C_1	13.9	13.8	7.6	25.0	12.2	17.3	10.2	35.5	9.9	6	cosl
55-66+	C_2	23.0	14.8	7.2	24.5	11.0	15.2	4.3	33.4	8.9	14	lcos

Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, David Estrada,

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Rescue loam **LOCATION** El Dorado County, Calif.

SOIL FAMILY: Mollic Haploxeralf, fine-loamy, mixed, thermic

SOIL NOS. S61Calif-9-2 **LAB. NOS.** 6186 - 6192

DEPTH INCHES	HORIZON	1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-2½	A_1	8.4	16.2	10.1	16.9	8.0	27.8	12.6	28.0	17.0	10	sl
2½-8	A_3	11.3	14.5	9.2	14.1	7.6	27.7	15.6	25.3	18.1	7	cosl
8-12	B_1	10.6	16.1	8.7	12.9	7.2	25.6	18.9	23.4	16.6	7	cosl
12-24	B_2+	13.3	13.3	6.3	10.3	5.8	23.8	27.2	18.9	16.7	7	scl

RESCUE LOAM
S61Calif-9-2

Location: El Dorado County, California, 2.5 miles northwest of Bass Lake, SE 1/4 NE 1/4, Sec. 20, T. 10 N., R. 9 E. Date of Sampling: April 25, 1961.
Description by: Grant Kennedy. Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, David Estrada, Lothair Grant, and Lynn Brittan.
Classification: Mollic Haploxeralf, fine-loamy, mixed, thermic.
Vegetation: Filaree, soft chess, ripgut, wild oats, clover. Climate: Mean annual rainfall 30 inches - mean annual temperature is 62° F., the average January temperatures is 46.7° F., and the average July temperature is 78° F.
Parent material: Gabbro, diorite. Topography: Undulating uplands with a 5 percent slope to the east. Elevation: 1,380 feet. Drainage: Well drained; medium runoff; moderate over slow permeability.
Remarks: This area may have been farmed at one time and turkey manure may have been spread in recent years. C1 and C2 horizons are well weathered, but show original rock structure. There was a thin quartz ledge on the left side of the pit. Less than 5 percent of the are contained well-scattered rock outcrops. The dry color of the B2t is between hue 2.5YR and 5YR.

HORIZON	DESCRIPTION
A1. 0-21	Yellowish red (5YR 3/6) loam, dark reddish brown (5YR 3/2) when moist; [illegible]
[illegible]	[illegible]

CONSECUTIVE PEDON NUMBER C 68117 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES TRABUCO TAXADJUNCT SOIL NO. S68 CALI. 29-30 SAMPLE NO. 8 875-8 882 NEVADA COUNTY AREA

SOIL FAMILY MOLLIC HAPLOXERALF FINE-LOAMY,MIXED,THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SANDS VCS 2-1	CS 1-.05	MS .5-.25	FS .25-.10	VFS .1-.05	SILTS CSI .05-.02	FSI .02-.002	CLAYS CO. 2-.2U	FINE LT .2U	CARBONATE 3A1A	INT. II .2-.02 WHOLE SOIL	COARSE FRAGMENTS PCT. OF 20	I 75-20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75			
8 875	0- 10	AP	51.3	35.6	13.1	4.0	10.0	8.7	16.9	11.7	15.3	20.3				36.4	--A	--	--	--
8 876	10- 25	A12	49.4	33.8	16.8	3.3	8.6	8.4	17.0	12.1	13.1	20.7				34.9	--A	--	--	--
8 877	25- 38	B1	45.0	33.4	21.6	2.4	7.3	7.7	16.0	11.6	12.2	21.2				33.2	--A	--	--	--
8 878	38- 69	B21T	27.7	34.4	37.9	0.8	4.0	4.2	10.7	8.0	9.6	24.8				23.8	--A	--	--	--
8 879	69- 99	B22T	38.0	35.9	26.1	1.6	6.3	6.0	14.0	10.1	11.4	24.5				29.4	--A	--	--	--
8 880	99-130	B3 T	42.3	36.3	21.4	2.3	6.9	6.6	16.3	10.2	12.8	23.5				32.3	--A	--	--	--
8 881	130-170	C	66.0	21.2	12.8	5.9	19.6	11.7	19.8	9.0	8.7	12.5				27.3	--A	--	--	--
8 882	0- 10	AP1															--A	--	--	--

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	CARBONATE NITROGEN 6B1A	AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	DRY 4A1F	4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		PERCENT				PCT.NO.40		GRAMS PER CC				PERCENT			-PERCENT-	
8 875	0- 10	2.69				2.1	0.1			NP	NP		1.69	1.70	1.00		17.5	8.7	0.2	0.2
8 876	10- 25	0.72				2.2	0.2					1.60			1.00			8.5		
8 877	25- 38	0.44				2.4	0.2			29	22	1.58			1.00			10.3		
8 878	38- 69	0.26				2.8	0.1					1.57			1.00			19.4		
8 879	69- 99	0.12				2.0	0.1			44	27		1.57	1.73	1.00		25.6	17.2	3.3	3.3
8 880	99-130					1.8	TR					1.58			1.00			15.2		
8 881	130-170					1.1	0.1			32	27		1.59	1.71	1.00		22.5	9.7	2.5	2.5
8 882	0- 10	2.80										1.69			1.00			9.2		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACI-DITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CACL2 8C1E
		MEQ/100G									PERCENT					
8 875	0- 10	10.2	3.1	0.1	0.4	13.8			13.8	17.3	100+		80		5.8	5.3
8 876	10- 25	8.4	5.4	0.1	0.1	14.0			14.0	15.2	100+		92		6.2	5.7
8 877	25- 38	10.0	7.6	0.1	0.1	17.8			17.8	19.0	100+		94		6.3	5.8
8 878	38- 69	16.5	17.7	0.1	0.1	34.4			34.4	33.7	100+		100+		6.3	5.8
8 879	69- 99	18.5	16.6	0.2	0.1	35.4			35.4	36.0	100+		98		6.5	6.0
8 880	99-130	19.6	17.4	0.2	0.1	37.3			37.3	34.4	100+		100+		6.5	6.1
8 881	130-170	15.2	12.5	0.2	0.1	28.0			28.0	26.3	100+		100+		6.6	6.0
8 882	0- 10														6.0	5.4

CLAY MINERALOGY

SAMPLE NO	DEPTH CM.	LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 875	0- 10	KK	4	MI	3						KK KAOLINIT	1 TRACE
8 876	10- 25										MI MICA	2 SMALL
8 877	25- 38	KK	4	MI	3							3 MODERATE
8 878	38- 69										MT MONTMORT	4 ABUNDANT
8 879	69- 99	KK	4	MT	3							5 DOMINANT
8 880	99-130											6 INDETER.
8 881	130-170	KK	4	MT	2	MI	1					
8 882	0- 10											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.

TRABUCO LOAM (TAXADJUNCT) 1/
S68Calif-29-30

Location: Nevada County, California. 1,350 feet east of Pleasant Valley Road, 7.5 miles west of Grass Valley, California; 1,000 feet east and 1,100 feet south of W 1/4 corner sec. 21, T.16N., R. 7E. Date of Sampling: September 4, 1968.
Description by: L. A. Brittan, G. Staidl. Collectors: L. A. Brittan, G. Staidl, C. B. Goudey, T. M. Ryan, G. M. Kennedy.
Classification: Mollic Haploxeralf, fine-loamy, mixed, thermic.
Vegetation: This field has been cultivated recently. Irrigated pasture now, mostly annual grasses but some perennials. Ponderosa pine, Digger pine, blue oak. Climate: Average annual precipitation is 35 inches. Mean annual temperature 61°F. Parent Material: Granodiorite, coarse grained. Topography: Mountainous upland, strongly sloping, convex, 10 percent slope, southeast aspect. Elevation: 1,260 feet. Drainage: Well drained; runoff is medium; permeability is moderate to very slow. Soil Moisture: Dry to 10 inches, very slightly moist 10 inches to 36 inches.
Remarks: pH taken by Hellige method. This site was flood irrigated in past. See Ap and A12 horizons, which are mottled, probably due to past irrigation. Thickness and sequence of horizons in this profile very similar to S68Calif-29-29. An occasional quartz gravel occurs in the upper 15 inches of this profile. No remarks regarding fertilization can be made.

HORIZON	DESCRIPTION
Ap RSL 68875	(0-10 cm.) 0 to 4 inches, brown (7.5YR 5/4) loam, dark brown (7.5YR 3/2) moist; common fine distinct yellowish red and red (5YR 5/8, 2.5YR 4/6) mottles, yellowish red and red (5YR 5/8 and 2.5YR 4/6) moist; moderate very coarse platy structure; hard, friable, slightly sticky, slightly plastic; common very fine and few fine random roots; common very fine tubular and interstitial pores; strongly acid (pH 5.5); abrupt smooth boundary.
A12 RSL 68876	(10-25 cm.) 4 to 10 inches, brown (7.5YR 5/4) heavy loam, dark reddish brown (5YR 3/3) moist; common fine prominent red (2.5YR 4/8) mottles, dark red (2.5YR 3/6) moist; weak coarse prismatic structure; hard, friable, sticky, plastic; common very fine and fine vertical roots; common very fine tubular and interstitial pores; slightly acid (pH 6.2); clear wavy boundary.
B1 RSL 68877	(25-38 cm.) 10 to 15 inches, reddish brown (5YR 4/4) clay loam, dark reddish brown (5YR 3/4) moist; weak medium and coarse prismatic structure: hard, friable, sticky, plastic; few very fine vertical, medium and coarse horizontal roots; common very fine tubular and interstitial pores; few thin clay films line pores and as bridges; slightly acid (pH 6.2); clear wavy boundary.
B21t RSL 68878	(38-69 cm.) 15 to 27 inches, reddish brown (5YR 5/4) clay, dark reddish brown (5YR 3/4) moist; moderate coarse prismatic parting to moderate coarse angular blocky structure; very hard, firm, sticky, very plastic; few very fine, fine vertical, and medium and coarse horizontal roots; common very fine tubular pores; continuous moderately thick clay films on faces of peds and lining pores; light mineral grains apparent; slightly acid (pH 6.2); gradual wavy boundary.
B22t RSL 68879	(69-99 cm.) 27 to 39 inches, brown (7.5YR 5/4) clay, dark reddish brown (5YR 3/4) moist; moderate medium prismatic parting to moderate coarse angular blocky structure; very hard, firm, sticky, very plastic; few very fine vertical roots; few very fine tubular pores; many moderately thick dusky red (2.5YR 3/2 moist) clay films on faces of peds and continuous thick clay films line pores; mica and light mineral grains apparent; slightly acid (pH 6.2); gradual wavy boundary.
B3t RSL 68880	(99-130 cm.) 39 to 51 inches, pale brown (10YR 6/3) light clay, brown and dark brown (10YR 4/3) moist; weak medium prismatic parting to weak coarse angular blocky structure; hard, firm, sticky, very plastic; few very fine and fine vertical roots; few very fine tubular pores; many moderately thick dark red (2.5YR 3/6) clay films, dusky red (2.5YR 3/3) moist, on faces of peds and lining pores; mica and light mineral grains apparent; slightly acid (pH 6.2); gradual wavy boundary.
C RSL 68881	(130-170 cm.) 51 to 66 inches, light yellowish brown (10YR 6/4) heavy clay loam, brown and dark brown (7.5YR 4/4) moist; massive; hard, firm, sticky, very plastic; very few very fine random roots; few very fine tubular pores; common moderately thick red (2.5YR 4/6) clay films, dark red (2.5YR 3/6) moist, lining pores and as bridges; mica and other light mineral grains apparent; medium acid (pH 6.0).

1/ The Trabuco series is in a fine, mixed, thermic family of Mollic Haploxeralfs. This pedon has less than 35 percent clay in the upper 20 inches of the argillic horizon.

CONSECUTIVE PEDON NUMBER C 68111 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES SECCA SOIL NO. S68 CALI. 29- 7 SAMPLE NO. 8 832-8 838 NEVADA COUNTY AREA

SOIL FAMILY MOLLIC HAPLOXERALF FINE,MIXED,MESIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	CSI .05- .02	FSI .02- .002	CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A.02	1A2A INT. II .2- OF WHOLE SOIL	-COARSE FRAGMENTS- PCT. I 75-	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75		
8 832	0- 15	A1	23.5	58.8	17.7	2.4	3.9	2.8	6.6	7.8	18.7	40.1				30.6	61*	10	44	5
8 833	15- 38	A3	18.8	57.4	23.8	2.2	2.8	2.2	5.1	6.5	15.6	41.8				25.3	59*	9	25	2
8 834	38- 56	B1 T	17.7	50.7	31.6	2.7	3.5	2.0	4.2	5.3	13.5	37.2				21.4	45*	9	1	1
8 835	56- 91	B2 T	10.0	29.5	60.5	1.0	1.1	1.2	3.0	3.7	7.4	22.1				13.0	41*	--	TR	2
8 836	91-115	B3 T	22.4	35.3	42.3	3.8	4.8	2.8	6.1	4.9	6.1	29.2				14.7	35A	--	24	11
8 837	115-145	C	33.2	35.2	31.6	7.6	6.7	4.0	8.3	6.6	9.0	26.2				20.5	50A	--	22	28
8 838	0- 15	A1															41A	--	37	4

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR- BON 6A1A	NITRO GEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG L I M I T LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		-PERCENT-				PCT.NO.40		GRAMS PER CC				-PERCENT-			-PERCENT-	
8 832	0- 15	1.50	0.050			4.9	0.4			35	30	0.99	0.99	0.64		19.8	11.0			
8 833	15- 38	0.71	0.045			5.5	0.3					1.22	1.32	0.60		33.9	12.6	2.7	1.6	
8 834	38- 56	0.64				6.0	0.4			40	27	1.36	1.45	0.70		29.5	14.9	2.2	1.5	
8 835	56- 91	0.23				5.5	0.4			64	30	1.32	1.79	0.74		35.8	24.7	10.7	7.5	
8 836	91-115					3.3	0.2					0.98	1.27	0.83		60.4	25.6	9.0	7.3	
8 837	115-145					4.4	0.2			54	35	1.12	1.33	0.71		53.3	22.8	5.9	4.0	
8 838	0- 15	1.69	0.099								0.99			0.80			11.1			

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6Q2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI- DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4- OAC 5A6A	BASE SATURATION SUM+ ACI- DITY	SUM+ AL	NH4- OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA- CL2 8C1E
		MEQ/100G									PERCENT					
8 832	0- 15	10.4	6.8	0.1	0.2	17.5	7.9		25.4	20.8	69		84		6.2	5.7
8 833	15- 38	9.9	7.9	0.1	0.1	18.0	5.6		23.6	20.9	76		86		6.4	5.9
8 834	38- 56	10.2	10.7	0.1	0.1	21.1	6.4		27.5	23.5	77		90		6.3	5.8
8 835	56- 91	16.3	20.1	0.2	0.2	36.8			36.8	36.4	100+		100+		6.6	6.2
8 836	91-115	17.3	20.8	0.3	0.1	38.5			38.5	40.6	100+		95		6.7	6.3
8 837	115-145	12.6	15.6	0.3	0.1	28.6	4.6		33.2	31.7	86		90		6.3	6.1
8 838	0- 15														6.1	5.7

SAMPLE NO	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM MINERAL X RAY I	7A2E II	7A2E III	CODE	RELATIVE AMOUNT
8 832	0- 15	KK 4	MT 3								KK KAOLINIT	1 TRACE
8 833	15- 38										MT MONTMORT	2 SMALL
8 834	38- 56	KK 4	MT 2	MI 2								3 MODERATE
8 835	56- 91	KK 4	MT 3								MI MICA	4 ABUNDANT
8 836	91-115											5 DOMINANT
8 837	115-145	KK 4	MT 3									6 INDETER.
8 838	0- 15											

SAMPLE PREPARATION CODES
*=THE 75-250MM FRACTION IS INCLUDED.
A=STANDARD PREPARATION.

SECCA GRAVELLY SILT LOAM
S68Calif-29-7

Location: Nevada County, California, 1,300 feet east and 500 feet north of W 1/4 corner, sec. 16, T. 15 N., R.8E., 5 miles SSW of Grass Valley, California; 1.6 miles SSE of intersection of McCourtney and Wolf Creek Mountain Lookout Roads, 100 feet SE of Wolf Creek Mountain Lookout Road. Date of Sampling: September 3, 1968.

Description by: L. A. Brittan, C. B. Goudey. Collectors: G. M. Kennedy, L. A. Brittan, C. B. Goudey, T. M. Ryan, G. Staidl.

Classification: Mollic Haploxeralf, fine, mixed, mesic.

Vegetation: Manzanita, ceanothus, scattered Digger pine, blue oak, squirrel tail, very scattered ponderosa pine. Not cultivated. Climate: Average annual precipitation 45 inches. Mean annual temperature 56°F. Parent Material: Metabasic rock similar to diabase. C horizon (45-57") listed as weathered fractured metabasic rock or diabase on geology map (California Division of Mines and Geology.) Topography: Moderately steep, concave slope, 17 percent, aspect W., near top of slope. Elevation: 2,260 feet. Drainage: Well drained; runoff is medium; permeability is moderate over very slow in subsoil. Soil Moisture: Dry to 15 inches. Very slightly moist 15 inches to 57 inches.

Remarks: pH taken by Hellige method. pH approximately 1/2 unit lower in lower portions of profile S68Calif-29-7 than in S68Calif-29-5. About 10 to 25 percent of surface covered by stone fragments. Lower part of B1t, 15 to 22 inches, contained a stoneline, approximately 25 percent by volume of fragments greater than 3 inches in diameter. Gravel pavement on A1 horizon 1/2 to 1 inch thick. Few Mn shot in B3t horizon. The A1 horizon, 0 to 6 inches, specifically contains 2 percent by volume of fragments greater than 3 inches in diameter.

HORIZON	DESCRIPTION
	1 to 0 inches, gravel pavement.
	(0-15 cm.)
A1 RSL 68832	0 to 6 inches, brown (7.5YR 5/4) gravelly silt loam, dark reddish brown (5YR 3/4) moist; strong fine medium and coarse subangular blocky structure; slightly hard, friable, slightly sticky, plastic; many very fine and very few fine roots; common very fine tubular and interstitial pores; a few cobblestones and about 40 percent gravel by volume; medium acid (pH 6.0); clear wavy boundary.
	(15-38 cm.)
A3 RSL	6 to 15 inches, reddish brown (5YR 5/4) gravelly silt loam, dark reddish brown (5YR 3/4) moist; moderate medium and coarse subangular blocky structure; slightly

CONSECUTIVE PEDON NUMBER C 68116 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES TRABUCO SOIL NO. S68 CALI. 29-29 SAMPLE NO. 8 865-8 874 NEVADA COUNTY AREA

SOIL FAMILY MOLLIC HAPLOXERALF, FINE, MIXED, THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SANDS VCS 2-1	CS 1-.05	MS .5-.25	FS .25-.10	VFS .1-.05	CSI .05-.02	FSI .02-.002	SILTS CO. 2-.2U	FINE LT .2U	CLAYS CARBONATE 3A1A	INT. II .2-.02	COARSE FRAGMENTS PCT. OF WHOLE SOIL I 75-20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75		
8 865	0- 10	AP	38.1	40.4	21.5	2.5	7.6	7.5	14.4	6.1	14.7	25.7				27.7	--A	--	--	--
8 866	10- 25	A12	37.5	38.0	24.5	1.9	7.2	7.1	13.9	7.4	13.6	24.4				28.4	--A	--	--	--
8 867	25- 38	B1 T	33.2	35.4	31.4	1.5	6.3	6.2	11.9	7.3	11.1	24.3				24.7	--A	--	--	--
8 868	38- 74	B21T	23.4	27.3	49.3	0.5	4.4	4.7	8.8	5.0	8.1	19.2				17.6	--A	--	--	--
8 869	74-100	B22T	24.8	31.8	43.4	1.1	4.7	4.1	8.6	6.3	8.8	23.0				20.0	--A	--	--	--
8 870	100-140	B31T	35.4	32.9	31.7	2.5	7.0	5.0	11.5	9.4	11.0	21.9				27.3	--A	--	--	--
8 871	140-170	B32T	38.5	33.4	28.1	2.6	8.1	5.4	12.8	9.6	10.7	22.7				27.8	--A	--	--	--
8 872	170-205	C 1	42.6	34.3	23.1	4.4	9.0	5.7	14.5	9.0	9.8	24.5				27.4	--A	--	--	--
8 873	205-235	C 2	58.8	26.3	14.9	4.6	13.8	9.3	20.7	10.4	8.4	17.9				30.6	--A	--	--	--
8 874	0- 10	AP1															--A	--	--	--

SAMPLE NO.	DEPTH CM.	ORG. MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LIMIT LL 4F2	PL 4F1	1/3 BAR 4A1F	BULK DENSITY EST. 1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTMSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		-PERCENT-				PCT.NO.40		GRAMS PER CC				-PERCENT-			-PERCENT-	
8 865	0- 10	2.55	0.198			3.5	0.1			37	30		1.45	1.50	1.00		25.2	10.6	1.1	1.1
8 866	10- 25	0.73	0.068			3.8	0.2			32	9		1.47	1.53	1.00		23.2	10.4	1.3	1.3
8 867	25- 38	0.44				4.2	0.3						1.53	1.63	1.00		22.6	13.4	2.1	2.1
8 868	38- 74	0.29				4.9	0.3			47	27		1.45	1.63	1.00		29.3	20.1	4.0	4.0
8 869	74-100	0.20				4.4	0.2			48	23		1.40	1.61	1.00		30.8	22.2	4.8	4.8
8 870	100-140	0.14				3.8	0.3						1.48	1.62	1.00		28.6	20.6	3.1	3.1
8 871	140-170					3.2	0.1			41	17		1.49	1.64	1.00		28.2	18.8	3.2	3.2
8 872	170-205					2.1	0.1					1.51			1.00			15.3		
8 873	205-235					1.3	TR			32	27		1.54	1.55	1.00		23.0	10.8	0.2	0.2
8 874	0- 10	7.95	0.505									1.45			1.00			18.2		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACIDITY 6H2A	KCL EXT. AL 6G1E	CEC SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACIDITY	SUM+ NH4-AL OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT				
8 865	0- 10	9.4	2.9	0.1	0.5	12.9			12.9	18.1	100+	71		5.7	5.3
8 866	10- 25	7.5	2.7	0.1	0.3	10.6			10.6	14.5	100+	73		5.9	5.4
8 868	38- 74	10.8	7.0	0.1	0.1	18.0			18.0	22.6	100+	80		6.0	5.6
8 869	74-100	13.8	11.2	0.1	0.1	25.2			25.2	27.6	100+	91		6.1	5.8
8 870	100-140	14.1	12.7	0.2	0.1	27.1			27.1	27.9	100+	97		6.3	5.9
8 871	140-170	17.2	15.0	0.2	0.1	32.5			32.5	27.7	100+	100+		6.3	6.0
8 872	170-205	12.6	10.9	0.1	0.1	23.7			23.7	27.8	100+	85		6.5	6.1
8 873	205-235	14.0	11.0	0.1	0.1	25.2			25.2	23.1	100+	100+		6.6	6.2
8 874	0- 10													6.0	5.7

SAMPLE NO	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 865	0- 10	KK 3	MI 2								KK KAOLINIT	1 TRACE
8 866	10- 25										MI MICA	2 SMALL
8 867	25- 38											3 MODERATE
8 868	38- 74	KK 3										4 ABUNDANT
8 869	74-100											5 DOMINANT
8 870	100-140											6 INDETER.
8 871	140-170											
8 872	170-205											
8 873	205-235	KK 4	MI 2									
8 874	0- 10											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.

TRABUCO LOAM
S68Calif-29-29

Location: Nevada County, California; 600 feet north of N 1/4 corner sec. 9, T. 16 N., R. 7 E.; 1,400 feet NNW of intersection of Pleasant Valley and Beckman Hill Roads; 8½ miles NW of Grass Valley, California. Date of Sampling: September 4, 1968.
Description by: George Staidl and Lynn Brittan. Collectors: L. A. Brittan, G. Staidl, C. B. Goudey, T. M. Ryan, G. M. Kennedy.
Classification: Mollic Haploxeralf, fine, mixed, thermic.
Vegetation: Not cultivated. Oak-grass cover. Live oak, blue oak, Digger pine, ripgut, soft chess, annual grasses, forbs. Climate: Average annual precipitation is 35 inches. Mean annual temperature is 61°F. Parent Material: Granodiorite. Coarse grained granitic rock. Reaction slightly acid. C1 and C2 horizons weathered to loam. Light and dark mineral grains and mica particles apparent in the C1 and C2 horizons. These horizons were not sampled, but were augered. Topography: Moderately sloping, plain slope, 6 percent, west aspect near crest of slope. Elevation: 1,300 feet. Drainage: Well drained; runoff is medium; permeability is moderate over very slow. Soil Moisture: Dry to 15 inches. Very slightly moist 15 to 92 inches.
Remarks: pH taken by Hellige method. There is only an occasional gravel in the profile. Light and dark mineral grains and mica particles are quite apparent from 15 inches to 92 inches. The Ap and A12 horizons combined, B21t and B22t combined, and B32t horizon were sampled for the California Division of Highways. Moist chroma is one unit lower than characteristic of the series in all parts of the epipedon.

HORIZON	DESCRIPTION
Ap RSL 68865 (0-10 cm.)	0 to 4 inches, reddish brown (5YR 4/4) loam, dark reddish brown (5YR 3/3) moist; moderate medium and coarse prismatic structure; hard, friable, sticky and plastic; common very fine random roots; common very fine tubular and interstitial pores; medium acid (pH 6.0); abrupt wavy boundary.
A12 RSL 68866 (10-25 cm.)	4 to 10 inches, reddish brown (5YR 4/4) heavy loam, dark reddish brown (5YR 3/3) moist; weak medium and coarse prismatic structure; hard, friable, slightly sticky, plastic; common very fine and fine random, and medium oblique roots; common very fine tubular and interstitial pores; medium acid (pH 6.0); clear smooth boundary.
B1t RSL 68867 (25-38 cm.)	10 to 15 inches, reddish brown (5YR 4/4) clay loam, dark reddish brown (2.5YR 3/4) moist; weak medium and coarse prismatic structure; hard, friable, sticky, plastic; common very fine and fine random and medium oblique and horizontal roots; common very fine tubular and interstitial, few fine tubular pores; common thin clay films line pores; medium acid (pH 6.0); abrupt wavy boundary.
B21t RSL 68868 (38-74 cm.)	15 to 29 inches, dark red (2.5YR 3/6) clay, dark reddish brown (2.5YR 3/4) moist; moderate coarse prismatic parting to moderate coarse angular blocky structure; hard, firm, sticky, very plastic; common very fine, fine random, medium and coarse oblique roots, common medium and coarse horizontal roots at lower boundary; common very fine tubular pores; continuous moderately thick clay films on faces of peds and lining pores; medium acid (pH 6.0); clear wavy boundary.
B22t RSL 68869 (74-100 cm.)	29 to 40 inches, dark red (2.5YR 3/6) clay, dark reddish brown (2.5YR 3/4) moist; moderate coarse prismatic parting to moderate coarse angular blocky structure; hard, firm, sticky, very plastic; few very fine and fine random roots; common very fine tubular pores; continuous moderately thick clay films on faces of peds and lining pores; mica and other mineral grains apparent; medium acid (pH 6.0); clear irregular boundary.
B31t RSL 68870 (100-140 cm.)	40 to 55 inches, dark red (2.5YR 3/6) light clay, dark reddish brown (2.5YR 3/4) moist; moderate coarse angular blocky structure; hard, firm, slightly sticky and very plastic; few very fine and fine random and medium oblique roots; common very fine tubular pores; many moderately thick clay films on faces of peds and lining pores; mica and other mineral grains apparent; medium acid (pH 5.8); gradual wavy boundary.
B32t RSL 68871 (140-170 cm.)	55 to 67 inches, yellowish red (5YR 4/6) heavy clay loam, dark reddish brown (2.5YR 3/4) moist; moderate coarse angular blocky structure; hard, firm, slightly sticky, very plastic; few very fine and fine random and medium oblique roots; few very fine tubular pores; many moderately thick clay films on faces of peds and lining pores; mica and other mineral grains very apparent; medium acid (pH 5.8); gradual wavy boundary.
C1 RSL 68872 (170-205 cm.)	67 to 80 inches, variegated yellowish red, red (5YR 5/6, 2.5YR 5/6) loam, yellowish red, red (5YR 4/6, 2.5YR 4/6) moist; massive; very hard, firm, slightly sticky, slightly plastic; common thin clay films line pores and occur as bridges; slightly acid (pH 6.5); gradual wavy boundary.
C2 RSL 68873 (205-235 cm.)	80 to 92 inches, variegated yellowish red, red (5YR 5/6, 2.5YR 5/6) loam, yellowish red, red (5YR 4/6, 2.5YR 4/6) moist; massive; slightly hard, friable, slightly sticky, slightly plastic; few thin clay films line pores, many moderately thick clay films line larger pores and in fracture planes; slightly acid (pH 6.3).

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Argonaut gravelly loam **LOCATION** Amador County, Calif rnia

SOIL FAMILY: Mollic Haploxeralf, fine, montmorillonitic, thermic.

SOIL NOS. 59Calif-3-6 **LAB. NOS.** 59300 - 59305

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1	2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	
0-2	A_{11}	1.9	5.1	4.8	11.9	7.6	50.0	18.7	34.5	29.8	21	l-sil
2-6	A_{12}	2.8	4.7	4.7	11.6	8.1	46.7	21.4	32.0	29.7	20	l
6-10	B_1	2.2	4.3	4.4	11.2	8.2	44.9	24.8	31.6	28.1	15	l
10-14	B_{21}	1.8	3.8	4.2	10.2	7.2	42.3	30.5	27.7	27.9	8	cl
14-21	B_{22}	1.7	3.2	3.3	7.5	5.5	29.6	49.2	19.4	20.3	22	c
21-27+	C_2D_r	26.5	23.1	8.2	10.7	6.3	16.9	8.3	18.4	10.7	21	2/

pH 8C1a H_2O 1:1	pH 8C1c KCl 1:1	1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	"Free" Fe_2O_3 as % 6C1b	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	GYPSUM me./100g. SOIL	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.1	4.8		2.21	0.167	13.2	3.7						10.7
6.3	4.6		0.67	0.059	11.4	4.1						10.1
6.2	4.6		0.47	0.051	9.2	4.0						11.2
6.1	4.5		0.43	0.045	9.6	3.4						13.6
6.1	4.3		0.25	0.028	8.9	1.5						22.6
7.1	3.5		0.01	--	--	N.D.						8.8

5A2a CATION EXCHANGE	EXTRACTABLE CATIONS 5B1a	5B1a	6H2a	5B1a	BASE SAT. %	SATURATION EXTRACT SOLUBLE					MOISTURE AT

ARGONAUT GRAVELLY LOAM
S59Calif-3-6

Collectors: Grant Kennedy, Reuben Nelson, Bob Sketchley.
Date of Sampling: June 16, 1959. Description by: Grant Kennedy.
Location: Amador County, California, 1/2 mile west of Martel on Hwy. 88 in the SE1/4 of the NW1/4 of Section 19, T6N, R11E.
Elevation: 1,400 feet.
Relief: Gently sloping foothills with a 6 percent slope to the southwest.
Drainage: Moderately well drained; moderately rapid runoff when wet; moderately permeable over very slow.
Parent Material: Porphyritic meta-andesite.
Climate: Mean annual rainfall 27 inches.
Vegetation: Oak-grass with scrub oak, white oak, Digger pine, annual grasses, filaree, and sedge.
Remarks: This soil is a maximal Noncalcic Brown near the Brunizem soil zone. Some stones are present but mainly above the B_2. The sample site is included in a very stony mapping unit.
Classification: Mollic Haploxeralf, fine, montmorillonitic, thermic.

HORIZON		DESCRIPTION
A_{11}	0-2"	Brown (7.5YR 5/4) gravelly loam, near silt loam, dark reddish brown (5YR 3/4) when moist; very weakly platy; hard when dry, friable when moist; abundant very fine roots, few fine and medium; many fine and very fine pores; slightly acid, pH 6.1; abrupt smooth lower boundary.
A_{12}	2-6"	Yellowish red (5YR 5/6) heavy loam, yellowish red (5YR 3/6) when moist, nonsticky and slightly plastic when wet; massive; hard when dry, friable when moist; plentiful very fine, few fine and medium roots; plentiful very fine, common fine, few medium pores; occasional thin, discontinuous clay films in pores; slightly acid, pH 6.3; clear, smooth lower boundary.
B_1	6-10"	Yellowish red (5YR 4/6) heavy clay loam, yellowish red (5YR 3/6) when dry; massive; hard when dry, friable when moist, slightly sticky and plastic when wet; roots and pores as above; thin discontinuous clay films in most pores; slightly acid, pH 6.2; clear, smooth lower boundary.
B_{21}	10-14"	Similar to above but of clay texture, firm consistence, and clay films are thin and continuous in all pores; abrupt, wavy lower boundary.
B_{22}	14-21"	Brown (10YR 5/3) clay with brown (7.5YR 5/4) coatings, yellowish brown and brown (10YR 5/4 and 7.5YR 5/4) when moist; massive; very hard when dry, firm when moist, sticky and very plastic when wet; few very fine to coarse roots; few pores; thick, continuous clay films in pores; few slickensides; slightly acid, pH 6.1; abrupt, wavy lower boundary; 4 to 7 inches thick.
C_2/D_r	21-27"+	Light yellowish brown (2.5Y 6/4) deeply weathered meta-andesite with yellowish red (5YR 5/6) moderately thick, continuous clay films and occasional black (manganese?) stains along fracture planes, light olive brown (2.5Y 5/4) yellowish red (5YR 4/6) and black when moist; the weathered rock crumbles to gravelly coarse sandy loam when disturbed but becomes firmer with increasing depth; neutral, pH 7.1.

CONSECUTIVE PEDON NUMBER C 68115 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES DUBAKELLA VARIANT SOIL NO. S68 CALI. 29-36 SAMPLE NO. 8 860-8 864 NEVADA COUNTY AREA

SOIL FAMILY MOLLIC HAPLOXERALF FINE,SERPENTINITIC,MESIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SANDS VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A	INT. II .2- .02	1A2A COARSE FRAGMENTS PCT. OF WHOLE 20 I 75+	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														SOIL PCT. OF LT 75		
8 860	0- 5	A11	60.3	30.9	8.8	6.6	10.8	8.3	19.4	15.2	19.2	11.7	5.1	3.7		45.4	22A --	7	15
8 861	5- 15	A12	56.3	33.4	10.3	6.7	10.0	7.6	16.9	15.1	19.3	14.1	5.3	5.0		44.0	7A --	4	3
8 862	15- 41	B2 T	30.2	30.2	39.6	2.3	5.3	4.5	10.4	7.7	12.3	17.9	15.8	23.8		25.8	6A --	--	6
8 863	41- 66	C	37.6	28.8	33.6	3.2	7.3	6.1	13.1	7.9	9.5	19.3				24.7	13A --	1	12
8 864	0- 15	A1															5A --	1	4

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		-----PERCENT-------				PCT.NO.40		GRAMS PER CC				----PERCENT---			-PERCENT-	
8 860	0- 5	1.58	0.073			2.7	0.2			NP	NP		1.54	1.58	0.86		18.0	7.2	0.9	0.7
8 861	5- 15	0.62	0.031			2.9	0.3						1.47	1.54	0.96		18.7	7.2	1.6	1.5
8 862	15- 41	0.33	0.031			3.1	0.2			63	31		1.11	1.41	0.97		48.3	23.8	8.3	8.3
8 863	41- 66	0.20				2.7	0.2			66	37		0.76	0.86	0.96		91.0	27.8	4.2	4.2
8 864	0- 15	1.50	0.072									1.50			0.97			7.0		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACIDITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACIDITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CACL2 8C1E
		MEQ/100G									PERCENT					
8 860	0- 5	3.6	8.1	TR	0.2	11.9	4.7		16.6	15.3	72		78		5.8	5.3
8 861	5- 15	1.6	9.7	TR	0.1	11.4			11.4	13.2	100+		87		5.9	5.4
8 862	15- 41	1.3	43.1	0.1	TR	44.5			44.5	37.9	100+		100+		6.1	5.9
8 863	41- 66	1.5	44.8	0.1	TR	46.4			46.4	41.7	100+		100+		6.3	6.1
8 864	0- 15														6.0	5.5

SAMPLE NO.	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 860	0- 5	MT 4	KK 2	TL 2							MT MONTMORT	1 TRACE
8 861	5- 15										KK KAOLINIT	2 SMALL
8 862	15- 41	MT 5	KK 1								TL TALK	3 MODERATE
8 863	41- 66	MT 5	KK 1									4 ABUNDANT
8 864	0- 15											5 DOMINANT
												6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.

DUBAKELLA LOAM, SHALLOW VARIANT 1/
S68Calif-29-36

Location: Nevada County, California; 800 feet west and 750 feet north of E 1/4 corner, sec. 10, T. 16 N., R. 8 E. BLM property; dozer cut (pit) 60 feet west of dirt road and 650 feet south of Newtown Road. Date of Sampling: September 9, 1968.
Description by: L. A. Brittan and T. M. Ryan. Collectors: L. A. Brittan, G. Staidl, C. B. Goudey, T. M. Ryan and G. M. Kennedy.
Classification: Mollic Haploxeralf, fine, serpentinitic, mesic.
Vegetation: Digger pine, manzanita, yerba santa, ceanothus, live oak, blue oak, McNabb cypress, perennial and annual grasses, forbs. Not cultivated. Climate: Average annual precipitation 47 inches. Mean annual temperature 56°F. Parent Material: Ultrabasic rock. Topography: Moderately sloping, plain slope, 7 percent north northeast aspect; crest of slope. Elevation: 2,500 feet. Drainage: Well drained; runoff is medium; permeability is moderate over very slow in the subsoil. Soil Moisture: Dry to 6 inches. Very slightly moist 6 inches to 26 inches.
Remarks: pH taken by Hellige method. There is gravel pavement covering 50 to 60 percent of the surface. Of this pavement approximately 20 percent is 3/4 to 3 inches in diameter. There are few pebbles in the argillic horizon. The boundary between the A11 and A12 horizons is listed in the description as abrupt wavy; however, it is clear wavy in places. Most of these profiles are skeletal or nearly so. Vegetation is very poor. This profile was available because equipment could be gotten into the area. These profiles are usually lithic and skeletal.

HORIZON	DESCRIPTION
O1	½ to 0 inches, leaves, litter, undecomposed. (0-5 cm.)
A11 RSL 68860	0 to 2 inches, brown (7.5YR 5/4) loam, dark reddish brown (5YR 3/4) moist; moderate medium and coarse platy structure; slightly hard, friable, nonsticky and slightly plastic; common very fine roots; few very fine tubular and many very fine interstitial pores; medium acid (pH 6.0); abrupt wavy boundary. (5-15 cm.)
A12 RSL 68861	2 to 6 inches, brown (7.5YR 5/4) light loam, dark reddish brown (5YR 3/4) moist; moderate fine and medium granular structure; slightly hard, friable, slightly sticky, slightly plastic; common very fine, fine, medium and coarse roots; few very fine tubular and common very fine and fine interstitial pores; few thin clay films as bridges and as colloid stains on mineral grains; medium acid (pH 6.0); abrupt wavy boundary. (15-41 cm.)
B2t RSL 68862	6 to 16 inches, dark yellowish brown (10YR 4/4) clay, dark yellowish brown (10YR 3/4) moist; moderate medium and coarse prismatic structure; very hard, very firm, sticky, plastic; common very fine, fine, medium and coarse roots; very few very fine interstitial pores; continuous thick clay films on faces of peds and lining pores; some very fine and fine pores appear to be root channels; neutral (pH 6.7); clear wavy boundary. (41-66 cm.)
C RSL 68863	16 to 26 inches, variegated very pale brown, light brownish gray, and pale olive (10YR 7/4, 2.5Y 6/2 and 5Y 6/3) weathered basic rock, dark brown, very dark grayish brown and dark grayish brown (10YR 3/3, 2.5Y 3/2, 4/2) moist; massive; very hard, very firm; very few very fine and fine and few medium and coarse roots; common moderately thick clay films as bridges and as colloid stains on mineral grains; [illegible]
R	26 to 31 inches, slightly weathered ultrabasic rock.

1/ The Dubakella series is in a clayey-skeletal, serpentinitic, mesic family of Mollic Haploxeralfs. This pedon has less than 35 percent coarse fragments by volume.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Traver loamy fine sand **LOCATION** Riverside County, California

SOIL FAMILY: Natric Haploxeralf, coarse-loamy, mixed, thermic

SOIL NOS. S57Calif-33-14 **LAB. NOS.** 57549 - 57555

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0-6	Ap	0.3	2.9	8.3	47.5	22.1	15.5	3.4	64.8	5.0	0	lfs
6-13	A1	0.2	1.9	6.1	44.5	24.2	18.8	4.3	69.0	6.0	0	lfs
13-19	B21	0.1	1.0	4.1	39.5	23.7	20.2	11.4	65.9	7.4	0	fsl
19-24	B22	0.1	0.6	2.2	39.9	23.3	21.2	12.7	67.5	7.7	0	fsl
24-38	B23	0.1	0.6	2.5	36.1	24.3	23.6	12.8	65.8	10.0	0	fsl
38-53	Bca	0.3	0.6	1.3	22.2	21.9	37.6	16.1	59.6	18.3	0	l
53-60+	C1	0.0	0.4	1.0	21.1	27.6	35.8	14.1	65.7	15.3	0	l

8C1b pH SATURATED PASTE 1:1	CO_3 meq/l.	8C1a pH 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	6J1a * HCO_3 meq./l.	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM 25°C	6E1b $CaCO_3$ equivalent %	4A3a Bulk Density* g/cc	MOISTURE TENSIONS 4B 1/10 ATMOS. %	4B 1/3 ATMOS. %	4B2 15 ATMOS. %

TRAVER LOAMY FINE SAND
S57Calif-33-14

Location: Riverside County, California. 2200 feet east of the southwest corner of Sec. 30, T. 4 S., R. 1 W. On the Clayton Record Ranch, about 4 miles west of San Jacinto. Date of Sampling: June 6, 1957.
Description by: Fred Eggers. Collectors: George Borst, R. E. Nelson, Cliff Henry, Arnold Knecht, and Fred Eggers.
Classification: Natric Haploxeralf, coarse-loamy, mixed, thermic.
Vegetation: Barley, good stand, was pastured, weeds present including spike weed, alkali lettuce and Russian thistle. Precipitation: 12 inches. Parent Material: Granitic alluvium. Physiography: Alluvial fan, lower end approaching basin.
Elevation: 1,495 feet. Drainage: Well drained at present.

HORIZON	DESCRIPTION
S57Calif-33-14-1 Ap 0-6"	Gray (10YR 5/1 dry, very dark gray 3/1 moist) loamy fine sand; weakly defined medium subangular blocky breaks to weakly defined fine granular; soft when dry, very friable when moist; nonsticky and nonplastic when wet; moderately rapid permeability; occasional plant roots; many fine pores; pH 8.0-; noncalcareous; clear smooth lower boundary.
S57Calif-33-14-2 A1 6-13"	Gray (10YR 5/1 dry, very dark gray 3/1 moist) loamy fine sand; weakly defined medium blocky breaking to weakly defined fine granular; slightly hard when dry, very friable when moist; nonsticky and nonplastic when wet; moderately permeable; many plant roots; many fine pores; pH 8.0-; noncalcareous; a definite cultivation pan; abrupt wavy lower boundary.
S57Calif-33-14-3 B21 13-19"	Gray to dark gray (10YR 5/1-4/1 dry, very dark gray 3/1 moist) fine sandy loam; moderately defined medium angular blocky; very hard when dry, friable when moist; nonsticky and nonplastic when wet; moderate permeability; occasional plant roots; many pores; pH 8.0-; gradual smooth lower boundary.
S57Calif-33-14-4 B22 19-24"	Grayish brown (10YR 5/2 dry, dark grayish brown 4/2 moist) fine sandy loam; moderately defined medium angular blocky; very hard when dry, friable when moist; nonsticky and nonplastic when wet; occasional roots; occasional small pores; moderate permeability; pH 8.4; slight effervescence; small pockets of soft lime present; clear smooth lower boundary.
S57Calif-33-14-5 B23 24-38"	Light brownish gray (10YR 6/2 dry, dark grayish brown 4/2 moist) fine sandy loam; massive, very hard when dry, firm when moist; slightly sticky and plastic when wet; moderately permeable; few roots; occasional pores; pH 9.0; strong effervescence; lime throughout; gradual irregular lower boundary.
S57Calif-33-14-6 Bca 38-53"	Light brownish gray (10YR 6/2 dry, dark grayish brown 4/2 moist) loam; massive; slightly hard when dry, friable when moist; slightly sticky and plastic when wet; moderate permeability; no roots; very few pores; pH 9.2; violent effervescence; calcareous, nodular concretions, most of them soft; gradual irregular lower boundary.
S57Calif-33-14-7 C1 53-60"+	Light brownish gray (10YR 6/2 dry, dark grayish brown 4/2 moist) loam; massive; slightly hard when dry, friable when moist; slightly sticky and slightly plastic when wet; moderate permeability; no roots; very few pores; pH 9.2; strong to violent effervescence.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Traver loamy fine sand **LOCATION** Riverside County, California

SOIL FAMILY: Natric Haploxeralf, coarse-loamy, mixed, thermic

SOIL NOS. S57Calif-33-15 **LAB. NOS.** 57556 - 57563

DEPTH INCHES	HORIZON	1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
		PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)										
0-11	Ap	1.0	4.0	8.7	49.5	23.8	9.1	3.9	63.9	2.8	0	lfs
11-18	B21	1.0	3.0	5.0	34.2	23.6	17.2	16.0	54.8	11.0	0	fsl
18-22	B22	1.1	3.8	5.1	27.3	20.6	22.5	19.6	52.6	10.0	0	fsl
22-29	B23ca	1.4	3.5	4.7	25.0	20.1	27.9	17.4	53.2	13.0	0	fsl
29-39	Bcs	1.1	5.9	8.2	23.5	15.9	28.9	16.5	44.3	15.6	0	fsl
39-47	C1	2.6	9.7	12.7	30.9	12.9	20.6	10.6	40.9	10.3	0	fsl
47-62	C2	1.2	4.7	6.9	31.3	19.8	26.3	9.8	54.3	13.0	0	fsl
62-70+	C3	1.6	4.1	4.6	19.6	18.1	39.7	12.3	51.4	20.4	0	l

8C1b pH SATURATED PASTE 1:1	6J1a HCO_3 meq/l.	8C1a pH 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	6I1a CO_3 meq./l.	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	4A3a* Bulk Density g/cc	MOISTURE TENSIONS 4B1a 1/10 ATMOS. %	4B1a 1/3 ATMOS. %	4B2 15 ATMOS. %
7.0	3.5	7.4	0.25	0.025	10.0	0.0	0.8	0	1.45	10.1	4.9	2.0
7.8	1.5	8.5	0.19	0.028		0.7	6.0	0	1.78			7.0
7.8	1.2	9.0	0.12	0.018		0.7	10.0	Tr	1.62			8.9
7.9	1.7	9.5	0.10			0.0	12.5	<1	1.46			8.6
7.9	0.8	8.2	0.09			0.0	16.0	2.3	1.50			8.0
8.2	1.3	9.5	0.06			0.0	14.0	1.9	1.51			5.5
8.3	1.2	9.7	0.06			0.0	11.0	3.1	1.51			5.6
8.3	1.4	9.5	0.10			0.0	9.5	3.9	1.47			7.4

5A2a CATION EXCHANGE CAPACITY ←(Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	5B1b Na 6P2a	5B1b K 6Q2a	5D1 EXCH. Na %	SATURATION EXTRACT SOLUBLE 6P1a Na	6Q1a K	6N1a	6O1a	6K1a	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
5.0	2.2	2.1	0.5	0.3	0.4	6	3.1	1.0	1.9	1.5	1.1	29.9
12.6	4.4	4.2	0.4	4.0	0.3	32	44.5	0.2	1.9	2.2	17.0	36.3
17.5	7.0	16.9	0.3	7.1	0.3	41	91.0	0.3	5.1	5.5	28.3	43.7
18.8	12.2	5.3		8.6	0.4	46	122.0	0.1	8.4	9.2	31.2	44.0
17.3	24.0	30.0		9.6	0.3	55	167.0	0.3	22.4	17.2	31.3	40.3
12.7	15.1	3.9		7.0	0.3	55	133.0	0.2	11.1	10.8	23.4	33.1
12.9	15.2	4.0		6.9	0.1	53	105.0	0.1	3.0	4.5	21.4	38.8
15.8	13.2	5.0		9.9	0.2	63	91.0	0.1	2.0	3.6	19.9	48.5

* no gypsum found

TRAVER LOAMY FINE SAND
S57Calif-33-15

Location: Riverside County, California. 650 feet west and 1,600 feet south of the northeast corner of Sec. 9, T. 4 S., R. 2 W. Along fence 300 feet north of road. On the Mott Bros. Ranch, about 2 miles east of Lakeview. Date of Sampling: June 7, 1957.
Description by: Fred Eggers. Collectors: George Borst, R. E. Nelson, Cliff Henry, Arnold Knecht, and Fred Eggers.
Classification: Natric Haploxeralf, coarse-loamy, mixed, thermic.
Vegetation: Fallow. Has been in dry farmed grain. Precipitation: 12 inches. Parent material: Granitic alluvium. Physiography: Lower part of alluvial fan approaching basin. Elevation: 1,435 feet. Drainage: Well drained at present.

HORIZON	DESCRIPTION
Ap RSL No. 57556	0-11 inches, gray to grayish brown (10YR 5/1-5/2 dry, dark gray to dark grayish brown 4/1-4/2 moist) loamy fine sand; weakly defined fine subangular blocky breaking to single grain; soft when dry, very friable when moist; nonsticky and nonplastic when wet; moderately rapid permeability; many roots; many fine pores; pH 8.0; noncalcareous; abrupt smooth lower boundary.
B21 RSL No. 57557	11-18 inches, light olive brown (2.5Y 5/4 dry, olive brown 4/3 moist) loam; strongly defined medium angular blocky breaking to moderately defined fine granular; hard when dry, firm when moist; slightly sticky and plastic when wet; moderate permeability; many roots; many pores; pH 8.4; noncalcareous; weak fragmentary clay skins; clear wavy lower boundary.
B22 RSL No. 57558	18-22 inches, light yellowish brown (2.5Y 6/3, dark grayish brown 4/2 moist) loam; strongly defined medium angular blocky breaking to moderately defined fine granular; hard when dry, firm when moist; slightly sticky and plastic when wet; many roots; many fine pores; pH 8.4; noncalcareous; prominent fragmentary clay skins; clear wavy lower boundary.
B23ca RSL No. 57559	22-29 inches, grayish brown (2.5Y 5/2 dry, dark grayish brown 4/2 moist) loam; moderately defined medium subangular blocky breaking to weakly defined fine granular; slightly hard when dry, firm when moist; slightly sticky and plastic when wet; moderate permeability; occasional roots; many pores; pH 8.4; violent effervescence; gradual wavy lower boundary.
Bcs RSL No. 57560	29-39 inches, light brownish gray (2.5Y 6/2 dry, dark grayish brown 4/2 moist) loam; massive; slightly hard when dry, firm when moist; slightly sticky and plastic when wet; moderate permeability; occasional roots; many pores; pH 8.2; strong effervescence; gypsum in seams; gradual wavy lower boundary.
C1 RSL No. 57561	39-47 inches, light olive gray (5Y 6/2 dry, olive gray 4/2 moist) loam; massive; slightly hard when dry, firm when moist; slightly sticky and plastic when wet; moderate permeability; few roots; occasional pores; pH 8.4; violent effervescence; gradual wavy lower boundary.
C2 RSL No. 57562	47-62 inches, light olive gray (5Y 6/2 dry, olive gray 4/2 moist with slight mottling) very fine sandy loam; massive; slightly hard when dry, friable when moist; nonsticky and slightly plastic when wet; moderate permeability; few roots; occasional pores; pH 8.6; violent effervescence; clear wavy lower boundary.
C3 RSL No. 57563	62-70 inches +, light gray to pale yellow (5Y 7/2-7/3 dry, olive gray to olive 5/2-5/3 moist) silt loam (very fine sandy loam with silt lenses); stratified; slightly hard when dry, friable when moist; slightly sticky and plastic when wet; no roots; occasional pores; pH 8.6; violent effervescence.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Natric Haploxeralf, fine-loamy, mixed, thermic.

SOIL Bonsall sandy loam (taxadjunct) SOIL Nos. S64Calif-37-5 LOCATION San Diego County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 6440 - 6449

Depth (In.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	* Clay <.002	** Clay <.002	1/ Coarse fragments 1A2a: > 2 Pct.	2 - 19	19 - 76
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm														Pct. of < 76 mm		
0-10	Ap	60.2	30.7	9.1	7.2	13.4	9.1	17.8	12.7	16.8	13.9	38.9	47.5	8.4	8.1	3	3	0
10-15	A2	58.9	30.3	10.8	8.1	13.2	8.6	16.8	12.2	16.4	13.9	37.9	46.7	10.7	10.4	3	3	0
15-20	B1	57.7	28.2	14.1	9.2	12.6	8.5	15.9	11.5	15.3	12.9	35.3	46.2	14.6	13.9	5	5	0
20-29	B21t	52.3	23.3	24.4	9.6	12.1	7.8	13.7	9.1	11.0	12.3	27.2	43.2	25.6	23.7	8	8	0
29-41	B22t	52.8	24.5	22.7	8.8	11.6	7.5	14.9	10.0	11.8	12.7	30.1	42.8	22.4	21.5	4	4	0
41-59	B23t	59.4	24.2	16.4	11.3	12.4	7.4	16.1	12.2	11.8	12.4	33.4	47.2	17.2	15.3	11	11	0
59-73	B24t	55.8	26.3	17.9	12.0	12.0	7.0	14.2	10.6	12.1	14.2	30.4	45.2	21.1	18.3	11	11	0
73-85	B25t	48.7	29.7	21.6	7.7	10.4	6.6	13.3	10.7	13.2	16.5	31.6	38.0	22.3	20.6	8	8	0
85-92	B3	45.8	33.2	21.0	9.4	10.3	5.8	11.3	9.0	12.2	21.0	27.5	36.8	21.8	18.7	14	14	0
92-100+	C	52.5	33.6	13.9	15.7	11.7	5.7	11.0	8.4	13.8	19.8	28.3	44.1	15.3	11.0	28	28	0

Depth (In.)	6A1a * Organic carbon Pct.	6B1a * Nitrogen Pct	C/N	* 6C2a Ext. Iron as Fe Pct.	6E1b ** Carbonate as $CaCO_3$ Pct.		Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 C'	Water content: Pct.	4B1c 1/3 bar Pct.	4B2* 15 bar Pct	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-10	0.94	0.080	12	0.8			1.53	1.54		0.98		11.5	4.7	0.002	0.002	6.7	6.8	
10-15	0.26			0.8			1.59	1.60		0.98		13.0	5.2	0.002	0.002	7.0	7.2	
15-20	0.15			1.1			1.78	1.83		0.97		12.9	6.4	0.009	0.009	6.7	6.9	
20-29	0.16			1.2			1.75	1.85		0.95		16.2	10.6	0.019	0.018	6.8	7.1	
29-41	0.06			1.1									10.2			7.6	8.0	
41-59	0.04			1.0			1.66	1.71		0.93		21.9	8.5	0.010	0.009	7.8	8.1	
59-73	0.03			0.9									10.2			7.8	8.3	
73-85	0.04			1.1	<1		1.77	1.85		0.95		16.8	11.7	0.015	0.014	7.7	8.1	
85-92	0.01			1.2									13.5			7.6	8.0	
92-100+	0.01			1.5			1.65	1.71		0.81		19.5	12.0	0.012	0.010	7.7	8.1	

Depth (In.)	Extractable bases 6N2a * Ca	6O2a* Mg	6P2a* Na	6Q2a* K	5B1a Sum of bases	6H2a* Ext. Acidity	Cation Exch. Capacity: 5A2a* NaOAc	5A3a* Sum	Water extract from saturated paste 8A1 **: 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-10	5.0	2.2	0.2	0.3	7.7	1.6	9.7	9.3	2.9	2.2	1.5	0.3		-	5.4	0.3		0.58
10-15	5.3	2.4	0.4	0.1	8.2	1.4	10.7	9.6	1.4	1.0	3.1	tr.		-	2.8	1.0		0.52
15-20	5.9	3.4	0.8	0.1	10.2	1.9	12.6	12.1	1.8	1.8	5.6	0.1		-	1.2	5.4		1.00
20-29	8.7	6.0	1.3	0.1	16.1	2.3	19.6	18.4	1.5	2.6	6.5	0.1		-	1.0	5.9		1.11
29-41	10.2	6.4	1.6	0.1	18.3	2.0	19.8	20.3	2.5	2.3	7.9	0.1		-	3.2	6.8		1.33
41-59	9.7	5.8	1.9	0.1	17.5	1.0	19.0	18.5	2.3	2.3	10.4	tr.		-	2.9	9.2		1.55
59-73	13.4	7.5	3.7	0.1	24.7	1.2	25.3	25.9	1.9	1.6	13.5	tr.		-	2.9	11.0		1.78
73-85	14.4	8.4	4.5	0.1	27.4	1.0	27.1	28.4	1.8	1.7	14.1	0.1		-	2.6	11.6		1.90
85-92	17.6	9.7	6.6	0.1	34.0	1.4	34.7	35.4	1.2	1.4	13.4	tr.		-	2.0	11.6		1.74
92-100+	18.1	8.8	7.1	0.1	34.1	1.2	33.6	35.3	1.0	0.8	11.1	tr.		-	1.8	9.2		1.46

	8A * Mineral	5D2	5F	Base	Base	Clay mineralogy (<0.002 mm) 7A			

1/ From characterization sample; determined

BONSALL SANDY LOAM (taxadjunct) 1/
S64Calif-37-5

Location: San Diego County, California. A soil profile pit located 1.5 miles south and 1.5 miles east of Fallbrook in the SW1/4 of the NW1/4 of section 29, T. 9 S., R. 3 W. Date of Sampling: February 19, 1964.
Description By: David C. Estrada and Gerald Kester. Collectors: Klaus Flach, Gerald Kester, George Borst, Clifford Henry, Roy Bowman, and Gerald Anderson.
Classification: Natric Haploxeralf, fine-loamy, mixed, thermic.
Vegetation: Idle at present (few annual weeds and grasses). Climate: The annual rainfall is 15 inches. The mean annual temperature is 60° F., the mean January temperature 54° F., and the mean July temperature 70° F. Parent Material: Tonalite grus. Topography: East-facing, smooth concave 7 percent slopes in lower backslope positions on rolling upland. Elevation: 665 feet above sea level. Drainage: Moderately well drained; very slowly permeable. Soil Moisture: Profile dry.
Remarks: The parent material is deeply weathered. In the B23t and B25t horizons the roots are concentrated in the ped faces and are predominantly very fine. Manganese staining increases from the B23t to B25t horizon.

HORIZON	DESCRIPTION
Ap	0 to 10 inches, brown (10YR 5/3) sandy loam, dark brown (10YR 3/3) moist; moderate

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Natric Haploxeralf, fine-loamy, mixed, thermic.
SOIL Redding sandy loam (taxadjunct) SOIL Nos. S66Calif-37-4 LOCATION San Diego County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66250 - 66258

Depth (In)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)			1/ Coarse fragments 1A2a > 2 Pct.	2-19	19-76
		Pct. of < 2 mm															Pct of < 76mm	
0- 4	A1	56.0	38.7	5.3	2.2	11.6	17.0	18.6	6.6	20.2	18.5	34.5	49.4	0	5	18	12	6 2/
4-10	A2	53.6	41.5	4.9	2.3	10.6	16.1	18.4	6.2	19.1	22.4	32.8	47.4	0	5	15	11	4 2/
10-17	B11	46.6	35.6	17.8	1.7	8.9	13.6	16.1	6.3	16.4	19.2	29.4	40.3	0	18	19	14	5 2/
17-30	B12	42.9	33.0	24.1	1.6	7.7	12.6	15.0	6.0	14.8	18.2	27.0	36.9	0	24	15	11	4 2/
30-36	B13	39.9	31.5	28.6	1.6	6.8	11.6	14.1	5.8	14.3	17.2	26.0	34.1	0	29	25	13	6 3/
36-45	B2t	27.5	20.9	51.6	1.9	6.0	7.6	8.5	3.5	8.8	12.1	15.9	24.0	0	52	13	5	0 3/
45-49	B31sim	53.5	12.1	34.4	15.1	16.1	9.0	9.9	3.4	5.8	6.3	13.4	50.1	0	34	40	35	0 3/
49-59	B32sim	85.2	8.2	6.6	24.3	25.1	14.8	16.1	4.9	6.1	2.1	17.6	80.3	0	7	44	39	0 3/
59-72	IIB33sim	86.7	9.4	3.9	13.9	18.5	23.8	24.5	6.0	6.7	2.7	20.9	80.7	0	4	73	15	58 4/

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct	Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content: Pct	4B1c 1/3 bar Pct.	4B2 15 bar Pct	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1a H_2O 1:1		
0- 4	2.30					1.29	1.31		0.90		10.7	4.1	0.007	0.007	5.3		
4-10	0.26					1.81	1.81		0.89		9.0	2.0	0.000	0.000	5.5		
10-17	0.29					1.79	1.86		0.86		12.9	6.5	0.012	0.011	4.6		
17-30	0.25					1.71	1.78		0.90		15.1	8.6	0.014	0.012	4.5		
30-36	0.28					1.68	1.77		0.83		17.0	10.9	0.018	0.015	4.3		
36-45	0.30					1.53	1.87		0.92		25.7	20.1	0.070	0.064	4.1		
45-49	0.24					1.36	1.48		0.74		29.8	18.4	0.029	0.021	4.2		
49-59	0.13					1.43	1.43		0.71		20.8	9.8	0.001	0.000	4.5		
59-72	0.10														4.5		

Depth (In.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1a NH_4OAc	5A3a Sum	6G1d KCl Ext. Al^{+++}	Water extract from saturated paste 8A1: 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K	6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1b SO_4	8A1a Electrical conductivity
	meq/100 g									meq/liter								mmho/cm
0- 4	3.6	0.9	0.2	0.4	5.1	5.0	7.9	10.1	0.2	2.4	1.6	1.0	0.4	-	2.4	1.5		0.52
4-10	1.2	0.8	0.3	0.1	2.4	1.3	3.0	3.7	0.1	1.0	0.4	3.0	0.1	-	1.0	1.0		0.39
10-17	1.9	3.3	0.8	0.2	6.2	4.9	9.6	11.1	1.3	0.4	0.3	3.8	tr.	-	0.4	1.0		0.33
17-30	2.4	4.9	2.4	0.2	9.9	5.1	12.2	15.0	0.9	1.0	2.2	19.4	0.1	-	0.2	17.3		2.43
30-36	2.9	6.5	4.7	0.2	14.3	6.8	14.9	21.1	0.7	4.5	11.6	53.0	0.2	-	0.4	61.6		6.2
36-45	5.6	5.9	7.9	0.3	19.7	9.0	27.9	28.7	1.1	7.4	20.4	69.0	0.3	-	1.9	85.9	10.6	8.2
45-49	5.9	6.6	10.3	0.3	23.1	10.4	35.2	33.5	1.5	8.9	23.2	86.5	0.3	-	0.6	103	11.6	10.3
49-59	4.3	9.8	7.2	0.2	21.5	7.9	28.7	29.4	2.2	3.1	5.7	52.5	0.2	-	0.7	51.8	8.8	5.67
59-72	4.2	10.2	6.6	0.2	21.2	8.1	25.4	29.3	2.2	1.6	3.7	39.4	0.2	-	0.7	37.4		4.68

Depth (In)	8A Water at Saturation Pct.	5D2 Exchangeable Na Pct.	Clay mineralogy (<0.002 mm.) 7A: Mont.	Verm.	Mica	Kaolin.	Base saturation: 5C3 Sum Pct.	5C1 NH_4OAc Pct.
			X-ray					
0- 4	30.8	2.5					50	65
4-10	12.4	10.0	t	-	xx	xx	65	80
10-17	21.4	7.3					56	65
17-30	31.3	14.8	x	-	-	xx	66	81
30-36	39.8	17.4					68	96
36-45	66.7	11.8	xxx	-	-	xxx	69	71
45-49	50.7	17.0	xxx	-	-	xx	69	66
49-59	41.7	17.4					73	75
59-72	39.9	19.6	xxxx	-	-	x	72	83

1/ From characterization sample (5 kg)(except as noted).
2/ 2-19 mm. from characterization sample, 19-76 mm. from field weight (>50 kg).
3/ 2-19 mm. from characterization sample, 19-76 mm. from field weight; >2 mm. includes volume estimate of >76 mm. fraction.
4/ Calculation of % >2 mm. of RSL 66258 differs from that of the horizons above only in that a 1/3 bar Bd of 1.92 g./cc. was assumed.

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

REDDING SANDY LOAM (taxadjunct) 1/
S66Calif-37-4

Location: San Diego County, California; Miramar Transect No. 2, station 9.0; 100 feet east and 50 feet north of the southwest corner of section 19, T. 15 S., R. 2 W. Date of sampling: June 14, 1966.
Description by: George Borst, Roy Bowman, David Estrada, and Gerald Kester. Samples collected by: Leo Klameth and Dennis Nettleton.
Classification: Natric Haploxeralf, fine-loamy, mixed, thermic.
Vegetation: Sparse chamise and flattop buckwheat. Climate: Weather record for San Diego, near the sampling site, is mean winter air temperature 55° F., mean summer air temperature 68° F., mean annual precipitation 10 inches. Parent Material: Stream deposit over Pleistocene marine terrace.
Topography: Flat marine terrace of less than 1 percent slope. Elevation: About 425 to 450 feet.
Drainage: Moderately well drained; runoff is medium; permeability is very slow. Soil Moisture: Dry.

HORIZON	DESCRIPTION
A1 RSL No. 66250	0 to 4 inches, yellowish brown (10YR 5/4) sandy loam, brown to dark brown (10YR 4/3) when moist; weak fine granular structure; loose, very friable, slightly sticky, nonplastic; plentiful fine and medium roots; many very fine interstitial and common fine tubular pores; strongly acid (pH 5.3); abrupt smooth boundary.
A2 RSL No. 66251	4 to 10 inches, light gray (10YR 7/2) fine sandy loam, brown (10YR 5/3) when moist; weak medium subangular blocky structure; slightly hard, friable, slightly sticky, nonplastic; few fine to medium roots; common very fine and fine tubular pores; strongly acid (pH 5.3); [redacted]
B11 RSL No. 66252	10 to 17 inches, brown (7.5YR 5/4) sandy clay loam, brown to dark brown (7.5YR 4/4) when moist; moderate medium angular blocky structure; hard, friable, sticky, slightly plastic; few fine to medium roots; few very fine and fine tubular pores; common thin clay skins in pores; very strongly acid (pH 4.8); abrupt wavy boundary.
B12 RSL No. 66253	17 to 30 inches, brown (7.5YR 5/4) sandy clay loam, brown to dark brown (7.5YR 4/2) when moist; moderate medium angular blocky structure; hard, friable, sticky, slightly plastic; very few fine roots; few very fine and fine tubular pores; common thin clay skins in pores; very strongly acid (pH 4.5); gradual smooth lower boundary.
B13 RSL No. 66254	30 to 36 inches, brown to dark brown (7.5YR 4/4) heavy clay loam, brown to dark brown (7.5YR 4/2) when moist; moderate medium angular blocky structure; hard, friable, sticky, plastic; very few fine roots; few very fine and fine tubular pores; common thin clay skins in pores; very strongly acid (pH 4.5); abrupt smooth boundary.
B2t RSL No. 66255	36 to 45 inches, brown to dark brown (7.5YR 4/2) clay, dark brown (7.5YR 3/2) when moist; strong medium to coarse angular blocky structure; very hard, firm, sticky, plastic; very few fine roots; many fine interstitial and very few fine tubular pores; extremely acid (pH 4.2); abrupt wavy boundary.
B31sim RSL No. 66256	45 to 49 inches, grayish brown (10YR 5/2) ped exteriors and brown to dark brown (7.5YR 4/2) ped interiors both of which are of clay texture and dark gray (10YR 4/1) and brown to dark brown (7.5YR 4/2) when moist; moderate medium to coarse angular blocky structure; hard to extremely hard, firm to extremely firm, sticky, plastic; very few fine roots; very few fine tubular pores; many thin to moderately thick clay skins on ped faces and in pores; extremely acid (pH 4.2); abrupt irregular boundary.
B32sim RSL No. 66257	49 to 59 inches, grayish brown (10YR 5/2) ped exteriors and dark brown (7.5YR 3/2) ped interiors of clay texture, dark gray (10YR 4/1) and dark brown (7.5YR 3/2) respectively when moist; moderate coarse platy structure; strongly indurated ped interiors; very few fine roots; very few fine tubular pores; common thick clay skins on ped faces; clear smooth boundary.
IIB33sim RSL No.	59 to 72 inches +, color pattern same as the B32sim; massive; weakly indurated; very common fine interstitial pores; common moderately thick clay skins on ped faces;

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Natric Haploxeralf, fine, mixed, thermic

SOIL San Miguel sandy loam variant SOIL Nos. S66Calif-37-5 LOCATION San Diego County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66259 - 66260

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: (0.05-0.02)	Silt: Int. III (0.02-0.002)	Silt: Int. II (0.2-0.02)	Silt: (2-0.1)			1/ Coarse fragments 1A2a > 2 Pct.	2-19 (Pct. of < 76mm)	19-76 (Pct. of < 76mm)	
		Pct. of < 2 mm (Size class and particle diameter (mm) 3A1)																	
35-45	B21t	25.8	26.1	48.1	0.9	4.6	7.7	9.0	3.6	10.8	15.3	18.0	22.2	0	48	3	2	1	2/
45-57	B22t	27.4	28.3	44.3	1.0	4.8	7.9	9.7	4.0	12.0	16.3	19.7	23.4	0	44	3	3	0	2/

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content Pct.	4B1c 1/3 bar Pct	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1a H_2O 1:1		
35-45	0.35						1.62	1.86		0.98		22.5	16.8	0.047	0.046	4.2		
45-57	0.21						1.54	1.79		0.98		24.5	16.8	0.052	0.051	4.1		

Depth (in.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1a NH_4OAc	5A3a Sum		6G1d KCl Ext. Al^{+++}						Base saturation: 5C3 Sum	5C1 NH_4OAc
	meq/100 g															Pct.	Pct.
35-45	4.2	4.5	6.8	0.3	15.8	8.5	22.7	24.3		1.2						65	70
45-57	4.6	4.8	8.0	0.2	17.6	8.5	24.2	26.1		1.1						67	73

Depth (in.)	Clay mineralogy (<0.002 mm) 7A: Mont.	Verm.	Mica	Kaolin.
	X-ray			
35-45	xx	x	-	xxx
45-57	xxx	-	-	xx

1/ From characterization sample (5Kg) (except as noted).
2/ 2-19 mm. from characterization sample, 19-76 mm. from field weight (>50 Kg).

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

SAN MIGUEL SANDY LOAM variant 1/
S66Calif-37-5

Location: San Diego County, California; Miramar Transect No. 2, station 21.1; 100 feet east and 50 feet north of the southwest corner of section 19, T. 15 S., R. 2 W. Date of sampling: June 14, 1966.

Description by: George Borst, Roy Bowman, David Estrada, and Gerald Kester. Samples collected by:

Classification: Natric Haploxeralf; fine, mixed, thermic family.

Vegetation: Sparse chamise and flattop buckwheat. Climate: Weather record for San Diego, near the sampling site, is mean winter air temperature 55° F., mean summer air temperature 68° F., mean annual precipitation 10 inches. Parent Material: Stream deposit over Pleistocene marine terrace. Topography: Flat marine terrace of less than 1 percent slope. Elevation: About 425 to 450 feet. Drainage: Moderately well drained; runoff is medium; permeability is very slow. Soil Moisture: Dry.

HORIZON	DESCRIPTION
A1	0 to 3 inches, grayish brown (10YR 5/2) sandy loam, very dark grayish brown (10YR 3/2) when moist; weak fine granular structure; slightly hard, very friable, slightly sticky, nonplastic; plentiful very fine to fine roots; common fine tubular and very fine interstitial pores; medium acid (pH 5.9); abrupt smooth boundary.
A21	3 to 14 inches, pale brown (10YR 6/3) loam, brown to dark brown (10YR 4/3) when moist; massive; hard, very friable, slightly sticky, slightly plastic; few fine to medium roots; common fine tubular and many fine vesicular pores; medium acid (pH 5.8); clear smooth boundary.
A22	14 to 24 inches, very pale brown (10YR 7/3) loam, brown (10YR 5/3) when moist; massive; hard, very friable, slightly sticky, slightly plastic; few fine to medium roots; common fine tubular and many fine vesicular pores; very strongly acid (pH 4.8); gradual smooth boundary.
B1	24 to 35 inches, pinkish gray (7.5YR 6/2) and pale brown (10YR 6/3) clay loam, dark grayish brown (10YR 4/2) when moist; massive; very hard, friable, slightly sticky, slightly plastic; few fine roots; few very fine tubular pores; common thin clay skins in pores; very strongly acid (pH 4.8); clear smooth boundary.
B21t RSL No. 66259	35 to 45 inches, dark brown to brown (7.5YR 4/2) clay, dark brown (7.5YR 3/2) when moist; weak medium prismatic breaking to coarse angular blocky structure; very hard, firm, sticky, plastic; few fine roots; few very fine tubular pores; common thin to thick clay skins on ped faces; very strongly acid (pH 4.5); gradual smooth boundary.
B22t RSL No. 66260	45 to 57 inches, grayish brown (10YR 5/2) clay, dark grayish brown (10YR 4/2) when moist; weak medium to coarse angular blocky structure; extremely hard, firm, sticky, plastic; very few fine roots; very few very fine tubular pores; very strongly acid (pH 4.5); gradual smooth boundary.
B23t	57 to 92 inches, gray (10YR 5/1) clay, dark gray (10YR 4/1) when moist; weak coarse angular blocky structure; extremely hard, firm, sticky, plastic; very few fine roots; very few very fine tubular pores; very strongly acid (pH 4.5); clear smooth boundary.
B3	92 to 98 inches +, grayish brown (10YR 5/2) clay loam to sandy clay, dark grayish brown (10YR 4/2) when moist; massive; very hard, friable, slightly sticky, slightly plastic;

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Psammentic Haploxeralf, coarse-loamy, mixed, thermic

SOIL Garey sandy loam SOIL Nos. S62Calif-42-3 LOCATION Santa Barbara County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 6286 - 6294

Depth (In.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: (0.05-0.02)	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)			Coarse fragments 2A2: > 2 Pct.	2-19	19-76
		Pct. of < 2 mm															Pct. of < 76mm	
0-4	Ap1	51.2	42.1	6.7	0.0	5.3	24.3	12.2	9.4	27.6	14.5	40.4	41.8			0		
4-8	Ap2	48.9	43.2	7.9	0.0	5.5	23.7	11.9	7.8	28.0	15.2	39.0	41.1			0		
8-16	A13	47.5	42.5	10.0	0.1	4.6	23.2	11.8	7.8	26.4	16.1	37.5	39.7			0		
16-27	A3	47.7	42.6	9.7	0.1	5.0	22.8	11.9	7.9	26.7	15.9	37.9	39.8			0		
27-36	B21	53.0	39.1	7.9	0.0	5.4	27.3	13.7	6.6	24.0	15.1	34.3	46.4			0		
36-47	B22	52.0	40.3	7.7	0.0	6.1	23.4	16.7	5.8	28.0	12.3	38.2	46.2			0		
47-58	B31	68.8	25.0	6.2	0.0	7.8	38.2	18.5	4.3	14.7	10.3	23.6	64.5			0		
58-70	B32	75.4	19.9	4.7	0.0	8.3	40.3	22.4	4.4	13.0	6.9	26.7	71.0			0		
70-83+	C	84.4	12.3	3.3	0.1	9.2	48.5	23.5	3.1	7.2	5.1	16.1	81.3			0		

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.	6C2a Ext. Iron as Fe Pct.	Bulk density: 4A1f 1/3 bar clods g/cc	4A1h Oven-dry clods g/cc	g/cc		Water content: Pct.	4B1c 1/3 bar clods Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.		pH: 8C1a H_2O 1:1	8C1c 1 N KCl 1:1	(1:1)
0-4	0.43	0.048	9			0.6	1.42	1.44				10.6	3.2	0.003		5.9	4.3	
4-8	0.34					0.7	1.64	1.64				10.4	3.5	0.000		5.5	3.8	
8-16	0.25					0.8	1.49	1.50				10.0	4.2	0.003		5.6	3.9	
16-27	0.16					0.8	1.58	1.61				12.1	4.1	0.007		6.2	4.4	
27-36	0.08					0.7	1.64	1.66				10.1	3.5	0.003		6.5	4.7	
36-47	0.03					0.7	1.85	1.89				9.3	3.6	0.007		6.8	5.0	
47-58	0.01					0.6		1.85*					3.0			6.3	4.6	
58-70	0.01					0.4							2.3			7.0	5.0	
70-83+	0.00					0.3	1.61	1.61				3.5	1.5	0.007		7.3	5.1	

Depth (In.)	Extractable bases 5B1a: 6N2b Ca	6O2c Mg	6P2a Na	6Q2a K		6H2a Ext. Acidity	Cation Exch. Capacity 5A3a Sum									Base saturation 5C3 Sum Pct.	Pct.
	meq/100 g																
0-4	2.1	0.9	0.1	0.2		2.9	6.2									59	
4-8	1.8	0.8	0.2	0.2		3.6	6.6									45	
8-16	2.6	1.1	0.2	0.1		3.3	7.3									55	
16-27	2.8	1.8	0.4	0.1		2.2	7.3									70	
27-36	2.4	2.2	0.4	0.1		1.3	6.4									80	
36-47	2.3	2.3	0.6	0.1		0.6	5.9									90	
47-58	1.5	1.7	0.4	Tr.		0.6	4.2									86	
58-70	1.3	1.2	0.4	Tr.		0.4	3.3									88	
70-83+	0.9	0.8	0.3	Tr.		0.2	2.2									91	

Depth (In.)				

* Lamella only

GAREY SANDY LOAM
S62Calif-42-3

Location: Santa Barbara County, California. SW¼ Sec. 9, T.9N., R.33W.; 5 miles east of Orcutt on Clark Road; 10,600 feet east of Highway 101, 600 feet north, 40 feet west. Date of Sampling: March 14, 1962

Description by: L. C. Leifer. Collectors: R. E. Nelson, B. R. Brasher, G. E. Shipman, C. J. Hilliker, L. C. Leifer.

Classification: Psammentic Haploxeralf, coarse-loamy, mixed, thermic.

Vegetation: Abandoned alfalfa field; annual weeds and sparse stand of alfalfa. Climate: Precipitation 14-15 inches. Mean annual temperature 57°F., January 50°F., July 63°F. Parent Material: Aeolian sandy material. Topography: Rolling terrace-like, locally slightly convex 6 percent south slope. Elevation: 740 feet. Drainage: Well drained. Runoff is medium. A horizons have moderately rapid and B horizons moderately slow to slow permeability. Soil Moisture: Moist to 72 inches, dry below.

Remarks: Field determinations of pH by Hellige-Truog triplex indicator. Paired site sample numbers are Garey S62Calif-42-4-1 through 9.

HORIZON	DESCRIPTION
Ap1 RSL 6286	0 to 4 inches, brown (10YR 5/3) sandy loam, dark brown (10YR 4/3) when moist; weak fine subangular blocky and weak medium granular structure; slightly hard, very friable, nonsticky and slightly plastic; abundant very fine roots; many very fine interstitial pores; medium acid (pH 5.7); clear smooth boundary.
Ap2 RSL 6287	4 to 8 inches, brown (10YR 5/3) sandy loam, dark brown (7.5YR 4/3) when moist; slightly finer than above; massive; slightly hard, friable, slightly sticky and slightly plastic; few very fine roots with abundant very fine roots in joints; many very fine interstitial pores; very few fine and medium tubular pores; strongly acid (pH 5.4); clear wavy boundary.
A13 RSL 6288	8 to 16 inches, brown (10YR 5/3) sandy loam, dark brown (7.5YR 3/3) when moist; weak fine and medium subangular blocky structure; slightly hard, friable, slightly sticky and slightly plastic; very few very fine roots with plentiful very fine roots in joints, very few woody roots 3 to 10 mm; many very fine interstitial and common fine and very few medium tubular pores; strongly acid (pH 5.4); gradual smooth boundary.
A3 RSL 6289	16 to 27 inches, light brown (7.5YR 6/3) sandy loam, dark brown (7.5YR 3/3) when moist; slightly more clay than above; massive; hard, friable, slightly sticky and slightly plastic; very few very fine roots, mostly in joints; many very fine interstitial and common fine and very few medium tubular pores, common thin clay films as bridges and in pores; medium acid (pH 6.0); gradual wavy boundary.
B21 RSL 6290	27 to 36 inches, light brown (7.5YR 6/3) heavy sandy loam, dark brown (7.5YR 3/3) when moist; massive; hard, firm, slightly sticky and slightly plastic; very few very fine roots; many very fine interstitial, common fine and very fine and very few medium tubular pores; common thin clay films as bridges in pores; indistinct dark reddish brown (5YR 3/3m) lamellae 1/2 inch thick, slightly brittle; medium acid (pH 6.0); gradual smooth boundary.
B22 RSL 6291	36 to 47 inches, light brown (7.5YR 6/4) sandy loam, reddish brown (5YR 4/4) when moist; massive; hard, soil mass friable, lamellae firm, slightly sticky and slightly plastic; very few very fine roots; many very fine interstitial and common very fine and very few medium tubular pores; common thin clay films as bridges and in pores; several indistinct discontinuous dark reddish brown (5YR 3/2 3/3m) lamellae that are weakly brittle; slightly acid (pH 6.2); clear wavy boundary.
B31 RSL 6292	47 to 58 inches, pinkish gray (7.5YR 7/2) loamy sand, dark brown (7.5YR 4/3) when moist; massive; slightly hard, very friable, nonsticky and nonplastic; lamellae are very hard, dry, firm when moist; very few very fine roots, many very fine interstitial pores with very few very fine tubular pores in lamellae only; distinct reddish brown (5YR 5/3d) dark reddish brown (5YR 3/3) when moist continuous lamellae about 1/2 inch thick; slightly acid (pH 6.2); clear wavy boundary.
B32 RSL 6293	58 to 70 inches, pinkish gray (7.5YR 7/3) loamy sand, brown (7.5YR 5/4) when moist; massive; slightly hard, very friable, nonsticky and nonplastic; no roots; many very fine interstitial pores; no clay films except thin weak bridges in bands; lamellae 1/4 to 1/2 inch thick, continuous and wavy, thinner in lower part of horizon, similar to those in horizon above; medium acid (pH 6.0); clear wavy boundary.
C RSL 6294	70 to 83 inches +, very pale brown (10YR 7/3) sand, brown (7.5YR 5/4) when moist; massive; soft, very friable, nonsticky and nonplastic; no roots; many very fine interstitial pores; weak lamellae 1/4 to 1/2 inch thick, 5 to 7 inches apart, continuous and wavy, reddish brown (5YR 4/3d) a few pendants of lamellae 1 to 1½ inches long; medium acid (pH 6.0), lamellae essentially same reaction as other soil of same horizon.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Psammentic Haploxeralf, coarse-loamy, mixed, thermic

SOIL Garey sandy loam SOIL Nos. S62Calif-42-4 LOCATION Santa Barbara County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 6295 - 62103

Depth (in.)	Horizon	1B1b Total Sand (2–0.05)	Total Silt (0.05–0.002)	Total Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	3A1 Int. II (0.2–0.02)	(2–0.1)			Coarse fragments 2A2 > 2 Pct.	2 – 19	19 – 76
		Pct. of < 2 mm															Pct. of < 76mm	
0–8	Ap	52.8	40.8	6.4	0.0	6.3	25.7	13.5	7.3	29.1	11.7	39.8	45.5			0		
8–19	A12	51.9	39.7	8.4	0.0	5.9	26.0	12.8	7.2	26.6	13.1	37.2	44.7			0		
19–27	A3	51.4	40.7	7.9	0.0	5.8	26.4	13.0	6.2	28.2	12.5	37.9	45.2			0		
27–33	B21	54.9	37.6	7.5	0.0	5.5	27.7	13.8	7.9	24.8	12.8	36.0	47.0			0		
33–47	B22	60.3	33.3	6.4	0.1	6.0	32.2	16.4	5.6	22.5	10.8	32.3	54.7			0		
47–54	B31	69.4	25.4	5.2	0.1	6.9	38.5	18.9	5.0	16.2	9.2	25.5	64.4			0		
54–64	B32	75.6	21.1	3.3	0.1	7.3	42.8	21.0	4.4	14.0	7.1	23.0	71.2			0		
64–75	C1	79.8	16.5	3.7	0.1	7.1	45.6	23.6	3.4	10.8	5.7	19.4	76.4			0		
75–96+	C2	85.5	11.3	3.2	0.0	8.3	51.5	23.2	2.5	7.6	3.7	14.7	83.0			0		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.		Bulk density: 4A3a Field moist cores g/cc	4A1h Oven-dry clods g/cc	g/cc		Water content: 4B1d 1/10 bar Pct.	4B1d 1/3 bar Pct.	4B2 15 bar Pct.			pH: 8C1a H_2O 1:1	8C1c 1 N KCL 1:1	(1:1)
0–8	0.54	0.046	12				1.60				14.7	9.2	2.8			5.8	4.3	
8–19	0.24						1.52				13.3	9.2	3.4			6.1	4.1	
19–27	0.17						1.46				13.2	9.1	3.3			6.7	4.8	
27–33	0.13						1.66				13.1	9.4	3.4			7.0	5.1	
33–47	0.02						1.87				12.1	9.0	3.0			7.3	5.5	
47–54	0.00						1.82				10.0	7.4	2.4			7.3	5.0	
54–64	0.00						1.82				9.6	6.2	1.8			7.1	5.0	
64–75	0.00						1.72				8.2	5.2	1.9			7.1	5.3	
75–96+	0.00						1.50	1.82*			5.9	3.7	1.9			7.2	5.2	

Depth (in.)	Extractable bases 5B1a: 6N2b Ca	602c Mg	6P2a Na	6Q2a K		6H2a Ext. Acidity	Cation Exch. Capacity 5A3a Sum									Base saturation 5C3 Sum Pct.	Pct.
	meq/100 g																
0–8	1.8	0.8	0.2	0.3		2.7	5.8									53	
8–19	2.9	1.1	0.2	0.1		2.2	6.5									66	
19–27	2.9	1.5	0.2	0.1		1.4	6.1									77	
27–33	2.3	1.9	0.4	0.1		1.0	5.7									82	
33–47	1.9	2.2	0.5	Tr.		0.4	5.0									92	
47–54	1.4	1.6	0.4	Tr.		0.3	3.7									92	
54–64	1.0	1.2	0.3	Tr.		0.2	2.7									93	
64–75	1.2	1.1	0.4	Tr.		0.2	2.9									93	
75–96+	1.0	1.0	0.4	Tr.		0.2	2.6									92	

Depth (in.)				

* lamella

GAREY SANDY LOAM
S62Calif-42-4

Location: Santa Barbara County, California. SW¼ Sec. 10, T9N, R33W; 1400 feet north on Dominion Road from Clark Avenue, 2000 feet east and north along road to oil well and 100 feet northwest into field; 6 miles east of Orcutt. Date of Sampling: March 14, 1962.
Description by: L. C. Leifer. Collectors: R. E. Nelson, B. R. Brasher, G. E. Shipman, C. J. Hilliker, L. C. Leifer.
Classification: **Psammentic Haploxeralf, coarse-loamy, mixed, thermic.**
Vegetation: Not cultivated for several years, used for range; small annual weeds, nit grass (Festuca megalora) and annual lupine. Climate: Precipitation 14-15 inches. Mean annual temperature 57°F., January 50°F., July 60°F. Parent Material: Aeolian sandy material. Topography: Low rolling terrace-like, locally plane 5 percent east slope. Elevation 600 feet. Drainage: Well drained. Runoff is medium. A horizons have moderately rapid permeability and B horizons have moderately slow to slow. Soil Moisture: Moist to 85 inches, dry below. A3 more moist than A12; soil above firm lamellae of B31, B32 and C1 is particularly moist.
Remarks: Field determinations of pH by Hellige-Truog triplex indicator. Paired site sample numbers are Garey S62Calif-42-3-1 through 9.

HORIZON	DESCRIPTION
Ap RSL 6295	0 to 8 inches, brown (10YR 5/3) sandy loam, dark brown (7.5YR 3/2) when moist; few thin yellowish brown (10YR 5/4m) plates due to tillage; massive; hard, firm, nonsticky, slightly plastic; plentiful very fine roots tend to be concentrated on joints; many very fine interstitial, few very fine and fine tubular pores; strongly acid (pH 5.2); clear smooth boundary.
A12 RSL 6296	8 to 19 inches, brown (10YR 5/3) sandy loam, dark brown (7.5YR 3/2) when moist; massive; hard, friable, slightly sticky and slightly plastic; plentiful very fine roots; many very fine interstitial and many fine and few medium tubular pores; strongly acid (pH 5.4); gradual wavy boundary.
A3 RSL 6297	19 to 27 inches, brown (7.5YR 5/3) sandy loam, darker brown (7.5YR 4/4) when moist; massive; hard, very friable, slightly sticky and slightly plastic; plentiful very fine roots; many very fine interstitial,many fine and few medium tubular pores; very few thin clay films in bridges; medium acid (pH 5.9); gradual wavy boundary.
B21 RSL 6298	27 to 33 inches, light brown (7.5YR 6/3) sandy loam, slightly finer than above, reddish brown (5YR 4/3) when moist; a few coarse indistinct reddish brown (5YR 5/3d) mottles; massive; very hard, friable, slightly sticky and slightly plastic; few very fine roots; many very fine interstitial, common fine and very few medium tubular pores; few thin clay films as bridges; several indistinct blotches similar to A1 material; slightly acid (pH 6.1); clear smooth boundary.
B22 RSL 6299	33 to 47 inches, light brown (7.5YR 6/3) slightly heavy sandy loam, reddish brown (5YR 4/3) when moist; a few indistinct darker lamellae 1/2 inch thick discontinuous, dark reddish brown (5YR 3/3m); massive; very hard, firm, somewhat brittle, slightly sticky and slightly plastic; few very fine roots; many very fine interstitial pores, many very fine few fine and very few medium tubular pores; few thin clay films in bridges; slightly acid (pH 6.5); clear wavy boundary, lower boundary of succeeding horizons coincides with a lamella. Otherwise boundaries are gradual.
B31 RSL 62100	47 to 54 inches, light brown (7.5YR 6/3) light sandy loam, reddish brown (5YR 4/3) when moist; massive; hard, very friable, nonsticky and nonplastic; very few very fine roots; many very fine interstitial, common very fine and very few fine tubular pores; no clay films except in lamellae; a prominent lamella bottom of horizon 1 to 1½ inch thick has common very fine and few fine tubular pores in addition to interstitial pores; some tubular pores have continuous moderately thick clay films, all of the lamella has thin or very thin clay films in bridges; slightly acid (pH 6.3); clear wavy boundary.
B32 RSL 62101	54 to 64 inches, pinkish gray (7.5YR 7/2) loamy sand, reddish brown (5YR 5/3) when moist; massive; slightly hard, very friable, nonsticky and nonplastic; no roots; many very fine interstitial and few fine tubular pores; bottom of horizon has lamellae 1 to 3½ inch thick, few tubular pores in addition to interstitial,most prominent lamella in profile, firm brittle when moist; slightly acid (pH 6.3); clear wavy boundary.

CONSECUTIVE PEDON NUMBER C 68 76 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES JABU SHALLOW VARIANT SOIL NO. S68 CALI. 9-29 SAMPLE NO. 8 614-8 626 TAHOE BASIN AREA

SOIL FAMILY ULTIC HAPLOXERALF COARSE-LOAMY,MIXED,FRIGID RIVERSIDE SOIL SURVEY LABORATORY

SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SANDS VCS 2-1	CS 1-.5	MS .5-.25	FS .25-.10	VFS .1-.05	SILTS CSI .05-.02	FSI .02-.002	CLAYS CO. 2-.2U	FINE LT .2U	CARBONATE 3A1A.02	INT. II .2-75	COARSE FRAGMENTS PCT. OF WHOLE SOIL	I 75-20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														1A2A	PCT. OF LT 75		
																	--A	--	--	--
8 614	3- 0	01																		
8 615	0- 4	A11	72.4	18.1	9.5	9.1	17.7	13.3	23.0	9.3	7.3	10.8				28.8	15A	--	15Z	N
8 616	4- 13	A12	74.9	16.2	8.9	9.5	17.7	14.2	23.7	9.8	6.3	9.9				28.9	17A	--	17Z	N
8 617	13- 33	C 1	74.0	16.4	9.6	6.6	17.1	14.8	24.8	10.7	7.0	9.4				31.1	16A	3	13Z	N
8 618	33- 56	C 2	74.8	15.5	9.7	11.9	16.2	13.3	23.6	9.8	6.1	9.4				28.0	12A	--	12Z	N
8 619	56- 76	2B1 TXB	68.7	18.3	13.0	8.1	16.1	12.8	22.8	8.9	6.8	11.5				27.2	10A	--	10Z	N
8 620	76-105	2B2 TXB	73.1	13.2	13.7	6.9	18.0	14.8	25.3	8.1	4.7	8.5				25.3	16A	--	16Z	N
8 621	105-125	2B31TXB	75.4	11.1	13.5	7.3	18.2	15.3	25.9	8.7	4.4	6.7				26.0	19A	--	19Z	N
8 622	125-140	2B32TXB	75.0	11.5	13.5	7.8	18.0	15.0	25.7	8.5	4.6	6.9				26.0	17A	--	17Z	N
8 623	140-165	2B33TXB	74.1	13.0	12.9	6.4	17.8	16.3	26.0	7.6	5.1	7.9				25.2	16A	--	16Z	N
8 624	165-195	2B34TB	76.0	11.1	12.9	8.3	17.1	15.2	26.3	9.1	4.1	7.0				26.2	20A	--	20Z	N
8 625	195-210	2C B	80.4	9.3	10.3	14.5	25.0	14.9	19.2	6.8	3.7	5.6				19.5	26A	--	26Z	TR
8 626	0- 13	A															19A	--	19Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A -PERCENT-	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B -PERCENT-	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B -PERCENT-	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LIMIT LL 4F2 PCT.NO.40	PL 4F1	BULK DENSITY EST. 1/3 4A1F GRAMS PER CC	DRY BAR 4A1F	CM 4A1H	WATER CONTENT 1/10 BAR 3B2	1/3 BAR 4B1C -PERCENT-	15 BAR 4B2	EXTNSBLTY LEF 4D1 -PERCENT-	LE 4D1
8 614	3- 0											0.40		1.00					
8 615	0- 4	6.56	0.355			1.1							0.70	0.70	0.96		14.0		
8 616	4- 13	1.94	0.097			1.3				NP	NP		1.10	1.10	0.92	11.7	6.3		
8 617	13- 33	0.66	0.042			1.2							1.20	1.21	0.92	11.5	5.5	0.3	0.3
8 618	33- 56	0.53				1.4				NP	NP		1.34	1.35	0.94	11.3	5.3	0.2	0.2
8 619	56- 76	0.21				1.2							1.61	1.64	0.94	15.2	4.2	0.6	0.6
8 620	76-105	0.14				1.1				NP	NP		1.58	1.61	0.90	13.2	6.2	0.6	0.6
8 621	105-125	0.11				1.1							2.14	2.22	0.84	10.3	5.8	1.2	1.0
8 622	125-140	0.13				1.1							1.52	1.54	0.89	12.4	6.3	0.4	0.4
8 623	140-165	0.13				1.1							1.52	1.54	0.90	13.4	5.9	0.4	0.4
8 624	165-195	0.15				1.0							1.54	1.54	0.87	12.3	6.3		
8 625	195-210	0.06				0.9				NP	NP		1.41	1.41	0.84	13.8	4.8		
8 626	0- 13	2.18	0.097									1.05			0.91		5.7		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	CEC SUM 5A3A	NH4-DAC 5A6A	BASE SUM+ ACI-DITY	SATURATION SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 614	3- 0															
8 615	0- 4	21.2	1.8	TR	0.5	23.5	14.8		38.3	38.1	61		62		5.9	5.5
8 616	4- 13	5.6	0.5	TR	0.2	6.3	8.4		14.7	12.7	43		50		6.2	5.5
8 617	13- 33	3.6	0.4	TR	0.2	4.2	6.3		10.5	8.3	40		51		6.3	5.6
8 618	33- 56	3.5	0.4	0.1	0.2	4.2	6.7		10.9	8.3	39		51		6.2	5.4
8 619	56- 76	4.4	0.7	0.1	0.2	5.4	3.9		9.3	7.9	58		68		6.2	5.6
8 620	76-105	4.5	0.7	0.1	0.2	5.5			5.5	7.4	100+		74		6.1	5.5
8 621	105-125	4.2	0.8	0.1	0.2	5.3	3.1		8.4	7.5	63		71		6.2	5.7
8 622	125-140	3.8	0.8	0.1	0.1	4.8	3.3		8.1	6.9	59		70		6.2	5.7
8 623	140-165	4.5	0.8	0.1	0.1	5.5	3.9		9.4	7.7	59		71		6.1	5.6
8 624	165-195	4.0	0.7	0.1	0.1	4.9	3.3		8.2	7.0	60		70		6.1	5.5
8 625	195-210	3.4	0.6	0.1	0.1	4.2			4.2	5.9	100+		71		6.1	5.6
8 626	0- 13	5.6	0.7	TR	0.3	6.6	9.0		15.6	12.7	42		52		6.1	5.5

CLAY MINERALOGY

SAMPLE NO	DEPTH CM.	LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM MINERAL X RAY 7A2E I	7A2E II	7A2E III	CODE		RELATIVE AMOUNT
8 614	3- 0												1 TRACE
8 615	0- 4										VR	VERMICLT	2 SMALL
8 616	4- 13	VR 3	KK 3								KK	KAOLINIT	3 MODERATE
8 617	13- 33												4 ABUNDANT
8 618	33- 56												5 DOMINANT
8 619	56- 76												6 INDETER.
8 620	76-105												
8 621	105-125												
8 622	125-140												
8 623	140-165												
8 624	165-195												
8 625	195-210	KK 2	VR 1										
8 626	0- 13												

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

JABU COARSE SANDY LOAM, SHALLOW VARIANT
S68Calif-9-29

Location: Tahoe Basin, El Dorado County, California, 1/2 mile northeast of the southwest corner, sec. 21, T. 12 N., R. 18 E., in swale 10 feet west of dirt road along moraine crest, 1.3 miles northeast of Meyers.
Date of Sampling: August 9, 1968. Description by: L. N. Langan. Collectors: J. H. Rogers, G. M. Kennedy, L. I. Larsen, D. Palmquist, B. R. Brasher, W. D. Nettleton.
Classification: Ultic Haploxeralf, coarse-loamy, mixed, frigid.*
Vegetation: Not cultivated; Jeffrey pine, white fir, mountain whitethorn, greenleaf manzanita. Climate: Precipitation is 30 to 35 inches. Mean annual temperature is 45° F., January 27° F., July 61° F. Parent Material: Mixed morainal material, dominantly granitic. Topography: Swale on crest of lateral moraine, pre-Tahoe? Two percent slope. Elevation: 6,500 feet. Drainage: Moderately well drained; runoff is slow; permeability is moderately rapid over slow. Soil Moisture: Dry.
Remarks: Field determination for pH by Hellige-Truog triplex indicator. Composite A taken from 10 satellite samples. C2-IIB1txb horizon contact contains a stoneline with cobbles and stones. *Fragixeralf group not provided for in current classification system. Fragipan properties not strongly and clearly expressed.

HORIZON	DESCRIPTION
O1 RSL 68614	1 to 0 inches (3-0 cm.), pine litter.
A11 RSL 68615	0 to 1½ inches (0-4 cm.), dark grayish brown (10YR 4/2) silt loam, very dark brown (10YR 2/2) moist; moderate fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine roots; many very fine and fine interstitial pores; 5 percent 3/8 inch gravel; strongly acid (pH 5.5); clear wavy boundary.
A12 RSL 68616	1½ to 5 inches (4-13 cm.), brown (10YR 5/3) light sandy loam, dark brown (7.5YR 3/2) moist; dark grayish brown (10YR 4/2) spots; moderate very fine and fine granular structure; soft, very friable, nonsticky and nonplastic; many very fine, fine and medium roots; many very fine and fine interstitial pores; 5 percent 3/8 inch gravel; medium acid (pH 5.8); clear wavy boundary.
C1 RSL 68617	5 to 13 inches (13-33 cm.), pale brown (10YR 6/3) light sandy loam, brown (7.5YR 4/3) moist; weak fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine, fine, medium and coarse roots; many very fine and fine interstitial pores; 5 percent 3/8 inch gravel; medium acid (pH 6.0); clear wavy boundary.
C2 RSL 68618	13-22 inches (33-56 cm.), pale brown (10YR 6/3) loamy sand, brown (7.5YR 4/3) moist; massive; soft, very friable, nonsticky and nonplastic; many very fine, medium and coarse, common fine roots; many very fine and fine interstitial pores; 5 percent 3/8 inch gravel; medium acid (pH 5.8); abrupt smooth boundary.
IIB1txb RSL 68619	22 to 30 inches (56-76 cm.), pale brown and light yellowish brown (10YR 6/3, 6/4) sandy loam, dark yellowish brown (10YR 4/4) moist; massive; very hard, very friable, nonsticky and slightly plastic; common very fine, few fine and medium roots; few fine tubular and common very fine interstitial pores; common thin clay films in pores and bridges and many thin clay films coating sand grains; many 1/8 to 3/8 inch vertical and horizontal light gray (10YR 7/2) seams, brown (10YR 4/3) moist, containing common clean sand grains and many silt bridges; 10 percent gravel; strongly acid (pH 5.5); clear wavy boundary.
IIB2txb RSL 68620	30 to 41 inches (76-105 cm.), pale brown and light yellowish brown (10YR 6/3, 6/4) sandy loam, dark yellowish brown (10YR 4/4) moist; massive; very hard, very friable, nonsticky and slightly plastic; common very fine, few fine and medium roots; few fine tubular and common very fine interstitial pores; few moderately thick, many thin clay films in bridges and many thin clay films coating sand grains; common 1/8 to 3/8 inch vertical and horizontal light gray (10YR 7/2) seams, brown (10YR 4/3) moist, containing common clean sand grains and many silt bridges; 10 percent gravel; strongly acid (pH 5.5); clear wavy boundary.
IIB31txb RSL 68621	41 to 49 inches (105-125 cm.), pale brown and light yellowish brown (10YR 6/3, 6/4) sandy loam, dark yellowish brown (10YR 4/4) moist; massive; hard, very friable, nonsticky and slightly plastic; few very fine, fine, medium and coarse roots; many very fine, common fine interstitial pores; common thin clay films in bridges and many thin clay films coating sand grains; common 1/8 to 3/8 inch vertical and horizontal light gray (10YR 7/2) seams, brown (10YR 4/3) moist, containing many clean sand grains and thin silt bridges; 10 percent ½ inch minus gravel; strongly acid (pH 5.5); gradual wavy boundary.
IIB32txb RSL 68622	49 to 56 inches (125-140 cm.), pale brown and light yellowish brown (10YR 6/3, 6/4) sandy loam, dark yellowish brown (10YR 4/4) moist; massive; slightly hard, very friable, nonsticky and slightly plastic; very few fine, medium and coarse roots; many very fine, few fine interstitial pores; common thin clay films in bridges and many thin clay films coating sand grains; common 1/8 to 3/8 inch vertical and horizontal light gray (10YR 7/2) seams, brown (10YR 4/3) moist, containing many clean sand grains and thin silt coatings; 10 percent ½ inch minus gravel; strongly acid (pH 5.5); gradual wavy boundary.
IIB33txb RSL 68623	56 to 64 inches (140-165 cm.), pale brown and light yellowish brown (10YR 6/3, 6/4) sandy loam, dark yellowish brown (10YR 4/4) moist; common fine distinct dark brown (7.5YR 3/3) moist mottles; massive; slightly hard, very friable, nonsticky and slightly plastic; very few fine, medium and coarse roots; many very fine, few fine interstitial pores; common thin clay films in bridges and many thin clay films coating sand grains; few 1/8 to 3/8 inch vertical and horizontal light gray (10YR 7/2) seams; brown (10YR 4/3) moist; 10 percent ½ inch minus gravel; strongly acid (pH 5.5); gradual wavy boundary.
IIB34tb RSL 68624	64 to 77 inches (165-195 cm.), very pale brown (10YR 7/4) light sandy loam, dark yellowish brown (10YR 4/4) moist; common fine distinct brownish yellow (10YR 6/6) mottles; massive; slightly hard, very friable, nonsticky and nonplastic; very few fine and medium roots; many very fine, common fine interstitial pores; few thin clay films in bridges and common thin clay film coatings; 10 percent ½ inch minus gravel; medium acid (pH 5.8); clear wavy boundary.
IICb RSL 68625	77 to 82 inches (195-210 cm.), white (10YR 8/2) loamy sand, dark yellowish brown (10YR 4/4) moist; massive; soft, very friable, nonsticky and nonplastic; very few fine and medium roots; many very fine and fine interstitial pores; few fine and medium dark brown and very dark brown (7.5YR 3/3, 2/2) moist, iron and manganese stains on gravel; 5 percent ½ inch minus gravel; medium acid (pH 6.0).

CONSECUTIVE PEDON NUMBER C 68 77 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES JABU SHALLOW VARIANT SOIL NO. S68 CALI. 9-30 SAMPLE NO. 8 627-8 642 TAHOE BASIN AREA

SOIL FAMILY ULTIC HAPLOXERALF COARSE-LOAMY,MIXED,FRIGID RIVERSIDE SOIL SURVEY LABORATORY

SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 1A2A

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SANDS VCS 2-1	CS 1- .5	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A.02	INT. II .2- OF WHOLE 20 SOIL	COARSE FRAGMENTS I PCT. 75-	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM													PCT. OF LT 75			
8 627	3- 0	01														--A	--	--	--
8 628	0- 8	A11	74.7	19.1	6.2	14.6	20.8	12.5	18.3	8.5	7.2	11.9			25.3	16A	--	16Z	N
8 629	8- 20	A12	71.3	18.9	9.8	9.5	18.5	13.1	20.0	10.2	8.9	10.0			29.8	16A	--	16Z	N
8 630	20- 28	B11TX	70.7	19.1	10.2	8.7	18.0	13.2	20.2	10.6	8.6	10.5			30.1	11A	--	11Z	N
8 631	28- 36	B12TX	72.9	17.2	9.9	16.3	19.2	12.1	16.8	8.5	7.6	9.6			24.9	18A	--	18Z	N
8 632	36- 48	B13TX	69.3	19.8	10.9	11.2	18.3	12.3	18.8	8.7	8.7	11.1			27.5	11A	--	11Z	N
8 633	48- 74	B14TX	62.3	20.6	17.1	7.0	12.8	11.0	20.3	11.2	9.3	11.3			31.8	9A	--	9Z	N
8 634	74-110	B21TX	67.5	16.9	15.6	11.3	16.2	12.1	18.9	9.0	7.0	9.9			26.1	12A	--	12Z	N
8 635	110-165	B22TX	75.4	11.8	12.8	13.9	19.1	13.1	20.4	8.9	6.1	5.7			25.6	25A	--	25Z	N
8 636	165-190	B31TX	76.8	12.0	11.2	16.8	21.3	13.7	18.1	6.9	5.5	6.5			20.8	33A	--	33Z	N
8 637	190-230	B32TX	76.1	15.5	8.4	16.1	20.2	13.1	18.0	8.7	7.1	8.4			24.5	18A	--	18Z	N
8 638	230-290	2C 1X	77.1	20.1	2.8	12.9	18.1	12.9	22.0	11.2	9.9	10.2			32.6	11A	--	11Z	N
8 639	290-445	2C 2X	81.5	14.0	4.5	16.7	22.2	13.5	20.5	8.6	6.8	7.2			25.5	16A	--	16Z	N
8 640	445-495	2C 3	79.8	14.5	5.7	19.1	20.8	9.9	20.0	10.0	7.4	7.1			28.3	8A	--	8Z	N
8 641	495-510	2C 4	80.6	14.5	4.9	15.4	21.7	13.8	21.5	8.2	7.0	7.5			25.5	12A	--	12Z	N
8 642	0- 20	A1														13A	--	13Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR-BON 6A1A	NITRO-GEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	DRY 4A1F	1/3 BAR 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		PERCENT		PERCENT		PERCENT				PCT.NO.40		GRAMS PER CC				PERCENT			PERCENT	
8 627	3- 0											0.40		1.00						
8 628	0- 8	2.58	0.092			1.2							0.94	0.98	0.94	12.5	6.3	1.4	1.3	
8 629	8- 20	1.43	0.045			1.4				NP	NP		1.36	1.36	0.91	10.8	4.4			
8 630	20- 28	1.05				1.4							1.47	1.47	0.93	12.6	2.8			
8 631	28- 36	0.97				1.4				NP	NP		1.46	1.47	0.89	12.8	4.6	0.2	0.2	
8 632	36- 48	0.27				1.5							1.76	1.76	0.92	12.0	4.7			
8 633	48- 74	0.19				1.5							1.72	1.72	0.94	11.5	7.3			
8 634	74-110	0.18				1.2				22	18		1.86	1.91	0.91	13.2	6.8	0.9	0.8	
8 635	110-165	0.12				1.4							1.71	1.78	0.82	16.9	6.4	1.3	1.1	
8 636	165-190	0.09				1.5							1.78	1.82	0.76	13.7	3.6	0.7	0.6	
8 637	190-230					0.9							1.70	1.74	0.87	13.9	4.8	0.8	0.7	
8 638	230-290					0.5							1.74	1.75	0.93	12.9	1.5	0.2	0.2	
8 639	290-445					0.5							1.74	1.74	0.89	13.5	2.0			
8 640	445-495					1.0							1.78	1.78	0.94	12.0	3.7			
8 641	495-510					0.7				NP	NP	1.78			0.92		3.2			
8 642	0- 20	1.80	0.065									1.20			0.94		5.1			

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 584A 6N2E	MG 584A 6O2D	NA 584A 6P2A	K 584A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACI-DITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 627	3- 0															
8 628	0- 8						10.8	0.8	10.8						5.2	4.6
8 629	8- 20						8.3	0.8	8.3						5.6	5.0
8 630	20- 28	2.4	0.2	TR	0.2	2.8	7.5		10.3	8.3	28		34		5.9	5.1
8 631	28- 36	2.0	0.3	0.1	0.2	2.6	8.3		10.9	8.6	24		30		5.8	5.1
8 632	36- 48	2.6	0.4	0.1	0.2	3.3	4.1		7.4	6.7	45		49		5.7	5.2
8 633	48- 74	5.2	1.1	0.1	0.1	6.5			6.5	8.2	100+		79		5.7	5.2
8 634	74-110	4.4	1.0	0.1	0.1	5.6			5.6	7.6	100+		74		5.7	5.2
8 635	110-165	6.7	1.5	0.1	TR	8.3	3.3		11.6	10.3	72		81		5.6	5.1
8 636	165-190	4.9	1.1	0.1	TR	6.1			6.1	7.9	100+		78		5.7	5.3
8 638	230-290	2.9	0.7	0.1	TR	3.7			3.7	4.0	100+		93		6.7	6.3
8 639	290-445	3.8	1.0	0.1	TR	4.9			4.9	5.3	100+		93		6.8	6.2
8 640	445-495	4.7	1.3	0.1	0.1	6.2			6.2	6.7	100+		93		6.7	6.1
8 641	495-510	4.1	1.1	0.1	TR	5.3			5.3	6.0	100+		89		6.7	6.2
8 642	0- 20														5.9	5.1

CLAY MINERALOGY

SAMPLE NO	DEPTH CM.	LESS THAN 0.002 MM X RAY I 7A2E	II 7A2E	III 7A2E	IV 7A2E	DTA -PERCENT- I 7A3	II 7A3	L.T. 0.0002 MM X RAY I 7A2E	II 7A2E	III 7A2E
8 627	3- 0									
8 628	0- 8									
8 629	8- 20	VR	3	KK	3					
8 630	20- 28									
8 631	28- 36	VR	3	KK	3					
8 632	36- 48									
8 633	48- 74									
8 634	74-110	MI	3	KK	3	VM	2			
8 635	110-165									
8 636	165-190									
8 637	190-230									
8 638	230-290	MI	3	KK	2					
8 639	290-445									
8 640	445-495									
8 641	495-510	MI	3	KK	2	VR	1			
8 642	0- 20									

MINERAL CODE		RELATIVE AMOUNT
		1 TRACE
VR	VERMICLT	2 SMALL
KK	KAOLINIT	3 MODERATE
MI	MICA	4 ABUNDANT
VM	VM/MICA	5 DOMINANT
		6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED

JABU COARSE SANDY LOAM, SHALLOW VARIANT
S68Calif-9-30

Location: Tahoe Basin, El Dorado County, California; 2,700 feet east and 200 feet south of northwest corner sec. 28, T. 12 N., R. 18 E.; 100 feet south of dirt road extension of Oneidas Street, 1.3 miles northeast of Meyers, 10 feet east of sample site S67Calif-9-13. Date of Sampling: August 15, 1968.
Description by: L. N. Langan. Collectors: J. H. Rogers, G. M. Kennedy, L. C. Leifer, L. I. Larsen, D. Palmquist, B. R. Brasher, W. D. Nettleton.
Classification: Ultic Haploxeralf, coarse-loamy, mixed, frigid.*
Vegetation: Not cultivated. Jeffrey pine, lodgepole pine, squaw carpet. Climate: Precipitation is 35 inches. Mean annual temperature is 42° F., January 27° F., July 60° F. Parent Material: Mixed, dominantly granitic alluvium. Topography: Glacial terrace outwash, 1 percent northwest slope. Elevation: 6,420 feet. Drainage: Moderately well drained; runoff is slow; permeability is very slow. Soil Moisture: Moist below 75 inches.
Remarks: Field determinations for pH by Hellige-Truog triplex indicator. Composite A taken from 10 satellite samples. All bleached seams are soft and very friable. *Fragixeralfs group not provided for in current classification system. Fragipan properties not strongly and clearly expressed.

HORIZON	DESCRIPTION
01	1-0 inches (3-0 cm.), pine needle litter.
RSL 68627	
A11	0-3 inches (0-8 cm.), brown (7.5YR 5/4) coarse sandy loam, dark brown (7.5YR 3/2) moist; weak fine and medium platy [illegible]
68628	and fine tubular pores; 8 percent ¼ inch minus gravel; medium acid (pH 5.7); abrupt wavy boundary.
A12 RSL 68629	3-8 inches (8-20 cm.), brown (7.5YR 5/4) coarse sandy loam, dark brown (7.5YR 3/2) moist; weak medium and coarse subangular blocky structure; soft, very friable, nonsticky and nonplastic; common very fine, few fine and medium roots; very few very fine interstitial, very few fine tubular pores; 5 percent 1 inch minus gravel; medium acid (pH 5.8); clear wavy boundary.
B11t RSL 68630	8-11 inches (20-28 cm.), reddish yellow (7.5YR 6/6) coarse sandy loam, reddish brown (5YR 4/4) moist; weak medium and coarse subangular blocky structure; soft, very friable, nonsticky and nonplastic; many fine, medium and coarse roots; very few medium interstitial, very few fine tubular pores; few thin clay films in bridges and pores; 10 percent 1 inch minus gravel; medium acid (pH 5.8); abrupt wavy boundary.
B12tx RSL 68631	11-14 inches (28-36 cm.), reddish yellow (7.5YR 6/6) coarse sandy loam, reddish brown (5YR 4/4) moist; 1/8 inch white (10YR 8/2) seams; massive; hard, very friable, nonsticky and nonplastic; many fine, medium and coarse roots; very few very fine interstitial and fine tubular pores; many strong brown (7.5YR 5/6) thin clay films in bridges and pores; 10 percent 1 inch minus gravel; medium acid (pH 5.6); abrupt wavy boundary.
B13tx RSL 68632	14-19 inches (36-48 cm.), light brown (7.5YR 6/3) coarse sandy loam, brown (7.5YR 4/4) moist; strong brown (7.5YR 5/6) spots, few very pale brown (10YR 7/3) seams, moist; massive; hard, friable, nonsticky and nonplastic; few very fine, fine and medium roots; many very fine and few fine interstitial pores; common very thin, few thin clay films in bridges; 10 percent gravel; medium acid (pH 5.6); clear wavy boundary.
B14tx RSL 68633	19-29 inches (48-74 cm.), light yellowish brown (10YR 6/4) sandy loam, strong brown (7.5YR 4/6) moist; very pale brown (10YR 8/3) seams; massive; hard, friable, sticky and plastic; few very fine and fine roots; common very fine, few fine interstitial pores; many strong brown and reddish yellow (7.5YR 5/6, 6/6) thin and very few moderately thick clay films in bridges; few stones; 10 percent gravel; medium acid (pH 5.8; abrupt wavy boundary.
B21tx RSL 68634	29-44 inches (74-110 cm.), pale brown (10YR 6/3) coarse sandy loam, brown (10YR 4/3) moist; very pale brown (10YR 7/3) seams and few fine and medium reddish yellow (7.5YR 6/8) iron stains; massive; very hard, friable, sticky and plastic; few very fine and fine roots; common very fine, few fine interstitial pores; many thin, common moderately thick brown (7.5YR 4/4) clay films as bridges; 10 percent gravel; medium acid (pH 5.8); clear wavy boundary.
B22tx RSL 68635	44-64 inches (110-165 cm.), strong brown (7.5YR 5/6) crushed, gravelly coarse sandy loam, dark yellowish brown(10YR 4/4) moist; many 1/8 to 1/4 inch white (10YR 8/2) bleached and unoriented seams, few coarse prominent yellowish red and reddish brown (5YR 5/8, 4/4) iron stains moist; massive; very hard, friable, slightly sticky and plastic; few fine and

CONSECUTIVE PEDON NUMBER C 68 79 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES JABU SOIL NO. S68 CALI. 9-34 SAMPLE NO. 8 652-8 663 TAHOE BASIN AREA

SOIL FAMILY ULTIC HAPLOXERALF COARSE-LOAMY,MIXED,FRIGID RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 VCS 2-1	CS 1-.05	MS .5-.25	FS .25-.10	VFS .1-.05	CSI .05-.02	FSI .02-.002	CO. 2-.2U	FINE LT .2U	CARBO NATE 3A1A	INT. II .2-.02	1A2A -COARSE FRAGMENTS- PCT. OF WHOLE 20 I 75-	II 20-5	III 5-2
						PERCENT OF LESS THAN 2MM											SOIL PCT. OF LT 75		
8 652	0- 5	A1	77.7	14.5	7.8	17.5	19.8	12.4	19.5	8.5	6.9	7.6				25.7	13A --	13Z	N
8 653	5- 15	A3	80.6	12.3	7.1	31.5	18.4	9.7	15.2	5.8	5.4	6.9				19.1	14A --	14Z	N
8 654	15- 25	B1 T	77.4	13.4	9.2	15.9	22.0	12.4	18.7	8.4	6.3	7.1				24.5	15A --	15Z	N
8 655	25- 51	B2 T	77.0	12.8	10.2	11.4	21.3	13.6	22.1	8.6	5.8	7.0				25.0	15A --	15Z	N
8 656	51- 74	2B21TB	78.8	10.6	10.6	12.6	17.9	12.8	26.4	9.1	4.7	5.9				27.4	27A 8	19Z	N
8 657	74- 94	2B22TB	81.1	7.6	11.3	9.5	20.2	15.7	28.4	7.3	3.6	4.0				24.7	25A 15	10Z	N
8 658	94-115	2B23TB	81.3	8.3	10.4	12.8	23.1	14.9	24.2	6.3	4.1	4.2				21.3	45A 37	8Z	N
8 659	115-130	3B31TXB	83.8	10.6	5.6	11.2	23.7	14.2	24.6	10.1	6.3	4.3				29.1	7A --	7Z	N
8 660	130-170	3B32TXB	72.2	21.8	6.0	10.4	17.3	10.6	22.1	11.8	10.4	11.4				33.6	10A --	10Z	N
8 661	170-185	4C 1B	57.3	29.0	13.7	8.9	15.5	10.1	16.5	6.3	7.3	21.7				21.5	14A --	14Z	TR
8 662	185-255	5C 2B	81.2	12.0	6.8	20.0	21.6	11.7	18.9	9.0	6.4	5.6				25.2	9A --	9Z	N
8 663	0- 15	A															18A --	18Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR-BON 6A1A	NITRO-GEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LIMIT LL 4F2	PL 4F1	EST. 1/3 BAR 4A1F	BULK DENSITY 1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		-PERCENT-				PCT.NO.40		GRAMS PER CC				-PERCENT-			-PERCENT-	
8 652	0- 5	2.73	0.115			1.5							0.90	0.95	0.95		20.2	6.2	1.8	1.7
8 653	5- 15	1.02	0.052			1.5					NP	NP	1.36	1.36	0.92		13.0	5.0		
8 654	15- 25	0.71				1.8							1.31	1.32	0.92		11.4	5.5	0.3	0.2
8 655	25- 51	0.47				1.9					NP	NP	1.39	1.39	0.91		9.0	5.6		
8 656	51- 74	0.30				2.9							1.57	1.57	0.82		14.3	6.2		
8 657	74- 94	0.22				3.5							1.45	1.46	0.84		17.4	6.5	0.2	0.2
8 658	94-115	0.28				2.6					NP	NP	1.50	1.50	0.69		17.2	6.9		
8 659	115-130	0.18				1.6							1.64	1.64	0.95		9.5	4.0		
8 660	130-170	0.10				0.6							1.92	1.92	0.93		9.7	3.4		
8 661	170-185					1.3							1.60	1.62	0.91		19.3	7.9	0.4	0.4
8 662	185-255					0.5					NP	NP	1.40		0.95			3.5		
8 663	0- 15	1.87	0.082										1.30		0.90			9.4		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4-OAC 5A6A	BASE SUM+ ACI-DITY	SATURATION SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 652	0- 5	6.1	0.4	TR	0.4	6.9	9.0		15.9	12.8	44		54		6.0	5.5
8 653	5- 15	3.4	0.3	TR	0.3	4.0	5.5		9.5	9.9	42		41		6.2	5.6
8 654	15- 25	2.1	0.3	TR	0.2	2.6	5.7		8.3	7.9	32		33		5.7	5.3
8 655	25- 51	2.0	0.3	0.1	0.1	2.5	4.9	0.5	7.4	7.1	34	83	35		5.4	4.7
8 656	51- 74	2.3	0.4	0.1	0.1	2.9	4.0	0.3	6.9	6.5	42	91	45		5.4	4.7
8 657	74- 94	1.9	0.3	0.1	0.1	2.4	4.9	0.2	7.3	5.0	33	92	48		5.5	4.9
8 658	94-115	2.2	0.4	0.1	0.1	2.8	5.9	0.3	8.7	5.9	32	90	47		5.4	4.8
8 659	115-130	2.0	0.3	0.1	0.1	2.5	2.8	0.2	5.3	5.1	47	93	49		5.4	4.7
8 660	130-170	2.8	0.4	0.1	0.1	3.4	1.4	0.2	4.8	4.5	71	94	76		5.6	4.8
8 661	170-185	9.2	1.7	0.2	0.1	11.2	2.3	0.4	13.5	14.8	83	97	76		5.4	4.8
8 662	185-255	2.5	0.5	0.1	0.1	3.2			3.2	3.8	100+		84		6.1	5.6
8 663	0- 15														6.1	5.4

CLAY MINERALOGY

SAMPLE NO	DEPTH CM.	LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 652	0- 5											1 TRACE
8 653	5- 15	VR	3	KK	3						VR VERMICLT	2 SMALL
8 654	15- 25										KK KAOLINIT	3 MODERATE
8 655	25- 51	VR	3	KK	3						MI MICA	4 ABUNDANT
8 656	51- 74											5 DOMINANT
8 657	74- 94											6 INDETER.
8 658	94-115	KK	3	MI	1							
8 659	115-130											
8 660	130-170											
8 661	170-185											
8 662	185-255	MI	3	KK	3							
8 663	0- 15											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

JABU COARSE SANDY LOAM
S68Calif-9-34

Location: Tahoe Basin, El Dorado County, California; 0.2 mile east of southwest corner sec. 23, T. 12 N., R. 18 E., 15 feet north of Fountain Place Road. Date of Sampling: August 14, 1968.
Description by: G. M. Kennedy. Collectors: J. H. Rogers, D. Palmquist, B. R. Brasher, W. D. Nettleton.
Classification: Ultic Haploxeralf, coarse-loamy, mixed, frigid.
Vegetation: Not cultivated; White fir, Jeffrey pine, squaw carpet, greenleaf manzanita. Climate: Precipitation is 35 to 40 inches. Mean annual temperature is 41°F., January 26°F., July 60°F. Parent Material: Mixed alluvium. Topography: Glacial outwash over moraine, 10 percent north-facing slope. Elevation: 6940 feet. Drainage: Well drained; runoff is slow; permeability is slow. Soil Moisture: Dry to 72 inches.
Remarks: Field determinations for pH by Hellige-Truog triplex indicator. Composite A taken from 10 satellite samples.

HORIZON	DESCRIPTION
	(3-0 cm.)
01	1 to 0 inch, fir needles.
	(0-5 cm.)
A1 RSL 68652	0 to 2 inches, brown (7.5YR 5/4) coarse sandy loam, dark brown (7.5YR 3/2) moist; weak coarse platy structure; soft, very friable, nonsticky and nonplastic; many very fine, few fine roots; common very fine interstitial pores; 13 percent gravel; medium acid (pH 5.6); clear smooth boundary.
	(5-15 cm.)
A3 RSL 68653	2 to 6 inches, brown (7.5YR 5/4) coarse sandy loam, reddish brown (5YR 4/4) moist; weak medium subangular blocky structure; soft, very friable, nonsticky and nonplastic; many very fine, common fine, medium and coarse roots; common very fine interstitial, few fine tubular pores; 14 percent gravel; medium acid (pH 6.0); clear smooth boundary.
	(15-25 cm.)
B1t RSL 68654	6 to 10 inches, brown (7.5YR 5/4) coarse sandy loam, reddish brown (5YR 4/4) moist; weak medium subangular blocky structure; slightly hard, very friable, nonsticky and nonplastic; many very fine, fine, medium, and coarse roots; common very fine interstitial and few fine tubular pores; common thin clay films in bridges and coating sand grains; 14 percent gravel; medium acid (pH 5.8); clear smooth boundary.
	(25-51 cm.)
B2t RSL 68655	10 to 20 inches, strong brown (7.5YR 5/6) coarse sandy loam; dark reddish brown (5YR 3/4) moist; massive; hard, friable, nonsticky and nonplastic; many very fine, fine, medium and coarse roots; common very fine tubular and interstitial pores; many thin clay films in bridges and coating sand grains, clay films moderately thick in places; 14 percent gravel; strongly acid (pH 5.5); abrupt wavy boundary.
	(51-74 cm.)
IIB21tb RSL 68656	20 to 29 inches, strong brown (7.5YR 5/6) gravelly coarse sandy loam, yellowish red (5YR 4/6) moist; brown (7.5YR 5/4) coatings; massive; hard, firm, slightly sticky and slightly plastic; few very fine, fine, medium and coarse roots; common very fine tubular and interstitial, few fine tubular pores; many thin, few moderately thick clay films in bridges and coating sand grains; few cobbles resting on upper boundary of horizon; 27 percent gravel; strongly acid (pH 5.5); gradual smooth boundary.
	(74-94 cm.)
IIB22tb RSL 68657	29 to 37 inches, strong brown (7.5YR 5/6, 5/8) gravelly coarse sandy loam, yellowish red (5YR 5/6) moist; brown (7.5YR 5/4) coatings; massive; hard, firm, slightly sticky, slightly plastic; very few fine, medium and coarse roots; common very fine interstitial and tubular, few fine tubular pores; continuous thin, few moderately thick clay films in bridges and coating sand grains; 26 percent gravel; strongly acid (pH 5.5); gradual wavy boundary.
	(94-115 cm.)
IIB23tb RSL 68658	37 to 45 inches, strong brown (7.5YR 5/6, 5/8) gravelly coarse sandy loam, yellowish red (5YR 4/8) moist; massive; hard, firm, slightly sticky and slightly plastic; very few fine, medium and coarse roots; common very fine interstitial pores; continuous moderately thick yellowish red (5YR 4/6) clay films in bridges and coating sand grains; 44 percent gravel; strongly acid (pH 5.5); abrupt wavy boundary.
	(115-130 cm.)
IIIB31tkb RSL 68659	45 to 52 inches, variegated reddish yellow and strong brown (7.5YR 6/6, 5/6) loamy coarse sand, strong brown (7.5YR 5/6, 5/8) moist; light gray (10YR 7/2) bleached areas, reddish yellow (7.5YR 6/6) moist; massive; hard, firm, nonsticky and nonplastic; no roots; common very fine interstitial pores; common thin clay films line pores; 7 percent gravel; strongly acid (pH 5.5); clear wavy boundary.
	(130-170 cm.)
IIIB32tkb RSL 68660	52 to 66 inches, light gray (2.5Y 7/2) loamy coarse sand, olive gray (5Y 5/2) moist; dark grayish brown (2.5Y 4/2) bands; massive; extremely hard, extremely firm, nonsticky and nonplastic; no roots; few very fine interstitial pores; 10 percent gravel; strongly acid (pH 5.5); clear wavy boundary.
	(170-185 cm.)
IVC1b RSL 68661	66 to 72 inches, white (N 8/) silty clay, light gray (N 7/) moist; black coatings dry and moist; strong medium platy structure; slightly hard, firm, nonsticky and plastic; whole horizon has horizontal bands of clay and silt with black coatings between the plates; 6 percent gravel; strongly acid (pH 5.5).

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Ahwahnee sandy loam (taxadjunct) **LOCATION** El Dorado County, California

SOIL FAMILY: Ultic Haploxeralf, coarse-loamy, mixed, mesic.

SOIL NOS. S61Calif-9-6 **LAB. NOS.** 61114 - 61120

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1	2A2	TEXTURAL CLASS
		VERY COARSE SAND	COARSE SAND	MEDIUM SAND	FINE SAND	VERY FINE SAND	SILT	CLAY				

AHWAHNEE SANDY LOAM (taxadjunct) 1/
S61Calif-9-6

Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, David Estrada, Lothair Grant, Lynn Brittan
Date of Sampling: April 28, 1961 Description by: Grant Kennedy.
Location: El Dorado County, 1.5 miles southwest of Fairplay, California, NW cor. SE1/4NW1/4 sec. 5, T. 8 N., R. 12 E., El Dorado County, California.
Elevation: 2,390 feet
Relief: Rolling uplands with a 12 percent slope to the south.
Drainage: Well drained; medium runoff; rapid over moderate permeability
Parent Material: Granodiorite
Climate: Mean annual rainfall 37 inches. Mean annual temperature 57° F.
Vegetation: Cleared and formerly cultivated field, now heavily grazed by sheep. Present cover filaree, annual forbs, bur clover, wild oats, soft chess, ripgut brome, scattered ponderosa pine, live oak and black oak.
Classification: Ultic Haploxeralf, coarse-loamy, mixed, mesic.
Remarks: The profile has considerable mica which is most noticeable in the B horizon. The C1 is intermittent, in places being interrupted by an irregular [illegible]

HORIZON		DESCRIPTION
Ap1	0- 3"	Dark brown (10YR 4/3) sandy loam, dark brown (10YR 3/3) when moist; massive; slightly hard when dry, friable when moist, nonsticky and nonplastic when wet; abundant very fine roots; many very fine and fine tubular and interstitial pores; medium acid, pH 6.0; clear smooth boundary.
Ap2	3-15"	Dark brown (10YR 4/3) sandy loam, dark brown (10YR 3/3) when moist; massive; slightly hard when dry, very friable when moist, nonsticky and nonplastic when wet; abundant very fine and fine roots; many very fine tubular and interstitial, common fine tubular, few medium tubular pores; medium acid, pH 6.0; clear wavy boundary.
B1	15-26"	Yellowish brown (10YR 5/4) sandy loam near sandy clay loam, brown (7.5YR 4/4) when moist; massive; hard when dry, friable when moist, slightly sticky and slightly plastic when wet; plentiful very fine and fine roots; many very fine tubular and interstitial, common fine and medium tubular pores; few thin clay films in pores and bridges; medium acid, pH 6.0; clear smooth boundary.
B21t	26-38"	Yellowish brown (10YR 5/4) sandy clay loam near sandy loam, brown (7.5YR 4/4) when moist; massive; extremely hard when dry, friable when moist, slightly sticky and plastic when wet; few very fine and fine roots; many very fine tubular, common fine, few medium pores; many thin clay films in pores and bridges; medium acid, pH 6.0; gradual smooth boundary.
B22t	38-47"	Yellowish brown (10YR 5/4) sandy clay loam, brown (7.5YR 4/4) when moist; massive; extremely hard when dry, firm when moist, sticky and plastic when wet; few very fine and fine roots; common very fine tubular and interstitial, common fine tubular pores; many [illegible]

CONSECUTIVE PEDON NUMBER C 68 78 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES JABU SHALLOW VARIANT SOIL NO. S68 CALI. 9-33 SAMPLE NO. B 643-B 651 TAHOE BASIN AREA

SOIL FAMILY ULTIC HAPLOXERALF FINE-LOAMY,MIXED,FRIGID RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SANDS VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A	INT. PCT. II .2- .02	COARSE FRAGMENTS OF WHOLE I 75- 20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														SOIL PCT. OF LT 75		
																	--A --	--Z --	
8 643	3- 0	01																	
8 644	0- 15	A1	71.1	18.4	10.5	17.4	17.4	10.0	17.1	9.2	8.3	10.1				26.1	38A --	38Z	N
8 645	15- 41	A3	68.3	19.6	12.1	10.6	15.9	10.3	19.9	11.6	9.7	9.9				32.1	12A --	12Z	N
8 646	41- 69	2B1 TXB	61.9	16.0	22.1	14.5	15.5	9.0	14.6	8.3	6.2	9.8				22.5	13A --	13Z	N
8 647	69-105	2B21TXB	56.7	15.7	27.6	8.5	13.5	9.5	16.3	8.9	6.4	9.3				24.2	11A --	11Z	N
8 648	105-175	2B22TXB	71.0	18.6	10.4	14.4	17.4	12.1	18.8	8.3	7.7	10.9				25.4	13A --	13Z	N
8 649	175-205	2B3 TB	74.5	13.6	11.9	20.1	19.4	10.9	16.4	7.7	5.2	8.4				21.7	17A --	17Z	N
8 650	205-220	2C B	84.7	10.2	5.1	23.5	31.0	11.7	13.5	5.0	4.9	5.3				16.1	24A --	24Z	N
8 651	0- 41	A															19A --	19Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR-BON 6A1A	NITRO GEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6F1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP EXT FE 6C5A	AL 6G5A	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		-PERCENT-				PCT.NO.40		GRAMS PER CC				-PERCENT-				
												0.40			1.00					
8 643	3- 0																			
8 644	0- 15	2.83	0.119			3.3				NP	NP		1.22	1.24	0.78		13.4	6.7	0.5	0.4
8 645	15- 41	1.48	0.067			3.7							1.31	1.31	0.94		14.7	6.6		
8 646	41- 69	0.50				7.5				32	19		1.44	1.47	0.92		23.9	10.5	0.7	0.6
8 647	69-105	0.18				7.3				39	25		1.54	1.60	0.93		19.2	12.4	1.3	1.2
8 648	105-175	0.09				5.1							1.64	1.65	0.91		16.3	5.8	0.2	0.2
8 649	175-205	0.08				5.7							1.62	1.64	0.89		19.1	5.7	0.4	0.4
8 650	205-220					2.6				NP	NP		1.68	1.68	0.83		8.4	3.5		
8 651	0- 41	2.51	0.104									1.28			0.90			7.0		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACI-DITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 643	3- 0															
8 644	0- 15	5.2	0.6	TR	0.3	6.1	11.9		18.0	14.3	34		43		5.8	5.2
8 645	15- 41	4.7	0.5	TR	0.2	5.4	9.8		15.2	10.7	36		51		6.0	5.3
8 646	41- 69	3.3	0.9	0.1	0.1	4.4	7.2		11.6	8.5	38		52		5.8	5.4
8 647	69-105	4.5	1.4	0.1	0.1	6.1	5.3		11.4	9.1	54		67		5.9	5.6
8 648	105-175	3.0	0.8	0.1	TR	3.9			3.9	5.2	100+		76		5.9	5.6
8 649	175-205	3.0	0.8	0.1	TR	3.9			3.9	5.3	100+		74		5.9	5.5
8 650	205-220	2.3	0.6	0.1	TR	3.0			3.0	3.6	100+		85		6.0	5.5
8 651	0- 41														5.8	5.3

SAMPLE NO	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 643	3- 0											1 TRACE
8 644	0- 15	VR	3	KK	3						VR VERMICLT	2 SMALL
8 645	15- 41										KK KAOLINIT	3 MODERATE
8 646	41- 69	KK	3	VR	1						MI MICA	4 ABUNDANT
8 647	69-105	MI	3	KK	3							5 DOMINANT
8 648	105-175											6 INDETER.
8 649	175-205											
8 650	205-220	MI	3	KK	3	VR	1					
8 651	0- 41											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.
N=NOT DETERMINED.

JABU COARSE SANDY LOAM, SHALLOW VARIANT 1/
S68Calif-9-33

Location: Tahoe Basin, El Dorado County, California, 0.6 miles south and 0.4 miles east of the northwest corner of sec. 22, T. 12 N., R. 18 E., 200 feet south of Fountain Place Road. Date of Sampling: August 14, 1968.
Description by: G. M. Kennedy and P. G. Nazar. Collectors: J. H. Rogers, G. L. Anderson, B. R. Brasher, W. D. Nettleton.
Classification: Ultic Haploxeralf, fine-loamy, mixed, frigid.
Vegetation: Not cultivated; Jeffrey pine, big sagebrush, perennial and annual grasses. Climate: Precipitation is 35 inches. Mean annual temperature is 43°F., January 27°F., July 61°F. Parent Material: Mixed alluvium. Topography: Glacial outwash terrace. 5 percent north-northwest facing slope. Elevation: 6400 feet. Drainage: Well drained; runoff is slow; permeability is very slow. Soil Moisture: Dry.
Remarks: Field determination for pH by Hellige-Truog triplex indicator. Composite A taken from 10 satellite samples.

HORIZON	DESCRIPTION
01 RSL 68643	(3-0 cm) 1 to 0 inches, loose mat of pine needles.
A1 RSL 68644	(0-15 cm.) 0 to 6 inches, reddish brown (5YR 4/4) sandy loam, dark reddish brown (5YR 3/4) moist; weak medium platy structure parting to moderate fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine and fine, common medium roots; common very fine tubular and interstitial pores; 3 percent gravel; medium acid (pH 6.0); clear smooth boundary.
A3 RSL 68645	(15-41 cm.) 6 to 16 inches, yellowish red (5YR 5/6) heavy sandy loam, yellowish red (5YR 4/6) moist; weak medium subangular blocky structure; slightly hard, friable, slightly sticky and slightly plastic; many very fine, fine, medium and coarse horizontal roots; common very fine interstitial and few very fine tubular pores; few thin clay films in pores and on ped faces; 5 percent gravel and 15 percent 5 inch minus cobbles; slightly acid (pH 6.2); abrupt wavy boundary.
IIB1txb RSL 68646	(41-69 cm.) 16 to 27 inches, strong brown (7.5YR 5/6) sandy clay loam, strong brown (7.5YR 4/6) moist; massive; hard, firm, brittle, sticky and plastic; common fine, medium and coarse roots; common very fine tubular and interstitial and few fine tubular pores; continuous thin clay films in bridges and coating sand grains; 10 percent gravel and 10 percent cobbles; slightly acid (pH 6.2); gradual wavy boundary.
IIB21txb RSL 68647	(69-105 cm.) 27 to 42 inches, strong brown (7.5YR 5/6) clay loam, strong brown (7.5YR 4/6) moist; massive; very hard, very firm, brittle, sticky and plastic; few medium and coarse roots; few very fine tubular and interstitial pores; continuous thick clay films in bridges and coating sand grains; 5 percent gravel and 15 percent cobbles; slightly acid (pH 6.2); gradual smooth boundary.
IIB22txb RSL 68648	(105-175 cm.) 42 to 68 inches, strong brown and brown (7.5YR 5/6, 4/4) gravelly sandy clay loam, strong brown (7.5YR 4/6) moist; massive; very hard, very firm, brittle, slightly sticky and slightly plastic; very few fine and medium, few coarse roots; common very fine tubular pores; continuous thin clay films in bridges and coating sand grains; 20 percent gravel and 25 percent cobbles; slightly acid (pH 6.2); clear wavy boundary.
IIB3tb RSL 68649	(175-205 cm.) 68 to 80 inches, variegated strong brown (7.5YR 5/8, 5/6) gravelly sandy clay loam, strong brown (7.5YR 5/6) moist; massive; very hard, firm, brittle, slightly sticky and slightly plastic; no roots; common very fine tubular pores; continuous thin clay films in bridges and coating sand grains; 20 percent gravel and 20 percent cobbles; slightly acid (pH 6.2); clear wavy boundary.
IICb RSL 68650	(205-220 cm.) 80 to 86 inches, variegated strong brown and brown (7.5YR 5/6, 4/4) coarse sand, brown (7.5YR 4/4) moist; massive; slightly hard, friable, nonsticky and nonplastic; no roots; common very fine interstitial pores; common thin clay films coating sand grains; 30 percent gravel; medium acid (pH 6.0).

1/ The Jabu series is in a coarse-loamy, mixed, frigid family of Ultic Haploxeralfs. This pedon is in a fine-loamy, mixed, frigid family of the same subgroup.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Boomer stony loam **LOCATION** San Diego County, California

SOIL FAMILY: Ultic Haploxeralf, fine-losmy, mixed, mesic.

SOIL NOS. S60Calif-37-9 **LAB. NOS.** 60446 - 60451

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 3	A11	2.6	5.5	4.1	13.0	11.8	41.0	22.0	36.8	24.6	10	l
3- 8	A12	2.1	5.5	4.3	13.0	11.9	40.3	22.9	36.0	24.7	15	l
8-16	B1	1.2	4.4	3.8	14.9	13.2	39.4	23.1	38.4	23.7	3	l
16-31	B21	2.1	5.4	3.7	13.4	11.3	35.5	28.6	34.9	21.0	4	cl
31-46	B22	2.2	7.0	4.9	13.7	10.4	29.5	32.3	31.3	17.5	7	cl
46-57+	B3	3.2	13.0	8.8	20.7	10.7	22.7	20.9	33.4	12.4	6	scl-fsl

8C1a pH H_2O 1:1	8C1c pH KCl 1:1	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	Free Fe_2O_3 6C1b %	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.7	5.6	3.81	0.241	15.8				3.8			15.9
6.5	5.4	3.01	0.187	16.1				3.8			10.1
5.7	4.5	1.33	0.092	14.5				4.1			14.3
6.0	4.6	0.54	0.049	11.0				4.2			17.4
5.9	4.7	0.47	0.040	11.8				3.9			22.4
6.0	4.8	0.37	0.033					3.3			16.5

5A2a CATION EXCHANGE CAPACITY ←(Na)	EXTRACTABLE CATIONS 6N2b Ca	6O2c Mg	6H2a H	6P2a Na	6Q2a K	BASE SAT. % 5C3	Sum of Cations me./100 soil 5A3a	g. K	MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						milliequivalents per liter →		
34.9	20.8	5.6	10.1	0.2	1.4	73	38.1		
32.0	17.4	4.9	10.3	0.2	0.9	69	33.7		
27.6	13.1	4.4	11.2	0.2	0.4	62	29.3		
24.2	11.8	5.4	8.9	0.2	0.3	67	26.6		
26.3	11.2	5.7	9.0	0.4	0.1	66	26.4		
21.3	8.5	3.8	8.9	0.6	0.1	59	21.9		

1/ Percent of total soil

BOOMER STONY LOAM
S60Calif-37-9

Description: George Borst
Date of Sampling: November 18, 1960
Collectors: Reuben Nelson, James Lockard, Cliff Henry, and George Borst
Location: Approximately 75 feet east of Engineer's Road, approximately 0.3 mile northwest of its junction with Old Mill Truck Trail in SE1/4 sec. 30, T. 13 S., R. 3 E. (projected) of Rancho Cuyamaca, San Diego County, California.
Elevation: 4,800 feet
Relief: Steep, 33 percent
Use: Woodland
Vegetation: Cedar, oak, and pure forest
Drainage: Well drained
Parent Material: Deeply weathered basic intrusive rocks
Climate: This soil occurs under an average annual rainfall of about 38", a mean annual temperature of 53°, a mean January temperature of 39°, and a mean July temperature of 70°.
Remarks: This soil is relatively stony or rocky throughout, especially in the lower part of the B horizon. This profile is typical of Cohasset as to size and degree of development of the solum although it is less acid than is typical for the series. It was sampled as Aiken stony loam.
Reaction: pH was determined in the field with a Truog (Hellige) soil reaction kit
Classification: **Ultic Haploxeralf, fine-loamy, mixed, mesic.**

HORIZON		DESCRIPTION
O1 & O2	1- 0"	Partially decomposed and undecomposed leaves and litter, neutral (pH 7.0).
A11	0- 3"	Dark grayish brown (10YR 4/2) loam, very dark grayish brown (10YR 3/2) when moist; moderate fine crumb structure; soft, very friable, slightly sticky and plastic; abundant very fine random roots; many fine interstitial pores; slightly acid (pH 6.5); abrupt smooth boundary.
A12	3- 8"	Dark brown (10YR 4/3) loam, dark brown (10YR 4/3) when moist; moderate medium granular structure; slightly hard, friable, slightly sticky and plastic; plentiful fine random roots; many fine inter-

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Elkhorn fine sandy loam (taxadjunct) **LOCATION** San Mateo Co., Calif.

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, mesic

SOIL NOS. S55Ca1.41-5 **LAB. NOS.** 55125-55130

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0- 6	A_{1p}	0.3	5.2	19.7	36.8	7.8	21.5	8.7	34.8	9.0	0	fsl
6-14	A_{12}	0.7	6.3	20.2	33.9	7.3	21.6	10.0	33.3	8.4	0	fsl
14-26	A_2	0.3	5.9	22.1	37.1	7.7	22.1	4.8	34.1	9.3	0	fsl
26-40	B_{21}	0.2	5.2	18.8	29.9	6.2	17.9	21.8	26.9	7.8	0	scl
40-50	B_{22}	0.2	4.8	18.3	30.1	6.5	19.4	20.7	28.9	8.1	1	scl
50-60	C1	0.4	5.1	18.0	29.4	6.0	19.8	21.3	27.9	8.6	0	scl

8C1b pH SATURATED PASTE 1:1	8C1a pH 1:5	8C1a pH 1:10	6A1a ORGANIC MATTER ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	Free Fe_2O_3 Pct. 6C1b	4A3a Bulk Density g/cc	4B2 MOISTURE TENSIONS 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
5.3		5.9	1.07	.092	11.6		0.5	0.8			9.5	4.0
5.4		5.9	0.97	.086	11.3		0.5	0.8	1.52		10.5	4.0
5.5		5.9	0.14				0.6	0.6			8.1	2.0
6.0		6.4	0.15				0.5	2.7	1.64			9.5
6.1		6.5	0.13				0.4	2.0				9.0
6.2		6.6	0.11				0.4	2.0	1.62			9.2

5A2a CATION EXCHANGE CAPACITY (+ Na)	5B1a EXTRACTABLE CATIONS Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	6P1a SATURATION EXTRACT SOLUBLE Na	6Q1a K	6N1a 6O1a Ca+Mg	6J1a 6I1a CO_3+HCO_3 1/	6K1a 8A1 Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
7.9	3.0	0.8	5.4	0.3	0.3	45	1.5	0.5	2.2	0.6	0.9	22.7
7.9	3.6	0.9	5.2	0.2	0.2	49	1.7	0.1	3.0	0.5	0.7	21.7
3.3	1.6	0.7	1.8	0.1	Tr	57	1.8	0.1	3.4	0.2	1.3	17.4
11.3	2.9	4.2	3.5	0.4	0.1	68	2.5	<0.1	1.6	0.2	1.4	41.3
11.4	2.3	3.8	3.6	0.4	0.1	65	2.5	<0.1	1.3	0.2	1.3	40.8
11.4	4.9	4.7	3.4	0.6	0.1	75	2.5	Tr	0.7	0.2	1.2	44.3

1/ only HCO_3^- found

ELKHORN FINE SANDY LOAM (taxadjunct) 1/
S55Calif-41-5

Location: San Mateo County, California. Two miles southwest of Pescadero, 1/2 mile north of junction of Highway No. 1 and Old Coast Highway on east side of Old Coast Highway. See Sheet 40, Soil Survey of San Mateo Area, California, May, 1961.
Date of Sampling: February 25, 1955.
Description by: W. M. Johnson. Collectors: Reuben E. Nelson, Ralph E. Nelson, Richard Wagner, Melvin Mello, and W. M. Johnson.
Classification: Ultic Haploxeralf, fine-loamy, mixed, mesic.
Vegetation: Cultivated, used for production of Brussels sprouts. Original vegetation assumed to be grass and shrubs. Climate: Rainfall about 20 inches annually. June, July, and August are very dry. Fogginess is high in many parts of the year, particularly during summer and fall. Average January temperature is about 48° F., average July temperature about 60° F. On-shore winds carrying salt spray are common.
Parent Material: Sandy coastal plains sediments. Relief: Undulating to rolling marine terrace. On second terrace above ocean level. Slope about 5 percent, plane, toward the west. Elevation: About 65 feet. Drainage: Runoff medium, Permeability is moderate. Affected by laterally flowing underground water. At time of collection of sample, water flowed into the pit at a depth of 56 inches, and rose to within 50 inches of the surface overnight.
Remarks: Profile moist in A and B horizons, wet below.

HORIZON		DESCRIPTION
A_{1p}	0 - 6"	Gray (10YR 5/1.5, dry; 3/2, moist) light fine sandy loam. Weak medium and fine granular. Slightly hard, very friable, nonplastic, nonsticky. Roots very numerous. Peds slightly porous. pH 4.8. Abrupt, wavy lower boundary.
A_{12}	6 - 14"	Light fine sandy loam of about the same color as the A_{1p}. Very weak medium and fine subangular blocky; nearly massive. Slightly hard, very friable, nonplastic, nonsticky. Roots very numerous. Peds are slightly porous. pH 5.0. Abrupt wavy lower boundary.
A_2	14 - 26"	Light gray (1Y 7/2.5, dry; 5/3, moist) light fine sandy loam. Massive. Slightly hard, very friable, nonplastic, nonsticky. Roots are common. Peds are moderately porous. There are lumps of B_2 in this horizon, and "tongues" of A_2 penetrate to a depth of about 4 feet in this profile. pH 5.8. Abrupt, irregular lower boundary.
B_{21}	26 - 40"	Brownish yellow (10YR 6/6, dry; 5/5, moist) sandy clay loam with common fine mottles of black and dark brown (dry). Moderate very coarse prismatic, breaking to weak coarse and medium subangular blocky. Moderate continuous clay skins. Very hard, friable, plastic, sticky. Few roots. Peds are nonporous. pH 6.0. Clear, irregular lower boundary.
B_{22}	40 - 50"	Brownish yellow (10YR 6/6, dry; 5/6, moist) light sandy clay loam with common fine mottles of black, very pale brown and dark brown (dry). Weak coarse and medium subangular blocky. Very hard, friable, plastic, sticky. Moderate, nearly continuous clay skins. Very few roots. Peds are nonporous. pH 6.2. Gradual, wavy lower boundary.
C1	50 - 60"	Finely mottled very pale brown and brownish yellow (10YR 7/3 and 6/6, dry; 5/3 and 4/6, moist) light sandy clay loam with a few small black spots. Massive. Hard, friable, plastic, sticky. No roots observed. Slightly porous. A few thin patches of clay skins along tubular pores and cracks. pH 6.4.

1/ The Elkhorn series is in a fine-loamy, mixed, mesic family of Pachic Argixerolls. This pedon has an epipedon with a base saturation (by sum of cations) of less than 50.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Elkhorn sandy loam (taxadjunct) **LOCATION** San Mateo County, California

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, mesic.

SOIL NOS. S55Cal. 41-6 **LAB. NOS.** 55131-55137

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 2/	TEXTURAL CLASS
0-13	A_{1p}	2.3	21.7	28.4	20.4	4.3	15.9	7.0	20.1	7.6	1	sl
13-17	A_{12}	2.6	19.3	26.7	20.3	4.5	17.5	9.1	21.1	8.3	2	sl
17-24	A_{21}	2.5	19.4	27.5	20.2	4.5	16.6	9.3	21.0	7.5	1	sl
24-27	A_{22}	2.5	20.7	26.6	19.0	4.0	15.9	11.3	19.6	7.2	1	sl
27-40	B_2	1.3	16.3	26.0	18.7	3.9	13.4	20.4	18.1	5.8	1	scl
40-55	B_3	3.2	17.2	31.6	21.9	3.3	10.1	12.7	15.7	4.9	1	sl
55-60	C_g or C_{ug}	19.1	16.2	10.6	11.6	4.2	16.3	22.0	18.9	7.6	14	scl

pH 8C1b SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C	Free Fe_2O_3 6C1b %	4A3a Bulk Density g/cc	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
5.1		5.6	1.13	.087	13.0		0.6	0.8			8.4	3.5
5.4		6.0	0.72	.069	10.4		0.6	1.0	1.57		9.7	3.9
5.5		6.0	0.58				0.7	1.1			9.7	3.8
5.6		6.0	0.39				0.6	1.3			10.4	4.7
5.9		6.0	0.25				0.6	1.8	1.75			8.6
6.0		6.2	0.10				0.8	1.7				6.2
5.2		5.9	0.07				0.4	1.7	1.84			10.0

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a 6O1a	6J1a 6I1a $CO_3^{=}$ HCO_3^- 1/	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
7.8	3.1	0.7	5.5	0.2	0.3	44	2.1	0.6	4.7	1.3	0.9	20.1
8.5	2.5	0.7	5.8	0.3	0.2	39	2.2	0.3	2.8	0.4	1.8	19.9
7.5	3.2	0.7	5.3	0.3	0.2	45	2.6	0.4	3.5	0.4	1.9	19.5
7.7	2.3	0.6	4.6	0.2	0.2	42	2.1	0.2	3.6	0.3	1.9	20.9
9.8	2.1	3.1	4.3	0.4	0.1	57	2.7	0.2	4.6	0.2	1.9	28.0
7.5	1.3	2.8	3.8	0.4	0.1	55	4.1	0.1	3.5	0.2	2.5	24.3
16.4	2.4	6.8	4.0	0.8	0.2	72	3.2	0.1	0.8	0.1	1.6	48.9

1/ only HCO_3^- found

2/ Percent of whole soil

ELKHORN SANDY LOAM (taxadjunct) 1/
S55Calif-41-6

Location: San Mateo County, California. 115 yards east of California Highway No. 1, measured from a point 4/10 mile north of Pigeon Point Lighthouse. In a cultivated field about 10 feet north of a field road. Date of Sampling: February 25, 1955.
Description by: W. M. Johnson. Collectors: Reuben Nelson, Ralph Nelson, Richard Wagner, Melvin Mello, and W. M. Johnson.
Classification: Ultic Haploxeralf, fine-loamy, mixed, mesic.
Vegetation: Cultivated, irrigated. Land Use: Used for Brussels sprouts. Climate: Rainfall about 20 inches annually. June, July, and August are very dry. Fogginess is high in many parts of the year, particularly during summer and fall. Average January temperature is about 48° F., average July about 60° F. On-shore winds carrying salt spray are common. Parent Material: Sandy coastal plains sediments. Relief and Slope: Eroded low marine terrace. Gently undulating. About 3 percent, plane, toward the northwest. Elevation: About 55 feet. Drainage: Runoff, medium; permeability, medium. Affected by laterally moving ground water. Water seeped into hole at depth of 43 inches. Soil Moisture: Profile moist to depth of 40 inches, wet below.

HORIZON		DESCRIPTION
A_{1p}	0 - 13"	Dark grayish brown (10YR 4.5/1.5; 3/1.5, moist) light sandy loam. Cloddy due to tillage. Very weak coarse and medium granular. Slightly hard, friable, nonplastic, nonsticky. Porous. pH 5.0. Abrupt, smooth lower boundary.
A_{12}	13 - 17"	Dark grayish brown (10YR 4.5/1.5; 2.5/1.5, moist) light sandy loam. Weak medium and fine granular. Slightly hard, friable, nonplastic, nonsticky. Porous. pH 5.3. Clear, wavy lower boundary.
A21	17 - 24"	Mottled dark grayish brown and brown (10YR 4.5/1.5 and 4/3, dry; 10YR 3/1.5 and 3/3, moist) light sandy loam. Weak medium subangular blocky. Slightly hard, friable, nonplastic, nonsticky. Porous. pH 5.5. Clear, wavy lower boundary.
A22	24 - 27"	Mottled gray and brown (10YR 5/1 and 5/3, dry; 3/1 and 3/3, moist) with a few darker brown spots. Sandy loam, with slightly more clay than the horizon above. Moderate medium subangular blocky. Hard, friable, slightly plastic, nonsticky. Porous. Few small patches of thin clay skins on vertical surfaces. pH 5.6. "Interfingered" with horizon below.
B_2	27 - 40"	Yellowish brown (10YR 5/6, dry; 4/6, moist) with common medium and fine mottles of black, very dark brown, and dark brown (dry). Light sandy clay loam. Moderate medium subangular blocky. Very hard, friable, plastic, sticky. Thin, nearly continuous clay skins. Few pores. pH 6.0. Indistinct lower boundary.
B_3	40 - 55"	Yellowish brown (10YR 5/6, dry; 4/6, moist) with many medium and coarse mottles of black. Sandy loam. Very weak coarse and medium subangular blocky. Hard, friable, slightly plastic, nonsticky. Moderate number of fine pores. There are thin, patchy clay skins. pH 6.2. Clear, wavy lower boundary.
C_g or C_{ug}	55 - 60"+	Mottled light gray and yellowish brown (2.5Y 7/1 and 10YR 5/6, dry; 2.5Y 6/2 and 10YR 4/6, moist) with many black spots and streaks. Sandy clay loam. Massive. Very hard, friable, plastic, sticky. Contains more coarse fragments than horizons above. There are a few patches of thin clay skins on vertical surfaces. pH 5.3.

1/ The Elkhorn series is in a fine-loamy, mixed, mesic family of Pachic Argixerolls. This pedon has an epipedon with a base saturation of less than 50 percent (by sum of cations).

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Holland sandy loam **LOCATION** El Dorado County, California

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, mesic

SOIL NOS. S61Calif-9-5 **LAB. NOS.** 61106 - 61113

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	3A1 2A2 > 2	TEXTURAL CLASS
0- 2	A_{11}	7.6	19.6	9.2	18.4	7.6	26.4	11.2	28.2	16.3	0	cosl
2-10	A_{12}	7.0	18.5	8.5	20.0	8.2	25.4	12.4	28.5	16.6	0	cosl-sl
10-21	B_1	4.5	19.5	8.2	19.3	8.0	25.1	15.4	27.5	16.6	0	sl
21-30	B_{21t}	6.3	17.7	6.8	16.6	6.8	24.1	21.7	24.4	16.0	0	scl
30-36	B_{22t}	5.1	17.6	7.5	17.1	6.5	21.7	24.5	22.5	15.4	0	scl
36-45	B_{3t}	6.2	18.7	8.8	18.6	7.2	20.4	20.1	23.8	15.0	0	scl-cosl
45-54	C_1	8.8	21.3	10.6	20.7	7.3	17.7	13.6	24.4	12.5	0	cosl
54+	R											

pH 8C1a H_2O 1:1	8C1c KCl (1 N) 1:1	Organic Matter Organic Carbon % 6A1a	Nitrogen % 6B1a	C/N			6C1a Free Fe_2O_3 %	4A3a Bulk Den. gm/cc			4B2 Soil Water 15 ATMOS. %
6.0	5.0	2.77	0.203	13.6			0.7				7.0
5.9	4.6	1.08	0.085	12.7			0.9	1.57			5.8
6.0	4.5	0.36	0.033	10.9			1.1	1.54			6.8
6.1	4.6	0.18	0.019				1.5	1.80			9.9
6.3	4.7	0.10					1.5	1.36			10.5
6.5	4.7	0.07					1.0	1.89			10.6
6.6	4.7	0.06					0.5				7.8

5A2a CATION EXCHANGE	EXTRACTABLE CATIONS 6N2b	6O2c	6H2a	6P2a	5B1a 6Q2a	BASE SAT. %	5A3a Sum of	Calc. Sum of	Calc. Sum of

HOLLAND SANDY LOAM
S61Calif-9-5

Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, David Estrada, Lothair Grant, Lynn Brittan
Date of Sampling: April 28, 1961 **Description by: Grant Kennedy.**
Location: 600 feet north of Slug Gulch Road, 1-1/4 mile northeast of Fairplay.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Holland loam (taxadjunct) **LOCATION** San Diego County, California

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, mesic.

SOIL NOS. S61Calif-37-2 **LAB. NOS.** 6110 - 6116

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 3	A11	1.7	5.7	5.3	33.0	21.7	23.9	8.7	58.5	11.7	18	fsl
3-10	A12	1.8	5.2	4.4	32.3	23.1	24.4	8.8	60.9	12.1	16	fsl
10-20	A3	1.4	4.6	4.2	29.1	25.6	25.3	9.8	61.2	12.6	15	fsl-vfsl
20-29	B21	1.2	2.8	2.6	20.0	27.7	24.8	20.9	57.0	11.9	5	cl-vfsl
29-35	B22	0.3	1.1	1.0	14.0	32.0	31.3	20.3	61.1	14.6	0	l
35-50	B3	0.2	0.9	1.1	19.1	35.7	31.6	11.4	69.2	15.4	0	vfsl
50-64+	C1	3.2	4.4	3.1	21.4	35.5	23.6	8.8	67.5	9.6	3	vfsl

pH 8C1a H_2O 1:1	8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	Free Fe_2O_3 6C1b %	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
5.7	4.9		4.21	0.187	22.5				1.6			6.7
5.8	4.8		1.14	0.057	20.0				1.8			4.9
5.7	4.5		0.30	0.024					1.7			5.2
5.7	4.6		0.26	0.022					2.1			10.0
5.5	4.6		0.30	0.018					1.9			12.7
5.6	4.4		0.18						2.6			10.7
5.7	4.2		0.13						2.2			7.4

5A2a CATION EXCHANGE CAPACITY (+ Na)	EXTRACTABLE CATIONS 6N2b Ca	6O2c Mg	6H2a H	6P2a Na	6Q2a K	BASE SAT. % 5C3	Sum of Cations me./100g. soil 5A3a	K				MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						milliequivalents per liter →					

HOLLAND LOAM (taxadjunct) 1/
S61Calif-37-2

Description: George Borst
Date of Sampling: January 18, 1961
Collectors: Reuben Nelson, Cliff Henry, and George Borst
Location: About 700 feet east of Boulder Peak Road, 1,425 feet south-southeast of its junction with Eagle Way, in Pine Hills, in the SW1/4 sec. 13, T. 13 S., R. 3 E., **San Diego County, California.**
Elevation: 4,100 feet
Relief: Strongly sloping, 16 percent
Use: Woodland range
Vegetation: Open forest of pine and oak, with occasional manzanita
Drainage: Well drained, moderately slow subsoil permeability
Parent Material: Deeply weathered micaceous schist
Climate: This soil occurs under an average annual rainfall of about 36", a mean annual temperature of 53°, a mean January temperature of 39°, and a mean July temperature of 70°.
Remarks: The A horizon of this soil is relatively thick, and erosion relatively slight, as compared with most soils in this series in San Diego County.
Reaction: pH was determined in the field with a Truog (Hellige) soil reaction kit.
Classification: **Ultic Haploxeralf, fine-loamy, mixed, mesic.**

HORIZON		DESCRIPTION
A11	0- 3"	Dark brown (10YR 4/3) loam, dark brown (10YR 3/3) when moist; weak fine granular structure; soft, very friable, nonsticky and slightly plastic; abundant very fine random roots; common very fine random tubular pores; highly micaceous; medium acid (pH 6.0); abrupt smooth boundary.
A12	3-10"	Yellowish brown (10YR 5/4) stony fine sandy loam, dark yellowish brown (10YR 4/4) when moist; weak medium subangular blocky structure; slightly hard, friable, nonsticky and slightly plastic; plentiful fine random roots; common fine random pores; highly micaceous; medium acid (pH 6.0); clear smooth boundary.
A3	10-20"	Yellowish brown (10YR 5/4) stony fine sandy loam, dark yellowish brown (10YR 4/4) when moist; weak medium subangular blocky structure; slightly hard, friable, nonsticky and slightly plastic; plentiful medium random roots; many medium random tubular pores; highly micaceous; medium acid (pH 6.0); abrupt wavy boundary.
B21	20-29"	Brown (7.5YR 5/4) clay loam, dark brown (7.5YR 4/4) when moist; strong coarse angular blocky structure; hard, firm, slightly sticky and plastic; few medium to coarse random roots; many medium to coarse random tubular pores; many moderately thick clay films on ped faces; highly micaceous; medium acid (pH 5.8); clear wavy boundary.
B22	29-35"	Brown (7.5YR 5/4) clay loam, dark brown (7.5YR 4/4) when moist; strong coarse angular blocky structure; hard, firm, sticky and very plastic; few medium to coarse random roots; many medium random tubular pores; many thick clay films on ped faces; highly micaceous; medium acid (pH 6.0); clear irregular boundary.
B3	35-50"	Brownish yellow (10YR 6/6) fine sandy loam, yellowish brown (10YR 5/4) when moist; massive; hard, firm, slightly sticky and plastic; few medium random roots; common medium random tubular pores; common thin clay films on ped faces; highly micaceous; medium acid (pH 6.0); gradual irregular boundary.
C1	50-64"+	Light gray (2.5Y 7/2) deeply weathered highly micaceous schist of very fine sandy loam texture, dark yellowish brown (10YR 4/4) on cleavage planes, light brownish gray (2.5Y 6/2) when moist; few medium random roots; medium acid (pH 6.0).

1/ **The Holland series is in a fine-loamy, mixed, mesic family of Ultic Haploxeralfs. This pedon has a solum slightly thinner than permitted for the series (greater than 50 inches).**

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Josephine loam variant **LOCATION** Amador Co., California

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, mesic

SOIL NOS. 59Calif-3-4 **LAB. NOS.** 59286 - 59293

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE	COARSE	MEDIUM	FINE	VERY FINE						

JOSEPHINE LOAM variant 1/
859Calif-3-4

Collectors: Grant Kennedy, Reuben Nelson, Bob Sketchley, John Rogers.
Date of Sampling: June 17, 1959. Description by: Grant Kennedy.
Location: Amador County, California, Bennett property, 1/4 mile E. of the center of Section 21, T6N, R12E, about 4 miles E of Jackson.
Elevation: 2,180 feet.
Relief: Gently sloping uplands on a plain north-facing slope of 7 percent below a ridge.
Drainage: Well drained; slow runoff; moderately slow permeability in the B_2.
Parent Material: Fine grained schist of the Calaveras formation.
Biological Activity: Worms and burrowing rodents active in these soils.
Climate: Mean annual rainfall 40".
Vegetation: Ponderosa pine, black oak, manzanita, poison oak.
Remarks: A medial developed Reddish-Brown Lateritic soil. Bureau of Public Roads samples were obtained for the A_{12}, B_2, and C_1 horizons.
Classification: Ultic Haploxeralf, fine-loamy, mixed, mesic.

HORIZON		DESCRIPTION
A_{00} & A_0 Not sampled	1-0"	Fresh and partly decomposed litter of leaves and pine needles.
A_{11}	0-2½"	Brown (7.5YR 5/4) loam, dark reddish brown (5YR 3/3) when moist; massive in place crushing to moderate very fine granular; slightly hard when dry, friable when moist, nonsticky and nonplastic when wet; abundant very fine and fine roots; many very fine and fine pores, few medium; few shot less than 5 mm. in diameter (magnetic); medium acid, pH 5.8; abrupt, smooth lower boundary.
A_{12}	2½-7"	Reddish brown (5YR 5/4) loam, dark reddish brown (5YR 3/4) when moist; massive in place crushing to moderate medium and and fine granular structure; hard when dry, friable when moist, nonsticky and nonplastic when wet; abundant very fine and fine roots, few medium and coarse; many very fine and fine pores, few medium; few shot; slightly acid, pH 6.1; gradual, smooth lower boundary.
A_3	7-13"	Reddish brown (5YR 5/4) loam, dark reddish brown (5YR 3/4) when moist; essentially massive in place breaking to subangular blocks when disturbed; hard when dry, friable when moist, slightly sticky and nonplastic; roots and pores as above except common medium pores; thin, discontinuous clay films in pores; occasional shot; medium acid, pH 5.9; gradual, smooth lower boundary.
B_1	13-27"	Reddish brown (5YR 5/4) light clay loam, dark reddish brown (5YR 3/4) when moist; massive; hard when dry, firm when moist, slightly sticky and slightly plastic when wet; abundant very fine and fine, occasional medium and coarse roots; many very fine and fine, common medium pores; thin, continuous clay films in pores; occasional shot; medium acid, pH 5.9; gradual, smooth lower boundary.
B_2	27-44"	Yellowish red (5YR 5/6) light clay, yellowish red (5YR 4/6) when moist; massive; hard when dry, very firm when moist, sticky and plastic when wet; few very fine and fine roots; common very fine, fine, and few medium pores; moderately thick, continuous clay films in pores and bridges; medium acid, pH 5.9; gradual, wavy lower boundary.
B_3	44-62"	Yellowish red (5YR 5/6) with some coatings of red (2.5YR 5/6) clay loam, yellowish red (5YR 4/6) when moist; massive; hard when dry, very firm when moist, slightly sticky and plastic when wet; roots, pores, and clay films as above but clay films are not as conspicuous; medium acid, pH 5.7; gradual, wavy lower boundary.
C_1	62-72"	Strong brown (7.5YR 5/6) loam with yellowish red (5YR 5/6) streaks, strong brown (7.5YR 4/6) when moist with red (2.5YR 4/8) streaks; massive; slightly hard when dry, firm when moist, slightly sticky and slightly plastic when wet; occasional root; few very fine and fine pores; thin, continuous clay films in some pores; strongly acid, pH 5.5; clear, somewhat irregular lower boundary.
D_r	72-84"+	Dark bluish gray somewhat weathered fine-grained schist with reddish brown and yellowish brown staining on fracture planes; strongly acid, pH 5.4

1/ The Josephine series is in a fine-loamy, mixed, mesic family of Typic Haploxerults. This pedon has an argillic horizon with a base saturation of more than 35 percent.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Jos phine loam (taxadjunct) **LOCATION** Amador County, California

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, mesic.

SOIL NOS. S60Calif-3-29 **LAB. NOS.** 60134 - 60142

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)							3A1		2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	
0- 1	A11	2.0	5.2	2.9	8.7	9.7	57.6	13.9	37.5	35.6	14	sil
1- 5	A12	1.8	5.8	2.9	8.9	10.8	54.4	15.4	36.5	34.7	17	sil
5-12	B1	5.6	6.6	3.8	8.3	10.0	46.1	19.6	30.0	31.4	23	l
12-25	B21	6.1	7.9	4.4	7.8	10.1	44.0	19.7	31.5	27.1	7	l
25-38	B22t	6.0	7.6	4.2	6.3	11.0	42.5	22.4	30.0	26.6	13	l
38-50	B23t	4.7	8.0	4.4	9.1	9.4	42.8	21.6	32.5	25.5	18	l
50-63	B24t	6.1	9.7	4.7	8.9	8.5	38.3	23.8	29.4	23.0	20	l
63-69	B25t	7.4	10.4	4.6	8.4	9.1	32.3	27.8	28.5	18.1	26	cl-l
69+	B3	Rock										

pH 8C1a H_2O 1:1	pH 8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	6C1b Free Fe_2O_3 Pct.	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.2	5.2		4.97	0.217	22.9				4.3			12.2
6.3	5.2		1.74	0.078	22.3				5.1			9.1
6.0	4.9		1.19	0.057	20.9				6.4			10.3
6.1	5.0		0.57	0.036	15.8				6.7			10.5
5.9	4.8		0.38	0.024					6.6			11.2
5.9	4.8		0.36	0.023					6.9			14.9
5.9	4.8		0.31						7.9			11.9
5.8	5.1		0.25									13.1

5A2a CATION EXCHANGE CAPACITY (+ Na)	EXTRACTABLE CATIONS 6N2b Ca	6O2c Mg	6H2a H	6P2a Na	5B1a 6Q2a K	BASE SAT. % 5C3	5A3a Sum of Cations eq/100g. soil	MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →							
27.7	11.1	5.4	13.1	<0.1	0.8	57	30.5	
16.1	5.5	3.3	7.2	0.1	0.6	57	16.7	
13.9	3.2	2.4	7.0	<0.1	0.5	47	13.2	
11.0	3.0	2.0	5.7	0.1	0.4	49	11.2	
10.8	2.5	2.1	4.8	0.1	0.3	51	9.8	
10.5	2.6	2.2	5.1	0.1	0.3	50	10.3	
11.8	3.4	2.8	5.2	0.1	0.3	56	11.8	
11.4	3.3	3.2	4.4	0.1	0.2	61	11.2	

1/ Percent of whole soil

JOSEPHINE LOAM (taxadjunct) 1/
S60Calif-3-29

Location: Amador County, California. 0.15 mile east of W1/4 corner of sec. 24, T. 7 N., R. 11 E., 100 feet north of Shake Ridge Road; 1/4 mile northwest of Golden Bare Health Ranch turnoff.
Date of Sampling: May 6, 1960.

Henry, Paul Nazar, and Jack McClelland.
Classification: Ultic Haploxeralf, fine-loamy, mixed, mesic.
Vegetation: Ponderosa pine, sugar pine, black oak, poison oak, mountain misery; herbs. Climate: Mean annual precipitation 32 inches; average temperatures (Tiger Creek Station): January - 40° F., July - 70° F.; mean annual - 55°F. Parent Material: Schists of the Calaveras formation with vertical cleavage. Relief: Steep uplands on a north-facing slope of 35 percent. Elevation: 2,090 feet.
Drainage: Well drained; moderate runoff; moderate permeability.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, mesic

SOIL Josephine loam (taxadjunct) SOIL Nos. S66Calif-47-1 LOCATION Siskiyou County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66334 - 66342

1B1b	Size class and particle diameter (mm) 3A1						3A1a	1/ Coarse fragments 1A2a	[redacted]
Total		Sand		Silt					

JOSEPHINE CLAY LOAM (taxadjunct) 1/
S66Calif-47-1

Location: Siskiyou County, California. A pit located below Trestle Road, and west of Hilt, California. From Hilt travel the road along the W. Fork of Cottonwood Creek. Turn right (N.) on to Trestle Rd. Site is located 1250 feet north and 750 feet west of the Southeast corner of Sec. 20, T48N, R7W. Date of Sampling: May 17, 1966.
Description by: J. J. Newlun. Collectors: K. W. Flach and J. J. Newlun.
Classification: Ultic Haploxeralf, fine-loamy, mixed, mesic.
Vegetation: Near pure stand of Douglas Fir with some ponderosa pine, sugar pine, and incense cedar. Hazel brush and serviceberry plus a small amount of perennial grass and forbs make up about 3 percent of the vegetative cover. The canopy cover is about 85 percent. The area has been both logged and burned in the past. Climate: The average annual precipitation is approximately 30 inches with much of it in the form of snow. The average January temperature is 32°F., the average July temperature 70°F., and the mean annual temperature 50°F. Parent Material: Residium from the sedimentary rock of the Chico Formation. This soil has developed from the sandstone in this formation. The sandstone has formed from mixed rock sources that were dominantly basic. Topography: The site is located on 26 percent slope, east aspect in mountainous upland. Elevation: 3520 feet. Drainage: Well drained, moderately slow permeability and medium runoff. Soil Moisture: Soil was near field capacity at time of description. Moisture was the same throughout the profile.
Remarks: Reactions were determined by the Truog soil reaction test. The profiles S66Calif-47-4 and S66Calif-47-5 are paired profiles. S66Calif-47-2, S66Calif-47-3 are similar soils. This profile has an ochric epipedon and an argillic horizon. The soil varies primarily in depth and colors. It is dominantly deep (36-60") with an average depth of about 40 inches.

HORIZON	DESCRIPTION
O1 & O2	1½ to 0 inches, new partially decomposed organic debris from coniferous trees. Abrupt smooth boundary.
A11 RSL 66334	0 to 2 inches, dark brown (10YR 4/3) light clay loam, very dark brown (10YR 2/2) moist; moderate medium to fine platy breaking to moderate fine granular structure; slightly hard, friable, slightly sticky and slightly plastic; many medium, fine and very fine roots; many random interstitial pores; strongly acid (pH 5.5); abrupt smooth boundary.
A12 RSL 66335	2 to 4 inches, dark brown (10YR 4/3) light clay loam, very dark brown (10YR 2/2) moist; moderate medium platy breaking to moderate medium subangular blocky structure; slightly hard, friable, sticky and plastic; many medium and fine and common very fine roots; common random interstitial and few tubular pores; strongly acid

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, mesic
SOIL Josephine loam (taxadjunct) SOIL Nos. S66Calif-47-2 LOCATION Siskiyou County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66343 - 66349

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand Very coarse (2-1)	Sand Coarse (1-0.5)	Sand Medium (0.5-0.25)	Sand Fine (0.25-0.1)	Sand Very fine (0.1-0.05)	Silt 0.05-0.02	Silt Int III (0.02-0.002)	Silt Int II (0.2-0.02)	Silt (2-0.1)			1/ Coarse fragments 1A2a > 2 Pct.	2-19 (Pct. of < 76mm)	19-76 (Pct. of < 76mm)
		Pct. of < 2 mm																
0-2	A11															5	5	-
2-9	A12															5	5	-
9-17	B21t															3	3	-
17-27	B22t															3	3	-
27-35	B3t															6	5	1
35-55+	C															45	28	17
0-4	A1 2/															4	4	-

Depth (In.)	6A1a Organic carbon Pct	6B1a Nitrogen Pct	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc		Water content Pct	4B1c 1/3 bar Pct	4B2 15 bar Pct	4D1 Extensibility COLE in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1		
0-2	1.59	0.075	21										7.8			6.1		
2-9	0.78	0.055	14										8.8			5.9		
9-17	0.43	0.042	10										9.3			5.7		
17-27	0.26												10.6			5.7		
27-35	0.12												7.4			5.7		
35-55+	0.20												7.5			5.8		
0-4	1.29	0.060	21										8.2			6.2		

Depth (In.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	5B1a Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A1a NH_4OAc	5A3a Sum		6G1d KCl Ext. Al^{+++}						Base saturation 5C1 NH_4OAc	5C3 Sum
	meq/100 g															Pct.	Pct.
0-2	8.5	1.9	0.1	0.8	11.3	7.8	15.3	19.1								74	59
2-9	7.5	1.9	0.1	0.9	10.4	6.2	13.9	16.6								75	63
9-17	6.7	2.6	0.1	0.4	9.8	6.1	13.8	15.9								71	62
17-27	8.2	3.7	0.1	0.4	12.4	5.0	15.9	17.4								78	71
27-35	7.1	3.3	0.1	0.3	10.8	3.9	13.4	14.7								81	73
35-55+	10.2	4.8	0.2	0.3	15.5	3.6	18.3	19.1								85	81
0-4	8.8	2.0	0.2	0.2	11.2	6.3	15.7	17.5								71	64

Depth (In.)		Clay mineralogy (<0.002 mm) 7A Mont.	Verm.	Mica	Kaolin	
		X-ray				

1/ From characterization sample (5 kg).

2/ Composite of 20 subsamples taken in a 15x12' grid adjacent to the master pedon.

JOSEPHINE LOAM (taxadjunct) 1/
S66Calif-47-2

Location: Siskiyou County, California. Below Trestle Road, and west of Hilt, California. From Hilt travel west on the road along the West Fork of Cottonwood Creek, then turn right on to Trestle Road. Site is located 1375 feet north and 300 feet west of the SE

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, mesic
SOIL Josephine loam (taxad.junct) SOIL Nos. S66Calif-47-3 LOCATION Siskiyou County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66350 - 66351

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand Very coarse (2-1)	Sand Coarse (1-0.5)	Sand Medium (0.5-0.25)	Sand Fine (0.25-0.1)	Sand Very fine (0.1-0.05)	Silt (0.05-0.02)	Silt Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)			1/ Coarse fragments 1A2a > 2 Pct.	2 - 19 (Pct. of < 76mm)	19 - 76 (Pct. of < 76mm)
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm																
0-4																14	14	-
17-26																7	7	-

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	Bulk density 4A1h Oven dry g/cc	g/cc		Water content Pct.	Water content 4B1c 1/3 bar Pct.	Water content 4B2 15 bar Pct	4D1 Extensibility COLE in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1		
0-4	1.80	0.074	24										6.1			6.6		
17-26	0.22												7.3			5.7		

Extractable bases 6N1a | 6H2a Cation Exch | 6G1d | Base saturation

JOSEPHINE LOAM (taxadjunct) 1/
S66Calif-47-3

Location: Siskiyou County, California. A pit located below Trestle Road west of Hilt, California. From Hilt travel west along road adjacent to the West Fork Cottonwood Creek and turn right (north) on to Trestle Road. The site is located 2400 feet east and 1125 feet south of NW corner Sec. 28, T48N, R7W. Date of Sampling: May 17, 1966.
Description by: J. J. Newlun. Collectors: K. W. Flach, S. B. Johnson, J. J. Newlun.
Classification: Ultic Haploxeralf, fine-loamy, mixed, mesic.
Vegetation: In order of dominance: Douglas fir, black oak, ponderosa pine, sugar pine, Oregon grape, snowberry, vetch and a few grasses and other forbs. Tree canopy is about 75 percent. The area has been logged and burned in the past. The trees are small and growth is slow. Climate: The average annual precipitation is approximately 30 inches with much of it in the form of snow. The average January temperature is 32°F., the average July temperature is 70°F., and the mean annual temperature is 50°F. Parent Material: Residium from sedimentary rocks of the Chico Formation. This soil has developed in the sandstone in this formation. The sandstone is fairly fine grained and weathers to heavy sandy clay loams in most cases. The sandstone has formed from mixed rock sediments but is dominantly basic. Topography: The site is located on mountainous upland with a single convex slope, a southeast aspect, a 23 percent gradient and is about midslope. Microrelief is negligible. Elevation: 3300 feet plus or minus 40 feet. Drainage: Well drained, moderately slow permeability, medium runoff. Soil Moisture: Soil was about 50 percent of field capacity throughout the profile at the time it was described.
Remarks: Only two horizons were sampled of this profile--the A11 (0-4") and B2t (17-26"). This profile represents a variation in the Josephine soils. It has less strongly developed horizons than the modal series in this County. The soil is also less deep to the C horizon which has only rock structure. The soil has a paralithic contact at 39 inches. The soil has a lower site index for timber than other profiles sampled because of the aspect and rooting depth limitation.

HORIZON	DESCRIPTION
	boundary.
A11 RSL 66350	0 to 4 inches, pale brown (10YR 6/3) loam, dark brown (10YR 3/3) moist; moderate fine granular structure; slightly hard, friable, slightly sticky and slightly plastic; many fine, very fine and medium roots; medium acid (pH 5.8); 5 percent gravels; clear smooth boundary.
A3	4 to 10 inches, pale brown (10YR 6/3) loam, dark brown (7.5YR 4/4) moist both rubbed and ped faces; weak medium subangular blocky breaking to moderate fine subangular blocky structure; hard, friable, slightly sticky and slightly plastic; many fine very fine and medium roots; many very fine tubular pores; medium acid (pH 6.0); clear smooth boundary.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, mesic

SOIL Josephine loam (taxadjunct) SOIL Nos. S66Calif-47-4 LOCATION Siskiyou County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66352 - 66361

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)		3A1a Non-carbonate clay	1/ Coarse fragments 1A2a > 2	2-19	19-76
		Size class and particle diameter (mm) 3A1																
		Pct. of < 2 mm														Pct.	Pct. of < 76mm	
0-2	A11	60.0	33.3	6.7	2.5	9.1	11.0	22.7	14.7	17.0	16.3	43.9	45.3		7	4	4	-
2-6	A12	61.2	31.0	7.8	2.5	9.0	11.3	22.3	16.1	14.8	16.2	43.1	45.1		8	4	4	-
6-10	A3	59.7	31.2	9.1	2.7	9.0	11.1	22.0	14.9	15.4	15.8	42.4	44.8		9	3	3	-
10-25	B1t	55.7	28.1	16.2	3.1	8.2	10.3	20.0	14.1	13.2	14.9	38.9	41.6		16	2	2	-
25-36	B21t	51.5	26.9	21.6	1.9	6.6	9.0	19.2	14.8	13.7	13.2	39.7	36.7		22	1	1	-
36-50	B22t	50.8	28.8	20.4	1.3	7.3	9.3	19.5	13.4	16.2	12.6	41.0	37.4		20	-	-	-
50-68	B23t	53.2	29.9	16.9	2.1	7.8	9.3	19.5	14.5	17.2	12.7	43.1	38.7		17	1	1	-
68-91	B3t	59.0	27.7	13.3	1.9	7.2	9.9	23.5	16.5	15.4	12.3	45.4	42.5		13	1	1	-
91-94+	C	72.2	22.1	5.7	1.3	13.5	16.0	26.9	14.5	11.2	10.9	40.2	57.7		6	-	-	-
0-6	A1 2/															4	4	-

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 C'	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1		
0-2	1.69	0.072	23				1.46	1.47		0.98		14.1	6.0	0.001	0.001	5.8		
2-6	0.68	0.043	16				1.52	1.52		0.98		12.9	5.3	0.001	0.001	5.7		
6-10	0.49	0.038	13				1.50	1.51		0.98		11.9	5.5	0.001	0.001	5.5		
10-25	0.20						1.60	1.62		0.99		12.5	8.2	0.003	0.003	5.6		
25-36	0.14						1.65	1.74		0.99		17.2	10.6	0.018	0.018	5.4		
36-50	0.12						1.66	1.77		1.00		17.7	10.6	0.021	0.021	5.4		
50-68	0.06						1.62	1.73		0.99		18.0	9.7	0.022	0.022	5.6		
68-91	0.12						1.59	1.66		0.99		14.7	8.6	0.014	0.014	5.8		
91-94+	0.29						1.82	1.87		1.00		13.6	5.9	0.010	0.010	5.9		
0-6	1.43	0.055	26										5.4			5.8		

Depth (in.)	Extractable bases 5B1a 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A1a NH_4OAc	5A3a Sum		6G1d KCl Ext. Al^{+++}						Base saturation 5C1 NH_4OAc	5C3 Sum
	meq/100 g															Pct	Pct.
0-2	2.4	0.6	0.1	0.6	3.7	17.2	11.8	20.9		0.1						31	18
2-6	4.3	1.0	0.1	0.8	6.2	4.7	9.6	10.9		0.2						64	57
6-10	3.6	1.0	0.1	0.6	5.3	4.5	8.7	9.8		0.2						61	46
10-25	6.3	2.3	0.1	0.4	9.1	3.7	12.5	12.8		0.3						73	71
25-36	9.2	3.4	0.1	0.6	13.3	5.3	17.5	18.6		0.4						76	72
36-50	11.8	4.1	0.1	0.6	16.6	5.0	20.6	21.6		0.3						81	77
50-68	11.7	4.0	0.2	0.4	16.3	4.8	23.6	21.1		0.4						69	77
68-91	14.1	4.2	0.2	0.2	18.7	3.8	22.4	22.5		0.3						83	83
91-94+	9.0	2.2	0.3	0.1	11.6	2.8	13.1	14.4		0.2						89	81
0-6	4.2	1.2	0.2	0.1	5.7	5.9	11.0	11.6		0.1						52	49

Depth (in.)	Clay mineralogy (<0.002 mm) 7A Mont.	Verm.	Illite	Kaolin	
	X-ray				
0-2			xx	t	
2-6					
6-10					
10-25					
25-36					
36-50		xxx		xx	
50-68					
68-91					
91-94+		xx		x	
0-6					

1/ From characterization sample (5 kg).

2/ Composite of 20 subsamples taken in a 15x12' grid adjacent to the master pedon.

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

JOSEPHINE LOAM (taxadjunct) 1/
S66Calif-47-4

Location: Siskiyou County, California. Pit located above Trestle Rd. near its junction with the road following the West Fork of Cottonwood Creek, 1750 feet east and 1200 feet south of the NW corner of Sec. 34, T48N, R7W. Date of Sampling: May 18, 1966.
Description by: J. J. Newlun. Collectors: K. W. Flach, S. B. Johnson, L. Mann.
Classification: Ultic Haploxeralf, fine-loamy, mixed, mesic.
Vegetation: In order of dominance: Ponderosa pine, sugar pine, black oak, white oak, Douglas fir, Incense cedar and various grasses and forbs. Very little understory vegetation. This is a fully stocked stand of trees which show good growth rate. Canopy is approximately 65 percent. The area has been logged and burned in the past. Climate: Average annual precipitation approximately 22 inches with much of it in the form of snow. Average January temperature 33°F., average July temperature 71°F., and mean annual temperature 51°F.
Parent Material: Residium from sedimentary rock of Chico Formation. This soil has developed from the sandstone of this formation. The sandstone is fairly fine textured and weathers to heavy sandy clay loam or silty clay loam in most cases. Sandstone has formed from mixed rock sources that were dominantly basic. Topography: The site is located on the upland foothills of the Klamath Mountains. Slope is single convex with a gradient of 16 percent and a north aspect. No significant microrelief is present. Elevation: 3040 feet plus or minus 40 feet. Drainage: Well drained, moderately slow permeability, medium runoff. Soil Moisture: About 75 percent of field capacity in A horizons and approximately 100 percent in B and C horizons.
Remarks: Site is representative of the soils developed at near minimum precipitation for this series. It was originally thought to be a member of the Parrish series but was correlated Josephine at the field level. Soil reaction determined by Truog soil reaction test. Profile has an ochric epipedon and an argillic horizon. Soil varies primarily in depth and colors. They are dominantly deep (36-60") with an average depth of about 40 inches. There may be a few quartzite gravels and cobbles throughout the profile.

HORIZON	DESCRIPTION
O1 & O2	1 to 0 inches, new and partially decomposed organic debris; abrupt smooth boundary.
A11 RSL 66352	0 to 2 inches, brown (10YR 5/3) loam, very dark grayish brown (10YR 3/2) moist and rubbed, moist ped faces are dark brown (10YR 3/3); weak fine granular structure; slightly hard, friable, slightly sticky and nonplastic; many fine and very fine and common medium roots; many random interstitial and common fine tubular pores; strongly acid (pH 5.5); abrupt smooth boundary.
A12 RSL 66353	2 to 6 inches, light brown (7.5YR 6/4) loam, dark brown (10YR 3/3) moist and rubbed, moist ped faces are dark brown (7.5YR 3/4); massive, appears to have traffic compaction; slightly hard, friable, slightly sticky and nonplastic; many fine and medium roots; common very fine and few fine tubular pores; strongly acid (pH 5.3); abrupt smooth boundary.
A3 RSL 66354	6 to 10 inches, light brown (7.5YR 6/4) loam, dark brown (10YR 3/3) moist and rubbed, ped faces moist are dark brown (7.5YR 3/4); weak medium subangular blocky to massive; slightly hard, friable, slightly sticky and slightly plastic; common fine and medium roots; many random interstitial and few fine tubular pores; few thin clay films in pores; strongly acid (pH 5.3); clear smooth boundary.
B1t RSL 66355	10 to 25 inches, light yellowish brown (10YR 6/4) heavy sandy clay loam or silty clay loam, dark brown (7.5YR 4/4) moist and rubbed, moist ped faces are dark yellowish brown (10YR 4/4); very weak medium prismatic breaking to weak medium subangular blocky structure; hard, friable, sticky and plastic; common fine, medium and coarse roots; many very fine and fine tubular pores; common thick and moderately thick clay films in pores; few thin clay films on ped faces; strongly acid (pH 5.3); clear smooth boundary.
B21t RSL 66356	25 to 36 inches, reddish yellow (7.5YR 6/6) heavy silty clay loam, dark yellowish brown (10YR 4/4) moist and rubbed with (10YR 4/4) and yellowish brown (10YR 5/4) moist ped faces; weak medium prismatic breaking to moderate medium subangular blocky structure; very hard, slightly firm, sticky and very plastic; many medium and coarse and common fine roots; common fine tubular pores; common thick and many moderately thick clay films on ped faces and in pores; strongly acid (pH 5.5); gradual wavy boundary.
B22t RSL 66357	36 to 50 inches, yellowish brown (10YR 5/6) silty clay loam, dark yellowish brown (10YR 4/4) moist, rubbed and ped faces; moderate medium subangular blocky structure; very hard, slightly firm, sticky and plastic; many medium and common fine roots; many fine tubular pores; many moderately thick clay films in pores and on ped faces and common thick films in pores; strongly acid (pH 5.5); gradual wavy boundary.
B23t RSL 66358	50 to 68 inches, same colors, structure, consistence, roots, and clay films as B22t. Texture is a light silty clay loam, medium acid (pH 5.8); and boundary clear and wavy. Few fine but distinct purplish black stains.
B3 RSL 66359	68 to 91 inches, yellowish brown (10YR 5/4) light loam, dark yellowish brown (10YR 4/4) moist rubbed and on ped faces; rock structure to massive; very hard, slightly firm, slightly sticky and nonplastic; few fine and common medium roots; many fine tubular and interstitial pores; very few thin clay films causing some bridging of sand grains; medium acid (pH 6.0); abrupt smooth boundary. Few distinct purplish black stains.
C RSL 66360	91 to 94 inches, essentially well weathered sandstone that is only slightly firm moist and very hard dry; common medium roots; few thin clay films on structural faces; medium acid (pH 5.8).

1/ The Josephine series is in a fine-loamy, mixed, mesic family of Typic Haploxerults. This series has a base saturation greater than 35 percent in the argillic horizon.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, mesic

SOIL Josephine loam (taxad.junct) SOIL Nos. S66Calif-47-5 LOCATION Siskiyou County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66362 - 66370

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)			1/ Coarse fragments 1A2a > 2 Pct.	2-19 (Pct. of < 76mm)	19-76 (Pct. of < 76mm)
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm																
0-2	A11															5	5	-
2-6	A12															4	4	-
6-18	A3															3	3	-
18-35	B1t															2	2	-
35-52	B21t															1	1	-
52-67	B22t															1	1	-
67-84	B31t															-	-	-
84-91+	B32t															1	1	-
0-6	A1 2/															7	7	-

Depth (In.)	6A1a Organic carbon Pct	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc		Water content: Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1a H_2O 1:1		
0-2	2.10	0.071	30										7.4			6.1		
2-6	2.43	0.065	37										7.3			6.0		
6-18	0.49	0.032	15										7.1			5.9		
18-35	0.27												8.0			5.7		
35-52	0.18												8.3			5.8		
52-67	0.12												8.2			5.8		
67-84	0.13												8.0			5.9		
84-91+	0.11												9.0			5.8		
0-6	2.19	0.083	26										7.3			6.2		

Depth (In.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1a NH_4OAc	5A3a Sum		6G1d KCl Ext. Al^{+++}						Base saturation: 5C1 NH_4OAc Pct	5C3 Sum Pct.
	meq/100 g																
0-2	7.3	1.3	0.1	0.8	9.5	7.2	13.4	16.7								71	57
2-6	6.9	1.3	0.1	0.8	9.1	7.0	12.8	16.1								71	57
6-18	4.2	1.3	0.1	0.6	6.2	4.7	8.7	10.9								71	57
18-35	4.2	1.7	0.1	0.4	6.4	3.7	9.1	10.1								70	63
35-52	5.0	2.0	0.1	0.3	7.4	3.7	10.1	11.1								73	67
52-67	4.6	2.0	0.1	0.3	7.0	3.6	9.3	10.6								75	66
67-84	4.5	2.0	0.1	0.3	6.9	3.4	9.4	10.3								73	67
84-91+	5.7	2.5	0.2	0.2	8.6	3.4	10.7	12.0								80	72
0-6	8.5	1.3	0.1	0.7	10.6	7.8	13.9	18.4								76	58

Depth (In.)	Clay mineralogy (<0.002 mm) 7A: Mont.	Verm.	Mica	Kaolin

1/ From characterization sample (5 kg).

[redacted]

JOSEPHINE LOAM (taxadjunct) 1/
S66Calif-47-5

Location: Siskiyou County, California, below Trestle Road, west of Hilt, California. From Hilt, travel the road along the west fork of Cottonwood Creek to Trestle Road, then turn right onto it. Site is 2000 feet east and 1700 feet south of the NW corner, Sec. 27, T48N, R7W. Date of Sampling: May 18, 1966.
Description by: J. J. Newlun. Collectors: K. W. Flach, S. B. Johnson, and L. Mann.
Classification: Ultic Haploxeralf, fine-loamy, mixed, mesic.
Vegetation: In order of dominance: Ponderosa pine, Douglas fir, black oak and a few forbs and grasses. The tree canopy is about 80 percent. Very little understory vegetation. Trees show good growth rate and size. Site area has been burned and logged in the past.
Climate: Average annual precipitation is approximately 22 inches with much of it in the form of snow. Average January temperature is 33°F., average July temperature 71°F., and mean annual temperature 51°F. Parent Material: Residium from sedimentary rock of the Chico Formation. Soil has developed from the sandstone in this formation. The sandstone is fairly fine textured and weathers to heavy sandy clay loams or silty clay loams. The sandstone has formed in sediments from mixed rock sediments that were dominantly basic. Topography: The site is located on a mountain foothill. The slope is single convex, 23 percent gradient, and aspect is 25° east of north. Elevation: 3040 feet plus or minus 40 feet. Drainage: Well drained, moderately slow permeability and medium runoff. Soil Moisture: Soil moisture was about 50 percent of field capacity throughout the profile.
Remarks: Soil reactions determined by Truog soil reaction test. Profile has an ochric epipedon and an argillic horizon. The soils are dominantly deep (36-60") with an average depth of about 40 inches. There are few quartzite gravels throughout the profile.

HORIZON	DESCRIPTION
01 & 02	1 to 0 inch, new and partially decomposed organic debris. Abrupt smooth boundary.
A11 RSL 66362	0 to 2 inches, brown (7.5YR 5/4) loam, dark brown (7.5YR 3/2) moist; weak medium platy structure; slightly hard, very friable, slightly sticky and nonplastic; many fine and very fine roots; many very fine and fine tubular pores; medium acid (pH 5.8); abrupt smooth boundary.
A12 RSL 66363	2 to 6 inches, light reddish brown (5YR 6/4) heavy loam, dark reddish brown (2.5YR 3/4) moist and rubbed, with dark reddish brown (5YR 3/4) moist ped faces; massive; slightly hard, friable, slightly sticky and slightly plastic; common fine and medium roots; many very fine and common fine tubular pores; medium acid (pH 5.8); abrupt smooth boundary.
A3 RSL 66364	6 to 18 inches, light reddish brown (5YR 6/4) heavy clay loam, dark reddish brown (5YR 3/3) moist and rubbed, with moist reddish brown (5YR 4/4) ped faces; moderate medium subangular blocky structure; hard, friable, sticky and plastic; many medium and common fine roots; common fine and very fine tubular pores; few thin clay films in pores and on ped faces; medium acid (pH 6.0); clear smooth boundary.
B1t RSL 66365	18 to 35 inches, reddish brown (5YR 5/4) heavy sandy clay loam, reddish brown (5YR 4/4) moist and rubbed with 60 percent yellowish red (5YR 4/6) and 30 percent dark reddish brown (5YR 3/4) on moist ped faces; weak medium prismatic structure breaking to moderate medium subangular blocky; hard, friable, sticky and very plastic; many medium and common fine roots; many fine tubular pores; few thin clay films in pores and on ped faces; strongly acid (pH 5.5); gradual wavy boundary. Clay films in this horizon occur mainly in pockets of peds.
B21t RSL 66366	35 to 52 inches, same colors as B1t, sandy clay loam; weak medium prismatic structure breaking to moderate medium subangular blocky; very hard, slightly firm, sticky and very plastic; many medium and coarse and common fine roots; many very fine, fine and medium tubular pores; few moderately thick clay films in pores; strongly acid (pH 5.5); clear irregular boundary. A few purplish black stains and concretions are present throughout the horizon.
B22t RSL 66367	52 to 67 inches, same colors as B1t, sandy clay loam; weak medium prismatic structure breaking to moderate medium subangular blocky; very hard, slightly firm, sticky and plastic; few fine roots; many fine, very fine and medium tubular pores; few thin clay films in pores; strongly acid (pH 5.5); gradual wavy boundary. Some purplish black stains and shot-like to irregular shaped concretions throughout horizon.
B31 RSL 66368	67 to 84 inches, same colors as B1t, heavy sandy loam; massive; very hard, slightly firm; common fine and medium roots; many very fine interstitial and common fine tubular pores; few thin and moderately thick clay films in pores; strongly acid (pH 5.5); abrupt wavy boundary.
B32 RSL 66369	84 to 91 inches, light reddish brown (5YR 6/4) heavy sandy loam, dark brown (7.5YR 4/4) moist and rubbed with yellowish red (5YR 4/6) moist ped faces; massive; very hard, slightly firm, sticky and plastic; common medium roots; common fine tubular and many very fine interstitial pores; few thin and moderately thick clay films in pores; strongly acid (pH 5.5); this horizon retains structure and fabric of the original rock.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Mariposa gravelly silt loam (taxadjunct) **LOCATION** Amador County, California

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, mesic

SOIL NOS. S60Calif-3-26 **LAB. NOS.** 60114 - 60119

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 >2 1/	TEXTURAL CLASS
0- 1	A11	7.5*	11.5*	4.1	8.1	5.6	54.1	9.1	29.0	35.5	34	sil
1- 5	A12	5.8*	10.3*	3.7	7.3	5.6	55.3	12.0	27.5	37.7	30	sil
5-10	B1t	4.1*	10.1*	3.9	7.1	5.4	54.8	14.6	25.2	39.1	25	sil
10-16	B21t	4.9*	8.9*	3.3	6.1	4.8	54.9	17.1	23.7	39.6	40	sil
16-22	B22t	2.1	3.8	1.8	1.6	3.8	65.7	21.2	13.4	56.4	6	sil
22-36+	R	Rock										

8C1a H_2O 1:1	pH 8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	4A3a Bulk Density g/cc	$CaCO_3$ equivalent %	6C1b "Free" Fe_2O_3 %	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
5.8	4.8		6.87	0.284	24.2				2.9			11.0
5.7	4.5		1.81	0.104	17.4		1.4		3.5			7.8
5.6	4.4		1.51	0.099	15.3		1.33		3.4			8.4
5.5	4.3		1.02	0.089	11.4		1.4		3.6			9.0
5.1	3.9		0.23						4.2			10.3

5A2a CATION EXCHANGE CAPACITY (+ Na)	EXTRACTABLE CATIONS 6N2b Ca	6O2c Mg	6H2a H	6P2a Na	6Q2a K	5B1a BASE SAT. % 5C3	5A3a Sum of Cations meq/100g. soil	MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →							
32.0	12.6	2.8	17.6	0.2	1.0	49	34.2	
15.4	4.7	1.2	8.9	0.1	0.6	43	15.5	
14.4	3.8	1.6	9.0	0.1	0.5	40	15.0	
13.8	3.3	2.3	7.8	0.1	0.4	44	13.9	
12.5	1.1	3.2	7.4	0.1	0.1	38	11.9	

Many iron and manganese concretions in these fractions.

1/ Percent of the whole soil

MARIPOSA GRAVELLY SILT LOAM (taxadjunct) 1/
S60Calif-3-26

Location: 0.2 mile NW of the S1/4 corner of sec. 36, T. 8 N., R. 12 E., 300 feet W of Markley Mine Road Junction with the Fiddletown-Silver Lake Road, 150 feet S of the Fiddletown-Silver Lake Road, Amador County, California.
Date of Sampling: May 6, 1960
Collectors: Grant Kennedy, Reuben Nelson, Robert Sketchley, Clifford Henry, Paul Nazar, Jack McClelland Description by: Grant Kennedy.
Vegetation: Manzanita locally with black oak, mountain misery, and ponderosa pine
Climate: Mean annual precipitation - 38 inches; average temperatures (Tiger Creek Station): January - 40° F.; July - 70° F.; mean annual 55° F.

Relief: Sloping uplands near a ridge crest, on a south-facing slope of 10 percent
Elevation: 2,960 feet
Drainage: Well drained; moderate runoff; moderate permeability
Remarks: This is a weakly developed Red-Yellow Podzolic soil. Many krotovena are evident. A few rock outcrops occur nearby. Roots are mainly horizontal. About 20 percent gravel was present to 16 inches. The soil was moist when described.
Classification: Ultic Haploxeralf, fine-loamy, mixed, mesic.

HORIZON		DESCRIPTION
01 & 02 NOT sampled	1- 0"	Mainly loose leaf litter, the lower part being partly decomposed.
A11 RSL No. 60114	0- 1"	Yellowish brown (9YR 5/4) gravelly loam, dark brown (7.5YR 3/4) when moist although upper 1/4" is very dark gray (N 3/) moist; massive; soft; friable, slightly sticky and slightly plastic; abundant very fine roots; many very fine interstitial and tubular pores; strongly acid (pH 5.5); abrupt smooth boundary.
A12 RSL No. 60115	1- 5"	Yellowish brown (9YR 5.5/4) gravelly silt loam near loam, dark brown (7.5YR 3/4) when moist; weak fine subangular blocky structure; soft, friable, slightly sticky and slightly plastic; abundant very fine, few fine and medium roots; many very fine interstitial and tubular, few fine and occasional medium pores; strongly acid (pH 5.5); clear smooth boundary.
B1t RSL No. 60116	5-10"	Light yellowish brown (9YR 6/4) gravelly silt loam near loam, dark brown (7.5YR 3/4) when moist; weak fine subangular blocky structure; slightly hard, friable, slightly sticky and slightly plastic; roots and pores as above; few thin discontinuous clay films in pores; strongly acid (pH 5.2); clear smooth boundary.
B21t RSL No. 60117	10-16"	Similar to above except gravelly clay loam; sticky and plastic; thin continuous clay films in pores; very strongly acid (pH 5.0); abrupt irregular boundary.
B22t RSL No. 60118	16-22"	Light yellowish brown (10YR 6/4) silty clay loam between stones and rock (80 percent), brown (7.5YR 4/4) when moist; massive;

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Musick sandy loam **LOCATION** Amador County, California

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, mesic

SOIL NOS. 59Calif-3-13 **LAB. NOS.** 59325 - 59331

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	3A1 0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
2-0	A_{oo} & A_o	Not	sampled									
0-9	A_1	4.7	15.0	11.1	18.3	9.6	26.8	14.5	32.0	14.8	3	sl
9-20	A_3	3.5	14.0	9.9	17.2	10.3	27.6	17.5	32.2	15.3	0	fsl
20-27	B_1	4.2	14.3	8.2	14.5	7.8	23.9	27.1	26.0	13.7	0	scl
27-39	B_{21}	3.5	13.5	8.0	15.7	7.6	16.0	35.7	24.9	7.6	0	sc-scl
39-58	B_{22}	5.4	15.3	9.0	18.6	9.0	19.8	22.9	29.9	9.7	0	scl
58-78	B_{23}	5.6	10.9	5.9	13.1	8.2	24.3	32.0	27.2	13.2	0	cl
78-100+	B_3	11.8	18.6	8.9	19.2	10.4	21.3	9.8	33.4	9.4	0	cosl

pH 8C1a H_2O 1:1	pH 8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	6C1b "Free" Fe_2O_3 %	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	Bulk Density 4A3a g/cc	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
5.8	4.6		3.92	0.152	25.8	2.2			1.09			10.0
5.9	4.7		0.50	0.032	15.6	2.9			1.22			8.9
5.8	4.9		0.27	0.019	14.2	4.3			1.36			12.9
6.0	5.5		0.12			5.7			1.43			16.2
5.8	5.2		0.08			4.3			1.54			12.7
5.6	5.1		0.13			5.4			1.30			17.0
5.4	4.2		0.04			3.5			1.31			8.7

MUSICK SANDY LOAM
S59Calif-3-13

Collectors: Grant Kennedy, Reuben Nelson, Bob Sketchley.
Date of Sampling: June 18, 1959. Description by: Grant Kennedy.
Location: Amador County, Calif., 1/2 mile south of the junction of Highway 88, and Westpoint Road in the NE1/4 of the NW1/4 of Section 1, T6N, R12E. (Stout Realty lot 6, subdivision next to the McFadden fence.)
Elevation: 2,680 feet.
Relief: Sloping uplands on a convex slope of 12 percent facing north.
Drainage: Well drained; slow runoff; moderate permeability.
Parent Material: Granodiorite.
Climate: Mean annual rainfall - 34 inches.
Vegetation: Ponderosa pine, black and live oak, blue wild rye.
Remarks: This is a medial developed Reddish-Brown Lateritic soil which in some areas includes small areas with a weakly developed A_2 horizon (Red-Yellow Podzolic?). The profile is micaceous throughout, particularly below the B_1. Samples were obtained for the Bureau of Public Roads of the following horizons: 9-20", 39-58", 78-100", the A_3, B_{22}, and B_3 horizons respectively.
Classification: Ultic Haploxeralf, fine-loamy, mixed, mesic.

HORIZON		DESCRIPTION
$A_{oo}A_o$ Not sampled	2-0"	Fresh litter of pine needles and leaves, becoming partially decomposed in the lower portion.
A_1	0-9"	Brown (7.5YR 4/4) sandy loam, dark brown (7.5YR 3/4) when moist; moderate fine and very fine granular structure; soft although slightly hard in upper 4 inches when dry, friable when moist; abundant very fine, fine, few medium and coarse roots; many very fine, fine, medium, and coarse pores; few very fine and fine shot (essentially nonmagnetic); medium acid, pH 5.8; clear, smooth lower boundary.
A_3	9-20"	Yellowish red (5YR 5/6) fine sandy loam, reddish brown (5YR 4/4) when moist; massive; slightly hard when dry, friable when moist, slightly plastic and nonsticky when wet; roots and pores as above; thin discontinuous clay films in pores; few very fine shot; medium acid, pH 5.9; gradual, smooth lower boundary.
B_1	20-27"	Yellowish red (5YR 5/6) sandy clay loam, yellowish red (5YR 3/6) when moist; similar to above except clay films are thin and continuous in pores; gradual, smooth lower boundary.
B_{21}	27-39"	Yellowish red (5YR 5/8) sandy clay near sandy clay loam, dark red (2.5YR 3/8) when moist; massive; hard when dry, firm when moist, slightly plastic, slightly sticky when wet; plentiful very fine, fine and medium, few coarse roots; many very fine, fine and medium, few coarse pores; moderately thick continuous clay films in pores and bridges; moderately thick in some pores; medium acid, pH 6.0; gradual, smooth lower boundary
B_{22}	39-58"	Yellowish red (5YR 4/8) sandy clay loam, dark red (2.5YR 3/6) when moist; similar to above horizon except fewer roots and pores are present, clay films are thinner, and the pH is 5.8; gradual, smooth lower boundary.
B_{23}	58-78"	Light red (2.5YR 6/8) clay loam, red (2.5YR 4/8) when moist; massive; slightly hard when dry, firm when moist, slightly plastic and slightly sticky when wet; few very fine, fine, and medium roots; many very fine and fine, few medium and coarse pores; thin continuous clay films in pores and bridges, moderately thick in some pores; medium acid, pH 5.6; gradual, smooth lower boundary.
B_3	78-100"+	Reddish yellow (5YR 7/8) coarse sandy loam, red (2.5 R 5/8) when moist; massive; soft when dry, firm when moist, slightly plastic and nonsticky when wet; few very fine to medium roots and pores; thin, continuous clay films in most pores; strongly acid, pH 5.4.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Musick loam **LOCATION** Amador County, California

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, mesic

SOIL NOS. 59Calif-3-14 **LAB. NOS.** 59332 - 59339

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE	COARSE	MEDIUM	FINE	VERY FINE						

MUSICK LOAM
S59Calif-3-14

Collectors: Grant Kennedy, Reuben Nelson, Bob Sketchley.
Date of Sampling: June 18, 1959. Description by: Grant Kennedy.
Location: Amador Co., California in the SE1/4 of the SE1/4 of Section 19, T7N, R13E, 1/4 mile NE of Pioneer, follow black top across from the Post Office for 1/4 mile, then turn left on Schafer Road for 1/4 mile.
Elevation: 2,960 feet.
Relief: Steep uplands just below a ridgetop on a convex slope of 20 percent facing NNE.
Drainage: Well drained; slow runoff; moderate permeability.
Parent Material: Granodiorite.
Climate: Mean annual rainfall 37".
Vegetation: Ponderosa pine, incense cedar, black oak, madrona, sweet birch, mountain misery, ripgut, lupine.
Remarks: This is a medial Reddish Brown Lateritic soil, containing a few cobbles in the profile. In this area a few rock outcrops occur. The site has been cultivated in the past. Samples were obtained for the Bureau of Public Roads of the following horizons: 0-14", 29-47" and 74-97" (A_p, B_{22}, and C_1 horizons).
Classification: Ultic Haploxeralf, fine-loamy, mixed, mesic.

HORIZON		DESCRIPTION
A_p	0-14"	Reddish brown (5YR 4/4) loam, dark reddish brown (5YR 3/4) when moist; moderate very fine and fine granular structure; soft when dry, friable when moist; abundant very fine, fine and medium, plentiful coarse roots; many very fine, fine and medium, common coarse pores; common very fine and fine shot (now magnetic); medium acid, pH 6.0; gradual wavy lower boundary.
B_1	14-23"	Yellowish red (5YR 4/6) heavy loam, dark red (2.5YR 3/6) when moist; massive; slightly hard when dry, friable when moist, slightly plastic and nonsticky when wet; roots and pores as in horizon above; few thin discontinuous clay films, mainly in pores; medium acid, pH 5.7; clear, smooth lower boundary.
B_{21}	23-29"	Red (2.5YR 4/8) heavy clay loam, dark red (2.5YR 3/8) when moist; massive; hard when dry, firm when moist, plastic and sticky when wet; common very fine, fine, and medium roots, few coarse; many very fine and fine, occasional medium and coarse roots; many moderately thick continuous clay films in pores and in bridges; strongly acid, pH 5.5; gradual, smooth lower boundary.
B_{22}	29-47"	Similar to above except moist color is red (10R 4/6); gradual smooth lower boundary.
B_{31}	47-60"	Light red (2.5YR 6/8) sandy loam, red (2.5YR 4/8) when moist; massive; slightly hard when dry, friable when moist; slightly sticky and nonplastic when wet; few roots; common very fine, fine, and medium pores; few discontinuous clay films in pores and in bridges; medium acid, pH 5.6; diffuse lower boundary.
B_{32}	60-74"	Similar to above but coarse sandy loam texture and pH 5.5; clear, irregular lower boundary.
C_1	74-97"	Reddish yellow (5YR 6/6) fine sandy loam with light red (2.5YR 6/8) staining on fracture planes, red and dark red (2.5YR 4/8 & 3/8) when moist; massive; slightly hard when dry, friable when moist; few roots; many very fine and fine pores; strongly acid, pH 5.4; abrupt irregular lower boundary.
D_r	97"+	Speckled white, reddish yellow, and very dark gray weathered

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Boomer loam (taxadjunct) **LOCATION** El Dorado County, California

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, thermic.

SOIL NOS. S61Calif-9-3 **LAB. NOS.** 6193 - 6198

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 5	A1	6.9	14.1	5.7	9.0	5.1	39.4	19.8	23.0	26.5	26	l
5-13	A3	5.0	11.8	4.9	8.5	5.4	39.5	24.9	22.4	27.4	21	l
13-24	B21t	5.1	13.5	5.1	7.6	4.3	31.4	33.0	18.4	21.4	31	cl
24-37	B22t	10.7	22.7	6.2	7.9	3.9	20.2	28.4	14.8	13.6	40	scl
37-52	B31	19.6	21.0	4.9	5.3	2.8	20.4	26.0	10.6	15.4	64	scl
52-63+	B32 & R	25.7	19.1	4.4	4.6	2.9	21.1	22.2	10.8	15.7	81	scl

pH 8C1b Saturated Paste 1:1	pH 8C1c KCl 1:1	Organic Matter 6A1a Organic Carbon Pct.	Organic Matter 6B1a Nitrogen Pct.	C/N	6C1b Free Fe_2O_3 %	4A3a Bulk Den. gm/cc	4B2 Soil water 15 atmos. Pct.
6.6	5.4	2.52	0.137	18.4	7.6	1.28	12.8
6.2	4.7	0.99	0.056	17.7	8.5	1.37	13.2
5.9	4.4	0.60	0.036	16.7	8.7	1.47	16.8
5.8	4.0	0.26	0.015		6.2	1.53	16.9
6.1	4.1	0.10			5.5		13.7
6.2	4.1	0.08			4.2		12.9

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6Q2c	6H2a H	Na 6P2a	K 6Q2a	BASE SAT. % 5C3	5A3a Sum of Cations soil	Calc. Sum of Cations clay	Calc. Sum of Cations clay + fine silt
	milliequivalents per 100g. soil						milliequivalents per 100g.		
26.9	11.9	6.1	8.2	0.2	0.9	70	27.3	138	46
20.4	9.7	6.0	9.1	0.2	0.6	65	25.6	103	40
27.7	11.7	7.1	8.7	0.2	0.3	69	28.0	85	43
36.0	7.2	12.6	7.7	0.2	0.2	72	27.9	98	57
24.3	14.3	11.7	5.9	0.2	0.2	82	32.3	124	70
29.2	13.6	11.8	4.9	0.2	0.1	84	30.6	138	71

1/ Percent of whole soil <1 inch

BOOMER LOAM (taxadjunct) 1/
S61Calif-9-3

Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, David Estrada, Lynn Brittan, Lothair Grant Description by: Grant Kennedy.
Date of Sampling: April 27, 1961

Location: El Dorado County, 6 miles northwest of Coloma, California, NE corner of NW1/4SW1/4 of sec. 33, T. 12 N., R. 9 E., El Dorado County.
Elevation: 1040 feet
Relief: Steep uplands with a 40 percent slope to the east.
Drainage: Well drained; medium runoff; moderate permeability
Parent Material: Basic Schist
Climate: Mean annual rainfall 35", T 60°, J 44°, J 77°
Vegetation: Ponderosa pine, black oak, blue wild rye, poison oak, lupine, and annual grasses
Classification: Ultic Haploxeralf, fine-loamy, mixed, thermic.
Remarks: The B31 and B32&R horizons are well-fractured schists, breaking to [illegible]

B32&R are the clay coatings. The color of the B horizons dry are between hue 2.5YR and 5YR. Horizons 1, 4, and 5 (A1, B22t, and B31) were sampled for the Bureau of Public Roads and the California State Division of Highways.

HORIZON		DESCRIPTION
O1 & O2	1- 0"	Pine needles and decomposed litter
A1	0- 5"	Yellowish red (5YR 5/6) loam, dark reddish brown (5YR 3/4) when moist; moderate fine granular; slightly hard when dry, friable when moist, and slightly sticky and slightly plastic when wet; abundant very fine, plentiful fine roots; many very fine tubular and interstitial, common fine tubular pores; medium acid, pH 6.0; clear smooth boundary.
A3	5-13"	Yellowish red (5YR 5/6) gravelly clay near loam, yellowish red (5YR 3/6) when moist; moderate medium subangular blocky breaking to weak fine to medium granular; slightly hard when dry, friable when moist, [illegible]
		tiful medium roots; many very fine and fine tubular and interstitial, few medium pores; few thin clay films in pores; medium acid, pH 6.0; clear smooth boundary.
B21t	13-24"	Red (2.5YR 4/6) gravelly clay loam, dark red (2.5YR 3/6) when moist; moderate fine to medium subangular blocky; slightly hard when dry, friable when moist, sticky and plastic when wet; plentiful very fine to coarse roots; many very fine tubular and interstitial pores; few fine tubular pores; many thin clay films in pores; medium acid, pH 6.0; clear smooth boundary.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Boomer loam (taxadjunct) **LOCATION** El Dorado County, California

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, thermic.

SOIL NOS. S61Calif-9-4 **LAB. NOS.** 6199 - 61105

DEPTH INCHES 1	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)							3A1		2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	
0- 8	A_1	6.3	10.3	4.8	8.4	6.0	40.7	23.5	23.7	28.0	30	l
8-15	A_3	5.8	9.7	5.0	8.3	6.4	39.1	25.7	22.7	27.8	34	l
15-23	B_{21t}	7.6	11.0	4.0	5.9	4.9	37.4	29.2	19.0	26.8	26	cl
23-36	B_{22t}	3.9	8.4	4.8	8.8	6.3	37.2	30.6	22.8	26.1	19	cl
36-46	B_{31}	4.4	11.1	7.0	14.6	8.7	35.6	18.6	31.8	21.5	1	l
46-58	B_{32} & R	0.8	7.0	8.2	20.2	10.5	37.2	16.1	38.7	21.0	0	l

BOOMER LOAM (taxadjunct) 1/
S61Calif-9-4

Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, David Estrada, Lynn Brittan, Lothair Grant **Description by: Grant Kennedy.**
Date of Sampling: April 28, 1961
Location: El Dorado County, California, 5 miles west of Coloma along Highway 49, SE1/4SW1/4 sec. 4, T. 11 N., R. 9 E.
Elevation: 900 feet
Relief: Steep uplands with a 38 percent slope to the northwest.
Drainage: Well drained, medium runoff; moderate permeability
Parent Material: Basic Schist
Climate: Mean annual rainfall 35 inches, T 60°, J 44°, J 77°
Vegetation: Ponderosa pine, black oak, Douglas fir, manzanita, poison oak, blue wild rye, lupine and other forbs.
Classification: **Ultic Haploxeralf, fine-loamy, mixed, thermic.**
Remarks: Some flattened medium and coarse roots extend to 70 inches in crevices and old fracture planes. The C1 horizon breaks into fractured pieces, gravel size, that are all coated with thin to moderately thick clay films, the R is similar but less weathered and shows less clay films. The dry color of the B horizons are between hue 2.5YR and 5YR. Horizons 1, 4, and 5 (A1, B22t, and B31) were sampled for the Bureau of Public Roads and the California State Division of Highways.

HORIZON		DESCRIPTION
O1 & O2	1-0"	Pine needles and decomposed litter.
A1	0-8"	Yellowish red (5YR 5/6) loam, dark reddish brown (5YR 3/4) when moist; moderate fine to medium granular; slightly hard when dry, friable when moist; slightly sticky and slightly plastic when wet; abundant very fine plentiful fine to medium roots; many very fine, common fine tubular and interstitial pores; medium acid, pH 6.0; gradual smooth boundary.
A3	8-15"	Yellowish red (5YR 5/6) loam near clay loam, dark reddish brown (2.5YR 3/4) when moist; moderate fine to medium granular; slightly hard when dry, friable when moist, slightly sticky and slightly plastic when wet; abundant very fine to medium and plentiful coarse roots; many very fine tubular and interstitial, common fine tubular and few medium tubular pores; few thin clay films in pores; slightly acid, pH 6.3; clear slightly wavy boundary.
B21t	15-23"	Red (2.5YR 4/6) clay loam, dark red (2.5YR 3/6) when moist; moderate very fine to fine subangular blocky; hard when dry, friable when moist; sticky and plastic when wet; plentiful very fine to coarse roots; many very fine tubular and interstitial, common fine tubular and few medium tubular pores; many thin clay films in pores; slightly acid, pH 6.3; clear slightly wavy boundary.
B22t	23-36"	Red 2.5YR 4/8) clay loam near clay, dark red (2.5YR 3/6) when moist; moderate very fine to fine subangular blocky; hard when dry, firm when moist, sticky and plastic when wet; few fine to coarse roots; common very fine tubular few fine to coarse tubular pores; continuous thin clay film in pores; slightly acid pH 6.3; clear slightly wavy boundary.
B31	36-46"	Reddish yellow (7.5YR 6/6) clay loam, yellowish red (5YR 4/6) with red (2.5YR 4/6) coatings; weak very fine to fine subangular blocky; hard when dry, firm when moist, and sticky and plastic when wet; few fine to coarse roots; common very fine pores; common thin clay films in pores; slightly acid pH 6.2; gradual smooth boundary.
B32 & R	46-50"	Reddish yellow (7.5YR 6/6), yellowish red (5YR 5/8) with red (2.5YR 4/6) coatings, when moist; massive, very hard when dry, firm when moist; see remarks on clay films; slightly acid, pH 6.2; gradual smooth boundary.
R	50-70"+	Strong brown (7.5YR 5/8) weathered basic schist, yellowish red (5YR 5/8) with black manganese stains, with red (2.5YR 4/6) coatings; massive; very hard when dry, firm when moist; see remarks on clay films.

1/ Labeled a taxadjunct because soil temperature is marginal to mesic and thermic. The Boomer series has a mesic temperature regime.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Ramona fine sandy loam (taxadjunct) **LOCATION** Riverside County, Calif.

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, thermic

SOIL NOS. S59Calif-33-1 **LAB. NOS.** 59548 - 59555

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-14	A1p	5.3	13.2	8.5	19.0	13.7	31.6	8.7	43.9	13.1	5	fsl
14-23	A12	5.4	11.8	7.0	16.0	12.6	33.3	13.9	40.4	15.4	3	fsl-l
23-29	B1	6.3	11.2	6.6	14.6	13.1	32.3	15.9	39.1	15.2	2	l-fsl
29-37	B21	6.0	10.7	6.5	15.4	12.8	30.5	18.1	38.9	14.3	5	l-fsl
37-46	B22	5.9	12.2	7.2	16.7	12.3	26.7	19.0	36.9	12.6	4	fsl
46-58	B23	5.3	10.5	7.7	19.5	13.2	25.6	18.2	38.7	12.4	4	fsl
58-68	B3	6.6	11.8	7.7	16.8	11.4	27.1	18.6	35.1	13.9	9	fsl
68-74	C	3.8	7.9	6.0	18.1	15.4	30.7	18.1	43.5	14.6	4	l-fsl

pH 8C1b SATURATED PASTE 1:1	pH 1:5	pH 8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	ORGANIC MATTER 6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. mmhos/cm. at 25° C.	$CaCO_3$ equivalent %	Ext. Iron % Fe 6C1b	6I1a CO_3 meq/l	6J1a HCO_3 meq/l	4B2 S il water 15 atmos. Pct.
5.9		6.6	0.40	0.046	8.7		1.1		1.1	-	0.7	4.4
6.4		6.7	0.45	0.049	9.2		0.3		1.3	-	0.6	6.2
6.6		6.9	0.34	0.044	7.7		0.4		1.3	-	0.6	6.8

RAMONA FINE SANDY LOAM (taxadjunct) 1/
S59Calif-33-1

Location: Approximately 1100 feet north, 500 feet west of the south quarter corner of section 31, T. 2 S., R. 1 W., Riverside County, California, about 3 miles northwest of the town of Banning.
Date: September 10, 1959
Description By: Arnold Knecht and George Borst
Samples Collected By: Arnold Knecht, Fred Eggers and Reuben Nelson
Classification: Ultic Haploxeralf, fine-loamy, mixed, thermic.
Vegetation: Grain stubble
Climate: Average annual precipitation 16 inches; average January temperature 50°F., average July temperature 70°F. and average annual temperature about 62° F.
Parent Material: Moderately coarse alluvium predominately from granitic sources.
Topography: Gently sloping alluvial terrace, with about 3 percent west-southwesterly slope. Elevation: 2,410 feet.
Soil Moisture: Moist below 12 inches Drainage: Well drained.
Remarks: This profile was described about 1300 feet southwest of the location at which surface soil samples were collected for nutrient level determination by the University of California. It is characterized by an ochric epipedon and an argillic horizon, according to the revised system of soil classification (7th Approximation).

HORIZON		DESCRIPTION
A1p RSL No. 59548	0-14"	Brown (10YR 5/3) fine loam, dark brown (10YR 3/3) moist; massive; hard, very friable, nonsticky and nonplastic; few fine roots; many very fine interstitial and tubular pores; medium acid (pH 6.0); clear, smooth boundary.
A12 RSL No. 59549	14-23"	Brown (10YR 5/3) fine sandy loam, dark brown (10YR 3/3) moist; massive, hard, very friable, slightly sticky and slightly plastic; few fine roots; common fine tubular pores; slightly acid (pH 6.5); clear, smooth boundary.
B1 RSL No. 59550	23-29"	Brown (7.5YR 5/4) loam, dark reddish brown (5YR 3/4) moist; moderate coarse angular blocky structure; hard, friable, slightly sticky and slightly plastic; few fine roots; many fine tubular pores; few thin clay films on ped faces and in pores; slightly acid (pH 6.5); clear smooth boundary.
B21 RSL No. 59551	29-37"	Reddish brown (5YR 4/4) clay loam, dark reddish brown (5YR 3/4) moist; moderate coarse prismatic structure; very hard, firm, sticky and plastic; few fine roots; common fine tubular pores; common thin clay films on ped faces and in pores; slightly acid (pH 6.5); gradual, smooth boundary.
B22 RSL No. 59552	37-46"	Yellowish red (5YR 5/6) clay loam, yellowish red (5YR 4/6) moist; moderate coarse prismatic structure; very hard, firm, sticky and plastic; very few fine roots; few fine tubular pores; many moderately thick clay films on ped faces and in pores; slightly acid (pH 6.5); gradual smooth boundary.
B23 RSL No. 59553	46-58"	Yellowish red (5YR 5/6) clay loam, yellowish red (5YR 4/6) moist; moderate coarse prismatic structure; very hard, firm, sticky and plastic; few fine tubular pores; many moderately thick clay films on ped faces and in pores; slightly acid (pH 6.5); gradual smooth boundary.
B3 RSL No. 59554	58-68"	Yellowish red (5YR 5/6) clay loam, yellowish red (5YR 4/6) moist; moderate coarse angular blocky structure; very hard, firm, sticky and plastic; few fine tubular pores; many moderately thick clay films on ped faces and in pores; neutral (pH 6.8); clear, irregular boundary.

OIL SURVEY LABORATORY Riverside, California

SOIL TYPE Red Bluff gravelly loam variant **LOCATION** Amador County, California

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, thermic

SOIL NOS. 59Calif-3-10 **LAB. NOS.** 59320 - 59324

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	3A1 0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-1	A_{11}	9.3	13.5	6.0	9.9	6.2	37.0	18.1	25.1	23.9	35	l
1-9	A_{12}	9.0	12.2	5.9	10.1	7.1	36.3	19.4	25.9	23.6	34	l
9-22	B_1	6.8	12.7	6.0	8.6	6.4	32.7	26.8	22.6	21.6	34	l-cl
22-31	B_2	13.4	17.2	4.6	4.5	4.4	24.7	31.2	14.6	17.1	33	cl-scl
31-42+	D*	11.3	7.3	2.7	5.3	14.8	33.4	25.2	36.8	15.0	3	l

pH 8C1a. H_2O 1:1	pH 8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	6C1b "Free" Fe_2O_3 %	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	GYPSUM me./100g. SOIL	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.4	5.7		4.86	0.268	18.1	5.0						10.7
6.7	5.8		1.89	0.150	12.6	5.2						9.1
6.0	5.0		0.60	0.058	10.3	6.9						10.7
5.2	4.2		0.21	0.028		7.8						13.0
4.5	3.2		0.24	0.018		2.8						13.0

RED BLUFF GRAVELLY LOAM variant 1/
S59Calif-3-10

Collectors: Grant Kennedy, Reuben Nelson, Bob Sketchley.
Date of Sampling: June 16, 1959. Description by: Grant Kennedy.
Location: Amador County, California near Buena Vista School, approximately 1/4 mile west of the NE corner of Section 18, T5N, R9E.
Elevation: 300 feet.
Relief: Gently rolling, high terrace on a W-facing slope of 10 percent.
Drainage: Well drained; moderate runoff; moderate over slow permeability.
Parent Material: Arroyo Seco gravels resting on the Ione formation.
Climate: 20 inches, mean annual rainfall.
Vegetation: Oak-grass-wedge leaf ceanothus.
Remarks: This is a medial developed Noncalcic Brown soil which is gravelly

The gravels & cobbles consist of slate, greenstone, quartz, etc.
Classification: Ultic Haploxeralf, fine-loamy, mixed, thermic.

HORIZON		DESCRIPTION
A_{11}	0-1"	Reddish brown (5YR 4/4) light gravelly loam, dark reddish brown (5YR 3/4) when moist; moderate fine and medium granular structure; soft when dry, friable when moist; abundant very fine roots; many fine and very fine pores; slightly acid, pH 6.4; abrupt, smooth lower boundary.
A_{12}	1-9"	Yellowish red (5YR 4/6) gravelly loam, dark reddish brown (5YR 3/4) when moist; massive or very weak very fine subangular blocky; slightly hard when dry, friable when moist; abundant very fine, fine and medium roots; many very fine, fine, common medium, few coarse pores; neutral pH 6.7; clear, smooth lower boundary.
B_1	9-22"	Reddish brown (2.5YR 4/5) gravelly loam near gravelly clay loam, dark reddish brown (2.5YR 3/5) when moist; massive; slightly hard when dry, friable when moist, slightly sticky and slightly plastic when wet; plentiful very fine to coarse roots; many very fine and fine, common medium, few coarse pores; thin continuous clay films in pores and bridges;

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Sierra coarse sandy loam **LOCATION** Amador County, California

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, thermic

SOIL NOS. 59Calif-3-15 **LAB. NOS.** 59355 - 59361

		1B1b	PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)	3A1	

SIERRA COARSE SANDY LOAM
S59Calif-3-15

Collectors: Grant Kennedy, Bob Sketchley, and John Rogers.
Date of Sampling: September 4, 1959. Description by: Grant Kennedy.
Location: Amador County, California on Jameson Ranch. 60 yards NE of intersection of Shenandoah Road. Shenandoah School Road in the NE1/4 SE1/4 of Section 19, T8N, R11E.
Elevation: 1,610 feet
Relief: On gently sloping uplands with a convex slope of about 9 percent to south
Drainage: Well drained, moderate runoff, moderate to moderately slow permeability.
Parent Material: Granodiorite.
Climate: Mean annual rainfall is 33 inches.
Vegetation: Annual grasses, mainly soft chess and wild oats. Field formerly cultivated over 8 years past, now reverted to annual grassland. Field pastured annually by sheep.
Remarks: Profile dry to 29 inches. Below 29" the moisture increases with depth to almost field capacity in the B_3. Occasional rock outcrops in the area. Some 2-inch Krotovinas in the B_{21}.
Classification: Ultic Haploxeralf, fine-loamy, mixed, thermic.
This soil is a medial developed Reddish Brunizem. Samples were obtained for the Bureau of Public Roads on the A_p, B_{22}, and C_1 horizons. Oriented ring samples were obtained on all but the A_p horizon.

HORIZON		DESCRIPTION
A_p	0-6"	Brown (7.5YR 5/4) coarse sandy loam, dark reddish brown (5YR 3/3) when moist; massive; slightly hard when dry, friable when moist, nonsticky and nonplastic when wet; abundant very fine roots; many very fine and common fine tubular pores; medium acid, pH 6.0; abrupt, smooth lower boundary.
A_3	6-15"	Yellowish red (5YR 5/6) loam, dark red (2.5YR 3/6) when moist; massive; hard when dry, friable when moist, slightly sticky and slightly plastic when wet; plentiful very fine roots; many very fine and common fine tubular pores; few thin discontinuous clay films in pores and bridges; medium acid, pH 5.9; clear, smooth lower boundary.
B_1	15-24"	Yellowish red (5YR 5/6) loam, dark red (2.5YR 3/6) when moist; massive; very hard when dry, friable when moist, slightly sticky and slightly plastic when wet; few very fine roots; many very fine, common fine tubular pores; few thin continuous clay films in pores and adjacent areas; medium acid pH 6.0; clear smooth lower boundary.
B_{21}	24-33"	Red (2.5YR 5/6) loam, dark red (2.5YR 3/6) when moist; massive; very hard when dry, firm when moist, sticky and slightly plastic when wet; very few fine roots; many very fine and common fine tubular pores; common thin continuous clay films in pores and adjacent areas; slightly acid, pH 6.2; gradual smooth lower boundary.
B_{22}	33-45"	Red (2.5YR 5/6) loam, dark red (2.5YR 4/6) when moist; massive; very hard when dry, firm when moist, sticky and plastic when wet, very few fine roots; few very fine tubular pores; common thin continuous clay films in pores and adjacent areas; slightly acid, pH 6.3; clear smooth lower boundary.
B_3	45-66"	Yellowish red (5YR 5/6) loam, dark red (2.5YR 4/6) when moist; massive; very hard when dry, firm when moist, sticky and plastic when wet; few very fine tubular pores; very few fine roots; common thin continuous clay films in pores and adjacent areas; slightly acid, pH 6.5; abrupt wavy lower boundary.
C_1	66-73"	Reddish yellow (7.5YR 7/6) coarse sandy loam, yellowish red (5YR 5/6) when moist; massive; hard when dry, friable when moist; slightly sticky and nonplastic when wet; few very fine pores; thin discontinuous clay films in pores and coatings and grains; medium acid, pH 6.0.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Sierra coarse sandy loam **LOCATION** Amador County, California

SOIL FAMILY: Ultic Haploxeralf, fine-loamy, mixed, thermic

SOIL NOS. 59Calif-3-16 **LAB. NOS.** 59349 - 59354

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1	2A2	TEXTURAL CLASS
		VERY COARSE SAND	COARSE SAND	MEDIUM SAND	FINE SAND	VERY FINE SAND	SILT	CLAY			> 2	

SIERRA COARSE SANDY LOAM
S59Calif-3-16

Collectors: Grant Kennedy, Reuben Nelson, Bob Sketchley, John Rogers.
Date of Sampling: June 20, 1959. Description by: Grant Kennedy.
Location: Amador County, California on the Ken Deaver Ranch, 1/2 mile south of Shenandoah School on the Shenandoah School Road in the SW1/4 of the SW1/4 of Section 29, T8N, R11E.
Elevation: 1,560 feet.
Relief: On gently sloping uplands with a convex slope of about 9 percent to the north.
Drainage: Well drained, slow runoff, moderate to moderately slow permeability.
Parent Material: Granodiorite.
Biological Activity: Burrowing rodents and earthworms are active in these soils.
Climate: Mean annual rainfall is 33 inches.
Vegetation: Cultivated, oat hay.
Remarks: The soil is a medial developed Reddish Brunizem. Samples were obtained for the Bureau of Public Roads on the A_p, B_2, and C_1 horizons.
Classification: Ultic Haploxeralf, fine-loamy, mixed, thermic.

HORIZON		DESCRIPTION
A_p	0-8"	Brown (7.5YR 5/4) coarse sandy loam, dark reddish brown (5YR 3/4) when moist; massive; slightly hard when dry, friable when moist, nonsticky and nonplastic when wet; abundant very fine roots; many very fine and fine pores; medium acid, pH 5.7; clear, smooth lower boundary.
A_3	8-18"	Reddish brown (5YR 5/4) loam, yellowish red (5YR 3/6) when moist; massive; hard when dry, friable when moist, slightly sticky and slightly plastic when wet; abundant very fine roots; many very fine, common fine, few medium and coarse pores; few thin discontinuous clay films in pores; medium acid, pH 5.9; gradual, smooth lower boundary.
B_1	18-27"	Yellowish red (5YR 5/6) heavy loam, dark red (2.5YR 3.6) when moist; massive; very hard when dry, firm when moist, slightly sticky and slightly plastic when wet; plentiful very fine roots; many very fine, common fine, few medium and coarse pores; thin, continuous clay films in pores; slightly acid, pH 6.2; gradual, smooth lower boundary.
B_2	27-48"	Red (2.5YR 4/6) clay loam, dark red (2.5YR 3/6) when moist; massive; very hard when dry, firm when moist, sticky and plastic when wet; plentiful very fine roots; common very fine, few fine pores; moderately thick continuous clay films in most pores and in bridges between sand grains; slightly acid, pH 6.5; gradual, smooth lower boundary.
B_3	48-68"	Yellowish red (5YR 5/8) loam, dark red (2.5YR 3/8) when moist; massive; very hard when dry, firm when moist, sticky and plastic when wet; few very fine roots; common very fine and few fine pores; thin continuous clay films in some pores and in bridges between sand grains; slightly acid, pH 6.5; clear irregular lower boundary.
C	68-78"+	Yellowish red (5YR 5/8) fine sandy loam with [illegible]

SOIL SURVEY LABORATORY Lincoln, Nebraska

SOIL TYPE Aiken loam (taxadjunct) **LOCATION** Amador County, California

SOIL FAMILY: Ultic Haploxeralf, fine, kaolinitic, mesic.

SOIL NOS. S59Calif-3-11 **LAB. NOS.** 11348 - 11355

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	* 2A2 > 2	TEXTURAL CLASS
0-15	A1	4.6a	5.1a	4.7a	11.0a	13.7a	41.5	19.4	36.7	25.1	8	l
15-24	A3	2.1a	2.9a	3.0a	8.6a	13.3a	42.4	27.7	34.3	26.8	8	cl
24-37	B11t	1.7a	3.4a	2.7a	7.7a	11.7a	40.7	32.1	31.8	25.4	7	cl
37-49	B12t	1.1a	2.1a	2.0a	6.0a	9.4a	38.8	40.6	26.4	25.6	7	c
49-59	B21t	1.9a	2.1a	1.6a	5.1a	8.9a	37.4	43.0	23.3	26.2	8	c
59-75	B22t	0.8b	1.4b	1.7b	5.1b	8.2b	37.0	45.8	21.5	26.9	19	c
75-84	B31t	1.2b	3.1b	3.0b	7.4b	9.0b	36.8	39.5	22.8	27.4	17	cl-c
84-92	B32t	1.5b	3.0b	2.8b	7.2b	9.4b	37.3	38.8	23.3	27.8	37	cl

pH 8C1a 1:1	pH 8C1a 1:5	pH 8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	ORGANIC MATTER 6B1a NITROGEN %	ORGANIC MATTER C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC-10³ MILLIMHOS PER CM @25°C.	6C1a Free Iron as % Fe_2O_3	4A3a Bulk Density g/cc	MOISTURE TENSIONS 1/10 ATMOS. %	MOISTURE TENSIONS 1/3 ATMOS. %	4B2 15 ATMOS. %
6.0	6.2	6.4	3.21	0.145	22			5.3	1.04			18.6
5.8	6.0	6.2	0.92	0.044	21			6.1	0.95			19.0
5.8	6.0	6.2	0.66					6.3	1.05			19.6
6.0	6.1	6.3	0.34					6.6	1.20			21.0
6.0	6.3	6.5	0.33					5.8	1.39			22.5
6.0	6.2	6.4	0.09					5.0	1.39			23.8
6.0	6.2	6.4	0.09					5.0	1.39			22.4
6.0	6.2	6.2	0.08					5.2	1.31			22.8

1/ CATION EXCHANGE CAPACITY 5A3a (Sum)	EXTRACTABLE CATIONS 6N2b Ca	6O2b Mg	6H2a H	6P2a Na	6Q2a K	5B1a BASE SAT. % 5C3	Clay mineralogy (<0.002 mm.) Mont.	Verm. X-ray	Mixed layer Mica-Verm. 7A2	Mica	% Kaolin DTA 7A3	% Gibb. DTA 7A3
	milliequivalents per 100g. soil →						←			→		
30.5	6.4	1.3	21.8	<0.1	1.0	28		xx			20	19
19.6	5.1	0.7	13.1	<0.1	0.7	33						
18.6	4.8	0.9	12.2	<0.1	0.7	34		xxx			27	27
16.6	4.6	1.4	10.0	<0.1	0.6	40						
13.5	3.7	1.2	8.0	<0.1	0.6	41		x			45	10
11.9	2.4	1.4	7.6	<0.1	0.5	36						
9.8	2.0	1.0	6.2	<0.1	0.6	37						
10.4	2.0	1.3	6.6	<0.1	0.5	36					45	tr.

a. Many (>50 percent) Fe? nodules
b. Few (5 - 25 percent) Fe? nodules
2 mm. - 3/4" fraction

1/ In acid soils CEC (NH_4OAc) is almost always lower than either CEC (Na) or CEC (sum).

AIKEN LOAM (taxadjunct) 1/
S59Calif-3-11

Location: On the Allen Ranch, 100 feet north of road east of Fiddletown to Highway 88 in the SW1/4 of the NE1/4 of sec. 34, T. 8 N., R. 13 E., Amador Co., Calif.
Date of Sampling: June 19, 1959 Description by: Grant Kennedy.
Collectors: Grant Kennedy, Reuben Nelson, and Bob Sketchley
Vegetation: Ponderosa pine, incense cedar, black oak, sweet birch, bracken fern
Climate: Mean annual rainfall - 45 inches
Parent Material: Andesitic tuff-breccia
Relief: Broad, gently sloping ridgetop with a plane or slightly convex slope of 8 percent, facing west.

Horizons 1, 5, and 8 (A1, B21t, and B32t) were sampled for the Bureau of Public Roads. Soil was moist when described.
Classification: Ultic Haploxeralf, fine, kaolinitic, mesic.

HORIZON		DESCRIPTION
O1	2- 0"	Organic layer consisting of fresh pine needles, leaves, and twigs, becoming more decomposed with depth (not sampled).
A1 LSL No. 11348	0-15"	Reddish brown (5YR 4/4) loam, dark reddish brown (5YR 3/3) when moist; strong very fine and fine granular structure; soft, very friable; abundant very fine, fine, and medium roots; many very fine and fine interstitial pores; common very fine and fine shot*; slightly acid (pH 6.3); clear smooth boundary.
A3 LSL No. 11349	15-24"	Reddish brown (5YR 4/5) heavy loam, dark reddish brown (5YR 3/4) when moist; moderate very fine and fine granular structure; soft, very friable, slightly plastic; abundant very fine and fine, few medium and coarse roots; many very fine, fine, and medium, few coarse pores; occasional thin,

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Hoda stony loam **LOCATION** San Diego County, California

SOIL FAMILY: Ultic Haploxeralf, fine, kaolinitic, mesic.

SOIL NOS. S60Calif-37-8 **LAB. NOS.** 60437 - 60445

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 3	A11	*2.9	*5.6	*3.1	7.8	8.9	51.2	20.5	34.6	30.7	4	sil
3-11	A12	3.1	6.1	4.7	11.2	10.5	44.4	20.0	35.0	27.0	10	l
11-17	B1	1.3	4.7	3.4	9.4	8.5	43.1	29.6	29.0	28.5	2	cl
17-25	B21	0.4	3.9	3.4	8.3	7.3	33.8	42.9	21.7	24.6	0	c
25-39	B22	0.6	4.1	3.3	8.1	7.2	34.7	42.0	23.8	23.2	0	c
39-53	B23	0.2	3.3	3.0	8.3	8.1	37.0	40.1	26.3	24.1	0	c-cl
53-66	B31	0.5	5.0	3.9	9.8	8.9	38.0	33.9	29.0	24.2	0	cl
66-75	B32	0.2	4.6	4.4	11.0	9.7	39.4	30.7	30.8	25.2	0	cl
75-88+	B33	0.4	5.2	5.4	11.7	10.7	37.0	29.6	29.8	25.1	0	cl

pH 8C1a H_2O 1:1	pH 8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	Free Fe_2O_3 6C1b %	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.6	5.7		9.22	0.377	24.5				7.0			25.4
6.0	4.8		2.84	0.127	22.4				7.8			17.1
6.2	4.9		0.72	0.042	17.1				7.8			21.0
5.9	4.9		0.60	0.031					8.8			24.8
5.6	4.6		0.29	0.019					10.5			26.8
5.4	4.4		0.20	0.013					8.6			26.9
5.6	4.5		0.40	0.012					7.9			25.6
5.6	4.4		0.17						7.7			25.1
5.7	4.5		0.16						8.8			25.2

5A2a CATION EXCHANGE CAPACITY ← (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Na 6P2a	5B1a K 6Q2a	BASE SAT. % 5C3	Sum of Cations me./100 g. soil 5A3a	g. K				MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						milliequivalents per liter →					
41.5	25.0	8.1	16.8	0.3	1.8	68	52.0					
27.7	9.9	3.6	16.5	0.2	1.2	48	31.4					
18.7	6.9	4.3	9.8	0.3	1.6	57	22.9					
21.1	6.3	5.1	10.1	0.4	1.7	57	23.6					
22.8	7.1	5.1	11.4	0.6	1.1	55	25.3					
22.9	7.5	5.1	10.8	0.9	0.6	57	24.9					
21.8	7.6	5.5	11.0	1.2	0.1	57	25.4					
22.8	7.6	6.0	10.7	1.4	0.1	59	25.8					
25.7	7.9	6.8	11.6	1.5	0.1	58	27.9					

* Sand is largely organic matter

1/ Percent of total soil

HODA STONY LOAM
S60Calif-37-S

Population: Greater Depot

CONSECUTIVE PEDON NUMBER C 68113 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES HODA SOIL NO. S68 CALI. 29-33 SAMPLE NO. 8 844-8 851 NEVADA COUNTY AREA

SOIL FAMILY ULTIC HAPLOXERALF FINE,KAOLINITIC,MESIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SANDS VCS 2-1	CS 1-.5	MS .5-.25	FS .25-.10	VFS .1-.05	SILTS CSI .05-.02	FSI .02-.002	CLAYS CO. 2-.2U	FINE LT .2U	CARBONATE 3A1A	INT. II .2-.02	COARSE FRAGMENTS PCT. OF WHOLE SOIL I 75-20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														1A2A PCT. OF LT 75		
8 844	0- 10	A11	67.0	25.5	7.5	16.4	19.2	9.2	13.8	8.4	11.1	14.4				26.4	11A --	10	1
8 845	10- 30	A12	52.2	32.5	15.3	8.9	13.4	8.2	13.4	8.3	12.2	20.3				27.4	10A --	2	8
8 846	30- 46	A3	44.4	36.8	18.8	11.2	11.2	5.9	10.0	6.1	10.6	26.2				22.0	8A --	2	6
8 847	46- 91	B21T	30.0	25.2	44.8	6.3	7.7	3.9	6.6	5.5	7.3	17.9				16.5	5A --	TR	5
8 848	91-115	B22T	33.9	21.4	44.7	6.8	8.1	4.1	8.4	6.3	6.3	15.1				17.3	3A --	TR	3
8 849	115-135	B3	52.9	17.6	29.5	9.9	16.2	7.7	12.4	6.7	5.9	11.7				18.9	3A --	TR	3
8 850	135-150	C	53.6	22.1	24.3	8.5	14.9	7.5	14.1	8.6	8.6	13.5				24.9	7A --	1	6
8 851	0- 10	A11												2.7			6A --	1	5

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LL 4F2	LIMIT PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		PERCENT				PCT.NO.40		GRAMS PER CC				PERCENT			-PERCENT-	
8 844	0- 10	2.03	0.093			2.4	0.4			NP	NP		1.23	1.24	0.95		13.1	7.3	0.3	0.3
8 845	10- 30	1.35	0.056			3.1	0.6			30	7	1.25			0.95			9.6		
8 846	30- 46	0.55	0.036			3.6	0.7					1.32			0.96			11.4		
8 847	46- 91	0.35				5.2	0.8			53	23	1.30			0.97			19.3		
8 848	91-115	0.17				5.5	0.7			NP	NP		1.27	1.33	0.99		35.4	21.1	1.6	1.5
8 849	115-135					3.7	0.6					1.25			0.98			14.5		
8 850	135-150					3.8	0.5			NP	NP		1.24	1.25	0.97		27.8	13.8	0.3	0.3
8 851	0- 10	3.39	0.149									1.23			0.97			9.6		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A	MG 5B4A	NA 5B4A	K 5B4A	SUM	EXT. ACI-DITY	KCL EXT. Al	C E C SUM	NH4-SUM	BASE SATURATION SUM+NH4-	SUM+Al	NH4-DAC	PH NAF 7 MIN	H2O 1/1	CA-CL2
		6N2E	6Q2D	6P2A	6Q2A		6H2A	6G1E	5A3A	5A6A	DITY		5C1		8C1D	8C1A	8C1E
		MEQ/100G									PERCENT						
8 844	0- 10	3.4	0.8	TR	0.5	4.7	9.2		13.9	11.1	34		43			5.8	5.1
8 845	10- 30	1.6	0.6	0.1	0.5	2.8	4.5	1.0	7.3	10.7	38	74	26			5.3	4.5
8 846	30- 46	2.4	1.1	0.1	0.5	4.1	4.2	0.5	8.3	9.5	49	89	43			5.2	4.6
8 847	46- 91	3.5	1.9	TR	0.6	6.0	13.6	0.1	19.6	11.7	31	98	52			5.5	5.2
8 848	91-115	3.0	1.8	0.1	0.5	5.4	12.0	0.1	17.4	10.6	31	98	51			5.5	5.3
8 849	115-135	2.0	1.1	TR	0.5	3.6	5.1	0.1	8.7	9.8	42	97	37			5.4	5.0
8 850	135-150	1.7	1.0	0.1	0.5	3.3	4.7	0.4	8.0	10.5	41	89	31			5.4	4.6
8 851	0- 10															6.0	5.4

CLAY MINERALOGY

SAMPLE NO.	DEPTH CM.	LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 844	0- 10	KK	4	VR	1						KK KAOLINIT	1 TRACE
8 845	10- 30										VR VERMICLT	2 SMALL
8 846	30- 46											3 MODERATE
8 847	46- 91											4 ABUNDANT
8 848	91-115	KK	4	VR	1							5 DOMINANT
8 849	115-135											6 INDETER.
8 850	135-150	KK	3	VR	1							
8 851	0- 10											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.

HODA SANDY LOAM
S68Calif-29-33

Location: Nevada County, California; 800 feet west and 500 feet south of the E 1/4 corner of sec. 30, T. 17 N., R.9E., 150 feet west of Rock Creek Road, 3 miles NNE of Nevada City, California. Date of Sampling: September 6, 1968.

Description by: G. Staidl, C. Goudey. Collectors: G. M. Kennedy, L. A. Brittan, G. Staidl, C. B. Goudey, T. M. Ryan.

Classification: Ultic Haploxeralf, fine, kaolinitic, mesic.

Vegetation: Ponderosa pine, incense cedar, black oak, madrone, manzanita, ceanothus, bear

CONSECUTIVE PEDON NUMBER C 68114 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES HODA SOIL NO. S68 CALI. 29-34 SAMPLE NO. 8 852-8 859 NEVADA COUNTY AREA

SOIL FAMILY ULTIC HAPLOXERALF FINE,KAOLINITIC,MESIC RIVERSIDE SOIL SURVEY LABORATORY

SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 1A2A

TOTAL FRACTION ---------- SANDS ---------- SILTS ---------- CLAYS ---------- INT COARSE FRAGMENTS

NO.	CM.		2- .05	.05- .002	LT .002	2-1	1- .05	.5- .25	.25- .10	.1- .05	.05- .02	.02- .002	2- .2U	LT .2U	NATE 3A1A	.2- .02	OF WHOLE SOIL	75- 20 PCT.	20-5 OF	5-2 LT 75
			---PERCENT OF LESS THAN 2MM---																	
8 852	0- 10	A11	58.6	27.1	14.3	16.9	18.4	7.2	10.5	5.6	11.0	16.1				22.0	2A	--	TR	2
8 853	10- 25	A12	56.6	27.8	15.6	18.7	15.4	7.0	9.8	5.7	10.2	17.6				21.0	3A	--	1	2
8 854	25- 46	A3	51.3	28.0	20.7	12.9	16.0	7.4	9.4	5.6	10.4	17.6	15.7	5.0		20.9	3A	--	TR	3
8 855	46- 69	B1 T	35.7	29.9	34.4	4.6	9.7	6.1	9.0	6.3	10.6	19.3	24.7	9.7		21.8	3A	--	--	3
8 856	69-100	B2 T	32.9	24.9	42.2	10.6	8.4	4.2	5.9	3.8	9.1	15.8	26.8	15.4		16.1	1A	--	--	1
8 857	100-140	B3 T	33.0	27.2	39.8	9.0	10.2	4.1	5.9	3.8	7.3	19.9				14.3	2A	--	--	2
8 858	140-170	C	39.4	24.6	36.0	11.8	10.9	4.8	7.2	4.7	7.4	17.2				16.0	2A	--	--	2
8 859	0- 10	A11															6A	--	TR	6

SAMPLE NO.	DEPTH CM.	ORG. CAR- BON	MATTER NITRO GEN	CARBONATE AS CACO3 LT 2	2-20	----SESQUIOXIDES---- DI-CIT FE	EXT AL	PYROP FE	EXT AL	ATTERBERG L I M I T LL	PL	BULK DENSITY EST. 1/3 BAR	1/3 BAR	CM DRY	WATER CONTENT 1/10 BAR	1/3 BAR	15 BAR	EXTNSBLTY LEF	LE
		[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]

HODA LOAM
S68Calif-29-34

Location: Nevada County, California; 1,150 feet S. and 600 feet E. of the N 1/4 corner of sec. 36, T. 17 N., R. 8 E., 2½ miles north of Nevada City, 1/2 mile south of Lake Vera, 200 feet west of Lake Vera Road. Date of Sampling: September 6, 1968.

Description by: C. B. Goudey and G. Staidl. Collectors: L. A. Brittan, G. Staidl, C. B. Goudey, T. M. Ryan, G. M. Kennedy.

Classification: Ultic Haploxeralf, fine, kaolinitic, mesic.

Vegetation: Ponderosa pine, incense cedar, black oak, bear clover, ferns. Not cultivated.

Climate: Average annual precipitation 47 inches. Mean annual temperature 56°F.

Parent Material: Granodiorite, coarse grained. Topography: Mountainous upland, strongly sloping, concave, 13 percent slope, NNE aspect. Top of slope. Elevation: 2,590 feet.

Drainage: Well drained; runoff is medium; permeability is moderate to moderately slow in the subsoil. Soil Moisture: Profile very slightly moist.

Remarks: pH taken by Hellige method. The pH is lower than normal.

HORIZON	DESCRIPTION
O1	2 to 0 inches, pine needles, oak leaves, undecomposed.
	(0-10 cm.)
A11 RSL 68852	0 to 4 inches, brown (10YR 5/3) light loam, very dark grayish brown (10YR 3/2) moist; strong coarse subangular blocky parting to strong fine and coarse crumb structure; slightly hard, friable, slightly sticky, slightly plastic; many very fine and fine random, common medium and coarse oblique roots; many very fine interstitial pores; strongly acid (pH 5.5); clear wavy boundary.
	(10-25 cm.)
A12 RSL 68853	4 to 10 inches, brown (10YR 5/3) loam, dark brown (7.5YR 3/2) moist; moderate coarse subangular blocky parting to strong fine and coarse crumb structure; slightly hard, friable, slightly sticky, slightly plastic; many very fine and fine random, common medium and coarse oblique and horizontal roots; many very fine interstitial pores; medium acid (pH 6.0); abrupt wavy boundary.
	(25-46 cm.)
A3 RSL 68854	10 to 18 inches, brown (7.5YR 5/4) heavy loam, brown and dark brown (7.5YR 4/4) moist; weak coarse subangular blocky structure; slightly hard, friable, sticky, plastic; common very fine, and fine random, common medium and coarse horizontal and oblique roots; many very fine interstitial, few fine tubular pores; few thin clay films line pores and as bridges; medium acid (pH 6.0); clear wavy boundary.
	(46-69 cm.)
B1t RSL 68855	18 to 27 inches, light brown (7.5YR 6/4) clay loam, reddish brown (5YR 4/4) moist; weak medium and coarse angular blocky structure; hard, friable, sticky, plastic; few very fine and fine random, common medium oblique roots; common very fine and fine interstitial, few fine tubular pores; few thin clay films on faces of peds and common moderately thick clay films line pores; strongly acid (pH 5.5); clear wavy boundary.
	(69-100 cm.)
B2t RSL 68856	27 to 40 inches, yellowish red (5YR 5/6) clay loam, yellowish red (5YR 3/6) moist; moderate medium and coarse angular blocky structure; hard, friable, sticky, plastic; few very fine, fine and medium oblique roots; many very fine and fine interstitial and few fine tubular pores; many moderately thick yellowish red (5YR 4/8) clay films on faces of peds and lining pores; very strongly acid (pH 4.8); gradual wavy boundary.
	(100-140 cm.)
B3t RSL 68857	40 to 56 inches, yellowish red (5YR 5/6) light clay loam, yellowish red (5YR 4/6) moist; weak medium and coarse angular blocky structure; slightly hard, friable, sticky, plastic; few very fine random roots; common very fine interstitial pores; many moderately thick dark red(2.5YR 3/6 moist)clay films on faces of peds and lining pores; individual weathered mineral grains present; very strongly acid (pH 4.8); gradual wavy boundary.
	(140-170 cm.)
C RSL 68858	56 to 66 inches, reddish yellow (7.5YR 6/6) loam, strong brown (7.5YR 5/6) moist; massive; slightly hard, very friable, slightly sticky, plastic; few very fine random roots; common very fine interstitial pores; common moderately thick red (2.5YR 4/8 dry, 4/6 moist) clay films line pores and as bridges; very strongly acid (pH 4.8).

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Ultic Haploxeralf, fine, kaolinitic, thermic

Red Bluff clay loam (taxadjunct) S4557-10 Tehama County, California

SOIL SURVEY LABORATORY _____ LAB. Nos. _____

Depth (In.)	Horizon	1B1b Total Sand (2–0.05)	Total Silt (0.05–0.002)	Total Clay (< 0.002)	Sand Very coarse (2–1)	Sand Coarse (1–0.5)	Sand Medium (0.5–0.25)	Sand Fine (0.25–0.1)	Sand Very fine (0.1–0.05)	Silt 0.05–0.02	Silt Int. III (0.02–0.002)	Int. II (0.2–0.02)	3A1 (2–0.1)			Coarse fragments > 2 Pct.	Coarse fragments 2 - 19	Coarse fragments 19 - 76
		Pct. of < 2 mm															Pct. of < 76mm	
0-8	Ap	42.5	29.7	27.8	9.7	7.8	4.0	8.6	12.4	11.9	17.8		30.1					
8-15½	A1	41.3	28.9	29.8	7.6	7.3	4.2	8.7	13.5	11.9	17.0		27.8					
15½-23	B1t	41.8	28.1	30.1	9.0	8.1	4.2	8.1	12.4	11.0	17.1		29.4					
23-32	B21t	30.7	23.9	45.4	7.2	5.0	2.7	5.9	9.9	10.2	13.7		20.8					
32-40	B22t	33.4	25.0	41.6	6.2	5.6	3.4	7.4	10.8	7.3	17.7		22.6					
40-54	B23t	36.6	26.9	36.5	5.8	6.6	4.2	8.9	11.1	11.1	15.8		25.5					
54-60	B24t	44.5	26.2	29.3	9.4	9.1	5.1	9.5	11.4	10.8	15.4		33.1					

Depth (In.)	6A2a Organic carbon Pct	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	Bulk density g/cc	Bulk density g/cc		Water content Pct.	Water content Pct	Water content Pct			pH		8C1a (1:1)
0-8	0.66	0.070																5.1
8-15½	0.26	0.039																4.7
15½-23	0.17	0.034																4.9
23-32	0.09	0.028																5.0
32-40	0.06	0.020																5.0
40-54	0.08	0.014																4.6
54-60	0.07	0.014																4.6

Depth (In.)	Extractable bases 5B1a Ca 6N2b	Mg 6O2b	Na 2/	K 1/	Sum of bases	6H2a Ext. Acidity	5A3a CEC Sum									Base saturation 5C3 Sum Pct.	Pct.
	meq/100 g																
0-8	1.6	0.7	0.01	0.41	2.7	5.9	8.6									32	
8-15½	1.2	0.7	tr.	0.38	2.3	5.9	8.2									28	
15½-23	1.6	1.2	tr.	0.31	3.1	4.9	8.0									39	
23-32	2.4	2.0	0.04	0.31	4.7	4.7	9.4									51	
32-40	2.5	2.1	0.07	0.25	4.9	4.2	9.1									54	
40-54	2.4	2.0	0.06	0.30	4.8	3.6	8.4									57	
54-60	2.3	1.8	0.07	0.21	4.4	2.7	7.0									62	

Depth (In.)					

1/ Precipitated as potassium sodium cobaltinitrite and titrated as potassium permanganate

2/ Gravimetrically as sodium uranyl magnesium acetate

RED BLUFF CLAY LOAM (taxadjunct) 1/
S4557-10

Location: Tehama County, California, NE¼ SW¼ Sec. 34, T27N, R4W, MDB&M. Date of Sampling: April 4, 1945.

Description by: R. C. Roberts and R. A. Gardner. Collectors: R. C. Roberts, F. O. Youngs, R. A. Gardner, L. T. Alexander, H. Jenny and R. E. Storie.

Classification: Ultic Haploxeralf, fine, kaolinitic, thermic.

Vegetation: Pasture - formerly cultivated. Cover consists of sparse annual plants

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Argonaut gravelly silt loam (taxadjunct) **LOCATION** Amador County, California

SOIL FAMILY: Ultic Haploxeralf, fine, montmorillonitic, thermic.

SOIL NOS. 59Calif-3-5 **LAB. NOS.** 59294 - 59299

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-2	A_{11}	2.8	3.8	3.4	8.9	7.6	56.8	16.7	38.3	31.5	24	sil
2-10	A_{12}	2.7	4.4	3.0	8.1	8.1	55.0	18.7	35.8	32.4	13	sil
10-18	B_1	2.1	3.9	3.0	7.9	8.3	52.0	22.8	34.1	31.3	8	sil
18-22	B_{21}	2.4	3.7	2.3	5.0	4.7	30.1	51.8	17.3	20.6	35	c

ARGONAUT GRAVELLY SILT LOAM (taxadjunct) 1/
S59Calif-3-5

Collectors: Grant Kennedy, Reuben Nelson, Bob Sketchley.
Date of Sampling: June 17, 1959. Description by: Grant Kennedy.
Location: Amador County, California, on the Garbarini Ranch, 1-1/4 miles west of Jackson on the Pardee Dam Road in the SW1/4 of SW1/4 of Section 20, T6N, R11E, HDB&M.
Elevation: 1,450 feet.
Relief: Uplands in the upper portion of a swale with a very slightly concave slope of 7 percent, facing north.
Drainage: Moderately well drained; moderately rapid runoff when wet; moderately permeable over very slow.
Parent Material: Metamorphosed andesite.
Climate: Rainfall 27"
Vegetation: Oak-grass including white and blue oak, scattered Digger pine, soft chess, wild oats, filaree, and tar weed.
Remarks: The soil is a maximal Noncalcic Brown near the Brunizem soil zone. At 10 to 18 inches about 25 percent of the solids consist of cobbles of over 1 inch in diameter; not sampled.
Classification: Ultic Haploxeralf, fine, montmorillonitic, thermic.

HORIZON		DESCRIPTION
A_{11}	0-2"	Brown (7.5YR 5/4) gravelly silt loam with common, fine yellowish red (5YR 5/8) organic stains, dark brown (7.5YR 3/4) when moist; massive; slightly hard when dry, friable when moist; abundant very fine roots; common very fine and fine, few medium and coarse pores; slightly acid, pH 6.3; clear, smooth lower boundary.
[illegible]	2-10"	[illegible] when moist, slightly sticky and slightly plastic when wet; abundant very fine and fine roots; many very fine and fine, common medium tubular pores; slightly acid, pH 6.2; gradual, smooth lower boundary.
B_1	10-18"	Yellowish red (5YR 5/6) silt loam, dark reddish brown (5YR 3/4) when moist; massive; slightly hard when dry, friable when moist, sticky and plastic when wet; abundant very fine and fine, few medium roots; many very fine and fine, few medium and coarse pores; thin, nearly continuous clay films in some pores; slightly acid, pH. 6.1; abrupt, wavy lower

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Natrixeralf, fine-loamy, mixed, thermic

SOIL Placentia sandy loam (taxad.junct) SOIL Nos. S65Calif-27-15 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65507 - 65517

Depth (in.)	Horizon	1B1b Total: Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Sand: Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt: 0.05–0.02	Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	* Coarse fragments 1A2a > 2 Pct	2 – 19	19 – 76
		Pct. of < 2 mm														Pct of < 76 mm		
0-12	Ap1	59.1	29.9	11.0	13.8	14.7	8.4	13.7	8.5	15.8	14.1	31.4	50.6	0	11	13	13	0 **
12-16	Ap2	59.8	27.1	13.1	11.7	14.8	8.8	14.7	9.8	13.8	13.3	31.4	50.0	0	13	16	16	0 **
16-24	B21t	50.6	13.0	36.4	9.8	11.8	7.8	14.2	7.0	6.6	6.4	20.9	43.6	0	36	18	18	0 **
24-31	B22t	58.0	21.3	20.7	9.8	11.8	8.4	17.9	10.1	11.1	10.2	31.4	47.9	0	21	17	17	0 **
31-40	B23t	34.1	27.8	38.1	4.4	5.8	4.0	10.0	9.9	13.4	14.4	29.5	24.2	0	38	13	13	0 **
40-57	B31t	22.7	40.6	36.7	2.0	2.4	1.8	7.6	8.9	15.1	25.5	29.2	13.8	0	37	7	7	0 **
57-71	B32t	40.4	33.6	26.0	8.6	7.9	3.9	10.8	9.2	14.5	19.1	30.5	31.2	0	26	20	20	0 **
71-83	B33t	43.4	33.5	23.1	3.4	6.4	5.4	16.0	12.2	16.0	17.5	38.0	31.2	0	23	11	11	0 **
83-89+	C1	59.3	16.5	24.2	13.7	14.7	7.8	14.6	8.5	7.8	8.7	24.5	50.8	0	24	19	19	0 **
115-120	C***	50.1	23.8	26.1	4.2	11.8	9.3	16.4	8.4	6.8	17.0	24.2	41.7	0	26	0	0	0 **
24'	C***	92.4	4.0	3.6	34.6	37.5	11.3	7.3	1.7	1.7	2.3	6.2	90.7	0	4	43	42	1

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	Water content: Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-12	0.60	0.054	11		-						4.8				7.2	
12-16	0.35	0.035	10		-						5.4				6.8	
16-24	0.45				-						14.2				7.6	
24-31	0.09				tr.						9.0				8.6	
31-40	0.07				1						15.7				8.5	
40-57	0.11				-						17.6				8.4	
57-71	0.08				-						12.7				8.4	
71-83	0.07				tr.						11.9				8.2	
83-89+	0.05				-						9.3				8.6	
115-120					-						12.0				8.1	
24'					-						2.3				6.6	

Depth (in.)	Extractable bases: 6N2a Ca	6O2a Mg	6P2a Na	5B1a 6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste: 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		8A1 6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-12	4.1	1.8	0.8	0.2	6.9	2.0		7.9										
12-16	4.7	2.7	1.0	0.1	8.5	2.8		9.6										
16-24	11.9	8.5	2.2	0.2	22.8			24.2										
24-31	11.5	5.6	3.1	0.3	20.5		19.5	16.1	1.6	1.4	16.1	0.1		-	3.0	12.8		2.00
31-40	17.9	9.6	6.0	0.4	33.9		30.7	28.3	1.1	0.8	16.4	0.1		-	3.3	11.8		1.93
40-57	16.8	11.9	6.1	0.8	35.6		38.3	36.9	1.1	0.8	17.2	0.1		-	2.6	12.4		1.96
57-71	11.9	7.8	6.6	0.3	26.6		28.7	25.6	0.4	0.8	15.2	0.1		-	1.4	14.4		2.04
71-83	11.5	7.1	5.7	0.2	24.5		26.3	23.9	0.6	0.5	17.5	0.1		-	1.0	12.7		1.81
83-89+	12.8	5.0	2.8	0.3	20.9		17.8	14.4	0.6	0.3	11.4	0.1		-	4.0	6.0		1.25
115-120	10.9	7.0	5.1	0.3	23.3		26.8	22.7	0.5	0.4	13.8	0.1		-	1.5	11.2		1.55
24'	3.0	2.0	1.6	0.1	6.7		7.5	5.0	12.6	10.6	55.8	0.1		-	0.5	57.4	0.4	6.3

8A 5D2 5B1b 5C1 Clay mineralogy (<0.002 mm.)/A [redacted]

PLACENTIA SANDY LOAM (taxadjunct) 1/
865Calif-27-15

Location: Monterey County, California, 85 feet NW from drainageway 280 feet SW into field from Iverson Road, 0.53 mile NW on Iverson from Gloria Road. Date of Sampling: May 21, 1965. Description by: Lewis C. Leifer. Collectors: W.D. Nettleton, B.R.Brasher, C.J. Weisel, R. D. Waggoner, F. J. Pecarich; site selection by T. D. Cook.
Classification: Typic Natrixeralf, fine-loamy, mixed, thermic. Noncalcic Brown (maximal).
Vegetation: Formerly irrigated alfalfa field, temporarily abandoned, soon to be plowed; alfalfa, barley, wild oats. Climate: Precipitation 12 inches; mean annual temperature 58°F., mean January 50°F., mean July 65°F. Parent Material: Alluvium from phaneritic, intermediate igneous rocks. Topography: Very slightly convex, smooth, 3 percent slope towards drainageway, west exposure. Elevation: 300 feet. Drainage: Well drained, A horizons have moderately rapid permeability, B2 horizons have very slow permeability, runoff is medium. Soil Moisture: Nearly dry.
Remarks: The field pH determinations by Truog Kit. Matched pair 865Calif-27-14-1 through 8. This site has had some rinning and leveling operations with considerable disruption of A1 and A2 horizons.

HORIZON	DESCRIPTION
Ap1 RSL 65507	0 to 12 inches, brown (10YR 5/3) sandy loam, dark brown (10YR 3/3) moist; massive; some clods due to tillage; hard, friable, slightly sticky and slightly plastic; plentiful very fine and medium roots; many very fine interstitial, common very fine and few fine tubular pores; slightly acid (pH 6.5); clear wavy boundary.
Ap2 RSL 65508	12 to 16 inches, brown (10YR 5/3) sandy loam, dark brown (10YR 3/3) moist; 10 to 30 percent of soil consists of discrete fragments of A2, very pale brown (10YR 7/4) dry, dark yellowish brown (10YR 4/4) moist; massive; hard, friable, slightly sticky and slightly plastic; plentiful very fine and medium roots; many very fine interstitial and common very fine and few fine tubular pores; slightly acid (pH 6.5); abrupt smooth boundary.
B21t RSL 65509	16 to 24 inches, reddish brown (5YR 4/4 interior of peds) clay, reddish brown (5YR 4/4) moist, brown (10YR 4/3) rubbed, dry exterior upper portion very dark grayish brown (10YR 3/2 dry) lower part exterior dark reddish gray (5YR 4/2); moderate coarse prismatic structure; extremely hard, firm, very sticky and plastic; plentiful very fine roots on ped exterior; few very fine interstitial pores; continuous thin clay films in bridges and continuous moderately thick films on ped faces; mildly alkaline (pH 7.5); clear wavy boundary.
B22t RSL 65510	24 to 31 inches, dark brown (7.5YR 3/4) heavy sandy clay loam, slightly lighter color (7.5YR 4/4) moist; massive; very hard, firm, sticky and plastic; few very fine roots; few very fine interstitial and few very fine tubular pores; common thin clay films in bridges and common moderately thick films in pores; films are dark reddish brown (5YR 3/4 dry); very few very fine lime segregations throughout horizon, effervescent in seams; moderately alkaline (pH 8.0); gradual wavy boundary.
B23t RSL 65511	31 to 40 inches, dark reddish brown (5YR 3/4) light clay, reddish brown (5YR 4/4) moist undisturbed or rubbed; weak medium prismatic and strong medium angular blocky structure; hard, firm, very sticky and plastic; plentiful very fine roots on ped exterior; very few very fine interstitial and very few very fine tubular pores; continuous moderately thick films on ped faces; common fine lime segregations and common black Mn stains on ped faces, effervescent in segregations; moderately alkaline (pH 8.0); gradual smooth boundary.
B31t RSL 65512	40 to 57 inches, dark reddish brown (5YR 3/4), brown (7.5YR 5/4 crushed) silty clay, dark reddish brown (5YR 3/4), brown (7.5YR 4/4 crushed) moist; strong fine and medium angular blocky structure; hard, firm, very sticky and plastic; few very fine roots on ped exteriors; common very fine tubular pores; continuous moderately thick clay films on ped faces, few Mn stains on ped faces, violently effervescent in few soft lime seams, moderately alkaline (pH 8.0); gradual smooth boundary.
B32t RSL 65513	57 to 71 inches, dark reddish brown (5YR 3/4), brown (7.5YR 5/4 crushed) heavy clay loam, dark reddish brown (5YR 3/4), brown (7.5YR 4/4 rubbed) moist; moderate medium angular blocky structure; hard, firm, sticky and plastic; few very fine roots on ped exteriors; common very fine interstitial, many very fine tubular pores; continuous thin clay films on ped faces, few Mn stains on peds; moderately alkaline (pH 8.0), noncalcareous; clear smooth boundary.
B33t RSL 65514	71 to 83 inches, strong brown (7.5YR 5/6) silt loam, dark yellowish brown (10YR 4/4), brown (7.5YR 4/4 rubbed) moist; massive; hard, very friable, sticky and plastic; no roots; common very fine interstitial and few very fine tubular pores; common thin clay films on joints of upper portion of horizon, considerable stratification, suggestion of weak clay bands; moderately alkaline (pH 8.0), noncalcareous; clear smooth boundary.
C RSL 65515	83 to 89 inches, strong brown (7.5YR 5/6) fine sandy loam, brown (7.5YR 4/4) moist; massive; slightly hard, very friable, slightly sticky and slightly plastic; many very fine interstitial and common fine and medium tubular pores; few thin clay films in tubular pores; augered to 100 inches without change; moderately alkaline (pH 8.0),

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Natrixeralf, fine-loamy, mixed, thermic
SOIL Solano loam SOIL Nos. S64Calif-48-4 LOCATION Solano County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64511 - 64519

Depth (cm)	Horizon	1B1b Size class and particle diameter (mm) 3A1: Total: Sand (2-0.05)	Total: Silt (0.05-0.002)	Total: Clay (<0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: (0.05-0.02)	Silt: Int. III (0.02-0.002)	Silt: Int. II (0.2-0.02)	Silt: 2-0.1			Coarse fragments 1A2a: > 2 Pct. of whole soil	2-19	19-76
		Pct. of <2 mm														Pct. of <76mm		
0-15	A2	41.6	45.9	12.5	tr.	0.8	2.6	18.4	19.8	25.0	20.9	56.4	21.8			0		
15-36	B21t	36.9	41.8	21.3	tr.	0.6	2.6	17.3	16.4	21.2	20.6	48.9	20.5			0		
36-66	B22t	35.4	40.6	24.0	tr.	0.7	2.4	16.3	16.0	20.2	20.4	47.1	19.4			0		
66-100	B23t	23.8	41.5	34.7	0.2	0.7	1.5	9.8	11.6	14.6	26.9	32.7	12.2			0		
100-125	B24t	13.3	44.5	42.2	1.0	1.4	1.1	4.5	5.3	11.8	32.7	20.0	8.0			0		
125-165	B3ca	32.8	37.7	29.5	5.4	5.3	3.1	9.3	9.7	14.2	23.5	30.7	23.1			0		
165-195	C1	36.5	33.0	30.5	0.8	1.7	2.5	17.5	14.0	14.7	18.3	40.1	22.5			0		
195-230	C2	39.4	44.5	27.7	tr.	0.1	0.9	19.9	18.5	28.4	16.1	61.0	20.9			0		
230-270	C3	39.3	36.9	23.8	tr.	0.1	0.8	18.1	20.3	20.8	16.1	54.3	19.0			0		

Depth (cm)	6A1a Organic carbon a/ Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. iron as Fe Pct.	Carbonate as $CaCO_3$: 6E1b <2 mm Pct.	6E1c Clay size Pct.	Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content: 4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4C1 1/3 to 15 bar cm/cm	4D1 Extensibility COLEf cm/cm	4D1 Extensibility COLE cm/cm	pH: 8C1b Saturated paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-15	1.13	0.105	11	0.8	-		1.62	1.66		1.00	14.3	4.8		0.008	0.008		5.5	6.0
15-36	0.41	0.054	8	0.9	-		1.71	1.80		1.00	14.4	10.4		0.018	0.018		7.4	8.0
36-66	0.15	0.026		0.9	-		1.60	1.79		1.00	18.0	11.7		0.037	0.037		8.4	9.1
66-100	0.11			1.2	3		1.41	1.68		1.00	26.0	16.4		0.061	0.061		8.4	9.4
100-125	0.10			1.4	8	tr.	1.44	1.71		1.00	25.6	18.5		0.060	0.060		8.4	9.5
125-165	0.06			1.0	22	1	1.70	1.80		1.00	18.2	13.9		0.019	0.019		8.7	9.8
165-195	0.05			1.0	8	tr.	1.66	1.81		1.00	19.9	13.8		0.028	0.028		8.7	9.7
195-230	0.02			1.1	tr.		1.75	1.85		1.00	18.5	12.7		0.019	0.019		8.5	9.4
230-270	0.02			1.0	tr.							11.1					8.4	9.0

Depth (cm)	Extractable bases: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	5B1a	6H2a Ext. acidity	5A2a CEC NaOAc	5D2 Exchangeable Sodium Pct.	5E Sodium adsorption ratio	8A Water at saturation Pct.	Water extract from saturated paste 8A1: 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K	6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1a SO_4	8A1a Electrical conductivity mmho/cm
	meq/100g										meq/liter								
0-15	3.6	4.6	0.8	0.2			12.6	6	5	33.7	0.6	0.6	3.5	0.2	-	tr.	2.6	1.1	0.52
15-36	1.6	9.4	0.5	0.2			17.5	2	2	48.2	0.5	1.0	18.6	0.1	-	0.9	16.9	3.3	2.33
36-66	1.6	10.4	11.8	0.2			17.3	43	35	71.7	0.5	5.4	60.4	0.6	-	2.0	34.3	37.0	7.0
66-100	4.3	17.4	17.7	0.3			24.7	42	35	89.2	0.9	10.1	82.7	0.1	-	2.7	47.5	52.6	9.4
100-125	4.5	18.5	17.6	0.3			28.1	40	35	101.5	0.5	5.8	63.5	0.3	-	2.6	36.1	36.3	7.1
125-165	8.5	15.2	10.6	0.3			23.6	32	33	85.0	0.6	1.5	34.2	0.2	-	3.1	20.7	11.6	3.62
165-195	15.3	15.2	6.4	0.3			25.6	20	18	67.9	0.3	1.6	16.9	0.2	-	3.5	12.4	3.1	2.10
195-230	8.0	14.2	3.4	0.3			24.3	12	10	56.9	0.6	1.2	9.7	0.1	-	3.0	7.8	1.0	1.32
230-270	6.7	13.5	1.7	0.3			21.9	7	6	48.0	0.5	1.0	5.0	tr.	-	2.6	2.2	1.3	0.69

Depth (cm)	Ratios to clay 8D1: CEC NaOAc	Ext. iron	15 bar water	Salt Pct.	6F1a Gypsum Pct.	Clay fraction mineralogy 3A1: 7A2 X-ray b/	Atterberg limits: Plastic limit Pct. of <0.4 mm.	Liquid limit	Plastic index
0-15	1.01	0.06	0.38			MT3, KK3	N.P.	21	N.P.
15-36	0.82	0.04	0.49						
36-66	0.72	0.04	0.49		-	MT5, KK2			
66-100	0.71	0.03	0.47		-		17	36	19
100-125	0.67	0.03	0.44		-				
125-165	0.80	0.03	0.49			MT4, KK2			
165-195	0.84	0.03	0.45			MT5, KK2	19	40	21
195-230	0.88	0.04	0.46						
230-270	0.92	0.04	0.47						

a/ 5.5 kg of organic carbon per square meter to a depth of 1 meter

b/ Mineral code
MT = montmorillonite
KK = kaolinite

Approximate weight fraction (X-ray)
5 = more than half
4 = one-third to half
3 = one-fifth to one-third
2 = one-twentieth to one-fifth

SOLANO LOAM
S64Calif-48-4

Location: Solano County, California. Art Brown Ranch, a pit located in NW corner of SW1/4 of NW1/4, Sec. 36, T. 6 N., R. 1 E., eight miles south of Dixon. Date of Sampling: October 6, 1964.
Description by: Grant Kennedy and Sam Johnson. Collectors: Reuben E. Nelson, Ben R. Brasher, Leland Bates, Sam Johnson, Grant Kennedy, and Wells Andrews.
Classification: Typic Natrixeralf, fine-loamy, mixed, thermic.
Vegetation: Not cultivated. Vegetation consists of diminished annual weeds and grasses, dominantly salt grass (Distichlis spicata), soft chess (Bromus mollis), and tarweed. Climate: 18 inches rainfall, 60° F. mean annual temperature. Parent Material: Alluvium from sedimentary rocks. Topograhy: Basin rim or low terrace, nearly level with hummocks up to 15 inches high. Elevation: 24 feet. Drainage: Runoff is very slow, permeability is very slow, drainage is imperfect. No water table encountered within 107 inches of the surface. Soil Moisture: Dry 0-12 inches, slightly moist 12-24 inches, moist below 24 inches.
Remarks: 1) pH determined by Hellige-Truog and Thymol blue. 2) Sample number of paired profile S64Calif-48-7. 3) Diagnostic horizons are ochric and natric. 4) Thickness of A horizons is greater on hummocks than in between. The thickness of the A horizons are about the same as the height of the hummocks. Between the hummocks the A horizon is very thin. Hummocks vary in height from almost nil to about 2 feet. Most hummocks are less than 15 inches in height, vary in diameter but average about 15-20 feet. Hummocks are almost continuous in some areas and fairly well scattered in other areas.

HORIZON	DESCRIPTION
	Colors are for the dry soil unless otherwise indicated.
A2 RSL No. 64511	0-6 inches (0-15 cm.), pale brown (10YR 6/3) light loam with common fine faint mottles of yellowish brown (10YR 5/4), dark grayish brown (10YR 4/2) with common fine faint mottles of brown (10YR 4/3) moist; weak coarse prismatic with capping of weak coarse platy structure; hard (dry), friable (moist), slightly sticky and slightly plastic (wet); plentiful micro and very fine random roots; common tubular micro pores; abrupt smooth boundary.
B21t RSL No. 64512	6-14 inches (15-36 cm.), brown (10YR 5/3) heavy loam, brown (10YR 4/3) moist; strong coarse columnar structure; extremely hard (dry), firm (moist), sticky and very plastic (wet); plentiful micro horizontal roots; common micro pores; common thin clay skins on ped faces and in pores; clear smooth boundary.
B22t RSL No. 64513	14-26 inches (36-66 cm.), pale brown (10YR 6/3) heavy loam, brown (10YR 4/3) moist; moderate coarse prismatic structure; hard (dry), firm (moist), sticky and plastic (wet); common very fine roots; many micro tubular pores; common thin clay skins on ped faces and in pores; gradual smooth boundary.
B23t RSL No. 64514	26-40 inches (66-100 cm.), pale brown (10YR 6/3) clay loam, dark yellowish brown (10YR 4/4) moist; weak coarse prisms breaking to moderate fine angular blocks; hard (dry), firm (moist), sticky and plastic (wet); few fine random roots; many micro tubular pores; many moderately thick clay skins on ped faces and in pores; strongly effervescent with lime in soft masses; clear wavy boundary.
B24t RSL No. 64515	40-49 inches (100-125 cm.), pale brown (10YR 6/3) silty clay with dark reddish brown (5YR 3/4) stains, dark yellowish brown (10YR 4/4) moist; weak coarse prisms breaking to moderate fine angular blocks; hard (dry), firm (moist), sticky and plastic (wet); few fine random roots; many micro, few very fine pores; many moderately thick clay skins on ped faces and in pores; strongly effervescent with carbonate in soft masses; clear wavy boundary.
B3ca RSL No. 64516	49-64 inches (125-165 cm.), mottled white (10YR 8/2) and light yellowish brown (10YR 6/4) light clay loam, dark yellowish brown (10YR 4/4) with light brownish gray (10YR 6/2) mottles (lime seams) moist; weak fine angular blocky structure; extremely hard (dry), very firm (moist), sticky and plastic (wet); few fine random roots; common micro and very fine pores; common moderately thick clay skins on ped faces and in pores; strongly effervescent with carbonate in seams and concretions; iron and manganese concretions; clear wavy boundary.
C1 RSL No. 64517	64-76 inches (165-195 cm.), yellowish brown (10YR 5/4) light clay loam, dark yellowish brown (10YR 4/4) moist; massive; extremely hard (dry), very firm (moist), sticky and plastic (wet); common micro and very fine pores; strongly effervescent with carbonate in soft masses and seams; iron and manganese concretions; clear smooth boundary.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Natrixeralf, fine, montmorillonitic, thermic
SOIL Antioch loam SOIL Nos. S64Calif-48-8 LOCATION Solano County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64537 - 64547

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	3A1 Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	Coarse fragments 1A2a: > 2	2 - 19	19 - 76
		Pct. of < 2 mm														Pct.	Pct. of < 76 mm	
0-5	Ap	39.2	45.4	15.4	0.3	1.8	4.9	19.6	12.6	22.5	22.9	45.7	26.6			0		
5-14	A12	39.8	43.1	17.1	0.2	1.9	5.0	20.1	12.6	21.1	22.0	44.9	27.2			0		
14-19	A2	41.8	42.5	15.7	0.4	2.3	5.2	20.4	13.5	20.5	22.0	45.0	28.3			0		
19-34	B21t	20.2	39.5	40.3	tr.	0.3	1.4	6.8	11.7	19.0	20.5	35.1	8.5			0		
34-37	B22t	19.1	44.7	36.2	tr.	0.2	0.8	4.9	13.2	23.6	21.1	40.2	5.9			0		
37-46	B23t	19.6	47.5	32.9	tr.	0.2	0.5	3.6	15.3	24.9	22.6	42.9	4.3			0		
46-60	B24t	11.7	54.4	33.9	0.8	0.6	0.8	3.7	5.8	21.5	32.9	29.7	5.9			0		
60-76	B3t	43.9	34.5	21.6	0.4	1.7	4.9	19.5	17.4	18.2	16.3	47.8	26.5			0		
76-81	C1	58.2	27.5	14.3	0.2	2.4	7.6	28.6	19.4	15.8	11.7	52.4	38.8			0		
81-87	IIC2	45.1	22.8	32.1	0.1	2.1	6.6	22.1	14.2	12.2	10.6	39.4	30.9			0		
100-111		50.9	24.5	24.6	1.0	14.0	10.9	15.0	10.0	11.8	12.7	30.3	40.9					

Depth (in.)	6A1a Organic carbon	6B1a Nitrogen	C/N	6C2a Ext. Iron as Fe	6E1b Carbonate as $CaCO_3$		4A1f Bulk density 1/3 bar	4A1h Bulk density Oven dry	Bulk density	3B2 Cm		Water content 4B1c 1/3 bar	4B2 15 bar	4D1 Extensibility COLE	4D1 Extensibility COLE	pH 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
	Pct.	Pct.		Pct.	Pct.		g/cc	g/cc	g/cc		Pct.	Pct.	Pct.	in./in.	in./in.			
0-5	1.35	0.127	11	1.4	-		1.59	1.63		1.00		17.8	6.6	0.010	0.010		5.8	6.2
5-14	0.54	0.064	8	1.4	-		1.60	1.64		1.00		16.3	6.5	0.008	0.008		6.2	6.4
14-19	0.28	0.042	7	1.5	-		1.68	1.72		1.00		14.3	6.5	0.006	0.006		6.5	6.8
19-34	0.28	0.047	6	2.0	-		1.56	1.87		1.00		24.1	19.5	0.062	0.062		6.2	7.1
34-37	0.10			2.0	-		1.52	1.84		1.00		28.5	20.3	0.066	0.066		7.5	8.4
37-46	0.09			1.9	-		1.61	1.78		1.00		21.9	18.3	0.035	0.035		8.1	8.9
46-60	0.07			1.9	2		1.62	1.78		1.00		22.0	19.7	0.033	0.033		8.4	9.2
60-76	0.04			1.1	1		1.76	1.83		1.00		14.8	9.7	0.013	0.013		8.6	9.3
76-81	0.03			1.0	tr.		1.72	1.78		1.00		14.5	6.6	0.013	0.013		8.5	8.8
81-87	0.01			1.5	tr.		1.70	1.86		1.00		17.5	13.9	0.031	0.031		8.1	8.8
100-111	0.02			2.1	tr.								11.8				8.4	9.2

Depth (in.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	5B1a 6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	8A1 6K1a Cl	6L1a SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-5	7.7	2.0	0.3	0.4	10.4	4.9	14.1		2.0	0.8	2.8	0.4		-	0.2	2.9		0.67
5-14	6.8	2.3	0.4	0.2	9.7	3.6	13.7		0.8	0.6	2.4	0.2		-	0.2	1.8		0.42
14-19	6.1	2.7	0.8	0.1	9.7	2.9	12.4		0.7	0.6	4.4	0.1		-	0.2	3.5		0.64
19-34	13.8	7.3	7.0	0.3	28.4	3.9	34.3		0.4	0.7	10.8	0.2		-	0.4	7.9		1.31
34-37	15.6	8.4	8.8	0.3	33.1		34.7		0.6	0.3	14.4	0.3		-	0.6	11.1		1.61
37-46	15.5	8.1	8.4	0.2	32.2		33.4		0.7	0.3	12.4	0.2		-	1.9	8.9		1.38
46-60	17.7	8.7	8.8	0.2	35.4		35.3		0.6	0.2	11.7	tr.		-	2.6	7.6		1.31
60-76	10.8	4.5	4.1	0.1	19.5		18.0		0.4	0.4	8.5	0.3		-	4.4	2.8		0.80
76-81	7.9	3.0	2.2	0.1	13.2		12.8		0.4	0.4	6.7	0.3		-	3.8	1.2		0.63
81-87	15.6	6.2	3.6	0.2	25.6		25.3		0.3	0.4	4.5	tr.		-	2.6	1.0		0.46
100-111	17.0	5.6	2.7	0.3	25.6		25.0		0.4	0.2	4.8	tr.		-	3.4	1.1		0.48

Depth (in.)	8A Water at Saturation	Atterberg limits: Plastic limit	Liquid limit	Plastic index	5D2 Exchangeable Na	5E SAR	5B1b Exchangeable Na	Clay mineralogy (<0.002 mm.) 8A2: Mont.	Verm.	Mica	Kaolin.	
	Pct.	Pct. of <0.4 mm.			Pct.		meq/100 g	X-ray 7A2				
0-5	40.1	20	23	3	1	2	0.2					
5-14	33.1				2	3	0.3	-	xx	x	x	
14-19	31.2				5	5	0.6					
19-34	63.8	18	43	25	18	15	6.3					
34-37	66.0				23	21	7.9					
37-46	59.0				23	18	7.7					
46-60	58.7				23	18	8.1	xxxx	-	-	x	
60-76	38.4	17	26	9	21	13	3.7					
76-81	29.7				16	11	2.0	xxx	-	-	xx	
81-87	46.8				13	8	3.4	xxx	-	tr	x	
100-111	44.0				10	9	2.5					

- = look for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

ANTIOCH LOAM
S64Calif-48-8

Location: Solano County, California. A pit located SW¼, SE¼, SW¼ of Sec. 29, T5N, R1W, MDB&M, 1½ miles east of Suisun, California. Date of Sampling: October 8, 1964.
Description by: Leland Bates, Sam Johnson. Collectors: Reuben Nelson, Ben Brasher, Leland Bates, Sam Johnson, Wells Andrews.
Classification: Typic Natrixeralf, fine, montmorillonitic, thermic.
Vegetation: Fallow. Climate: 20 inches rainfall, 60°F. mean annual temperature. Parent Material: Alluvium from sedimentary rocks. Topography: Old terrace. Elevation: 12 feet. Drainage: Runoff is slow, permeability is very slow, drainage is moderately well. No water table encountered within 111 inches of the surface. Soil Moisture: Dry 0-19", moist below 19".
Remarks: (1) pH determined by Hellige-Truog. (2) Sample number of paired profile 64 CAL 48-1-(1-8). (3) Diagnostic horizons are ochric, argillic and albic. (4) Thickness of the A2 varies from thin capping on B2 horizon to 5 inches in the general area of this soil.

HORIZON	DESCRIPTION
Ap RSL 64537	0 to 5 inches, light brownish gray (10YR 6/2) loam with common fine distinct yellowish brown (10YR 5/6) mottles, dark grayish brown (10YR 4/2) with common fine distinct strong brown (7.5YR 5/6) mottles moist; massive; hard, friable, slightly sticky, slightly plastic; plentiful micro roots; many very fine and medium tubular pores; medium acid (pH 5.6); clear smooth boundary.
A12 RSL 64538	5 to 14 inches, brown (10YR 5/3) loam with few fine distinct yellowish brown (10YR 5/6) mottles, dark brown (10YR 3/3) with few fine distinct strong brown (7.5YR 5/6) mottles moist; massive; hard, friable, slightly sticky, slightly plastic; few micro roots; many fine and medium tubular pores; medium acid (pH 6.0); clear wavy boundary.
A2 RSL 64539	14 to 19 inches, light gray (10YR 7/2) loam with common fine distinct yellowish brown (10YR 5/6) mottles, dark grayish brown (10YR 4/2) with common fine distinct strong brown (7.5YR 5/6) mottles moist; massive;hard, friable, slightly sticky, slightly plastic; very few micro roots; many fine and very fine pores; slightly acid (pH 6.5); manganese stains; abrupt smooth boundary.
B21t RSL 64540	19 to 34 inches, light yellowish brown (10YR 6/4) clay with common medium faint pale brown (10YR 6/3) mottles, dark yellowish brown (10YR 4/4) with common medium dark brown (10YR 3/3) mottles moist; moderate very coarse prismatic structure; extremely hard, very firm, sticky, very plastic; very few micro roots; common micro tubular pores; many moderately thick clay films on ped faces and in pores; medium acid (pH 6.0); few iron and manganese concretions; clear wavy boundary.
B22t RSL 64541	34 to 37 inches, yellowish brown (10YR 5/4) clay, dark brown (10YR 4/3) moist; weak medium angular blocky structure; extremely hard, very firm, sticky, plastic; common micro tubular pores; many moderately thick clay films on ped faces and in pores; moderately alkaline (pH 8.0); many iron and manganese concretions; clear wavy boundary.
B23t RSL 64542	37 to 46 inches, pale brown (10YR 6/3) clay, dark yellowish brown (10YR 4/4) with dark brown (7.5YR 3/2) and dark grayish brown (2.5Y 4/2) ped faces moist; weak medium angular blocky structure; hard, firm, sticky, plastic; common micro tubular pores; continuous moderately thick clay films on ped faces and in pores; moderately alkaline (pH 8.0); common iron and manganese concretions; diffuse boundary.
B24t RSL 64543	46 to 60 inches, pale brown (10YR 6/3) light clay, olive brown (2.5Y 4/4) with dark grayish brown (2.5Y 4/2) and dusky red (2.5YR 3/2) ped faces moist; weak medium angular blocky structure; hard, firm, sticky, plastic; common micro tubular pores; continuous moderately thick clay films on ped faces and in pores; moderately alkaline (pH 8.0); common iron and manganese concretions; clear wavy boundary.
B3t RSL 64544	60 to 76 inches, pale brown (10YR 6/3) loam, dark yellowish brown (10YR 4/4) moist; weak medium angular blocky structure; slightly hard, friable, slightly sticky, plastic; many micro tubular pores; common thin clay films on ped faces and in pores; moderately alkaline (pH 8.0); common iron and manganese stains; clear wavy boundary.
C1 RSL 64545	76 to 81 inches, dark yellowish brown (10YR 4/4) (moist) fine sandy loam; weak medium angular blocky structure; slightly hard, friable, nonsticky, nonplastic; many micro tubular pores; few thin clay films on ped faces and in pores; moderately alkaline (pH 8.0); common iron and manganese stains.
II C2 RSL 64546	81 to 87 inches. Composite sample taken for laboratory - no description.
	87 to 100 inches. No sample taken and no description made.
RSL 64547	100 to 111 inches. Composite sample taken for laboratory - no description.

Samples taken for Bureau of Public Roads: Ap, B21t, and B3t.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Natrixeralf, fine, montmorillonitic, thermic

SOIL Placentia sandy loam SOIL Nos. 865Calif-27-14 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65498 - 65506

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	3A1 Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	* Coarse fragments 1A2a ≥ 2 Pct.	2 - 19 (Pct. of < 76 mm)	19 - 76 (Pct. of < 76 mm)	
		Pct. of < 2 mm																	
0-13	Ap1	52.5	33.8	13.7	11.8	12.7	7.6	12.3	8.1	16.4	17.4	31.2	44.4	0	14	8	8	0	***
13-32	Ap2***	52.9	33.8	13.3	11.2	13.9	7.5	11.8	8.5	16.0	17.8	31.0	44.4	0	13	8	8	0	***
32-51	B21t	35.1	17.3	47.6	7.4	8.1	5.2	8.7	5.7	7.7	9.6	18.2	29.4	0	48	8	8	0	***
51-74	B22t	49.8	16.8	33.4	9.5	11.0	7.3	14.3	7.7	7.2	9.6	22.9	42.1	0	33	18	18	0	***
74-91	B23tca	54.6	17.1	28.3	10.3	14.3	8.7	14.2	7.1	7.4	9.7	21.8	47.5	0	28	20	20	0	***
91-110	B24t	54.3	20.3	25.4	12.1	14.9	7.7	12.0	7.6	9.9	10.4	23.4	46.7	0	25	11	11	0	***
110-150	B3t	61.1	20.1	18.8	12.3	17.1	9.7	14.4	7.6	10.4	9.7	25.3	53.5	0	19	15	15	0	***
150-175+	C *****	65.3	18.2	16.5	13.9	18.2	10.5	15.6	7.1	9.4	8.8	24.3	58.2	0	17	36	24	12	
240-255	Auger	70.5	11.5	18.0	10.7	16.8	12.1	22.6	8.3	6.1	5.4	25.6	62.2	0	18	15	15	0	***

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	Atterberg limits: Plastic limit (Pct. of <0.4 mm)	Liquid limit	Plastic index	Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	3B2 Cm	Water content 4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	8C1b Saturated Paste	pH 8C1a H_2O 1:1	8C1a H_2O 1:10
0-13	0.97	0.087	11	0.9	-	16	19	3	1.57	1.60	0.95	13.3	5.4	0.007	0.006		6.2	
13-32	0.86	0.072	12	0.8	-				1.58	1.63	0.95	11.4	5.2	0.011	0.010		5.9	
32-51	0.68			1.4	-				1.46	1.81	0.95	28.1	19.8	0.073	0.069		7.6	
51-74	0.33			1.2	1	25	58	33	1.56	1.89	0.89	24.1	15.0	0.066	0.058		8.5	
74-91	0.18			1.1	1				1.62	1.85	0.87	19.9	12.4	0.043	0.039		8.4	
91-110	0.11			0.8	-				1.73	1.87	0.93	15.0	10.4	0.026	0.024		8.4	
110-150	0.13			0.6	-				1.73	1.79	0.89	12.0	7.3	0.011	0.010		8.1	
150-175+	0.06			0.5	-				1.76	1.82	0.83	10.9	6.9	0.011	0.009		7.7	
240-255	0.16			0.7	-								8.2				7.2	

Depth (in.)	Extractable bases 6H2a Ca	6G2a Mg	6F2a Na	5B1a 6Q2a K	Sum of bases	6H2a Exr. Acidity	Cation Exch. Capacity 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	8A1 SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-13	4.4	2.5	0.4	0.4	7.7	4.4		10.0										
13-32	4.8	2.4	0.4	0.2	7.8	4.1		10.8										
32-51	12.8	13.7	5.5	0.3	22.3	3.1	36.3	32.5										
51-74	18.8	12.4	5.7	0.2	37.1		30.4		1.6	2.1	24.6	0.1		-	2.3	22.7	1.3	3.04
74-91	13.9	11.1	7.2	0.2	32.4		27.5	25.8	1.8	3.4	26.5	0.1		-	2.4	31.0	1.3	3.92
91-110	7.0	8.7	6.0	0.2	21.9		22.6	19.5	2.5	2.7	41.1	0.1		0.5	1.7	40.8	0.9	4.69
110-150	4.1	4.9	4.2	0.2	13.4		14.3	11.4	3.7	5.9	51.5	0.1		-	1.3	55.1	0.7	6.17
150-175+	3.3	3.9	3.5	0.1	10.8		12.0	9.1	4.4	6.6	51.2	0.2		-	0.9	57.9	0.4	6.85
240-255	5.4	5.8	4.7	0.2	16.1			14.3	8.2	11.6	61.2	0.2		-	0.7	78.0	0.1	8.33

Depth (in.)	8A Water at Saturation Pct.	5D2 Exchangeable Na Pct.	5B1b Exch. Na meq./100g.	Gypsum Pct.	5C1 Base Sat. NH_4OAc Pct.	Clay mineralogy (<0.002 mm.) 7A: Mont.	Verm.	Mica	Kaolin	
						X-ray				
0-13					77			xxxx	xx	
13-32					72					
32-51		15			69					
51-74	55.9	14	4.4		100+	xxxx		xx	x	
74-91	48.0	21	5.9		100+					
91-110	41.0	19	4.4		100+					
110-150	28.7	19	2.7		100+					
150-175+	26.8	18	2.1		100+					
240-255	31.6	18	2.6		100+	xxx		xxx	x	

* From characterization sample (5Kg) (except as noted).
** 2-19 mm. from characterization sample, 19-76 mm. from field volume estimate.
*** The A2 horizon (12.5-13") was not sampled.
***** Auger sample from below base of pit.

\- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

PLACENTIA SANDY LOAM
S65Calif-27-14

Location: Monterey County, California. 40 feet NW of fence, 300 feet SW from base of Reservoir 0.47 mile NE, .72 mile NW from intersection Iverson and Johnson Canyon Roads. Date of Sampling: May 21, 1965.

Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C.J. Weisel, R. D. Waggoner, F. J. Pecarich; site selection by T. D. Cook.

Classification: Typic Natrixeralf, fine, montmorillonitic, thermic. Noncalcic Brown (maximal).

Vegetation: Pasture - grass, ripgut brome, some soft chess, mustard, filaree. Climate: Precipitation is 12 inches; mean annual temperature 58°F., mean January 50°F., mean July is 65°F. Parent Material: Alluvium from phaneritic, intermediate igneous rocks. Topography: Gently sloping, slightly convex west-facing slope of 4 percent. Elevation: 300 feet. Drainage: Well drained, B2 horizons have very slow permeability, runoff is medium. Soil Moisture: Dry, wilting point or below throughout.

Remarks: Field pH determinations by Truog Kit. Paired samples are: S65Calif-27-15-1 through 9. Asterisks indicate horizons sampled for California Division of Highways.

HORIZON	DESCRIPTION
*Ap1 RSL 65498	0 to 5 inches, brown (7.5YR 5/4) sandy loam, dark brown (7.5YR 3/2) moist; massive; compacted by tillage; hard, very friable, slightly sticky and slightly plastic; plentiful very fine and few fine roots; many very fine interstitial and few very fine and medium tubular pores; medium acid (pH 6.0); clear smooth boundary.
Ap2 RSL 65499	5 to 12½ inches, brown (7.5YR 5/4) sandy loam, dark brown (7.5YR 3/2) moist; weak medium subangular blocky structure; hard, very friable, slightly sticky and slightly plastic; abundant very fine roots; many very fine interstitial and common very fine and fine tubular pores; medium acid (pH 6.0); horizon somewhat disturbed by many gopher holes filled with mixed granular material, roots much more numerous in the granular material; abrupt smooth boundary.
A2 (Not sampled)	12½ to 13 inches, brown (10YR 5/3) sandy loam, darker brown (7.5YR 4/4) moist; weak medium platy structure; hard, very friable, slightly sticky and slightly plastic; many very fine interstitial and few very fine tubular pores; medium acid (pH 6.0); very abrupt smooth boundary.
B21t RSL 65500	13 to 20 inches, dark reddish brown (5YR 3/4) moist or dry, clay; moderate coarse prismatic structure, prisms slightly rounded at top; extremely hard, very firm, very sticky and plastic; plentiful very fine roots on ped exterior; few very fine interstitial and few very fine tubular pores; continuous thin clay films as bridges and continuous moderately thick clay films on ped faces; moderately alkaline (pH 8.0), noncalcareous; gradual smooth boundary.
*B22t RSL 65501	20 to 29 inches, dark reddish brown (5YR 3/3) clay, dark brown (7.5YR 3/4) moist; strong coarse angular blocky structure; extremely hard, very firm, very sticky and plastic; plentiful very fine roots on ped exteriors; few very fine interstitial and few very fine tubular pores; continuous thin clay films in bridges and continuous moderately thick films on ped faces; moderately alkaline (pH 8.0), noncalcareous; few black Mn (?) stains on ped faces; clear wavy boundary.
B23tca RSL 65502	29 to 36 inches, reddish brown (5YR 4/4) moist or dry, heavy clay loam, same color rubbed; moderate medium angular blocky structure; very hard, firm, sticky and plastic; few very fine roots on ped exteriors; common very fine interstitial and common very fine tubular pores; many thin clay films as bridges and continuous thin films on ped faces; moderately alkaline (pH 8.0), strongly effervescent in medium lime seams; gradual smooth boundary.
B24t RSL 65503	36 to 42 inches, strong brown (7.5YR 5/6) heavy sandy clay loam, dark reddish brown (5YR 3/4) moist; moderate medium angular blocky structure; very hard, firm, sticky and plastic; very few very fine roots; many very fine interstitial pores; many thin clay films as bridges and continuous thin clay films on ped faces; moderately alkaline (pH 8.0); few lime seams trailing down from B23tca; clear smooth boundary.
B3t RSL 65504	43 to 58 inches, strong brown (7.5YR 5/6) light sandy clay loam, brown (7.5YR 4/4) moist; same moist color rubbed; massive; very hard, friable, sticky and plastic; very few very fine roots; many very fine interstitial and few very fine and few medium tubular pores; common thin clay films in bridges; moderately alkaline (pH 8.0), noncalcareous; this horizon somewhat brittle with weak duripan features; gradual smooth boundary.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Natrixeralf, fine, montmorillonitic, thermic

SOIL San Miguel stony silt loam SOIL Nos. S64Calif-37-18 LOCATION San Diego County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65602 - 65606

		1R1b	Size class and particle diameter (mm) 1A1

SAN MIGUEL STONY SILT LOAM
S64Calif-37-18

Location: The site is on the west side of Otay Mountain about 9.5 miles east and 1 mile north of San Ysidro, California. It is in the southwest 1/4 of the southwest 1/4 of Section 27. T. 19 S., R. 1 E., San Diego County, California. Date of Sampling: October 22, 1964.
Description by: Gerald Kester and George Borst. Collectors: K. W. Flach and W. D. Nettleton.
Classification: Typic Natrixeralf, fine, montmorillonitic, thermic.
Vegetation: Chamise, Ceonothis, sumac, Yerba Santa. Climate: Mean annual precipitation is about 15 inches and comes mostly as rain in the winter and early spring; mean annual air temperature is about 57° F., January about 40° F., and July about 74° F. Parent Material: Fine grained metabasic or metavolcanic rock. Topography: Mountainous, 38 percent southwest facing slope. Elevation: About 2,010 feet. Drainage: Well drained, slow permeability, high runoff, moderate erosion. Soil Moisture: Dry, ground water very deep.
Remarks: The A1 horizon is 12 inches thick on 35 to 45 percent of the area. Gravel, cobbles, and stones occur throughout the profile but are less than 35 percent of the soil volume. Rock outcrops are common. Color of the B21t horizon is mostly in hues of 5 and 7.5YR but ranges to 10YR.

HORIZON	DESCRIPTION
A1 RSL No. 65602	0 to 4 inches (0 to 10 cm.), light yellowish brown (10YR 6/4)* silt loam, brown to dark brown (7.5YR 4/4) moist; moderate fine to coarse granular structure; slightly hard, friable, nonsticky, nonplastic; plentiful very fine roots; common fine to medium tubular and common fine vesicular pores; medium acid (pH 6.0); abrupt wavy boundary.
A2 RSL No. 65603	4 to 8 inches (10-20 cm.), very pale brown (10YR 7/3) silt loam, yellowish brown (10YR 5/6) moist; massive; hard, friable, nonsticky, slightly plastic; plentiful fine to medium roots; many fine vesicular pores; strongly acid (pH 5.2); abrupt wavy boundary.
B21t RSL No. 65604	8 to 18 inches (20 to 45 cm.), strong brown (7.5YR 5/6) clay, yellowish red (5YR 4/6) moist, reddish yellow (7.5YR 6/6) crushed, yellowish red (5YR 4/6) crushed moist; strong coarse prismatic structure; extremely hard, firm, very sticky, very plastic; few medium and plentiful fine roots; few fine to medium pores; many pressure faces on peds and a few slickensides; strongly acid (pH 5.1); clear wavy boundary.
B22t RSL No. 65605	18 to 23 inches (45 to 58 cm.), yellowish brown (10YR 5/4) gravelly clay, brown to dark brown (7.5YR 4/4) moist; moderate medium subangular blocky structure; very hard, friable, very sticky, plastic; plentiful very fine and fine roots; few fine pores; common pressure faces on peds; strongly acid (pH 5.1); abrupt irregular boundary.
R RSL No. 65606	23 to 31 inches (58 to 78 cm.), light yellowish brown (2.5Y 6/4) metabasic rock, light olive brown (2.5Y 5/4) moist; many fine and medium roots in fractures in the rock.

* Colors are for dry condition unless otherwise noted.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aquic Natrixeralf, fine-loamy, mixed, thermic
SOIL Solano loam (taxadjunct) SOIL Nos. S64Calif-48-7 LOCATION Solano County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64520 - 64527

Depth (in)	Horizon	1B1b Total: Sand (2-0.05)	Total: Silt (0.05-0.002)	Total: Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	3A1 Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	Coarse fragments 1A2a > 2 Pct.	2 - 19	19 - 76
		Size class and particle diameter (mm)															Pct. of < 76 mm	
		Pct. of < 2 mm																
0-4	A21	31.8	53.3	14.9	0.2	1.1	3.0	12.8	14.7	24.0	29.3	46.9	17.1			0		
4-9	A22	31.4	52.7	15.9	0.2	1.0	2.5	12.2	15.5	22.7	30.0	46.1	15.9			0		
9-21	B21t	26.0	43.3	30.7	0.1	0.8	2.4	10.7	12.0	19.6	23.7	38.5	14.0			0		
21-32	B22t	36.4	32.6	31.0	tr.	1.0	4.0	15.8	15.6	15.3	17.3	40.3	20.8			0		
32-48	B23t	23.7	44.1	32.2	2.8	2.3	1.5	7.0	10.1	19.3	24.8	34.4	13.6			0		
48-62	B24t	17.4	50.4	32.2	2.5	2.9	1.3	3.7	7.0	18.0	32.4	27.5	10.4			0		
62-80	B25t	15.4	49.2	35.4	3.4	2.8	1.6	4.1	3.5	11.8	37.4	17.6	11.9			0		
80-92		42.3	34.5	23.2	0.1	0.4	2.9	20.6	18.3	19.3	15.2	50.6	24.0			0		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	6E1c Clay-size carbonate Pct.	Atterberg limits: Plastic limit	Liquid limit	Plastic index	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEf in./in.	4D1 Extensibility COLE in./in.	8C1b Saturated Paste	pH 8C1a H_2O 1:1	8C1a H_2O 1:10
							Pct. of <0.4 mm										
0-4	2.25	0.177	13		-		N.P.	30	N.P.			6.7				5.6	6.0
4-9	0.90	0.080	11		-							6.2				5.7	6.4
9-21	0.39	0.050	8		-		17	33	16			17.5				6.8	7.6
21-32	0.14				-							16.6				8.3	8.9
32-48	0.07				9	tr.						17.8				8.7	9.8
48-62	0.05				11	tr.	20	56	36			17.6				8.8	9.9
62-80	0.05				11	tr.						18.6				8.7	9.8
80-92	0.02				1							11.4				8.6	9.4

Depth (in.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	5B1a Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 8A1: 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K	6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1a SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter								mmho/cm
0-4	5.2	2.1	0.7	0.2	8.2	6.8	15.7		0.8	0.5	3.7	0.1	-	0.2	2.5	1.2	0.68
4-9	4.5	2.8	1.1	0.1	8.5	6.3	14.9		0.5	0.1	4.0	0.1	-	0.2	3.1	0.8	0.57
9-21	6.0	7.6	7.5	0.2	21.3	2.9	22.8		0.5	0.9	17.5	0.1	-	0.5	15.1	3.0	2.13
21-32	5.2	9.6	14.1	0.3	29.2		28.0		0.7	1.3	36.4	0.5	-	1.4	26.1	5.8	3.57
32-48	13.6	11.6	15.6	0.3	41.1		33.3		0.5	0.6	25.6	0.5	-	3.3	19.7	2.8*	2.86
48-62	16.9	12.3	11.9	0.3	41.4		34.1		0.5	0.3	12.6	tr.	-	4.3	8.3	0.9	1.49
62-80	21.0	12.7	9.3	0.3	43.3		37.8		0.3	0.1	9.2	tr.	-	4.3	4.4	0.4	0.95
80-92	18.4	8.3	3.0	0.3	30.0		25.5		4.8	4.2	20.9	0.5	-	2.2	6.7	22.0	2.83

Depth (in.)	8A Water at Saturation Pct.	Exch. bases 5B3b Ca	5B3b Mg	5B1b Na	5D2 Exchangeable Na Pct.	5E SAR	6F1a Gypsum Pct.	Clay mineralogy (<0.002 mm.) 3A1: Mont.	Verm.	Mica	Kaolin.	
		meq./100 g.						X-ray 7A2				
0-4	46.8			0.6	4	5		xx	-	-	x	
4-9	36.4			0.9	6	7						
9-21	60.9			6.4	28	21						
21-32	101.6			10.4	37	36		xxxx	-	-	x	
32-48	101.4			13.0	39	34		xxxx			tr.	
48-62	84.6			10.8	32	20						
62-80	75.5			8.6	23	21		xxxx	-	-	tr.	
80-92	49.1			2.0	8	10	-					

* Nitrites detected.

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

SOLANO LOAM (taxadjunct) 1/
S64Calif-48-7

Location: Solano County, California. A pit located at the NE corner of Sec. 3, T.4N., R.1W., M.D.B.&M., 5 miles east of Fairfield and 1 mile south of State Highway 12. Date of Sampling: October 8, 1964.
Description by: Wells Andrews. Collectors: Wells Andrews, Reuben Nelson, Ben Brasher, Leland Bates, Samuel Johnson.
Classification: Aquic Natrixeralf, fine-loamy, mixed, thermic.
Vegetation: Not cultivated. Natural vegetation consists of halophytic weeds and annual grasses. Climate: Average annual precipitation - 18 inches; mean annual temperature - 60°F. Parent Material: Alluvium from sedimentary rocks of the Coast Range. Topography: Convex basin rim or low terrace with an irregular slope of less than 1 percent. Elevation: 10 feet. Drainage: Very slow runoff, very slow permeability, and imperfect drainage. Free water was encountered at 68 inches. Soil Moisture: Dry to 12 inches, moist from 12 inches to 68 inches, and wet from 68 inches to 92 inches.
Remarks: Field pH determined by Hellige-Truog method and thymol blue. Paired profile No. Cal 48-4 (1-9). Diagnostic horizons are: Ochric epipedon, natric horizon.

HORIZON	DESCRIPTION
A21 RSL 64520	0 to 4 inches, light brownish gray (10YR 6/2) loam with few fine distinct yellowish brown (10YR 5/8) mottles; dark grayish brown (10YR 4/2) with few fine distinct yellowish red (5YR 4/8) mottles when moist; massive; hard, friable, nonsticky, slightly plastic; abundant micro and fine random roots; many micro and very fine tubular and common fine tubular pores; very strongly acid (pH 5.0); clear wavy boundary.
A22 RSL 64521	4 to 9 inches, light gray (10YR 7/2) loam with few fine distinct yellowish brown (10YR 5/6) mottles; dark grayish brown (10YR 4/2) with few fine distinct dark reddish brown (5YR 3/4) mottles when moist; massive; hard, friable, sticky, slightly plastic; abundant micro and very fine random roots; many micro and very fine tubular pores; strongly acid (pH 5.5); abrupt wavy boundary.
B21t RSL 64522	9 to 21 inches, brown (10YR 5/3) clay, brown (10YR 4/3) matrix and dark grayish brown (10YR 4/2) ped faces moist; strong coarse columnar structure; extremely hard, firm, sticky, plastic; plentiful micro and very fine random roots; many micro tubular pores; many thin clay films on ped faces and pores; neutral (pH 7.0); gradual wavy boundary.
B22t RSL 64523	21 to 32 inches, light yellowish brown (2.5Y 6/4) clay, olive brown (2.5Y 4/4) with dark reddish brown (5YR 2/2) stains moist; weak medium and coarse prismatic to weak medium angular blocky structure; extremely hard, firm, sticky, plastic; few micro random roots; common micro tubular pores; many moderately thick clay films on ped faces; moderately alkaline (pH 8.4); common Fe and Mn concretions; few slickensides; clear wavy boundary.
B23t RSL 64524	32 to 48 inches, light yellowish brown (10YR 6/4) clay, yellowish brown (10YR 5/6) matrix and light olive brown (2.5Y 5/4) ped faces moist with dark reddish brown (5YR 2/2) stains; weak medium and coarse prismatic to weak medium angular blocky structure; very hard, friable, slightly sticky, plastic; very few micro roots; common micro tubular pores; many moderately thick clay films on ped faces; strongly alkaline (pH 8.6); strongly effervescent medium soft masses and concretions of lime; many Fe and Mn concretions; few slickensides; gradual wavy boundary.
B24t RSL 64525	48 to 62 inches, light yellowish brown (10YR 6/4) clay, light olive brown (2.5Y 5/6) matrix and light olive brown (2.5Y 5/4) ped faces moist with dark reddish brown (5YR 2/2) stains; weak fine angular blocky structure; very hard, friable, sticky, plastic; no roots; common micro tubular pores; continuous moderately thick clay films on ped faces; strongly alkaline (pH 8.8); strongly effervescent large soft masses and concretions of lime; many Fe and Mn concretions; diffuse boundary.
B25t RSL 64526	62 to 80 inches, pale olive (5Y 6/3) light clay, olive (5Y 4/3) and yellowish brown (10YR 5/4) moist with dark reddish brown (5YR 2/2) stains; weak fine angular blocky structure; hard, friable, sticky, plastic; no roots; common micro tubular pores; continuous moderately thick clay films on ped faces; strongly alkaline (pH 8.8); strongly effervescent large soft masses and concretions of lime; many Fe and Mn concretions.
RSL 64527	80 to 92 inches, composite sample of which no description was made.

Samples taken for Bureau of Public Roads: A21, B21t, B24t

1/ The Solano series is in a fine-loamy, mixed, thermic family of Typic Natrixeralfs. This pedon has mottles within 30 inches of the soil surface.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aquic Natrixeralf, fine, montmorillonitic, thermic
SOIL Pescadero clay loam SOIL Nos. S64Calif-48-3 LOCATION Solano County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64451 - 64459

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	Coarse fragments 1A2a > 2 Pct.	2 - 19	19 - 76
		Pct. of < 2 mm															Pct. of < 76 mm	
0-4	A2	9.2	58.4	32.4	0.3	0.5	0.9	2.8	4.7	17.2	41.2	23.5	4.5			0		
4-14	B21t	19.3	41.5	39.2	0.1	0.3	1.0	7.3	10.6	18.3	23.2	34.1	8.7			0		
14-22	B22t	18.1	40.9	41.0	tr.	0.2	0.7	6.9	10.3	17.5	23.4	32.8	7.8			0		
22-34	B23t	20.0	40.4	39.6	0.2	0.4	0.9	7.5	11.0	16.7	23.7	33.2	9.0			0		
34-47	B3t	23.0	42.1	34.9	0.5	0.8	1.0	8.0	12.7	17.0	25.1	35.7	10.3			0		
47-58	C1	31.2	39.4	29.4	1.3	1.6	1.5	11.4	15.4	19.7	19.7	43.6	15.8			1		
58-69	C2	29.4	39.9	30.7	1.9	1.8	1.8	10.8	13.1	19.8	20.1	40.6	16.3			0		
69-85	C3	33.1	38.7	28.2	4.4	4.7	3.0	10.5	10.5	17.2	21.5	34.1	22.6			0		
109-119		24.3	44.0	31.7	tr.	0.1	0.6	7.1	16.5	24.7	19.3	46.8	7.8			0		

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	3B2 Cm	Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	Atterberg limits: Plastic limit Pct. of <0.4 mm	Liquid limit	Plastic index	Water content 4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLE in./in.	4D1 Extensibility COLE in./in.	pH 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-4	2.13	0.184	12	1.7	-	1.00	1.44	1.58	29	44	15	23.9	11.0	0.032	0.032		5.9	6.7
4-14	0.54	0.052	10	0.8	1	1.00	1.47	1.87				24.7	14.4	0.083	0.083		8.5	9.7
14-22	0.24	0.029	8	0.7	1	1.00	1.44	1.82	14	54	40	27.3	16.3	0.080	0.080		8.6	9.7
22-34	0.12			0.7	1	1.00	1.49	1.84				23.6	17.2	0.073	0.073		8.5	9.5
34-47	0.07			0.8	3	1.00	1.63	1.93				19.7	17.8	0.058	0.058		8.5	9.6
47-58	0.06			0.8	5	1.00	1.73	1.95	18	40	22	18.1	18.1	0.042	0.042		8.7	9.8
58-69	0.04			0.9	5	1.00	1.71	1.87				18.0	15.1	0.032	0.032		8.5	9.8
69-85	0.03			0.8	17	1.00	1.69	1.91				19.0	15.1	0.042	0.042		8.5	9.8
109-119	0.03			0.8	2								13.4				8.4	9.6

Depth (In.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	5B1a Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1a SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-4	8.2	10.8	3.0	0.5	22.5	7.9	32.3		0.3	0.2	6.6	0.1		-	0.5	5.6	1.8	0.91
4-14	11.3	12.3	18.3	0.6	42.5		28.6		0.5	1.9	91.2	0.2		-	4.2	15.4	30.2	5.33
14-22	9.9	13.7	22.1	0.5	46.2		29.3		1.0	2.9	68.5	0.3		-	2.9	30.0	49.0	7.8
22-34	3.6	13.2	22.5	0.3	39.6		27.5		0.8	2.0	71.5	0.3		-	2.4	27.5	44.6	7.1
34-47	3.9	13.8	20.2	0.2	38.1		26.4		0.6	2.0	57.2	0.3		-	3.0	24.8	37.9	6.3
47-58	6.5	12.0	16.5	0.2	35.2		23.7		0.5	1.2	40.5	0.3		-	3.2	18.2	24.6	4.58
58-69	7.2	12.9	14.2	0.2	34.5		25.8		0.6	1.5	40.0	0.3		-	3.1	19.6	20.6	4.30
69-85	10.7	14.4	10.3	0.2	35.6		25.7		0.7	1.7	30.4	0.1		-	2.7	17.9	14.3	3.67
109-119	10.0	15.0	5.6	0.2	30.8		27.7		0.6	1.0	12.7	0.2		-	2.7	6.4	5.6	1.68

Depth (In.)	8A Water at Saturation Pct.	Exch. bases 5B3b Ca	5B3b Mg	5B1b Na	5D2 Exchangeable Na Pct.	5E SAR	6F1a Gypsum Pct.	Clay mineralogy (<0.002 mm) 3A1 Mont.	Verm.	Mica	Kaolin.	
		meq./100 g.						X-ray 7A2				
0-4	51.0			2.6	8	13		xxxx	-	-	xx	
4-14	107.1			8.5	30	83	-					
14-22	116.0			14.2	48	49	-	xxxx	-	-	x	
22-34	135.6			12.8	46	60	-					
34-47	123.6			13.1	50	50	-					
47-58	114.1			11.9	50	44	-	xxxx	-	-	xx	
58-69	99.4			10.2	40	39	-					
69-85	81.7			7.8	30	28						
109-119	66.6			4.8	17	14						

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

PESCADERO CLAY LOAM
S64Calif-48-3

Location: Solano County, California. Art Brown Ranch, a pit located 200 feet south of NW corner of NE¼ NE¼ NW¼ Sec. 36, T6N, R1E, MDB&M. Date of Sampling: October 6, 1964
Description by: Grant Kennedy, Wells Andrews Samples Collected by: Reuben Nelson, Ben Brasher, Leland Bates, Sam Johnson, Grant Kennedy, Wells Andrews
Classification: Aquic Natrixeralf, fine, montmorillonitic, thermic.
Vegetation: Diminished annual grasses with some salt vegetation. Not cultivated. Climate: 18 inches of rainfall, 60 degrees mean annual temperature. Parent Material: Alluvium from sedimentary rocks. Topography: Basin with slope less than 1 percent. Elevation: 23 feet. Drainage: Runoff is slow, permeability is slow, drainage is imperfect. No water encountered within 119 inches of the surface. Soil Moisture: Dry 0-14", moist below 14".
Remarks: (1) pH determined by Hellige-Truog and thymol blue. (2) Sample number of paired profile 64 CAL 57-12-(1-8).(3) Diagnostic horizons are ochric and natric.(4) Inconspicuous intermittent gilgai is present.(5) Slickensides are present.(6) 1/2 inch cracks are present on the surface and extend downward 15 to 20 inches.

HORIZON	DESCRIPTION
A2	0 to 4 inches, light brownish gray (10YR 6/2) clay loam with common fine distinct

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aquic Natrixeralf, fine, montmorillonitic, thermic
SOIL Pescadero clay loam SOIL Nos. 864Calif-48-5 LOCATION Solano County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64460 - 64468

Depth (in.)	Horizon	1B1b Total: Sand (2–0.05)	Total: Silt (0.05–0.002)	Total: Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	3A1 Int. II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	Coarse fragments 1A2a > 2	2 – 19	19 – 76
		Size class and particle diameter (mm)														Pct.	Pct. of < 76 mm	
		Pct. of < 2 mm																
0-4	A2																	
4-14	B21t																	
14-21	B22t																	
21-30	B23t																	
30-34	B24tcs																	
34-49	B25t																	
49-62	B31tca																	
62-72	B32t																	
86-102																		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Pct.	Water content 4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEf in./in.	4D1 Extensibility COLE in./in.	pH 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-4													15.9					
4-14							1.49	2.00		1.00		22.8	19.0	0.105	0.105			
14-21													19.9					
21-30													18.5					
30-34							1.54	1.90		1.00		21.6	17.8	0.074	0.074			
34-49							1.59	1.88		1.00		22.3	17.1	0.056	0.056			
49-62													15.9					
62-72							1.70	1.87		1.00		18.6	14.5	0.033	0.033			
86-102													12.8					

Depth (in.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	5B1a Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 8A1 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1a SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-4	6.6	9.5	0.9	0.5	17.5		23.5		1.0	2.0	3.7	0.2		-	0.3	3.2	2.3	0.79
4-14	5.7	25.1	3.5	0.4	34.7		41.4		5.9	31.8	31.0	0.2		-	0.9	22.3	52.8	5.88
14-21	5.1	24.4	7.2	0.4	37.1		37.2		1.0	5.2	24.5	0.2		-	0.8	19.2	11.2	3.29
21-30	4.7	24.6	11.0	0.4	40.7		36.7		6.9	43.0	75.8	0.2		-	2.4	58.3	77.8	10.9
30-34	45.0	23.7	13.1	0.4	82.2		31.2		32.4	62.5	74.8	0.2		-	1.9	43.6	136	13.0
34-49	4.8	23.2	12.3	0.4	40.7		30.9		8.4	58.5	105	0.2		-	2.4	89.3	89.1	14.6
49-62	13.4	19.8	10.5	0.4	44.1		27.7		4.2	26.1	70.0	0.1		-	2.9	64.2	37.4	9.8
62-72	12.4	17.8	7.8	0.3	38.3		27.0		2.3	11.3	45.0	0.1		-	3.1	45.4	13.5	6.0
86-102	11.7	16.6	4.6	0.3	33.2		22.3		0.8	3.6	15.7	0.1		-	2.7	14.7	2.2	2.18

Depth (in.)	8A Water at Saturation Pct.	Exch. bases 5B3b Ca	5B3b Mg	5B1b Na	5D2 Exchangeable Na Pct.	5E SAR	6F1a Gypsum Pct.	Clay mineralogy (<0.002 mm.) Mont.	Verm.	Mica	Kaolin	
		meq./100 g.						X-ray				
0-4	41.3			0.8	3	3						
4-14	64.2			1.5	4	7	-					
14-21	67.0			5.5	15	14	-					
21-30	65.0			6.1	17	15	-					
30-34	60.0			8.6	28	11	2					
34-49	60.7			14.6	47	18	-					
49-62	61.5			6.2	22	18	-					
62-72	55.5			5.3	20	17						
86-102	53.8			3.7	17	11						

PESCADERO CLAY LOAM
S64-Calif-48-5

Location: Solano County, California. A pit on the Art Brown ranch about 9 miles south of Dixon - 300 feet NW of center of Sec. 1, T5N, R1E, MDB&M. Date of Sampling: October 7, 1964.
Description by: Grant Kennedy, Wells Andrews. Collectors: Reuben Nelson, Ben Brasher, [illegible]

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aquic Natrixeralf, fine, montmorillonitic, thermic

SOIL Pescadero clay loam SOIL Nos. S64Calif-48-6 LOCATION Solano County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64469 - 64479

Depth (In.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	Coarse fragments 1A2a: > 2 Pct.	2 - 19	19 - 76
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm															Pct. of < 76 mm	
0-5	A1																	
5-12	B21t																	
12-22	B22t																	
22-32	B23t																	
32-39	B24t																	
39-42	B25tca																	
42-51	B31tca																	
51-56	B32t																	
56-67	C1																	
67-80	IIC2																	
92-106																		

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc		Water content: Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLE in./in.	4D1 Extensibility COLE in./in.	pH: 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-5													12.9					
5-12													14.7					
12-22													15.5					
22-32													16.4					
32-39													18.8					
39-42													15.9					
42-51													15.1					
51-56													11.1					
56-67													10.6					
67-80													15.0					
92-106													14.3					

Depth (In.)	Extractable bases: 6N2a Ca	6O2a Mg	6P2a Na	5B1a 6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 8A1: 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1a SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-5	7.5	8.8	0.4	0.9	17.6		24.3		2.6	4.4	2.7	0.6		-	0.3	8.2	1.6	1.22
5-12	9.1	13.9	1.5	0.4	24.9		30.4		0.6	0.8	3.2	0.1		-	0.3	2.2	1.7	0.49
12-22	9.2	13.9	2.9	0.2	26.2		28.7		1.1	1.4	8.4	tr.		-	0.5	7.9	2.4	1.24
22-32	11.0	14.4	5.7	0.3	31.4		27.8		4.5	7.1	25.7	0.5		-	1.0	23.8	14.4	3.97
32-39	11.5	15.4	6.6	0.3	33.8		33.7		6.9	11.1	36.0	0.5		-	1.4	31.2	23.1	5.37
39-42	13.2	14.6	5.9	0.3	34.0		33.6		11.8	17.4	43.0	0.3		-	1.7	34.1	40.3	5.93
42-51	45.8	13.3	6.5	0.3	65.9		22.9		27.7	32.6	53.0	0.3		-	1.9	39.7	80.4	9.4
51-56	10.7	12.2	6.1	0.2	29.2		17.8		10.8	16.0	41.5	0.3		-	1.5	36.3	35.9	6.4
56-67	8.5	11.2	5.7	0.3	25.7		19.2		5.9	12.8	40.0	0.2		-	1.0	38.5	21.2	5.80
67-80	16.2	13.8	6.6	0.4	37.0		17.6		4.2	7.3	29.3	0.2		-	1.5	27.7	14.6	4.28
92-106	18.0	12.4	5.6	0.4	36.4		27.5		4.0	6.5	28.2	0.6		-	1.6	30.8	8.0	4.14

Depth (In.)	8A Water at Saturation Pct.	Exch. bases: 5B3b Ca	5B3b Mg	5B1b Na	5D2 Exchangeable Na Pct.	5E SAR	6F1a Gypsum Pct.	Clay mineralogy (<0.002 mm.): Mont.	Verm.	Mica	Kaolin	
		meq./100 g.						X-ray				
0-5	42.3			0.3	1	1						
5-12	40.8			1.3	4	4						
12-22	48.9			2.5	9	5						
22-32	54.9			4.3	16	11						
32-39	56.3			4.6	14	12	-					
39-42	53.2			3.6	11	11	-					
42-51	50.8			3.8	17	9	2					
51-56	48.2			4.1	23	11	-					
56-67	41.5			4.0	21	13	-					
67-80	58.6			4.8	27	12						
92-106	57.5			4.0	14	12						

PESCADERO CLAY LOAM
S64Calif-48-6

Location: Solano County, California. Schmeiser Ranch, a pit located NW¼SW¼NW¼ Sec. 30, T.5N, R.2E, MDB&M. 13 miles east of Fairfield and 1 mile north of State Highway 12. Date of Sampling: October 7, 1964.

Description by: Grant Kennedy and Wells Andrews. Collectors: Reuben Nelson, B. R. Brasher, Leland Bates, Samuel Johnson, Grant Kennedy, Wells Andrews.

Classification: Aquic Natrixeralf, fine, montmorillonitic, thermic.

Vegetation: Not cultivated. Diminished annual grasses. Climate: Average annual rainfall is 19 inches; mean annual temperature is 60°F. Parent Material: Alluvium from sedimentary rocks of the Coast Range. Topography: Basin. Elevation: 15 feet. Drainage: Runoff is slow, permeability is slow. Drainage is imperfect. No water table encountered within 106". Soil

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aquic Natrixeralf, fine, montmorillonitic, thermic

SOIL Pescadero clay SOIL Nos. 8640alif-57-12 LOCATION Yolo County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 6443 - 6450

Depth (in.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	Coarse fragments 1A2a: > 2 Pct.	2 - 19	19 - 76
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm															Pct. of < 76 mm	
0-3	A1	5.2	54.9	39.9	0.3	0.5	0.5	1.6	2.3	8.7	46.2	12.0	2.9			0		
3-13	B21t	2.0	46.0	52.0	0.1	0.1	0.1	0.5	1.2	6.6	39.4	8.1	0.8			0		
13-26	B22tca	2.1	49.8	48.1	0.3	0.3	0.1	0.5	0.9	6.2	43.6	7.4	1.2			0		
26-40	B23tca	8.5	51.2	40.3	0.5	0.8	0.4	1.8	5.0	12.0	39.2	18.3	3.5			0		
40-52	B31ca	16.7	48.6	34.7	1.0	1.1	0.6	3.8	10.2	19.2	29.4	32.3	6.5			0		
52-67	B32ca	17.7	46.2	36.1	0.2	0.5	0.4	4.1	12.5	21.0	25.2	36.8	5.2			0		
67-95	C1ca	22.4	46.0	31.6	0.2	0.2	0.3	5.0	16.7	21.7	24.3	42.6	5.7			0		
95-105	C2	20.8	56.5	22.7	tr.	0.1	0.2	4.9	15.6	23.1	33.4	43.0	5.2			0		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content: Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLE in./in.	4D1 Extensibility COLE in./in.		pH: 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-3	2.84	0.200	14	1.4	-		1.60	1.85		1.00		24.0	11.0	0.050	0.050			7.7	8.3
3-13	1.01	0.081	12	1.8	-		1.33	1.68		1.00		33.0	20.8	0.080	0.080			8.9	9.6
13-26	0.58	0.055	11	1.7	3		1.45	1.78		1.00		27.9	27.7	0.070	0.070			8.6	9.6
26-40	0.43	0.043	10	1.8	5		1.56	1.75		1.00		23.3	21.3	0.040	0.040			8.1	9.3
40-52	0.26			1.7	3		1.67	1.81		1.00		19.1	18.5	0.028	0.028			7.9	8.7
52-67	0.19			1.7	2		1.69	1.89		1.00		18.7	13.7	0.038	0.038			8.0	8.7
67-95	0.15			1.6	2		1.67	1.88		1.00		19.4	14.6	0.041	0.041			8.1	8.8
95-105	0.09			1.5	-		1.46	1.54		1.00		25.0	13.6	0.019	0.019			8.3	8.7

Depth (in.)	Extractable bases 5B1a: 6O2a Ca	6N2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 8A1: 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1a SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-3	4.8	9.5	12.6	1.8	28.7		28.9		1.2	2.2	61.2	0.7		-	4.1	50.2	6.6	6.6
3-13	3.5	9.6	34.4	0.4	47.9		38.0		0.5	0.8	73.8	0.2		5.4	2.0	35.5	25.6	7.0
13-26	4.7	15.0	32.2	0.3	52.2		33.3		0.4	4.9	136	0.7		-	6.8	64.7	66.3	13.0
26-40	10.2	18.8	21.6	0.3	50.9		29.7		4.4	25.8	170	0.7		-	6.5	65.0	81.8	15.8
40-52	15.5	18.8	5.6	0.5	40.4		25.1		13.2	35.1	49.8	0.4		-	1.9	69.4	15.8	9.4
52-67	18.5	17.7	3.9	0.4	40.5		28.1		10.6	22.8	31.8	0.2		-	2.2	51.8	12.2	6.2
67-95	16.4	16.0	1.4	0.4	34.2		25.1		3.3	5.4	6.4	0.1		-	2.3	10.3	2.5	1.64
95-105	8.8	14.1	0.9	0.4	24.2		22.8		0.4	0.6	2.7	0.1		-	2.6	1.3		0.49

Depth (in.)	8A Water at Saturation Pct.	Exch. bases 5B3b Ca	5B3b Mg	5B1b Na	5D2 Exchangeable Na Pct.	5E SAR	6F1a Gypsum Pct.	Atterberg limits: Plastic limit	Liquid limit	Plastic index		
		meq./100 g.						Pct. of <0.4 mm.				
0-3	55.0			9.2	32	47		31	48	17		
3-13	110.6			26.2	69	91	-					
13-26	106.8			17.7	53	84	-	30	56	26		
26-40	67.3			10.2	34	44	-					
40-52	47.7			3.2	13	10	-					
52-67	47.8			2.4	8	8	-					
67-95	44.6			1.2	5	3						
95-105	44.3			.0.8	4	4		18	24	6		

S64Calif-57-12

Location: Yolo County, California. A pit located about ½ mile east of Rd. 102 in the SE¼ NW¼ NW¼, Sec. 2, T.9N., R.2E. Area located about 2.5 miles east of Woodland. Photo ABB-41T-98. Date of Sampling: October 2, 1964.

Description by: G. J. Staidl. Collectors: R. E. Nelson, B. R. Brasher, G. M. Kennedy, W. F. Andrews, L. A. Bates, G. J. Staidl, C. B. Goudey.

Classification: Aquic Natrixeralf, fine, montmorillonitic, thermic.

Vegetation: Not cultivated, area is in permanent pasture, plants present are pickleweed and saltgrass. Climate: Average annual precipitation is 17.3 inches, mean annual temperature is 61°F. Parent Material: Fine textured alluvium from sedimentary rocks. Topography: Basin; plane (slightly hummocky); less than one percent slope. Elevation: 35 feet. Drainage: Runoff is slow: permeability is slow; drainage is imperfect to poor. Soil Moisture: The profile was moist below 8 inches.

Remarks: (1) Truog soil reaction test and Thymol Blue were used in field pH determination; (2) Paired sample No. 64 Cal. 48-3-1 through 9; (3) Diagnostic horizons-albic or ochric, and natric; (4) Soil temperature at about 100" was 22.5°C.; (5) BPR sample horizons: A2 - 0-3; B22t - 13-26; C1 - 95-105; (6) There is evidence of plant residue in cracks to a depth of 12 inches.

HORIZON	DESCRIPTION
A1 RSL 64443	0 to 3 inches, light gray (2.5Y 7/2) silty clay loam, dark grayish brown (2.5Y 4/2) when moist: weak medium platy structure; very hard, friable, sticky and plastic: plentiful micro random roots; common micro tubular pores; mildly alkaline (pH 7.7) on the immediate surface, moderately alkaline (pH 8.0) about an inch to three inches below the surface; abrupt wavy boundary.
B21t RSL 64444	3 to 13 inches, gray (5Y 5/1) silty clay, very dark gray (5Y 3/1) when moist: strong, very coarse prismatic structure; very hard, friable, sticky and plastic: plentiful micro and abundant medium roots; many micro tubular pores; many moderately thick clay films on ped faces and in pores; strongly alkaline (pH 8.9); gradual wavy boundary.
B22tca RSL 64445	13 to 26 inches, light olive gray (5Y 6/2) silty clay with common fine distinct strong brown (7.5YR 5/6) and gray (N 5/0) mottles, olive (5Y 4/3) with common fine distinct brown to dark brown (7.5YR 4/4) and very dark gray (N 3/0)mottles moist; moderate medium prismatic breaking to moderate medium to coarse angular blocky structure; very hard, firm, sticky and plastic; plentiful micro and abundant medium roots; common micro tubular pores; many moderately thick clay films on ped faces and in pores; slightly effervescent with segregated lime in soft masses; strongly alkaline (pH 8.6); clear wavy boundary.
B23tca RSL 64446	26 to 40 inches, light olive gray (5Y 6/2) silty clay with many fine distinct strong brown (7.5YR 5/6)mottles, olive gray (5Y 4/2) with many fine distinct yellowish red (5YR 4/6) and very dark grayish brown (2.5Y 3/2) mottles when moist; moderate medium prismatic breaking to moderate medium to coarse angular blocky structure; very hard, firm, sticky and plastic; few micro, plentiful medium roots; many micro tubular pores; many moderately thick clay films in pores and continuous moderately thick clay films on ped faces; slightly effervescent segregated lime in seams and soft masses; moderately alkaline (pH 8.1); gradual wavy boundary.
B31ca RSL 64447	40 to 52 inches, light olive gray (5Y 6/2) silty clay loam with many fine distinct strong brown (7.5YR 5/6) mottles, olive gray (5Y 4/2) with many fine distinct yellowish red (5YR 4/6)mottles when moist; moderate medium prismatic breaking to moderate medium to coarse angular blocky structure; hard, firm, sticky and plastic; few micro to fine roots; many micro tubular pores; continuous thin clay films on ped faces and many moderately thick clay films in pores; strongly effervescent with disseminated and segregated lime in seams; moderately alkaline (pH 7.9); gradual wavy boundary.

CONSECUTIVE PEDON NUMBER C 69 24 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES COMETA TAXADJUNCT SOIL NO. S69 CALI. 31-18 SAMPLE NO. 9 106-9 114 PLACER COUNTY AREA

SOIL FAMILY TYPIC PALEXERALF, FINE-LOAMY, MIXED, THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL SAND 2- .05	FRACTION SILT .05- .002	CLAY LT .002	SANDS VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A	INT. II .2- .02	1A2A PCT. OF WHOLE SOIL	COARSE FRAGMENTS I 75- 20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM															PCT. OF LT 75		
9 106	0- 20	AP	53.7	36.4	9.9	4.8	8.7	7.9	21.3	11.0	18.5	17.9				41.0	--A	--	--	--
9 107	20- 46	B21T	34.1	25.8	40.1	1.9	4.0	4.4	15.4	8.4	11.1	14.7				29.1	--A	--	--	--
9 108	46- 61	B22T	42.0	24.4	33.6	0.7	2.3	6.1	22.8	10.1	9.7	14.7				32.0	--A	--	--	--
9 109	61- 86	B31T	56.7	20.2	23.1	1.6	6.0	14.3	29.2	5.6	6.8	13.4				24.5	--A	--	--	--
9 110	86-115	B32T	71.7	10.4	17.9	3.0	13.4	23.1	28.6	3.6	3.8	6.6				17.0	--A	--	--	--
9 111	115-140	C1 SI	77.2	6.8	16.0	4.9	10.5	18.9	40.0	2.9	2.1	4.7				16.3	1A	--	--	1
9 112	140-165	C2 SI	79.4	7.8	12.8	9.9	10.6	14.8	42.1	2.0	2.8	5.0				18.4	5A	--	1	4
9 113	165-190	C3	80.7	8.6	10.7	15.8	19.6	16.7	25.0	3.6	4.3	4.3				17.5	17A	--	2	15
9 114	190-205	C4	84.2	6.6	9.2	20.7	30.3	17.3	13.5	2.4	3.8	2.8				11.4	14A	--	2	12

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR.	NITRO.	CARBONATE AS CACO3	FROM 1/3 BAR	ATTERBERG L.L.	P.L.	P.I.	BULK DENSITY EST.	1/3	DRY	CM	WATER CONTENT 1/10	1/3	15	EXTENSBLTY LES	LE	PH H2O	H2O	CACL2
9 106	0- 20																			
9 107	20- 46																			
9 108	46- 61																			
9 109	61- 86																			
9 110	86-115																			
9 111	115-140																			
9 112	140-165																			
9 113	165-190																			
9 114	190-205																			

SAMPLE NO.	DEPTH CM.	6A1A -PERCENT- CA	6B1A MG	6E1B -PERCENT- NA	6E1B K	6C2B PCT.	4F2 PCT.NO.40	4F1	4A1F GRAMS PER CC	4A1F	4AH	3B2	4B1C -PERCENT-	4B1C	4B2	4D1 -PERCENT-	4D1	8C1A	8C1A	8C1E
9 106	0- 20	0.46	0.042			0.7				1.79	1.85	1.00		13.7	4.1	1.1	1.1	5.9		5.0
9 107	20- 46	0.25	0.028			1.0				1.48	1.99	1.00		27.5	16.3	10.4	10.4	5.9		5.4
9 108	46- 61	0.16				1.0				1.49	1.83	1.00		25.6	16.7	7.1	7.1	6.4	7.4	6.1
9 109	61- 86	0.06				1.1				1.70	1.81	1.00		17.6	14.4	2.1	2.1	7.0		6.6
9 110	86-115					1.1				1.64	1.71	1.00		16.3	10.9	1.4	1.4	7.1	7.5	6.8
9 111	115-140					0.9				1.51	1.60	1.00		20.1	9.6	1.9	1.9	7.0	7.3	6.6
9 112	140-165					0.9				1.50	1.58	0.98		17.4	8.9	1.7	1.7	6.7	7.4	6.2
9 113	165-190					0.6				1.66	1.75	0.89		12.6	7.1	1.8	1.6	6.7		5.9
9 114	190-205									1.56	1.65	0.92		12.7	5.2	1.9	1.7	6.5		5.7

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT. PCT.	C E C NA- OAC 5A2A	NH4- OAC 5A6A	EC 8A1A	WATER EXTRACT FROM SATURATED PASTE CA 6N1B	MG 6O1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	8A1 SO4 6L1B	B 6T1A	PH SAT. PASTE 8C1B
		MEQ/100G						MEQ/100G		MMHO	MEQ/1000 ML								PPM	
9 106	0- 20	3.6	2.3	0.1	0.2	6.2	81		7.7											
9 107	20- 46	14.3	11.4	0.5	0.3	26.5	98		27.1											
9 108	46- 61	17.7	15.5	0.8	0.3	34.3	100+		28.4	0.57	1.8	1.9	3.3	0.1		2.5	0.6			6.1
9 109	61- 86	14.5	13.7	0.9	0.2	29.3	100+		27.6	0.43										
9 110	86-115	10.9	11.6	1.1	0.2	23.8	100+		22.0	2.07	5.9	6.9	7.1	0.1	TR	0.5	14.4	4.3		7.0
9 111	115-140	8.9	9.7	1.0	0.1	19.7	100+		16.5	2.85	9.0	10.1	8.6	0.1	TR	0.6	24.6	1.2		6.9
9 112	140-165	8.1	8.7	0.8	0.2	17.8	100+		15.7	2.36	7.2	8.1	6.8	0.1	TR	0.3	20.8			6.7
9 113	165-190	5.9	6.9	0.5	0.1	13.4	100+		11.7											
9 114	190-205	4.7	4.4	0.3	0.1	9.5	100+		8.8											

SAMPLE NO.	DEPTH CM.	WATER AT SAT. 8A PCT.	EXCHANGBLE NA 5B4B MEQ.	NA 5D2 PCT.	SAR	GYPSUM AMT. REQ. 6F1B PERCENT	6P3A	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 PCT.	L.T. .0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
9 106	0- 20		0.1	1				MI 3	KK 3	MT 1						MI MICA	1 TRACE
9 107	20- 46		0.5	2				MI 3	KK 3	MT 2						KK KAOLINIT	2 SMALL
9 108	46- 61	47.9	0.6	2	2											MT MONTMORT	3 MODERATE
9 109	61- 86		0.9	3													4 ABUNDANT
9 110	86-115	36.8	0.8	4	3	TR										MM MT/MICA	5 DOMINANT
9 111	115-140	36.8	0.7	4	3	TR											6 INDETER.

COMETA SANDY LOAM (TAXADJUNCT)* 1/
S69Calif-31-18

Location: Placer County, California. Approximately 2.2 miles southwest of Lincoln; Chris Story ranch. 1,700 feet north of SE corner of sec.20,T.12N.,R.6E. Date of Sampling: 5/8/69.
Description by: L. A. Brittan, T. M. Ryan. Collectors: K. W. Flach, W. D. Nettleton, G. M. Kennedy, T. M. Ryan, L. A. Brittan.
Classification: Typic Palexeralf, fine-loamy, mixed, thermic.
Vegetation: Soft chess, annual rye, rose clover, ripgut brome. Climate: Average annual precipitation 21 inches. Mean annual air temperature 61°F. Parent Material: Granitic sediments. Topography: Gently sloping terrace on a southwest slope of 5 percent. Elevation: 125 feet. Drainage: Moderately well drained; medium runoff; very slow permeability. Soil Moisture: Dry to 18 inches; slightly moist below.
Remarks: Roots stop at top of C1si horizon and are mainly in structural surfaces. Roots in B21t and B22t horizons mainly in cracks between prisms. Some pores in B22t horizon may be root channels. Small gravel occurs in C1si horizon and below. *Has very slightly less than 35 percent clay throughout the upper 50 cm of the argillic horizon and has less montmorillonitic clay than characteristic of the series.

HORIZON	DESCRIPTION
Ap RSL 69106	(0-20 cm.) 0 to 8 inches, brown (7.5YR 5/4) sandy loam, brown (7.5YR 4/4) moist; many medium distinct pale brown (10YR 6/3) mottles, common fine faint brown (7.5YR 5/4) mottles moist; massive; slightly hard, friable, nonsticky, nonplastic; common very fine vertical roots; common very fine tubular and interstitial pores; medium acid (pH 5.8); abrupt smooth boundary.
B21t RSL 69107	(20-46 cm.) 8 to 18 inches, brown (10YR 5/3 and 7.5YR 5/4) clay, dark brown (10YR 4/3) moist; strong medium and coarse prismatic structure; very hard, very firm, very sticky, very plastic; common very fine roots in structural planes; few very fine tubular pores; medium acid (pH 6.0); clear wavy boundary.
B22t RSL 69108	(46-61 cm.) 18 to 24 inches, light yellowish brown (10YR 6/4) clay, dark brown (7.5YR 4/4) moist; few fine distinct yellowish red (5YR 5/6) mottles; moderate medium prismatic structure; very hard, very firm, very sticky, very plastic; few very fine vertical and oblique roots; common very fine tubular pores; common moderately thick clay films line pores; slightly acid (pH 6.5); clear wavy boundary.
B31t RSL 69109	(61-86 cm.) 24 to 34 inches, light brown (7.5YR 6/4) sandy loam, variegated dark yellowish brown (10YR 4/4) and dark brown (7.5YR 4/4) moist; common medium distinct yellowish red (5YR 5/6) mottles, yellowish red (5YR 4/6) clay films moist; massive; hard, firm, nonsticky, nonplastic; few very fine vertical and oblique roots; many very fine interstitial pores; common moderately thick clay films line pores; mildly alkaline (pH 7.8); gradual wavy boundary.
B32t RSL 69110	(86-115 cm.) 34 to 46 inches, variegated yellowish red (5YR 5/6), pale brown (10YR 6/3) and light yellowish brown (10YR 6/4) sandy loam, reddish brown (5YR 4/4), brown (7.5YR 5/4) and strong brown (7.5YR 5/6) moist; massive; hard, firm, nonsticky, nonplastic; few very fine vertical and oblique roots; many very fine interstitial pores; many moderately thick clay films bridging and staining mineral grains; moderately alkaline (pH 8.0); clear smooth boundary.
C1si RSL 69111	(115-140 cm.) 46 to 55 inches, variegated yellowish red (5YR 5/6) light yellowish brown (10YR 6/4) gravelly sandy loam, reddish brown (5YR 4/4), brown (7.5YR 4/4), strong brown (7.5YR 5/6) moist; massive; hard, firm, nonsticky, nonplastic; many very fine interstitial pores; many moderately thick clay films bridging and staining mineral grains; mildly alkaline (pH 7.8); clear smooth boundary.
C2si RSL 69112	(140-165 cm.) 55 to 64 inches, variegated yellowish brown (10YR 5/8),strong brown (7.5YR 5/6) gravelly sandy loam, dark brown (7.5YR 4/4) moist; massive; hard, firm, nonsticky, nonplastic; many very fine interstitial pores; many moderately thick clay films bridging and staining mineral grains; slightly acid (pH 6.5); clear smooth boundary.
C3 RSL 69113	(165-190 cm.) 64 to 75 inches, variegated pale brown (10YR 6/3), light yellowish brown (10YR 6/4) gravelly sand, brown (7.5YR 5/4), dark brown (7.5YR 4/4) moist; massive; hard, firm, nonsticky, nonplastic; many very fine interstitial pores; few thin clay films bridging mineral grains; slightly acid (pH 6.5); gradual wavy boundary.
C4	(190-205 cm.) 75 to 80 inches, variegated gravelly sand; massive; hard, firm, nonsticky, nonplastic;

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Palexeralf, fine-loamy, mixed, thermic

SOIL Corning loam (taxadjunct) SOIL Nos. S4557-12 LOCATION Glenn County, California

SOIL SURVEY LABORATORY _____ LAB. Nos. _____

Depth (In.)	Horizon	7B1b Total Sand (2–0.05)	Total Silt (0.05–0.002)	Total Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	Int. II (0.2–0.02)	3A1 (2–0.1)			Coarse fragments > 2 Pct.	Coarse fragments 2 – 19 (Pct. of < 76mm)	Coarse fragments 19 – 76 (Pct. of < 76mm)
		Pct. of < 2 mm																
0-7	Ap	45.6	45.9	8.5	6.9	7.7	5.4	10.7	14.9	18.9	27.0		30.7					
7-13½	A1	43.3	44.7	12.0	5.1	6.0	5.0	12.1	15.1	18.0	26.7		28.2					
13½-21	B21t	36.0	24.5	39.5	1.6	2.7	5.1	16.9	9.7	8.8	15.7		26.3					
21-29	B22t	58.6	14.6	26.8	0.5	1.5	4.8	43.0	8.8	4.5	10.1		49.8					
29-35	IIC1	71.3	6.5	22.2	20.9	17.8	15.8	14.3	2.5	1.4	5.1		68.8					
35-41	IIC2	78.4	3.1	18.5	33.8	20.2	15.0	8.2	1.2	0.6	2.5		77.2					
41-47	IIC3	79.7	3.6	16.7	28.3	19.9	19.1	10.8	1.6	0.7	2.9		78.1					

Depth (In.)	6A2a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	Bulk density g/cc	Bulk density g/cc		Water content Pct.	Water content Pct.	Water content Pct.			pH	pH	8C1a (1:1)
0-7	0.48	0.058																5.7
7-13½	0.22	0.039																5.7
13½-21	0.32	0.058																6.4
21-29	0.14	0.036																7.1
29-35	0.09	0.025																7.6
35-41	0.08	0.024																7.6
41-47	0.08	0.020																7.6

Depth (In.)	Extractable bases 5B1a: Ca 6N2b	Mg 6O2b	Na 1/	K 2/	Sum of bases	6H2a Ext. Acidity	5A3a CEC Sum									Base saturation 5C3 Sum Pct.	Pct
	meq/100 g																
0-7	2.7	1.4	0.06	0.22	4.4	2.8	7.2									61	
7-13½	3.6	3.4	0.05	0.21	7.3	2.4	9.7									75	
13½-21	9.1	15.3	0.14	0.31	24.8	3.5	28.4									88	
21-29	6.1	11.0	0.11	0.33	17.5	1.4	18.9									93	
29-35	4.6	8.1	0.19	0.24	13.1	0.9	14.0									94	
35-41	5.4	7.5	0.19	0.31	13.4	1.0	14.4									93	
41-47	4.1	6.9	0.14	0.29	11.4	0.8	12.2									94	

Depth (In.)				

1/ Gravimetrically as sodium uranyl magnesium acetate

2/ Precipitated as potassium sodium cobaltinitrite and titrated as potassium permanganate

CORNING LOAM (taxadjunct) 1/
S4557-12

Location: Glenn County, California. Pit located in the NW¼ NE¼, Sec. 2, T22N, R3W, MDB&M, 0.6 miles east of the NW section corner. Date of Sampling: April 5, 1945.

Description by: R. C. Roberts and R. A. Gardner. Collectors: R. C. Roberts, F. O. Youngs, R. A. Gardner, L. T. Alexander, H. Jenny and R. E. Storie.

Classification: Typic Palexeralf, fine-loamy, mixed, thermic.

Vegetation: Sheep pasture--formerly cultivated. Cover consists of fairly dense stand of short annual grasses and associated herbaceous plants. Climate: Average annual precipitation is 18 inches. Average annual temperature is 63°F., January 45°F., and July 82°F. Parent Material: Mixed alluvium, mainly sedimentary. Topography: Nearly level older terrace. Elevation: 230 feet. Drainage: Well drained, slow permeability, slow runoff. Soil Moisture: Moist.

Remarks: This is a very good example of a soil that is only slightly gravelly in the upper part of the profile but very gravelly in the lower part. The outstanding characteristics of this soil are: (1) The abrupt boundary between the surface and subsoil; (2) the very compact subsoil; (3) the slight mottling probably due to imperfect drainage caused by occasional high water table in the lower part of the profile; (4) gravelly and rather friable condition of the profile below a depth of 30 inches.

Relief, climate, and land use of this site are very similar to those of profile S4557-13 of Corning loam. Vegetation and carrying capacity are much better here than at the site of profile S4557-3 of Corning loam.

HORIZON	DESCRIPTION
Ap	0 to 7 inches, reddish yellow (7.5YR 6/6) gravelly light loam containing a slight amount of gravel, reddish brown (5YR 4/4) moist; weak granular structure; slightly hard, very friable, slightly plastic; abundant small roots; medium acid (pH 5.7); distinct smooth boundary.
A1	7 to 13.5 inches, reddish yellow (7.5YR 6/6) loam containing a slight amount of gravel, reddish brown (5YR 4/4) moist; weak granular structure; hard, friable, slightly plastic; plentiful small roots; medium acid (pH 5.7); abrupt smooth boundary.
B21t	13.5 to 21 inches, yellowish red (5YR 5/6) heavy clay loam (near clay) containing a slight amount of gravel, red (2.5YR 4/6) moist; coarse blocky structure; very compact; extremely hard, extremely firm, very plastic; few roots; slightly acid (pH 6.4); distinct wavy boundary.
B22t	21 to 29 inches, reddish brown (5YR 5/4) sandy clay loam, reddish brown (5YR

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Palexeralf, fine, montmorillonitic, thermic

SOIL Corning loam SOIL Nos. 84557-3 LOCATION Tehama County, California

SOIL SURVEY LABORATORY _____ LAB. Nos. _____

Depth (in.)	Horizon	1B1b Total Sand (2–0.05)	Total Silt (0.05–0.002)	Total Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)			Coarse fragments ≥ 2 Pct.	2–19 Pct. of < 76mm	19–76 Pct. of < 76mm
		Pct. of < 2 mm																
0-7	Ap	43.8	46.9	9.3	3.8	4.9	4.4	12.1	18.6	23.1	23.8		25.2					
7-13	A11	43.5	45.9	10.6	4.6	4.8	4.3	11.9	17.9	22.3	23.6		25.6					
13-19	A12	43.4	43.8	12.8	7.3	5.2	4.4	11.1	16.2	21.2	22.6		27.2					
19-27½	B21t	38.8	19.3	41.9	5.8	4.8	6.8	14.1	7.3	6.4	12.9		31.5					
27½-35	B22t	31.3	24.3	44.4	6.8	5.1	4.5	7.6	7.3	9.7	14.6		24.0					
35-48	B23t	35.8	20.5	43.7	11.5	7.8	3.9	6.3	6.3	7.6	12.9		29.5					

Depth (in.)	6A2a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	Pct.			pH		8C1a (1:1)
0-7	0.53	0.056																5.7
7-13	0.28	0.038																6.1
13-19	0.18	0.029																6.5
19-27½	0.28	0.052																7.0
27½-35	0.12	0.030																7.6
35-48	0.08	0.024																7.8

Depth (in.)	Extractable bases: Ca 6N2b	Mg 602b	Na 1/	K 2/ (5B1a)	Sum of bases	6H2a Ext. Acidity	5A3a CEC Sum									Base saturation: 5C3 Sum Pct.	Pct.
	meq/100 g																
0-7	3.9	1.9	0.03	0.14	6.0	2.9	8.9									67	
7-13	4.4	2.5	0.02	0.09	7.0	2.4	9.4									75	
13-19	4.5	4.4	0.15	0.03	9.1	2.2	11.3									80	
19-27½	11.1	17.4	0.72	0.20	29.4	3.4	32.8									90	
27½-35	10.5	18.4	0.51	0.25	29.7	1.8	31.5									94	
35-48	10.1	17.5	0.46	0.25	28.3	1.7	30.0									94	

Depth (in.)

1/ Gravimetrically as sodium uranyl magnesium acetate

2/ Precipitated as potassium sodium cobaltinitrite and titrated as potassium permanganate

CORNING LOAM
S4557-3

Location: Tehama County, California. Pit located 0.05 miles south of the center of Sec. 29, T25N, R3W, MDB&M. Date of Sampling: April 1, 1945.

Description by: R. C. Roberts and R. A. Gardner. Collectors: R. C. Roberts, F. O. Youngs, R. A. Gardner, L. T. Alexander, H. Jenny and R. E. Storie.

Classification: Typic Palexeralf, fine, montmorillonitic, thermic.

Vegetation: Pasture--formerly cultivated. Vegetation is short annual grasses and weeds. Climate: Average annual precipitation is 19 inches. Average annual tem-

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Palexeralf, fine, montmorillonitic, thermic
SOIL Corning loam SOIL Nos. S4557-13 LOCATION Glenn County, California

SOIL SURVEY LABORATORY _____ LAB. Nos. _____

Depth (in.)	Horizon	1B1b Total: Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Size class and particle diameter (mm) 3A1 Sand: Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt: 0.05–0.02	Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)			Coarse fragments > 2 Pct.	2 – 19	19 – 76
		← Pct. of < 2 mm →															← Pct. of < 76mm →	
0-6	Ap	48.6	41.6	9.8	5.2	7.0	5.5	13.8	17.1	17.7	23.9		31.5					
6-12	A1	46.1	41.5	12.4	4.9	7.2	5.1	13.3	15.6	16.8	24.7		30.5					
12-18½	B21t	29.6	26.9	43.5	1.4	2.4	2.7	11.3	11.8	10.3	16.6		17.8					
18½-24	B22t	32.1	27.2	40.7	0.7	1.8	2.6	12.7	14.3	10.7	16.5		17.8					
24-35	B3	40.5	32.9	26.6	0.7	1.5	1.8	14.2	22.3	15.1	17.8		18.2					
35-44	C1	54.2	23.8	22.0	5.8	10.5	7.5	14.7	15.7	9.8	14.0		38.5					
44-54	C2	52.1	26.0	21.9	3.4	7.5	6.1	14.7	20.4	12.2	13.8		31.7					
54-70	IIC3	85.5	5.2	9.3	13.8	29.8	22.0	17.0	2.9	1.3	3.9		82.6					

Depth (in.)	6A2a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	Pct			pH		8C1a (1:1)
0-6	0.39	0.048																6.0
6-12	0.24	0.040																6.3
12-18½	0.30	0.046																6.9
18½-24	0.22	0.052																7.6
24-35	0.08	0.036																8.2
35-44	0.08	0.030																7.7
44-54	0.08	0.032																7.8
54-70	0.06	0.020																7.6

Depth (in.)	Extractable bases 5B1a: Ca 6N2b	Mg 6O2b	Na 1/	K 2/	Sum of bases	6H2a Ext. Acidity	5A3a CEC Sum									Base saturation 5C3 Sum Pct.	Pct.
	← meq/100 g →																
0-6	3.2	2.1	0.06	0.30	5.7	2.3	8.0									71	
6-12	4.4	3.5	0.05	0.18	8.1	2.0	10.1									80	
12-18½	11.1	16.4	0.21	0.37	28.1	3.5	31.6									89	
18½-24	9.8	15.3	0.16	0.46	25.7	1.5	27.2									95	
24-35	7.1	10.9	0.14	0.41	18.5	0.6	19.2									97	
35-44	5.2	8.4	0.08	0.35	14.0	0.7	14.7									95	
44-54	5.6	8.7	0.19	0.35	14.8	0.8	15.6									95	
54-70	2.4	3.5	0.03	0.17	6.1	0.6	6.7									91	

Depth (in.)

1/ Gravimetrically as sodium uranyl magnesium acetate

2/ Precipitated as potassium sodium cobaltinitrite and titrated as potassium permanganate

CORNING LOAM
S4557-13

Location: Glenn County, California. NW¼ SW¼ Sec. 2, T22N, R3W, MDB&M, 0.55 mile south of the section corner near Malton. Date of Sampling: April 5, 1945.
Description by: R. C. Roberts and R. A. Gardner. Collectors: R. C. Roberts, F. O. Youngs, R. A. Gardner, L. T. Alexander, H. Jenny and R. E. Storie.

Classification: Typic Palexeralf, fine, montmorillonitic, thermic.
Vegetation: Used for pasture. Cover consists of annual grasses and associated herbaceous plants. Climate: Average annual precipitation is 18 inches. Average annual temperature is 63°F., January 45°F., July 82°F. Parent Material: Mixed alluvium,

percent. Elevation: 250 feet. Drainage: Well drained, slow permeability, slow runoff. Soil Moisture: Moist.
Remarks: This is a representative profile of areas where the soils of the old terraces gradually grade into the soils of the basins. This soil probably has been influenced in the past by fluctuating ground-water table, and the higher pH of the lower part of the profile may be due to some deposition of calcium carbonate from the ground water. There appears to be some "greenish" or olive-colored mottling at depths between 2 and 3.5 feet.

HORIZON	DESCRIPTION
Ap	0 to 6 inches, strong brown (7.5YR 5/5) light loam containing a slight amount of gravel, reddish brown (5YR 4/3) moist; weak granular structure; slightly hard, friable, slightly plastic; abundant small roots; medium acid (pH 6.0); distinct smooth boundary.
A1	6 to 12 inches, strong brown (7.5YR 5/5) loam containing a slight amount of gravel; dark reddish brown (5YR 3/4) moist; structure and consistence similar to layer above; abundant small roots; slightly acid (pH 6.3); abrupt wavy boundary.
B21t	12 to 18.5 inches, yellowish red (5YR 5/6) clay, color of surface of dry aggregate red (2.5YR 5/6), red (2.5YR 4/6) moist; strong coarse blocky structure; very compact; extremely hard, extremely firm, very plastic; few roots; neutral (pH 6.9); gradual boundary.
B22t	18.5 to 24 inches, strong brown (7.5YR 5/5) light clay, yellowish red (5YR 4/5) moist; weak coarse blocky structure; compact; very hard, very firm, moderately

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Palexeralf, fine, montmorillonitic, thermic

SOIL Corning loam SOIL Nos. S4557-14 LOCATION Tehama County, California

SOIL SURVEY LABORATORY _____ LAB. Nos. _____

Depth (In.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)			Coarse fragments > 2 Pct.	2-19 (Pct. of < 76mm)	19-76 (Pct. of < 76mm)
		Pct. of < 2 mm																
0-8	Ap	47.5	42.3	10.2	10.5	9.4	5.6	9.5	12.5	19.7	22.6		35.0					
8-15	A11	45.9	41.5	12.6	8.4	8.9	5.3	10.0	13.3	19.4	22.1		32.6					
15-21	A12	43.8	41.8	14.4	7.9	8.4	5.2	9.5	12.8	19.6	22.2		31.0					
21-29	B21t	21.6	26.7	51.7	5.4	3.6	2.2	4.4	6.0	8.6	18.1		15.6					
29-36	B22t	36.8	28.6	34.6	9.9	6.8	3.8	7.9	8.4	7.2	21.4		28.4					
36-45	IIC1	56.1	20.9	23.0	16.8	15.9	5.6	9.0	8.8	4.3	16.6		47.3					
45-54	IIC2	51.8	15.3	32.9	17.2	16.8	7.1	6.1	4.6	3.6	11.7		47.2					

Depth (In.)	6A2a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	Pct.			pH		8C1a (1:1)
0-8	0.22	0.033																5.5
8-15	0.10	0.025																5.7
15-21	0.09	0.023																5.4
21-29	0.14	0.032																5.2
29-36	0.08	0.020																5.2
36-45	0.04	0.018																5.2
45-54	0.06	0.020																5.2

Depth (In.)	Extractable bases: Ca. 6N2b	Mg 6O2b	Na 1/	K 5B1a 2/	Sum of bases	6H2a Ext. Acidity	5A3a CEC Sum									Base saturation: 5C3 Sum Pct.	Pct
	meq/100 g																
0-8	1.2	1.1	<0.01	0.46	2.8	2.3	5.1									54	
8-15	1.4	1.9	<0.01	0.23	3.5	2.0	5.5									64	
15-21	1.6	2.6	tr.	0.17	4.4	2.3	6.7									65	
21-29	5.6	13.5	0.47	0.36	19.9	5.6	25.5									78	
29-36	8.0	18.6	0.64	0.27	27.5	6.1	33.6									82	
36-45	7.2	16.4	0.63	0.35	24.6	5.4	30.0									82	
45-54	7.4	16.7	0.51	0.37	25.0	5.0	30.0									83	

Depth (In.)				

1/ Gravimetrically as sodium uranyl magnesium acetate

2/ Precipitated as potassium sodium cobaltinitrite and titrated as potassium permanganate

CORNING LOAM
S4557-14

Location: Tehama County, California. Pit located just east of U. S. Highway 99W, in the SW¼ NW¼, Sec. 22, T23N, R3W, MDB&M. Date of Sampling: April 5, 1945.

Description by: R. C. Roberts and R. A. Gardner. Collectors: R. C. Roberts, F. O. Youngs, R. A. Gardner, L. T. Alexander, H. Jenny and R. E. Storie.

Classification: Typic Palexeralf, fine, montmorillonitic, thermic.

Vegetation: Used for sheep range. Vegetation is short annual grasses and associated herbaceous plants. Climate: Average annual precipitation is about 19 inches. Mean annual temperature is about 62°F., mean January temperature is about 45°F., and mean July temperature is about 82°F. Parent Material: Terrace alluvium. Topography: Older terrace on slope of about 2 percent. Elevation: 270 feet. Drainage: Well drained, very slow permeability, slow runoff. Soil Moisture: Moist throughout.

Remarks: This soil as observed in the pit and in nearby roadside cuts is very similar to the profile of Corning gravelly loam. One noticeable characteristic that is different is that this profile has little if any evidence of cementation below the claypan, and in this respect it is more typical of the Corning soils. Another difference is the rather abnormally low pH of the lower subsoil. There is evidence of more rodent activity throughout this profile than in the other Corning profile, and much of the rodent activity appears recent. It was quite evident that [illegible] grass was higher and denser above areas where rodents or insects had buried, at shallow depths, an assortment of grass, leaves, sticks, etc.

HORIZON	DESCRIPTION
Ap	0 to 8 inches, yellowish red (5YR 5/6) gravelly loam, yellowish red (5YR 4/6) moist; weak granular structure; hard, friable, slightly plastic; many small roots; strongly acid (pH 5.5); gradual boundary.
A11	8 to 15 inches, yellowish red (5YR 5/6) loam, red (2.5YR 4/6) moist; weak medium and fine blocky structure; hard, friable, slightly plastic; plentiful small roots; medium acid (pH 5.7); gradual boundary.
A12	15 to 21 inches, yellowish red (5YR 5/6) gravelly loam, red (2.5YR 4/6) moist; weak medium and fine blocky structure; hard, friable, slightly plastic; plentiful small roots; strongly acid (pH 5.4); boundary contains a very thin "bleached" layer just above the claypan that was not sampled. Abrupt wavy boundary.
B21t	21 to 29 inches, red (2.5YR 5/6) clay containing a slight amount of gravel; color of dry surface of aggregate is red (2.5YR 4/6), red (2.5YR 4/6) moist; strong coarse blocky structure; very compact; extremely hard, extremely firm, very plastic; very few roots; strongly acid (pH 5.2); distinct wavy boundary.
B22t	29 to 36 inches, yellowish red (5YR 5/6) gravelly clay loam, red (2.5YR 4/6) moist; weak blocky structure; compact; very hard, very firm, plastic; very few roots; strongly acid (pH 5.2); distinct wavy boundary.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Palexeralf, fine, montmorillonitic, thermic

SOIL San Ysidro fine sandy loam SOIL Nos. S64Calif-48-1 LOCATION Solano County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64528 - 64536

Depth (cm)	Horizon	1B1b Size class and particle diameter (mm) 3A1															
		Total			Sand					Silt					Coarse fragments 1A2a		
		Sand (2-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	(0.05-0.02)	Int. III (0.02-0.005)	Int. II (0.2-0.02)	2-0.1			> 2 Pct. of whole	2-19 / 19-76 Pct. of

SAN YSIDRO FINE SANDY LOAM
S64Calif-48-1

Location: Solano County, California. 3½ miles west and 1 mile south of Dixon on Forsgren Ranch, a pit located 300 yards south, 100 yards west of NE corner of NW1/4 sec. 29, T. 7 N., R. 1 E.
Date of Sampling: October 5, 1964.
Description by: Grant Kennedy, Sam Johnson, and Wells Andrews. Collectors: Reuben E. Nelson, Ben R. Brasher, Leland Bates, Sam Johnson, Grant Kennedy, and Wells Andrews.
Classification: Typic Palexeralf, fine, montmorillonitic, thermic.
Vegetation: Fallow. Climate: 22 inches of rainfall, 60° F. mean annual temperature. Parent Material: Alluvium from sedimentary rocks. Topgraphy: Level. Physiography: Old terrace. Elevation: 77 feet. Drainage: Moderately well drained; slow runoff; very slow permeability. No water table encountered within 114 inches of the surface. Soil Moisture: Dry 0-14 inches, slightly moist below 14 inches.
Remarks: pH determined by Hellige-Truog. Sample number of paired profile S64Calif-48-8. Diagnostic horizons are ochric and argillic. An A2 horizon rests on the B21t horizon and varies in thickness from zero to about 2 or 3 inches.

HORIZON	DESCRIPTION
Ap RSL No. 64528	0-7 inches (0-18 cm.), light brownish gray (10YR 6/2) fine sandy loam with few fine distinct mottles of brownish yellow (10YR 6/6), dark brown (10YR 3/3) moist; massive; hard (dry), friable (moist), nonsticky and slightly plastic (wet); abundant micro, plentiful fine and medium random roots; common very fine tubular and interstitial pores; clear smooth boundary.
A12 RSL No. 64529	7-14 inches (18-36 cm.), light brownish gray (10YR 6/2) fine sandy loam with few fine distinct mottles of brownish yellow (10YR 6/6), dark brown (10YR 3/3) moist; massive; hard (dry), friable (moist), nonsticky and slightly plastic (wet); abundant micro, plentiful fine and medium random roots; common micro, few very fine tubular pores; abrupt smooth boundary.
B21t RSL No. 64530	14-28 inches (36-71 cm.), dark yellowish brown (10YR 4/4) heavy clay loam, dark brown (7.5YR 4/4) moist; strong coarse prismatic structure; extremely hard (dry), very firm (moist), sticky and plastic (wet); few very fine and fine roots; common micro tubular pores; many stress cutans are on ped faces and in pores; iron and manganese concretions; gradual smooth boundary.
B22t RSL No. 64531	28-40 inches (71-100 cm.), yellowish brown (10YR 5/6) heavy sandy clay loam, dark yellowish brown (10YR 4/4) with dark brown (7.5YR 4/4) coatings moist; strong medium prismatic structure; extremely hard (dry), very firm (moist), sticky and plastic (wet); few very fine and fine exped roots; common micro tubular pores; many stress cutans are on ped faces and in pores; iron and manganese concretions; gradual smooth boundary.
B23t RSL No. 64532	40-54 inches (100-135 cm.), yellowish brown (10YR 5/4) light sandy clay loam, dark yellowish brown (10YR 4/4) moist; moderate medium prismatic structure; extremely hard (dry), very firm (moist), sticky and plastic (wet); few very fine exped roots; common micro tubular pores; many stress cutans are on ped faces and in pores; iron and manganese concretions; gradual wavy boundary.
IIB24t RSL No. 64533	54-68 inches (135-175 cm.), light yellowish brown (10YR 6/4) light clay loam, dark yellowish brown (10YR 4/4) with brown (7.5YR 4/4) coatings moist; strong medium prismatic structure; hard (dry), firm (moist), sticky and plastic (wet); few very fine exped roots; common micro tubular pores; continuous stress cutans are on ped faces and in pores; clear wavy boundary.
IIB3t RSL No. 64534	68-78 inches (175-200 cm.), yellowish brown (10YR 5/4) heavy sandy loam, dark yellowish brown (10YR 4/4) with brown (7.5YR 4/4) coatings moist; weak coarse prismatic structure; hard (dry), friable (moist), slightly sticky and plastic (wet); very few very fine roots; common micro and few very fine tubular pores; many stress cutans in pores; clear wavy boundary.

CONSECUTIVE PEDON NUMBER C 68110 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES SECCA TAXADJUNCT SOIL NO. S68 CALI. 29- 5 SAMPLE NO. 8 825-8 831 NEVADA COUNTY AREA

SOIL FAMILY MOLLIC PALEXERALF FINE,MIXED MESIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A	1A2A INT. .2- .02 II	COARSE FRAGMENTS PCT. OF WHOLE SOIL 20 75- 20-5 I	II	III 5-2 PCT. OF LT 75
			PERCENT OF LESS THAN 2MM																
8 825	0- 8	A1	37.1	49.5	13.4	6.9	7.7	4.7	8.9	8.9	24.2	25.3				38.1	43*	8	26	4
8 826	8- 23	A3	31.6	50.8	17.6	4.6	6.4	3.9	7.8	8.9	23.2	27.6				36.7	29*	7	10Z	N
8 827	0- 0	B21T	18.6	30.2	51.2	2.7	3.0	2.1	5.1	5.7	12.4	17.8				21.2	4A	--	4Z	N
8 828	58- 94	B22T	14.3	35.0	50.7	1.2	2.3	1.8	4.3	4.7	11.6	23.4				18.8	1A	--	TR	1
8 829	94-110	B3 T	16.1	40.8	43.1	1.0	2.7	1.9	4.7	5.8	10.2	30.6				18.8	4A	--	TR	4
8 830	110-125	C	35.5	47.0	17.5	4.5	8.8	4.7	9.4	8.1	12.6	34.4				25.9	40A	--	36	4
8 831	0- 8	A1															41A	--	33	8

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		PERCENT		PERCENT		PERCENT				PCT.NO.40		GRAMS PER CC			PERCENT			PERCENT	
8 825	0- 8	2.24	0.086			3.9	0.3			30	27		1.51	1.56	0.70		21.8	10.1	1.1	0.8
8 826	8- 23	1.02	0.048			4.3	0.3					1.50			0.81			9.7		
8 827	0- 0	0.32	0.022			4.2	0.3			57	37	1.45			0.98			18.1		
8 828	58- 94	0.21				3.0	0.2			56	26		1.44	1.82	0.99		29.1	21.3	8.1	8.0
8 829	94-110					0.9	0.1					1.38			0.98			22.7		
8 830	110-125		0.079			0.8	0.1			47	33		1.33	1.46	0.75		33.6	18.2	3.2	2.3
8 831	0- 8	1.58	0.090									1.51			0.71			8.6		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACIDITY 6H2A	KCL EXT. AL 6G1E	CEC SUM 5A3A	CEC NH4-OAC 5A6A	BASE SATURATION SUM+ ACIDITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CACL2 8C1E
		MEQ/100G									PERCENT					
8 825	0- 8	8.5	7.7	0.1	0.2	16.5	7.8		24.3	20.0	68		82		5.9	5.5
8 826	8- 23	6.5	7.2	0.1	0.1	13.9	6.4		20.3	17.7	68		79		5.9	5.3
8 827	0- 0	12.3	15.8	0.1	0.1	28.3			28.3	26.4	100+		100+		6.7	6.2
8 828	58- 94	11.1	22.5	0.2	0.1	33.9			33.9	32.3	100+		100+		6.9	6.7
8 829	94-110	12.3	26.8	0.3	0.1	39.5			39.5	37.6	100+		100+		7.1	6.7
8 830	110-125	11.6	24.0	0.3	0.1	36.0			36.0	36.2	100+		99		6.9	6.5
8 831	0- 8														6.0	5.6

SAMPLE NO	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 825	0- 8	KK 4	VR 3								KK KAOLINIT	1 TRACE
8 826	8- 23										VR VERMICLT	2 SMALL
8 827	0- 0											3 MODERATE
8 828	58- 94	KK 4	MC 3								MC MT/CHLOR	4 ABUNDANT
8 829	94-110										MT MONTMORT	5 DOMINANT
8 830	110-125	MT 4	KK 3									6 INDETER.
8 831	0- 8											

SAMPLE PREPARATION CODES
*=THE 75-250MM FRACTION IS INCLUDED.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.
A=STANDARD PREPARATION.

SECCA GRAVELLY LOAM (TAXADJUNCT) 1/
S68Calif-29-5

Location: Nevada County, California; 300 yards north of SW corner of SE¼ corner of sec. 4, T. 15 N., R. 8 E., 4 miles SW of Grass Valley, California; 300 yards north of intersection of McCourtney and Wolf Creek Mountain Lookout Roads—90 feet west of McCourtney Road. Date of Sampling: September 3, 1968.

Description by: G. M. Kennedy, T. M. Ryan, G. Staidl. Collectors: G. M. Kennedy, L. A. Brittan, G. Staidl, C. B. Goudey, T. M. Ryan.

Classification: Mollic Palexeralf, fine, mixed, mesic.

Vegetation: Manzanita, squirrel tail, igger pine, ceanothus. Not cultivated. Climate: Annual average precipitation 46 inches. Mean annual temperature 59°F. Parent Material: Amphibolite. Metabasic rock similar to diabase. These soils also occur in areas suggesting ultrabasic rock. Parent material at this site feels somewhat like talc. Possibly ultrabasic. Topography: Moderately sloping, plain slope, 8 percent near crest of slope. Southeast aspect. Elevation: 2,300 feet. Drainage: Well drained; runoff is medium; permeability is very slow. Soil Moisture: Profile dry to 9 inches, very slightly moist, 9 inches to bottom of profile.

Remarks: pH taken with Hellige kit. Sample S68Calif-29-7 and S68Calif-29-35 were also sampled as Secca soils. Thickness of clay varies from 14 to 34 inches. Reaction of Bt horizons varies from slightly acid to mildly alkaline. Reaction of surface seems uniform. Usually 10 to 25 percent of the surface is covered by stone fragments. Secca soils can occur under brush as well as grass cover. Eroded and open areas have gravel of 5 to 30 millimeters in diameter on surface. This layer is about 1/2 inch thick and covers 60 percent of the surface under the manzanita brush. The A1 horizon contains 3 to 5 percent gravel. There are 10 percent cobble-sized fragments in the A1 and A3 horizons. The A3 horizon contains 25 percent by weight fragments 3 to 10 inches in diameter. The A1 horizon (0-3"), B21t and B22t (9-23 and 23-37 combined) and C1 (43-50) were sampled for the California Division of Highways.

HORIZON	DESCRIPTION
O1	1/4 to 0 inches, manzanita leaves and bark. (0-8 cm.)
A1 RSL 68825	0 to 3 inches, brown (7.5YR 5/4) gravelly loam, dark reddish brown (5YR 3/3) moist; strong medium subangular blocky structure; slightly hard, friable, slightly sticky and slightly plastic; common very fine random roots; many very fine interstitial pores; medium acid (pH 6.0); abrupt smooth boundary. (8-23 cm.)
A3 RSL 68826	3 to 9 inches, brown (7.5YR 5/4) gravelly loam, dark reddish brown (5YR 3/4) moist; moderate medium subangular blocky structure; hard, friable, sticky, plastic; common very fine and fine oblique and common medium and coarse oblique roots; common very fine interstitial pores; medium acid (pH 5.8); clear wavy boundary. (23-58 cm.)
B21t RSL 68827	9 to 23 inches, brown (7.5YR 5/4) clay, brown and dark brown (7.5YR 4/4) moist; moderate coarse prismatic structure; hard, firm, sticky, very plastic; few very fine and fine oblique and few medium vertical roots; common very fine tubular pores; common moderately thick clay films on faces of peds and lining pores; few fine shot; mildly alkaline (pH 7.5); gradual wavy boundary. (58-94 cm.)
B22t RSL 68828	23 to 37 inches, brown (7.5YR 5/4) clay, brown and dark brown (7.5YR 4/4) moist; weak coarse prismatic structure; very hard, firm, sticky, very plastic; few very fine random and medium vertical roots; few very fine tubular pores; common moderately thick clay films on faces of peds and lining pores; pressure faces evident; mildly alkaline (pH 7.5); clear irregular boundary. (94-110 cm.)
B3t RSL 68829	37 to 43 inches, pale brown (10YR 6/3) clay, light olive brown (2.5Y 5/4) moist; massive; hard, firm, sticky, very plastic; few very fine tubular pores; continuous moderately thick reddish brown (5YR 4/4) clay films line pores; mildly alkaline (pH 7.5); clear wavy boundary. (110-125 cm.)
C RSL 68830	43 to 50 inches, white (5Y 8/1) gravelly loam, pink (5YR 7/4) moist; massive; very hard, very firm, slightly sticky, plastic; many moderately thick reddish yellow (5YR 6/6) clay films in fractures, red (2.5YR 4/8, 5/6) moist; neutral (pH 7.0).

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Millsap loam (taxadjunct) **LOCATION** San Luis Obispo County, California

SOIL FAMILY: Mollic Palexeralf, fin , montmorillonitic, thermic.

SOIL NOS. S55Cal. 40-4 **LAB. NOS.** 55465-470

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0- 5	A_{1p}	0.9	3.2	5.6	16.1	13.5	42.1	18.6	42.9	23.5	1	1
5-11	A_{12}	0.7	3.3	5.0	15.3	13.2	42.0	20.5	42.7	22.4	1	1
11-18	A_{13}	0.6	3.4	5.3	15.8	13.3	41.8	19.8	42.9	22.5	1	1
18-26	B_{21}	0.5	1.7	2.8	9.1	8.8	31.1	46.0	27.0	18.9	1	c
26-34	B_{22}	0.7	2.1	3.0	10.1	9.5	32.7	41.9	31.2	17.7	2	c
34-40	B_{23}	0.3	2.4	3.6	9.4	8.0	36.6	39.7	27.8	22.7	6	cl

pH 8C1b SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	6A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
5.8		6.5	1.95	0.154	12.7							11.4
6.0		6.8	1.34	0.109	12.3							11.8
6.1		7.2	0.74	0.070	10.6							10.8
7.1		8.7	0.62									22.5
7.8		9.1	0.45					<0.2				20.1
7.9		9.3	0.30					<0.2				18.8

MILLSAP LOAM (taxadjunct) 1/
S55Calif-40-4

Location: San Luis Obispo County, California. On Maino Ranch, about 3/8 mile northwest of San Bernardino Creek Road, 2.6 miles northeast of junction of Highway 101 and San Bernardino Creek Road. Date of Sampling: September 8, 1955.
Description by: J. M. Williams and Rudolph Ulrich. Collectors: W. M. Johnson, Rudolph Ulrich, Lewis Leifer, W. G. Harper, L. R. Wohletz, and J. M. Williams.
Classification: Mollic Palexeralf, fine, montmorillonitic, thermic.
Vegetation: Cultivated field, not irrigated. In barley stubble at time of sampling. Parent Material: Weathered, fractured, metamorphosed shale of the San Luis Obispo formation. Relief: Hilly, erosional, residual upland. Slope about 11 percent, slightly convex, toward the east. Elevation: About 300 feet. Drainage: Well drained. Runoff is moderately rapid; permeability is slow.

HORIZON		DESCRIPTION
A_{1p}	0-5"	Very dark grayish brown to grayish brown (10YR 5/2; 3/2 moist) loam, very hard, friable, very slightly plastic, nonsticky; essentially massive structure with weak coarse prismatic breakage due to extreme drying, probably weak very fine granular when moist, numerous fine roots and elongated porous openings due to fine root hairs, pH 6.0 (Truog). Abrupt, smooth lower boundary.
A_{12}	5-11"	Similarly colored loam, similar consistence, essentially massive with weak coarse prismatic breakage extending downward from surface above, numerous fine roots and elongated tubules due to root hair action. pH 6.2. Clear, smooth lower boundary.
A_{13}	11-18"	Very dark grayish brown (10YR 3/2, moist) loam; very hard; friable, very slightly plastic, nonsticky; essentially massive with very weak coarse prismatic breakage penetrating horizon from above, very numerous elongated fine pinhole openings or tubules due to fine root hair action (these openings have been described in numerous California reports as being vesicular, especially the lower surface or A_2 horizons of Planosolic profiles like Watsonville, Tierra, etc.), common fine root hairs. pH 6.8. Abrupt smooth lower boundary.
B_{21}	18-26"	Very dark grayish brown (2.5Y 3/2, moist) clay; extremely

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Positas gravelly loam **LOCATION** Alameda County, California

SOIL FAMILY: Mollic Palexeralf, fine, montmorillonitic, thermic

SOIL NOS. 59Calif-01-1 **LAB. NOS.** 59204 thru 59210

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)							3A1		2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	
0-8	A_p	3.1	4.1	3.6	9.7	10.3	53.2	16.0	47.7	21.9	21	sil
8-11	A_3	3.6	3.8	3.4	9.4	10.0	53.4	16.4	45.5	24.0	10	sil
11-20	B_{21}	1.9	2.3	1.8	5.3	5.8	29.9	53.0	25.3	13.8	14	c
20-29	B_{22}	2.5	2.9	2.9	10.9	3.7	34.4	42.7	27.7	16.5	4	c
29-39	B_{31}	2.9	4.3	4.6	15.4	11.0	34.9	26.9	38.2	17.3	8	1-cl
39-54	B_{32}	2.9	3.5	6.8	21.4	12.3	29.7	23.4	41.3	13.7	14	1
54-64+	C_1	13.8	10.8	9.6	18.4	8.2	18.1	21.1	27.4	8.9	40	scl
		Organic Matter			6B1h	8A1a						4B2

POSITAS GRAVELLY-LOAM
S59Calif-01-1

Location: Alameda County, California. 216 feet east of Greenville Road and 168 feet south of the northeast corner of Sec. 25, T. 3 S., R. 2 E. about 3 miles southeast of Livermore. Date of Sampling: May 4, 1959.
Description by: L. E. Welch. Collectors: R. G. Huff, L. E. Welch, L. H. Grant, and J. E. McClelland.
Classification: Mollic Palexeralf, fine, montmorillonitic, thermic.
Vegetation: Annual grasses including soft chess, filaree, and wild oats. Land Use: Not cultivated for about 5 years. Climate: About 15 inches rainfall. Parent Material: Old terrace deposits of mixed mineralogical origin from the Franciscan formation, probably of Livermore age. Relief: Gently sloping terrace, on a 4 percent southwest-facing slope. Elevation: About 600 feet. Drainage: Well drained, moderate runoff, very slow permeability.
Remarks: The soil is a maximal Noncalcic Brown with some stones and cobbles throughout the profile. The A3 horizon may be an A2, being slightly grayer (less than 1/2 chip) than the A1. Prisms have gray coatings on top and have slightly rounded edges, but have essentially flat upper faces. The A3 or A2 looks like degraded B2 in places with some clay films on small angular blocks. Few worm holes in surface.

HORIZON		DESCRIPTION
A_p	0-8"	Brown (10YR 5/3) gravelly loam, (7.5YR 3/2) when moist; essentially massive although surface few inches have weak horizontal cleavage; hard when dry, friable when moist; abundant fine roots; many fine pores; medium acid, pH 6.0; abrupt, smooth lower boundary.
A_3	8-11"	Similar to above in all respects except color values are nearly 1/2 chip higher; abrupt, smooth boundary.
B_{21}	11-20"	Dark reddish gray (5YR 4/2) clay, dark reddish brown (5YR 3/3) when moist; strong coarse prismatic structure; extremely hard when dry, extremely firm when moist, very plastic and sticky when wet; few fine roots along structure faces; few fine tubular pores; thick, continuous clay films on ped faces but colloid nearly fills pores; common slickensides; slightly acid, pH 6.5; gradual, smooth lower boundary.
B_{22}	20-29"	Reddish brown (5YR 4/4) dry and moist, clay; strong coarse prismatic structure; similar to B_{21} except has a little less clay; moderately alkaline, pH 8.0; abrupt, smooth lower boundary.
B_{31}	29-39"	Brown (7.5YR 5/5) heavy clay loam with many yellowish red (5YR 5/5) coatings and common fine (1 to 2 mm.) black stains, brown (7.5YR 4/4) and yellowish red (5YR 4/6) when moist; strong medium angular blocky structure; very hard when dry, firm when moist, plastic and sticky when wet; few fine roots; few fine tubular pores; moderately thick continuous clay films on peds and in pores; moderately alkaline, pH 8.0; very weakly calcareous; gradual, smooth lower boundary.
B_{32}	39-54"	Light yellowish brown (10YR 6/5) clay loam with few yellowish red (5YR 4/6) coatings and common fine black stains, brown (10YR 5/4) and yellowish red (5YR 4/6) when moist; strong medium angular blocky structure; very hard when dry, firm when moist, sticky and slightly plastic when wet; few fine roots; few fine tubular pores; moderately thick, continuous clay films on peds and in pores; moderately alkaline, pH 8.0; very weakly calcareous; gradual, smooth lower boundary.
C_1	54-64"+	Light yellowish brown (10YR 6/5) gravelly sandy loam with a few yellowish red (5YR 4/6) coatings, yellowish-brown (10YR 5/4) with yellowish red (5YR 4/6) coatings when moist; massive; slightly hard when dry, friable when moist; about 1 percent by volume of rocks over 3 feet in diameter and about 15 percent 1 to 3 inches in diameter, moderately alkaline, pH 8.0+.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Positas loam **LOCATION** Alameda County, California

SOIL FAMILY: Mollic Palexeralf, fine, montmorillonitic, thermic

SOIL NOS. 59Calif-01-2 **LAB. NOS.** 59211 thru 59217

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-8	A_{1p}	6.4	7.2	6.1	12.7	10.3	43.4	13.9	41.4	20.0	16	l
8-11	A_{12}	4.6	6.0	5.4	12.0	11.0	44.3	16.7	42.0	20.5	7	l
11-18	B_{21}	2.6	3.5	3.0	7.1	6.2	27.7	49.9	24.6	13.6	21	c
18-24	B_{22}	6.1	5.0	3.1	7.3	7.0	29.5	42.0	26.3	14.7	12	c
24-35	B_{31}	10.2	11.8	5.3	9.7	8.1	33.9	21.0	26.7	21.0	38	l
35-50	B_{32}	23.1	15.7	3.5	4.1	3.0	23.1	27.5	11.0	17.3	71	scl
50+	C_1	13.1	12.0	7.6	12.1	6.3	21.5	27.4	21.8	12.4	71	scl

pH 8C1b Saturated Paste 1:1	8C1a 1:10	Organic Matter 6A1a Organic Carbon Pct.	6B1a Nitrogen Pct.	C/N	6E1b $CaCO_3$ Equivalent Pct.	8A1a Elec. Conduc. mmhos/cm. at 25° C.	4B2 Soil water 15 atmos. Pct.
6.2	6.6	1.15	0.092	12.5		0.4	6.3
6.0	6.6	0.50	0.051	9.8		0.3	6.9
5.7	7.0	0.56	0.063	8.9		0.2	21.5
6.5	7.6	0.37	0.047	7.9		0.3	18.2
7.5	8.5	0.08				0.6	13.1
7.9	8.9	0.03				0.6	17.3
7.9	8.9	0.03				0.8	13.3

5A2a CATION EXCHANGE CAPACITY (+ Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a Ca	6O1a Mg	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
16.4	7.2	5.3	4.6	0.2	0.3	74	1.0	<0.1				28.5
17.3	6.6	6.0	4.6	0.3	0.2	74	1.4	<0.1				25.8
39.3	18.7	12.8	7.2	1.0	0.5	82	1.5	<0.1				57.4
38.4	20.2	12.1	4.4	1.2	0.4	89	2.4	<0.1				52.1
41.8	26.8	10.1	3.1	1.9	0.4	93	4.7	<0.1				41.2
44.3	26.8	11.4	2.3	3.6	0.6	95	5.1	<0.1				47.4
32.8	14.4	11.9	2.1	3.3	0.4	94	6.5	<0.1				41.3

1/ Stones and gravel >1" diameter discarded at time of sampling as follows:

Horizon B_{31} - 15%
Horizon B_{32} - 15%
Horizon C_1 - 20%

POSITAS LOAM
S59Calif-01-2

Location: Land grant - 169° east of true north of BM 448, 122° east of true north of BM 478, about 4 miles south of the center of Livermore in the Crowhare Ranch in Alameda County, California. Date of Sampling: May 4, 1959.
Description by: L. E. Welch. Collectors: R. C. Huff, L. E. Welch, L. H. Grant, and J. E. McClelland.
Classification: Mollic Palexeralf, fine, montmorillonitic, thermic.
Vegetation: Annual grasses and forbs, soft chess, filaree, wild oats, and clover. Land Use: Cultivated pasture. Climate: 15 inches rainfall. Parent Material: Old terrace deposits of the Livermore formation consisting of sediments of mixed mineralogical origin from the Franciscan formation. Relief: Gently sloping old terrace, slope 5 percent to the southwest. Elevation: 640 feet. Drainage: Well drained, moderate runoff, very slowly permeable.
Remarks: This is a maximal Noncalcic Brown soil with considerable gravel, cobbles and stones below the B22, few above. Very little indication of worm activity.

HORIZON		DESCRIPTION
A_{1p}	0-8"	Brown (10YR 5/3) heavy loam, dark brown (10YR 3/3); massive; very hard when dry, friable when moist; abundant fine roots; many fine tubular pores; medium acid, pH 6.0; clear, smooth lower boundary.
A_{12}	8-11"	Similar to above; abrupt, smooth lower boundary.
[redacted]		
		moist; strong coarse prismatic structure, nearly massive when moist; extremely hard when dry, extremely firm when moist, very plastic and sticky when wet; few fine roots along prism faces, very few in peds; nonporous except for cracks along ped faces; thick, continuous clay films on ped faces, colloid nearly fills all pores; slightly acid, pH 6.2; clear, smooth lower boundary.
B_{22}	18-24"	Brown (7.5YR 4/3) clay, brown (7.5YR 4/4) when moist; otherwise similar to B_{21} except pH 6.5; abrupt, irregular lower boundary.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Natric Palexeralf, fine, montmorillonitic, thermic.

SOIL Bonsall sandy loam SOIL Nos. S64Calif-37-3 LOCATION San Diego County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 6430 - 6439

Depth (In.)	Horizon	1B1b Total: Sand (2–0.05)	Total: Silt (0.05–0.002)	Total: Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)	* Clay <.002	** Clay <.002	1/ Coarse fragments 1A2a ≥ 2 Pct.	2 – 19 (Pct. of < 76 mm)	19 – 76 (Pct. of < 76 mm)
		Size class and particle diameter (mm)										3A1						
		Pct. of < 2 mm																
0- 6	A1	68.0	23.3	8.7	11.0	15.0	9.8	19.4	12.8	13.3	10.0	36.2	55.2	8.4	8.1	4	4	0
6-10	A2	62.9	24.7	12.4	9.3	13.6	9.0	18.6	12.4	13.9	10.8	36.2	50.5	11.0	10.7	3	3	0
10-14	B1	42.4	18.7	38.9	6.0	8.3	5.9	13.0	9.2	10.4	8.3	27.0	33.2	40.5	38.9	4	4	0
14-27	B21t	40.6	21.1	38.3	2.8	6.1	5.1	15.0	11.6	11.9	9.2	32.5	29.0	36.3	35.7	2	2	0
27-38	B22t	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]

BONSALL SANDY LOAM
S64Calif-37-3

Location: San Diego County, California. A soil profile pit located 2-1/4 miles south, 3/4 mile east of Fallbrook in the NE1/4 of the SW1/4 of section 31, T. 9 S., R. 3 W.
Date of Sampling: February 18, 1964.
Description By: David C. Estrada and Gerald Kester. Collectors: Klaus Flach, George Borst, Gerald Kester, David C. Estrada and Leo Klameth.
Classification: Natric Palexeralf, fine, montmorillonitic, thermic.
Vegetation: Annual grasses and annual weeds. Climate: This soil occurs under 15 inches of rainfall. The mean annual temperature is 60° F., the mean January temperature 54° F., and the mean July temperature 70° F. Parent Material: Tonalite grus. Topography: South-facing, smooth 8 percent slopes in lower backslope positions on rolling upland. Elevation: 545

Remarks: The texture of the C1 horizon is finer than the IIC2 or IIC3 horizons. This may reflect variation in grain size of the grus; differences in weathering of the horizons, or soil creep. This pedon could be classified as a Mollic Palexeralf.

HORIZON	DESCRIPTION
A1 RSL No. 6430	0 to 6 inches, brown (10YR 5/3) sandy loam, dark brown (10YR 3/3) moist; moderate fine to medium crumb structure; hard, friable, slightly sticky and nonplastic; abundant very fine and fine, and few medium roots; many very fine and fine tubular pores; slightly acid (pH 6.2); abrupt smooth boundary.
A2 RSL No. 6431	6 to 10 inches, same as above except the horizon is massive, roots are plentiful, and pores common; abrupt smooth boundary.
B1 RSL No. 6432	10 to 14 inches, brown (7.5YR 5/4) heavy clay loam, dark brown (7.5YR 4/4) moist; moderate medium angular blocky structure; very hard, firm, sticky, and plastic; plentiful very fine and fine, and few medium roots; few fine and medium tubular pores; slightly acid (pH 6.5); clear smooth boundary.
B21t * RSL No. 6433	14 to 27 inches, yellowish brown (10YR 5/4) light clay, dark yellowish brown (10YR 4/4) moist; strong coarse prismatic structure; extremely hard, very firm, sticky and plastic; few very fine, fine and medium roots; few fine and medium

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Sites gravelly loam (taxadjunct) **LOCATION** Amador County, California

SOIL FAMILY: Ultic Palexeralf, fine, kaolinitic, mesic

SOIL NOS. 59Calif-3-3 **LAB. NOS.** 59278 - 59285

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	
1/2-0	A_{oo}	Not sampled										
0-8	A_{11p}	2.5	3.4	2.2	10.1	14.6	42.8	24.4	41.2	24.0	28	l
8-15	A_{12}	1.4	2.1	1.7	9.8	15.3	42.0	27.7	42.2	22.8	28	cl- l
15-27	B_{11}	1.3	2.1	1.4	8.6	14.5	43.4	28.7	43.7	21.1	27	cl
27-35	B_{12}	1.5	2.2	1.4	8.1	13.8	40.7	32.3	40.2	20.8	35	cl
35-45	B_{21}	1.4	1.8	1.0	6.8	12.5	34.8	41.7	36.4	16.5	32	c
45-55	B_{22}	1.5	1.4	0.9	7.0	12.3	31.6	45.3	35.0	14.6	45	c
55-67	B_3	1.2	1.6	1.1	7.4	13.1	34.5	41.1	38.8	14.9	45	c
67-72	C_1	1.8	2.6	1.5	10.0	20.5	45.0	18.6	57.2	16.3	40	l
72+	D_r	Not sampled										

SITES GRAVELLY LOAM (taxadjunct) 1/
S59Calif-3-3

Collectors: Grant Kennedy, Reuben Nelson, Bob Sketchley, John Rogers.
Date of Sampling: June 17, 1959. Description by: Grant Kennedy.
Location: Amador County, California in the NE corner of Section 9, T6N, R12E, 100 yards north of the Ranger's residence, about 6 miles NE of Jackson.
Elevation: 2,800 feet.
Relief: Moderately steep uplands on a plain slope of 18 percent to the NW.
Drainage: Well drained; slow runoff; moderate permeability to 3 feet, moderately slow below.
Parent material; Metasedimentary rocks of the Calaveras formation.
Biological Activity: Krotovinas and worm holes common above 45 inches.
Climate: Mean annual rainfall about 45".
Vegetation: An abandoned black walnut orchard now with bracken fern, ponderosa pine, black oak, incense cedar, and sugar pine.
Remarks: A medial developed Reddish-Brown Lateritic soil. There are a few gravels above the B_{22} and considerable below. Bureau of Public Roads samples were obtained on samples 1, 3, and 6.
Classification: Ultic Palexeralf, fine, kaolinitic, mesic.

HORIZON		DESCRIPTION
A_{00}	1/2-0"	Pine needles. Not sampled.
A_{11m}	0-8"	Brown (7.5YR 5/4) gravelly loam, dark reddish brown (5YR 3/4) [illegible] slightly hard when dry, friable when moist, nonsticky and slightly plastic when wet; plentiful very fine, fine and medium roots, few coarse; many very fine and fine tubular and interstitial pores; slightly acid, pH 6.3; abrupt, smooth lower boundary.
A_{12}	8-15"	Reddish brown (5YR 4/5) heavy gravelly loam, dark reddish brown (5YR 3/4) when moist: moderate very fine and fine [illegible] pores, and pH as above except a few medium pores are present; gradual, smooth lower boundary.
B_{11}	15-27"	Reddish brown (5YR 4/5) gravelly clay loam, dark red (2.5YR 3/6) when moist; structure, consistence, roots, and pores as in horizon above except the soil is slightly sticky when wet; few, thin discontinuous clay films in pores; slightly acid, pH 6.4; gradual, smooth lower boundary.
B_{12}	27-35"	Similar to B_{11} except pH is 6.3; gradual, smooth lower boundary.
B_{21}	35-45"	Yellowish red (5YR 4/6) gravelly clay; dark red (2.5YR 3/7) when moist; weak very fine subangular blocky structure when crushed; slightly hard when dry, firm when moist, slightly sticky and slightly plastic when wet; few very fine and fine, very few medium roots; many fine and very fine, few medium tubular and interstitial pores; moderately thick, continuous clay films in pores and bridges; slightly acid; pH 6.2; gradual, smooth lower boundary.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Ultic Palexeralf, fine, kaolinitic, thermic

SOIL Red Bluff gravelly loam SOIL Nos. SL557-6 LOCATION Tehama County, California

SOIL SURVEY LABORATORY _____ LAB. Nos. _____

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	3A1 (2-0.1)			Coarse fragments > 2 Pct.	2-19 (Pct. of < 76mm)	19-76 (Pct. of < 76mm)
		Pct. of < 2 mm																
0-6	Ap	45.2	32.0	22.8	8.9	8.2	4.5	9.4	14.2	15.0	17.0		31.0					
6-10½	A11	44.1	30.7	25.2	7.7	8.9	4.8	9.0	13.7	13.3	17.4		30.4					
10½-19½	A12	43.3	31.3	25.4	5.5	7.7	4.9	10.1	15.1	14.1	17.2		28.2					
19½-27	B21tsi	36.1	29.7	34.2	6.9	6.2	3.7	7.4	11.9	13.4	16.3		24.2					
27-37	B22tsi	35.0	28.7	36.3	6.4	6.1	3.5	6.9	12.1	13.1	15.6		22.9					
37-45	B23tsi	34.5	28.8	36.7	6.7	5.8	3.4	6.9	11.7	12.9	15.9		22.8					
55-64	C1	39.6	28.4	32.0	8.7	7.1	3.6	7.8	12.4	12.9	15.5		27.2					
64-72	C2	43.3	27.3	29.4	11.5	7.0	3.9	8.3	12.6	12.4	14.9		30.7					

Depth (In.)	6A2a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	Pct			pH		8C1a (1:1)
0-6	0.61	0.066																5.0
6-10½	0.22	0.039																4.9
10½-19½	0.18	0.034																5.0
19½-27	0.11	0.034																5.0
27-37	0.09	0.029																5.0
37-45	0.07	0.027																5.2
55-64	0.06	0.019																5.2
64-72	0.04	0.020																5.4

Depth (In.)	Extractable bases 5B1a: Ca 6N2b	Mg 6O2b	Na 2/	K 1/	Sum of bases (meq/100 g)	6H2a Ext. Acidity	5A3a CEC Sum									Base saturation 5C3 Sum Pct.	Pct.
0-6	1.8	0.6	0.01	0.68	3.1	5.1	8.2									38	
6-10½	1.6	0.4	0.02	0.76	2.8	4.7	7.5									37	
10½-19½	1.6	1.0	0.02	0.72	3.3	4.3	7.6									44	
19½-27	2.3	1.7	0.03	0.62	4.6	4.2	8.8									53	
27-37	2.6	1.9	0.00	0.41	4.9	4.0	8.9									55	
37-45	2.9	2.1	0.02	0.31	5.3	3.5	8.8									61	
55-64	2.9	1.9	0.02	0.30	5.1	2.5	7.6									67	
64-72	2.9	2.0	0.07	0.16	5.1	2.4	7.5									68	

Depth (In.)

1/ Precipitated as potassium sodium cobaltinitrite and titrated as potassium permanganate

2/ Gravimetrically as sodium uranyl magnesium acetate

RED BLUFF GRAVELLY LOAM
S4557-6

Location: Tehama County, California. SW¼ NE¼ Sec. 36, T27N, R4W, MDB&M, 0.05 mile east of the center of Sec. 36. Date of Sampling: April 2, 1945.
Description by: R. C. Roberts and R. A. Gardner. Collectors: R. C. Roberts, F. O. Youngs, R. A. Gardner, L. T. Alexander, H. Jenny and R. E. Storie.
Classification: Ultic Palexeralf, fine, kaolinitic, thermic.
Vegetation: Sparse short annual plants dominated by alfilaria. Climate: Average annual precipitation is about 24 inches; average annual temperature about 62°F., January 45°F., July 82°F. Parent Material: Mixed alluvium. Topography: On nearly level to gently sloping very old terraces of mixed valley fill. Elevation: 340 feet. Drainage: Well drained, moderately slow permeability, slow runoff. Soil Moisture: Moist.
Remarks: One of the most noticeable characteristics of the soil as observed in the sampling pit, as well as in roadside cuts, is the seemingly high manganese oxide content between depths of about 12 and 20 inches. A high percentage of the manganese oxide appears to be contained in soft and medium hard small concretions. In road banks this zone is as conspicuous as the lime concentration zone of the arid regions. Another distinguishing characteristic of this profile is that the material below about 20 inches is brittle. It has the appearance of a degraded semiconsolidated hardpan.

Observations made in a deep ravine indicate that there are layers at a depth of about 9 or 10 feet that are more compact. These appear to intercept percolating water and force it to flow laterally to outlets in ravines or to hillside seeps. The material above the compact layer is acid and distinctly mottled.

HORIZON	DESCRIPTION
Ap	0 to 6 inches, reddish brown (5YR 4/4) gravelly loam, dark reddish brown (2.5YR 3/4) moist; strong granular structure; slightly hard, friable, slightly plastic; abundant fine roots; very strongly acid (pH 5.0); abrupt smooth boundary.
A11	6 to 10.5 inches, reddish brown (5YR 4/4) gravelly heavy loam; color dry and moist, structure and consistence similar to layer above; abundant fine roots; very strongly acid (pH 4.9); distinct wavy boundary.
A12	10.5 to 19.5 inches, reddish brown (5YR 4/4) heavy loam containing a slight amount of gravel, dark reddish brown (2.5YR 3/4) moist; strong granular structure; slightly hard, friable, slightly plastic; plentiful fine roots; very strongly acid (pH 5.0); abrupt irregular boundary.
B21tsi	19.5 to 27 inches, red (2.5YR 4/6) clay loam containing a slight amount of gravel, weak red (10R 4/4) moist; strong blocky structure; apparently weakly cemented with iron oxide; hard, firm and brittle, plastic; coatings of manganese oxide, particularly in lower part of layer, more in evidence than in any other layer; few fine roots; very strongly acid (pH 5.0); gradual boundary.
B22tsi	27 to 37 inches, red (2.5YR 4/6) clay loam containing a slight amount of gravel; color dry and moist, structure and consistence similar to layer above; apparently weakly cemented with iron oxide, as layer above, but contains somewhat less coatings of manganese oxide; few fine roots; very strongly acid (pH 5.0); diffuse boundary.
B23tsi	37 to 45 inches, red (2.5YR 4/6) gravelly clay loam, weak red (10R 4/4) moist; strong blocky structure; hard, firm and brittle, plastic; apparently weakly cemented but appers to be slightly less cemented than horizon above; contains manganese oxide; few fine roots; strongly acid (pH 5.2); distinct boundary.
C1	55 to 64 inches, red (2.5YR 4/6) gravelly clay loam, color dry and moist similar to layer above; weak blocky structure; compact; consistence similar to layer above; manganese oxide present; few fine roots; no distinct observable difference between this layer and underlying layer.
C2	64 to 72 inches, red gravelly clay loam (near loam); color dry and moist, structure and consistence similar to layer above; few roots present; strongly acid (pH 5.4).

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Ultic Palexeralf, fine, kaolinitic, thermic

SOIL Red Bluff loam SOIL Nos. S4557-8 LOCATION Shasta County, California

SOIL SURVEY LABORATORY _____ LAB. Nos. _____

Depth (In.)	Horizon	1B1b Total: Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Sand: Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt: 0.05–0.02	Int. III (0.02–0.002)	Int. II (0.2–0.02)	3A1 (2–0.1)			Coarse fragments > 2 Pct.	2 – 19	19 – 76
		Pct. of < 2 mm															Pct of < 76mm	
0–7	A11	40.7	34.5	24.8	10.4	7.5	4.8	8.9	9.1	11.1	23.4		31.6					
7–14	A12	34.2	32.5	33.3	7.0	6.3	4.1	8.2	8.6	10.4	22.1		25.6					
14–24	B21t	26.3	24.9	48.8	6.1	4.9	3.2	5.8	6.3	7.3	17.6		20.0					
24–33	IIB22t	20.5	23.0	56.5	4.8	3.8	2.3	4.6	5.0	5.5	17.5		15.5					
33–41	IIB23t	19.9	22.9	57.2	7.0	3.7	1.7	3.6	3.9	4.0	18.9		16.0					
41–51	IIB24t	24.6	23.8	51.6	9.5	4.9	2.2	3.7	4.3	4.4	19.4		20.3					

Depth (In.)	6A2a Organic carbon Pct.	6B1a Nitrogen Pct	C/N		Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	Pct			pH		8C1a (1:1)
0–7	1.25	0.077																6.1
7–14	0.62	0.048																5.5
14–24	0.23	0.035																4.9
24–33	0.17	0.037																4.9
33–41	0.15	0.029																4.8
41–51	0.16	0.030																4.4

Depth (In.)	Extractable bases 5B1a: Ca 6N2b	Mg 6O2b	Na 2/	K 1/	Sum of bases	6H2a Ext. Acidity	5A3a CEC Sum									Base saturation 5C3 Sum Pct.	Pct.
	meq/100 g																
0–7	6.2	0.8	0.02	0.44	7.5	4.4	11.9									63	
7–14	3.6	1.3	<0.01	0.52	5.4	4.9	10.3									53	
14–24	2.8	2.0	tr.	0.43	5.2	6.0	11.2									47	
24–33	2.8	2.6	0.05	0.55	6.0	6.4	12.4									48	
33–41	2.3	2.9	0.02	0.48	5.7	6.2	11.9									48	
41–51	0.6	2.8	0.04	0.53	4.0	6.6	10.6									37	

Depth (In.)					

1/ Precipitated as potassium sodium cobaltinitrite and titrated as potassium permanganate

2/ Gravimetrically as sodium uranyl magnesium acetate

* RED BLUFF LOAM
84557-8

Location: Shasta County, California, 1.35 miles E. of Olinda. Date of Sampling: April 4, 1945.
Description by: R. C. Roberts and R. A. Gardner. Collectors: R. C. Roberts, F. O. Youngs, R. A. Gardner, L. T. Alexander, H. Jenny and R. E. Storie.
Classification: Ultic Palexeralf, fine, kaolinitic, thermic.
Vegetation: Ceanothus, scrub oak, manzanita, digger pine, nitgrass, wild oats, alfilaria and associated herbaceous plants. Climate: Average annual precipitation is 33 inches. Average annual temperature is 62°F., January 45°F., and July 82°F. Parent Material: Mixed. Topography: Undulating to nearly flat tree- and grass-covered tops of old terraces of mixed valley-fill material. Elevation: 500 feet. Drainage: Well drained, moderate permeability, slow runoff. Soil Moisture: Moist to 41 inches, wet to 51 inches, water in bottom of pit.
Remarks: *This soil would be correlated to Red Bluff at the present time

annual precipitation of about 33 inches. The noticeable features of the profile are: (1) The rather intense reddish color of all layers, particularly below the surface; (2) the fairly good granular structure of the soil to a depth of about 2 feet; (3) the subsoil not being so distinct nor so compact a claypan as in the other Redding profiles; and (4) the material immediately below the claypan not being a distinct hardpan. There is not an abrupt change from surface to subsoil, but this soil appears to be quite lateritic judging from the granular structure, more rapid permeability, increasing acidity with depth, and the probable high iron, manganese and aluminum content. A moist clod of this soil has a tendency toward the characteristic soapy feel of a lateritic soil and may indicate that the soil is fairly high in aluminum. The lower part of the subsoil is more or less reticulately mottled, evidence of imperfect internal drainage or at least of uneven water per-

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Abruptic Xerollic Durargid, fine, montmorillonitic, mesic

SOIL Lovejoy loam SOIL Nos. S61Calif-32-21 LOCATION Sierra Valley, Plumas County, Calif.

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 61736 - 61741

Depth (in.)	Horizon	1B1b Total Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Sand: Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt: 0.05–0.02	Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.2)			Coarse fragments 2A2 > 2	2 – 19	19 – 76
		Pct. of < 2 mm														Pct.	Pct. of < 76 mm	
0–2½	A1	30.4	46.3	23.3	0.4	1.1	1.7	10.7	16.5	19.6	26.7	44.2	5.8			0		
2½–10	A21	42.1	30.3	27.6	0.1	0.6	1.7	18.3	21.4	14.2	16.1	49.6	6.7			0		
10–16	A22	52.2	30.6	17.2	1.7	4.1	5.0	21.8	19.6	14.5	16.1	49.0	17.7			0		
16–21	B2t	32.7	27.2	40.1	0.5	3.1	3.3	13.3	12.5	11.9	15.3	33.7	10.9			0		
21–41	C1m	52.0	35.5	12.5	2.5	6.4	6.1	17.5	19.5	14.7	20.8	46.2	20.5			0		
41–53+	C2	48.6	28.9	22.5	0.3	3.1	4.6	20.2	20.4	13.3	15.6	48.6	13.3			0		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		6E1b Carbonate as $CaCO_3$ Pct.	6C2a Ext. Iron as Fe Pct.	Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	4B2 15 bar Pct.			pH 8C1b Sat. Paste	8C1a 1:10
0–2½	1.20	0.097	12.4		-	0.6							10.4			5.8	6.5
2½–10	0.57	0.049	11.6		-	0.6							11.2			5.7	6.7
10–16	0.16				-	0.7							9.1			6.1	6.9
16–21	0.13				-	0.7							18.4			5.8	7.0
21–41	0.04				1	0.4							17.3			7.5	8.6
41–53+	0.03				-	0.4							17.4			7.1	7.9

Depth (in.)	Extractable bases 5B1a: 6N2c Ca	6O2c Mg	6P2a Na	6Q2a K	6H2a Ext. Acidity	Cat. Exch. Cap. 5A2a NaOAc	5B1b Exch. Na	Water extract from saturated paste 8A1: 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g							meq/liter									mmho/cm
0–2½	10.8	4.6	0.3	0.3	5.1	22.8	0.3	0.7	0.6	1.0	0.1		-	1.2	0.3		0.22
2½–10	12.6	5.2	0.6	0.1	4.5	25.0	0.5	0.4	0.4	1.3	-		-	1.2	0.2		0.13
10–16	8.6	3.8	0.7	0.1	3.0	18.5	0.6	0.4	0.3	1.9	-		-	0.9	0.5		0.26
16–21	18.1	8.6	1.8	0.2	4.5	33.5	1.6	0.9	0.9	3.2	-		-	0.9	1.5		0.37
21–41	27.9	16.0	2.4	0.1	0.7	37.5	2.2	1.6	1.2	5.3	-		0.3	2.8	3.7		0.85
41–53+	19.5	7.5	1.4	0.3	1.5	30.6	1.3	0.3	0.5	1.9	-		-	1.2	0.6		0.27

Depth (in.)	8A Water at Saturation Pct.	5C3 Base Sat. (Sum) Pct	Mineral Content (<0.002 mm.): Montmorillonite	Vermiculite	Mica	Kaolinite	Kaolinite % DTA
			X-ray				
0–2½	34.5	76	xx	t	-	x	12
2½–10	34.4	80	xxx	t	-	x	15
10–16	24.9	81	xxxx	t	-	x	15
16–21	46.2	86	xxxx	t	-	x	18
21–41	45.5	98	xx	t	-	t	5
41–53+	42.4	95					

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

LOVEJOY LOAM

S61Calif-32-21

Location: Sierra Valley, Plumas County, California. 0.6 mile north, and 0.1 mile west of the southeast corner of section 14, T. 22 N., R. 14 E. 100 feet north of farm road. Approximately four miles south of Beckwourth. Date of Sampling: September 21, 1961.
Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, James McLaughlin, Eugene Begg.
Vegetation: Low sage is predominant with some squirrel tail. Climate: Estimated annual precipitation 15 inches. Nearest weather station (Portola): Mean annual temperature 45.1° F.; January average 28.7° F.; July average 62.6° F.; mean annual precipitation 20.62 inches. Parent Material: Alluvium from mixed sources. Relief: Hummocky low terrace with a slope of less than 1 percent. Elevation: 4892 feet. Drainage: Moderately well drained; runoff is slow; permeability is moderately slow in the upper part of the profile and slow below. Soil Moisture: Slightly moist below 16 inches.
Remarks: This is a Planosol, with incipient Chestnut development in the A21. The surface has hummocks 3 to 4 inches high and 7 to 24 inches in diameter and the hummocks are matted with roots. Three inches below the bottom of the B2t in the top of the Clm there is a thin mat of roots spreading out over the underlying pan-like material. The A1 horizon (0-2½ inches), the B2t horizon (16-21 inches), and the C2 horizon (41-53 inches) were sampled for the Bureau of Public Roads and the California State Division of Highways. Texture of Clm cannot be determined satisfactorily in field because of difficulty of grinding up and wetting sample properly. **Classification: Abruptic Xerollic Durargid, fine, montmorillonitic, mesic**

HORIZON	DESCRIPTION
A1 RSL No. 61736	0 to 2½ inches, gray (10YR 6/1) loam (near clay loam), dark grayish brown (10YR 4/2) when moist; strong very fine platy structure; soft, very friable, slightly sticky and slightly plastic; abundant very fine and few fine roots; many very fine, and common fine interstitial, and common fine tubular pores; strongly acid (pH 5.3); clear smooth boundary.
A21 RSL No. 61737	2½ to 10 inches, light brownish gray (10YR 6/2) clay loam, dark grayish brown (10YR 4/2) when moist; moderate medium prismatic and moderate fine platy structure; hard, friable, slightly sticky and plastic; plentiful very fine and few fine to medium roots; many very fine mostly horizontal tubular pores and common very fine vertical tubular pores and common very fine interstitial pores; common thin clay films in pores; strongly acid (pH 5.5); clear, smooth boundary.
A22 RSL No. 61738	10 to 16 inches, light gray (10YR 7/1) loam (near sandy clay loam), dark grayish brown (10YR 4/2) when moist; moderate coarse prismatic structure; hard, friable, slightly sticky and slightly plastic; few very fine to medium mostly exped roots; common very fine tubular and interstitial pores; common thin clay films in pores; few black shot; medium acid (pH 5.9); noncalcareous; abrupt, smooth boundary.
B2t RSL No. 61739	16 to 21 inches, pale brown (10YR 6/3) clay, dark grayish brown (10YR 4/2) when moist; strong coarse columnar structure; extremely hard, extremely firm, very sticky and very plastic; few very fine to medium exped roots; few very fine tubular pores; continuous moderately thick clay films in pores and on ped faces; few black shot; medium acid (pH 6.0); abrupt, smooth boundary.
Clm RSL No. 61740	21 to 41 inches, light gray (2.5Y 7/2), olive brown (2.5Y 4/4) and pale yellow (2.5Y 7/4) when moist; massive, breaking to strong medium angular blocky structure; extremely hard, extremely firm, nonsticky and nonplastic; no roots except for a thin mat 3 inches below the B2; moderately alkaline (pH 8.0); strongly effervescent, lime is in distinct horizontal seams and the matrix is noncalcareous; pale yellow material is predominantly in horizontal seams that are noncalcareous in the upper part of the horizon; horizon does not slake in water; clear and smooth boundary.
C2 RSL No. 61741	41 to 53 inches +, pale yellow (2.5Y 7/4) clay loam (near loam), light olive brown (2.5Y 5/4) when moist; massive; very hard, firm, sticky and plastic; common very fine, few fine tubular and few very fine interstitial pores; common moderately thick clay films in pores; moderately alkaline (pH 7.9); noneffervescent.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Xerollic Durargid, fine-loamy, mixed, mesic

SOIL Lovejoy loam (taxadjunct) SOIL Nos. S61Calif-46-6 LOCATION Sierra Valley, Sierra County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 61742 - 61747

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.2)			Coarse fragments 2A2 > 2 Pct.	2 - 19 (Pct. of < 76 mm)	19-76 (Pct. of < 76 mm)
		Pct. of < 2 mm																
0-5	A1	32.6	39.4	28.0	1.9	5.6	4.6	10.3	10.2	14.0	25.4	30.7	15.9			1		
5-10	A21	55.1	26.8	18.1	3.1	14.9	10.0	19.4	7.7	13.3	13.5	31.8	36.6			1		
10-15	A22	68.5	17.8	13.7	7.2	20.9	14.3	19.5	6.6	7.0	10.8	22.9	52.6			5		
15-23	B2t	41.0	28.3	30.7	2.5	9.2	6.2	14.5	8.6	11.6	16.7	29.0	23.6			0		
23-42	C1m	36.8	44.4	18.8	0.1	1.2	1.7	15.1	18.7	17.8	26.6	48.1	6.5			0		
42-58+	C2g	41.0	38.4	20.6	0.1	1.0	1.4	15.2	23.3	20.5	17.9	56.0	5.5			0		

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		6E1b Carbonate as $CaCO_3$ Pct		Bulk density g/cc	Bulk density g/cc	Bulk density g/cc		Water content Pct.	Water content Pct.	4B2 15 bar Pct.			pH 8C1b Sat. Paste	pH 8C1a 1:10
0-5	2.32	0.191	12.1		-								13.8			5.7	6.5
5-10	0.78	0.072	10.8		-								8.7			5.9	6.5
10-15	0.23				-								6.9			6.1	6.6
15-23	0.17				-								14.7			6.6	7.4
23-42	0.04				1								19.5			7.4	8.5
42-58+	0.04				-								16.6			7.3	7.9

Depth (In.)	Extractable bases 6N2c Ca	6O2c Mg	6P2a Na	5B1a 6Q2a K	6H2a Ext. Acidity	Cat. Exch. Cap. 5A2a NaOAc		5B1b Exch. Na	Water extract from saturated paste 8A1 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6L1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-5	14.5	5.5	0.4	0.3	7.3	30.3		0.3	1.0	0.6	1.0	0.1		-	1.8	0.1		0.24
5-10	8.7	3.8	0.4	0.1	4.2	18.9		0.4	0.6	0.6	1.1	-		-	1.3	0.2		0.21
10-15	6.7	3.2	0.3	0.1	2.3	12.2		0.3	0.9	0.7	2.0	-		-	1.3	0.4		0.28
15-23	14.4	8.8	1.2	0.2	2.4	26.3		1.0	0.9	1.0	3.0	-		-	1.6	1.2		0.47
23-42	26.1	12.9	2.1	0.2	1.1	38.2		1.9	1.5	1.4	4.6	-		0.3	2.6	2.8		0.80
42-58+	18.0	9.0	1.6	0.2	2.4	30.1		1.5	0.5	0.9	3.0	-		-	1.6	1.2		0.43

LOVEJOY LOAM (taxadjunct) 1/

S61Calif-46-6

Location: Sierra Valley, Sierra County, California. 0.1 mile west of County Highway A-23. 0.25 mile north, 0.1 mile west of the southeast corner of section 9, T. 21 N., R. 14 E. Approximately 1½ miles northeast of Calpine. Date of Sampling: September 20, 1961.
Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, James McLaughlin, Eugene Begg.
Vegetation: Low sage and dry land sedges. Climate: Estimated annual precipitation 15 inches. Nearest weather station (Sierraville): Mean annual temperature 46.0° F.; January average 28.2° F.; July average 63.6° F.; mean annual precipitation 25.78 inches. Parent Material: Alluvium from mixed sources. Relief: Hummocky low terrace with a slope of 1 percent. Elevation: 4910 feet. Drainage: Moderately well drained; runoff slow; permeability moderately slow in the upper part of the profile and slow in the lower part. Soil Moisture: Slightly moist below 15 inches.
Remarks: This soil appears to be bisequal, with a Chestnut profile developing in the A horizon of an older Planosol or Solodized Solonetz soil. The surface has small hummocks 3 to 4 inches high and 3 to 24 inches in diameter which are very matted with roots. A one-half inch mat of roots spreads out at the bottom of the B2 over the pan below. The pan or Clm has the appearance of having strong medium angular blocky structure when broken and the peds all appear to have thin films. Horizons A1 (0-5 inches), B2t (15-23 inches), and C2g (42-58 inches) were sampled for the Bureau of Public Roads and the California Division of Highways. It is difficult to determine the texture of the Clm in the field because of difficulty in breaking up the pan and wetting satisfactorily.
Classification: Xerollic Durargid, fine-loamy, mixed, mesic.

HORIZON	DESCRIPTION
A1 RSL No. 61742	0 to 5 inches, gray (10YR 5/1 to 6/1) loam, very dark grayish brown (10YR 3/2) when moist; strong fine platy structure; soft, friable, slightly sticky and slightly plastic; abundant very fine, few fine to medium roots; plentiful very fine interstitial, few fine to medium tubular pores; neutral (pH 6.6); clear, smooth boundary.
A21 RSL No. 61743	5 to 10 inches, gray (10YR 5/1) clay loam with common medium faint brown (10YR 5/3) mottles, dark grayish brown (10YR 4/2) with common medium faint dark brown (10YR 4/3) mottles when moist; weak coarse platy and weak medium prismatic structure; hard, friable, sticky and plastic; plentiful very fine to medium roots; common very fine and fine tubular, and few very fine interstitial pores; many thin clay films in pores and on ped faces; medium acid (pH 6.0); clear, smooth boundary.
A22 RSL No. 61744	10 to 15 inches, light brownish gray (10YR 6/2) sandy clay loam, dark brown (10YR 3/3) when moist; moderate coarse prismatic structure; very hard, firm, slightly sticky and slightly plastic; plentiful very fine mainly exped roots; common very fine, few fine to medium tubular and interstitial pores; common thin clay films in pores and bridges; slightly acid (pH 6.4); abrupt, smooth boundary.
B2t RSL No. 61745	15 to 23 inches, grayish brown (10YR 5/2) sandy clay, very dark grayish brown (10YR 3/2) when moist; strong coarse columnar structure; extremely hard, extremely firm, sticky and plastic; plentiful very fine exped roots; few very fine tubular pores; continuous moderately thick clay films on ped faces and in pores; mildly alkaline (pH 7.8); abrupt, smooth boundary.
Clm	23 to 42 inches, light gray (2.5Y 7/2), variegated very dark grayish brown (2.5Y 3/2) and light

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Mohave sandy clay loam (taxadjunct) **LOCATION** San Bernardino County, California

SOIL FAMILY: Typic Haplargid, fine-loamy, mixed, thermic.

SOIL NOS. S55Calif-36-5 **LAB. NOS.** 5532 thru 5537

MOHAVE SANDY CLAY LOAM (taxadjunct) 1/
S55Calif-36-5

Sampled by: Fred Eggers, W. G. Harper, R. E. Nelson.
Description by: W. G. Harper Sampling date: January, 1955.
Location: 1000 feet W and 100 feet S of center Sec. 20, T8N, R4W, about 1 mile NW of Helendale, San Bernardino County, California.
Elevation: About 2500 ft. Annual precipitation about 4 inches
Classification: Typic Haplargid, fine-loamy, mixed, thermic.
Vegetation: Shadscale, inkweed, a few patches of saltgrass. About 70 percent of surface barren. Sample taken in barren area
Parent Material: Alluvium on high terrace of the Mojave River. Parent rocks are mixed but largely from granitic formations.
Physiography: Terrace - old river flood plain - slope less than 1 percent
Drainage: Runoff and permeability are slow.

HORIZON	DESCRIPTION
S55Calif-36-5-1 A_p 0- 4"	Light yellowish brown (10YR 6/4 yellowish brown 5/4 moist) loam; moderately defined medium to fine platy and vesicular; slightly hard when dry, friable when moist; very strongly calcareous; slowly permeable; moderately micaceous: contains a few plant roots; pH on 1 to 5 dilution with thymol blue 9.6; lower boundary abrupt and smooth.
S55Calif-36-5-2 B_1 4- 8"	Light brown (7.5YR 6/4 dry, brown 5/4 moist) silty clay loam; weak medium platy; soft when dry, very friable when moist; very strongly calcareous with some coarse prominent white mottling of lime and salts; moderately micaceous; slowly permeable; pH on dilution 9.0; moderate number of plant roots mainly between the plates; abrupt and smooth lower boundary.
S55Calif-36-5-3 B_2 8-12"	Light reddish brown (5YR 6/4 dry, reddish brown 5/4 moist) silty clay; moderately defined fine subangular blocky; extremely hard when dry, firm when moist. This is very strongly calcareous and has a few medium to fine white lime mottlings; contains a moderate amount of very fine flecks of mica; is very slowly permeable; contains very few plant roots; pH on dilution 9.2; abrupt and smooth lower boundary.
S55Calif-36-5-4 B_{2ca} 12-25"	Brown (7.5YR 5/4 dry, 4/3 moist) sandy clay loam; massive; extremely hard when dry, firm to friable when moist; moderately slow permeability. This is very strongly calcareous and contains many lime nodules that range up to 3/8 inch in diameter; pH on dilution 9.6; smooth lower boundary.
S55Calif-36-5-5 C_{ca} 25-44"	Pale brown (10YR 6/3 dry, brown 5/3 moist) fine sandy loam; massive; very hard when dry, friable when moist; very strongly calcareous with many white lime nodules ranging up to 3/8 inch in diameter; very strongly micaceous; moderately slow permeable; devoid of plant roots; pH on dilution 8.8; lower boundary abrupt and smooth.
S55Calif-36-5-6 C 44-52"	Same color as horizon above; fine sandy loam; massive; hard when dry, friable when moist; very mildly calcareous with an occasional small white lime fleck; moderate to moderately rapid permeability; pH on dilution 8.5; lower boundary abrupt.
S55Calif-36-5-7 D 52"+	Very gravelly loamy sand. Rounded gravel which is mainly from granite and quartzite range up to 2 inches in diameter.

REMARKS: Characteristics of this profile indicate that the soil probably developed under well drained conditions and subsequently became moderately to strongly impregnated with salts high in sodium. The sandy clay loam texture quoted in the name of this soil is taken from the soil survey of the Barstow area. The soil was mapped in that survey as strongly affected by alkali. Under present survey procedure, this soil would be a moderately saline-alkali phase.

1/ This pedon lacks the reddish colors and has more sodium than is characteristic of the series.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Mohave loamy coarse sand **LOCATION** San Bernardino County, Calif.

SOIL FAMILY: Typic Haplargid, fine-loamy, mixed, thermic

SOIL NOS. S59 Calif-36-1 **LAB. NOS.** 59680 - 59686

DEPTH INCHES	HORIZON	1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
		PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1		
0-5	A1	8.6	22.3	16.3	25.1	9.8	14.8	3.1	31.7	6.3	8	lcos
5-8	B1t	10.6	20.6	14.5	19.7	8.3	14.9	11.4	26.0	7.4	4	cosl
8-18	B2t	10.6	19.6	13.3	15.5	6.2	10.6	24.2	20.4	4.3	8	scl
18-32	B31t	10.6	22.5	14.5	19.1	7.6	9.3	16.4	23.2	3.7	9	cosl
32-40	B32t	13.8	23.0	12.3	15.9	6.9	11.7	16.4	21.2	6.0	24	cosl
40-45	C1	24.4	27.8	12.2	12.6	3.9	6.2	12.9	13.2	3.0	17	lcos
45-60+	C2	28.8	30.9	12.5	11.7	3.3	4.7	8.1	11.0	2.4	34	lcos

8C1b SATURATED PASTE 1:1	pH 8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	Ext. Iron % Fe 6C1b	6I1a CO_3 meq/l	6J1a HCO_3 meq/l	4B2 Soil water 15 atmos. Pct.
7.8		7.8	0.21	0.021			0.6	-	0.9	0.2	5.3	2.0
7.4		7.8	0.12	0.016			0.3	-	1.1	-	2.2	4.0
6.6		7.6	0.14	0.019			0.3	-	0.9	-	1.5	9.1
6.8		7.4	0.16	0.010			0.4	-	0.9	-	1.2	6.3
7.7		8.9	0.14	0.007			0.4	<0.5	0.7	-	2.7	6.5
7.8		8.9	0.10				0.4	-	0.7	-	3.0	5.1
8.0		8.9	0.07				0.4	-	0.8	-	2.9	3.6

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE SALTS 6P1a Na	6Q1a K	6N1a Ca	6O1a Mg	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
4.9	3.5	0.5	0.3	0.1	0.2	93	0.6	0.5	4.9	0.6	0.2	18.2
8.6	5.3	1.7	0.6	0.1	0.3	93	0.6	0.2	2.3	0.3	0.2	15.7
19.2	13.1	3.5	1.4	0.1	0.3	92	1.1	<0.1	2.1	0.2	0.5	29.9
15.5	9.7	2.1	0.7	0.2	0.2	95	0.8	<0.1	2.5	0.4	0.4	23.2
14.7	13.1	2.1	0.7	0.2	0.2	96	1.4	<0.1	3.1	<0.1	0.4	22.0
11.2	11.3	1.8	0.3	0.1	0.2	98	1.9	0.1	2.7	<0.1	0.4	19.3
8.0	6.0	1.1	0.0	0.1	0.1	100	1.8	0.1	1.1	<0.1	0.8	19.5

MOHAVE LOAMY COARSE SAND
S59Calif-36-1

Location: Approximately 400 feet north, 150 feet east of the south quarter corner of section 33, T. 5 N., R. 4 W., San Bernardino County, California, about 4 miles south of Victorville, California, 0.7 miles west of the intersection of Hesperia Road and Bear Valley Road.
Date: November 5, 1959
Description by: Fred C. Eggers and George Borst
[illegible]

and Reuben Nelson
Classification: **Typic Haplargid, fine-loamy, mixed, thermic.**
[illegible]

tea, desert needlegrass, and occasional Joshua trees.
Climate: Average annual precipitation about 6 inches; average January temperature 42°F., average July temperature 78°F., average annual temperature 66°F.
Parent Material: Coarse-texture alluvium, predominantly from granitic sources.
Topography: Nearly level alluvial terrace, with about 1 per cent northerly slope.
Elevation: 3030 feet above sea level.
Drainage: Well drained
Soil Moisture: None
Remarks: This profile is characterized by an ochric epipedon argillic horizon, according to the revised system of soil classification (7th Approximation). The relatively high pH of the A1 horizon may be due to contamination of calcareous dust from a cement plant about 5 miles north-northwest of this site.

HORIZON		DESCRIPTION
A1 RSL No. 59680	0-5"	Brown (10YR 5/3) loamy coarse sand, brown to dark brown (7.5YR 4/4) when moist; weak fine granular structure; loose, very friable, nonsticky and nonplastic; plentiful fine roots; mildly alkaline (pH 7.8); clear wavy boundary.
B1t RSL No. 59681	5-8"	Brown (7.5YR 5/4) coarse sandy loam, reddish brown (5YR 4/3) when moist; moderate fine subangular blocky structure; slightly hard, very friable, slightly sticky, and slightly plastic; plentiful fine roots; many fine tubular pores, slightly acid (pH 6.5); abrupt, smooth boundary.

SOIL FAMILY: Yamplicia Variant coarse-loamy mixed mesic

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 61712 - 61717

Depth (in.)	Horizon	1B1b Total: Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Sand: Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt: 0.05–0.02	Int. III (0.02–0.002)	3A1 Int. II (0.2–0.02)	(2–0.2)			Coarse fragments 2A2: > 2 Pct.	2 – 19	19 – 76
		Pct. of < 2 mm															Pct. of < 76 mm	
0–3	A11	88.4	5.8	5.8	2.1	17.9	20.7	41.0	6.7	2.5	3.3	29.6	61.3			1		
3–14	A12	90.5	4.8	4.7	2.1	12.3	21.4	46.9	7.8	2.4	2.4	34.1	58.8			1		
14–19	B1	88.1	4.9	7.0	1.5	14.4	19.7	45.4	7.1	4.7	2.2	33.5	57.3			0		
19–25	B2tca	81.3	7.7	11.0	1.0	12.5	19.5	42.0	6.3	3.2	4.5	31.3	53.2			1		
25–46	C1ca	88.1	5.9	6.0	1.8	12.9	21.6	45.6	6.2	2.8	3.1	32.4	58.5			1		
46–62+	C2g	90.3	4.4	5.3	4.3	27.5	25.5	29.8	3.2	2.3	2.1	17.7	74.9			5		

BECKWOURTH COARSE SANDY LOAM (taxadjunct) 1/

S61Calif-32-16

Location: Sierra Valley, Plumas County, California. 1350 feet north of Highway 40A at the center of section 30, T. 23 N., R. 16 E. Approximately three miles west of Vinton.
Date of Sampling: September 16, 1961.
Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, James McLaughlin, Eugene Begg.
Vegetation: Silver sage, dryland sedge, and shoestring. Climate: Estimated annual precipitation 15 inches. Nearest weather station (Portola): Mean annual temperature 45.1° F.; January average 28.7° F; July average 62.6° F.; mean annual precipitation 20.62 inches. Parent Material: Coarse textured alluvium dominantly from granitic sources. Relief: Flood plain with very slightly hummocky topography with a slope of less than 1 percent. Elevation: 4915 feet. Drainage: Moderately well drained; very slow runoff; rapidly permeable in the upper part of the profile and moderate in the lower part. Moisture: Profile moist below 14 inches.
Remarks: This is a Chestnut soil with a weakly developed Bt horizon. The A12 (3-14 inches), the B2tca (19-25 inches) and the C1ca (25-46 inches) were sampled for the Bureau of Public Roads and the California State Division of Highways. **Classification: Xerollic Haplargid, coarse-loamy, mixed, mesic.**

HORIZON	DESCRIPTION
A11 RSL No. 61712	0 to 3 inches, dark gray (10YR 4/1) loamy sand (near sandy loam), very dark brown (10YR 2/2) when moist; moderate very fine platy structure; soft, very friable, nonsticky and nonplastic; abundant very fine, few fine to medium roots; many very fine tubular and interstitial pores; slightly acid (pH 6.2); clear, smooth boundary.
A12 RSL No. 61713	3 to 14 inches, dark grayish brown (10YR 4/2) loamy sand (near sandy loam), dark brown (10YR 3/3) when moist; weak coarse platy structure; soft, very friable, nonsticky and nonplastic; plentiful very fine, few fine and medium roots; common very fine tubular and interstitial and few fine and medium tubular pores; slightly acid (pH 6.4); clear, smooth boundary.
B1 RSL No. 61714	14 to 19 inches, grayish brown (10YR 5/2) sandy loam, dark brown (10YR 3/3) when moist; massive; slightly hard, friable, nonsticky and nonplastic; few very fine and fine and medium roots; common very fine tubular and interstitial and few fine to medium tubular pores; few thin clay films in pores and bridges; strongly alkaline (pH 8.6)(pH 8.0 with thymol blue); slightly effer-
	vescent, lime is segregated; clear, smooth boundary.
B2tca RSL No. 61715	19 to 25 inches, light gray (10YR 7/2) sandy loam, variegated brown (10YR 4/3 and 5/3) and white (10YR 8/2) when moist; massive; hard, firm, slightly sticky and slightly plastic; few very fine, fine, and medium roots; common very fine tubular and interstitial, few fine and medium tubular pores; common thin clay films in pores and bridges; strongly alkaline (pH 8.6) (pH 8.0 with thymol blue); strongly effervescent, lime occurs in soft masses; clear, smooth boundary.
C1ca RSL No. 61716	25 to 46 inches, white (10YR 8/1) loamy sand, yellowish brown (10YR 5/4) when moist; massive; slightly hard, friable, nonsticky and nonplastic; very few fine roots; many very fine interstitial, common very fine tubular pores; moderately alkaline (pH 8.2); strongly effervescent, lime is disseminated; clear, smooth boundary.
C2g RSL No. 61717	46 to 62 inches +, light yellowish brown (10YR 6/4) loamy sand with many large prominent mottles, dark reddish brown (5YR 3/4) and light olive gray (5Y 6/2) when moist; massive, slightly hard, friable, nonsticky and nonplastic; very few fine roots; many fine interstitial and few very fine tubular pores; mildly alkaline (pH 7.4).

1/ This pedon has less organic carbon than required for the mollic epipedon of the Beckwourth series.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE McKay silt loam (taxadjunct) **LOCATION** Lassen County, California

SOIL FAMILY: Typic Nadurargid, fine-silty, mixed, mesic

SOIL NOS. S55-Cal. 18-2 **LAB. NOS.** 55488-497

		1B1b	PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)							3A1		
											2A2	

McKAY SILT LOAM (taxadjunct) 1/
855Calif-18-2

Location: About 10 miles east of Susanville. 53/100 mile west of the east quarter-corner of Section 1, T. 29 N. R. 13 E., Honey Lake Valley SCD, Lassen County, California. 240 feet north of an east-west graveled road.
Date of Sampling: August, 1955.

[illegible]

Classification: Typic Nadurargid, fine-silty, mixed, mesic.
Vegetation: Shrub-grass with many barren patches and an occasional patch of very light gray salt crust. Total density, 15 percent. Composition: Greasewood, 60 percent; big rabbitbrush, 25 percent, big sage, 10 percent; hop sage, 1 percent; horsebrush, trace; salt grass, 3 percent; giant and blue wild rye, 1 percent; pickleweed, trace. Forage production, 100-200 pounds per acre. Climate: At Susanville the January average temperature is 30.7° F., the July average is 69.5° F., and the frost-free period average 142 days. Summers are hot and dry, winters and cold and snowy. Mean annual rainfall at Susanville is 17.8 inches, most of which falls during the period October to May. At the site where samples were taken it is estimated that the mean annual rainfall is 8 to 10 inches. Parent Material: Probably alluvial sediments from Willow Creek, mixed with Lake Lahontan sediments. Material consists of calcareous silty and clayey earths. Use: Pasture in poor condition. Relief: Smooth, nearly level basin rim or old basin floor of ancient Lake Lahontan. There is slight (3-5 inches) microrelief, probably due to wind erosion. High spots are under shrubs; low spots are barren or sparsely covered with grass and herbs. Slope less than ½ percent, plane. Elevation: About 4080 feet. Drainage: Runoff, slow. Permeability, moderate or moderately slow. Soil seems to be imperfectly drained. Probably has high water table during part of the year. Other Features: The soil exists in complex with Canby and with a darker, friable, granular soil that seems to be higher in salt. The entire part of the basin is salt and sodium-affected, different units varying mainly in degree of salinity and alkalinity. Reaction: Free carbonates measured by observing reaction to dilute HCl. pH's determined with thymol blue in about 1:5 soil-indicator dilution.

HORIZON	DESCRIPTION
Av1 RSL No. 55488	0 to 1½ inches, light brownish gray (10YR 6.5/2, dry; 4.5/2.5, moist) silt loam. Very weak thick platy nearly massive. Soft, friable, plastic, slightly sticky. Has many roots, but they do not branch in this horizon. Very porous--strongly vesicular. Moderately calcareous. pH 9.1. Abrupt, smooth lower boundary.
Av2 RSL No. 55489	1½ to 3 inches, light brownish gray (10YR 6/2, dry; 4/2.5, moist) silt loam. Moderate medium and thin platy. Friable, plastic, slightly sticky. Many roots, but they do not branch in this horizon. Moderately porous, but not so strongly vesicular as the subhorizon above. Moderately calcareous. pH 9.4. Abrupt, smooth lower boundary.
B21 RSL No. 55490	3 to 4½ inches, brown (10YR 5/3, dry; 3.5/3, moist) silty clay loam. Weak medium prismatic, breaking to strong fine and very fine granular. Peds have moderate continuous clay skins. Very hard, firm, very plastic, very sticky. Roots are profuse; they "spread out" in this horizon. Peds are nonporous. Strongly calcareous. pH 9.4. Clear, smooth lower boundary.
B22 RSL No. 55491	4½ to 9 inches, pale brown (10YR 6/3, dry; 4/2.5, moist) silty clay loam with common faint fine mottles of very pale brown (dry). Weak medium prismatic breaking to weak medium and fine subangular blocky. Firm, plastic, very sticky. Peds have moderate continuous clay skins, but are moderately porous.

CASA GRANDE SAND (taxadjunct) 1/
S55Calif-36-7

Sampling Date: January, 1955. Description By: W. G. Harper
Collectors: Fred Eggers, W. G. Harper and Reuben E. Nelson.

Location: 0.2 mi. E of W1/4 cor., Sec. 26, T10N, R3W and 150 ft. N of railroad track, about 20 miles ENE of Barstow, San Bernardino County, Calif.
Elevation: About 2500 ft. Annual precipitation about 4 inches
Classification: Typic Natrargid, fine-loamy, mixed, thermic.
Vegetation: Desert sage and creosote bush. 98 percent of surface barren
Parent Material: Alluvium from mixed rocks but mainly from granitic formations
Physiography: Alluvial fan or old stream terrace - slope less than 1 percent
Drainage: Practically no runoff and moderately slow to slow permeability through B_2 horizon.

HORIZON		DESCRIPTION
S55Calif-36-7-1 A	0- 6"	Reddish brown (5YR 5/4 moist) sand; massive; soft when dry, loose when moist; noncalcareous; pH on dilution 8.0; rapidly permeable; devoid of plant roots; abrupt and smooth lower boundary.
S55Calif-36-7-2 B_{21}	6-12"	Reddish brown (5YR 4/4 moist) sandy clay loam; moderately to strongly defined coarse prismatic; hard when dry, slightly firm when moist; noncalcareous; pH on dilution 8.4; few plant roots; slowly permeable; abrupt lower boundary.
S55Calif-36-7-3 B_{22}	12-15"	Brown (7.5YR 5/4 dry, 4/4 moist) sandy clay loam; weakly to moderately coarse prismatic; very hard when dry, friable when moist; noncalcareous except where a few fine threads of lime occur; pH on dilution 8.4; practically devoid of plant roots; moderately permeable; abrupt lower boundary.
S55Calif-36-7-4 B_{2ca}	15-22"	Brown (7.5YR 5/4 moist) sandy clay loam; weak medium subangular blocky; hard when dry, friable when moist; moderate to moderately slow permeability; moderately calcareous with common medium to fine distinct white lime veins; pH on dilution 8.7; no plant roots; abrupt and smooth lower boundary.
S55Calif-36-7-5 C	22-26"	Pale brown (10YR 6/3 dry, brown 5/3 moist) fine sandy loam; massive; slightly hard when dry, friable when moist; moderately calcareous with few medium distinct white lime veins; moderately permeable; pH on dilution 8.6; no plant roots; lower boundary abrupt and wavy.
S55Calif-36-7-6 C_b	26-36"	Pink (7.5YR 7/4 dry, light brown 6/4 moist) sand; massive; soft when dry, loose when moist; mildly calcareous; very rapidly permeable; without plant roots; lower boundary abrupt and wavy.
S55Calif-36-7-7 B_{2b}	36-60"	Pink (7.5YR 7/4 dry, light brown 6/4 moist) sandy clay loam, moderately defined medium subangular blocky; moderately calcareous with common medium distinct white lime veins; moderately slow permeability; pH on dilution 8.9.

1/ The Casa Grande series is now in a hyperthermic family. This pedon has a mean annual soil temperature somewhat below 72° F.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Casa Grande sand (taxadjunct) **LOCATION** San Bernardino County, Calif.

SOIL FAMILY: Typic Natrargid, fine-loamy, mixed, thermic.

SOIL NOS. 855Calif-36-7 **LAB. NOS.** 5547 thru 5552

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 6	A	8.9	17.8	17.8	30.7	13.3	8.1	3.4	35.0	3.4	6	s
6-12	B21	11.2	14.1	11.0	18.8	8.4	12.8	23.7	25.0	6.6	10	scl
12-15	B22	13.3	15.2	9.0	16.9	7.3	13.0	25.3	23.2	6.5	9	scl
15-22	B2ca	9.3	13.8	7.3	16.0	8.9	19.4	25.3	29.0	9.0	5	scl
22-26	C	4.9	9.1	5.0	23.7	22.3	19.8	15.2	54.0	6.4	2	fsl
26-36	Cb	8.7	24.7	17.6	36.0	6.7	4.6	1.7	30.6	0.3	2	c s

pH 8C1b SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
8.0		7.8	0.10	0.007			0.67	0	0		3.8	1.8
8.2		9.0	0.15	0.016			11.0	0	0			9.3
8.1		9.3	0.07				16.4	0	0			11.2
8.0		9.3	0.06				17.6	2.7	0			10.0
8.1		9.5	0.05				15.2	3.3	0			5.5
8.3		9.3	0.01				8.9	0	0		4.7	1.7

5A2a CATION EXCHANGE CAPACITY (-(Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6Q2c	6H2a H	5B Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	5D1 EXCH. Na %	SATURATION EXTRACT SOLUBLE 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg	6O1a	6K1a	8A1 8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
3.6				0.6	0.6	17	5.1	0.8	1.6			19.8
17.0				7.5	0.6	44	99.0	1.1	8.0			34.3
19.0				7.9	0.6	42	153	2.3	13.7			49.7
20.5				9.3	0.3	45	160	2.3	23.3			38.5
12.3				4.7	0.3	38	140	2.5	16.9			38.3
4.2				1.8	Tr	43	80.0	0.7	7.2			22.7

1/ Percent of <1 inch fraction

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Casa Grande sand (taxadjunct) **LOCATION** San Bernardino County, California

SOIL FAMILY: Typic Natrargid, fine-loamy, mixed, thermic.

SOIL NOS. S55Calif-36-8 **LAB. NOS.** 5553 thru 5559

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 4	A	13.2	24.8	16.9	24.5	9.9	7.9	2.8	30.4	0.5	9	cos
4-11	B2	6.5	15.2	11.8	16.8	7.3	11.6	30.8	22.4	5.3	5	scl
11-18	B2ca	8.0	15.1	9.0	16.0	11.3	16.3	24.3	31.4	5.9	7	scl
18-24	B3	11.1	16.0	7.6	14.8	14.5	18.7	17.3	35.8	7.2	8	cosl
24-32	C	18.3	24.4	9.5	15.9	12.0	11.5	8.4	32.7	1.1	17	ls
32-44	B2cab	1.1	2.1	1.6	12.1	16.2	47.6	19.3	47.8	25.7	1	l
44-60	Cb	11.3	15.9	11.1	28.7	13.3	10.5	9.2	38.9	3.2	16	ls

8C1b pH SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	6A1a ORGANIC MATTER ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	4B2 MOISTURE TENSIONS 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
8.0		8.4	0.08	0.010			0.7	0	0		3.7	1.4
8.1		9.3	0.10	0.009			2.6	0	0			10.9
8.3		9.4	0.07	0.009			8.6	2.4	0			9.6
8.0		8.6	0.05				16.0	3.6	2.6			6.1
8.1		8.7	0.01				15.1	0	0		9.9	3.8
8.3		9.3	0.01				16.1	10.9	0			9.1
8.2		9.1	0.02				13.6	0	0		12.0	4.3

5A2a CATION EXCHANGE CAPACITY (-Na)	EXTRACTABLE CATIONS 5B 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	5D1 EXCH. Na %	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg	6O1a	6K1a	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
3.5				0.3	0.3	9	3.5	0.7	4.0			20.6
22.8				6.2	0.9	27	24.3	0.3	1.4			32.7
19.4				8.9	0.4	46	80.0	0.7	4.4			38.9
12.9				7.1	0.4	55	153	1.8	31.7			33.9
11.9				2.7	0.1	23	150	1.8	14.9			26.4
23.3				10.1	0.3	43	168	1.8	26.4			47.4
9.7				4.5	0.1	46	143	1.8	17.4			26.8

1/ Percent of <1 inch fraction

CASA GRANDE SAND (taxadjunct) 1/
S55Calif-36-8

Location: San Bernardino County, California. 200 feet north and 80 feet west of center Sec. 34, T. 10 N., R. 3 W., about 2 miles southwest of Hinkley.
Date of Sampling: January, 1955.
Description by: W. G. Harper. Collectors: Fred Eggers, W. G. Harper, and Reuben E. Nelson.
Classification: Typic Natrargid, fine-loamy, mixed, thermic.
Vegetation: Desert sage 24 inches high, creosote bushes 6 feet high. About 85 percent of surface barren. Climate: Annual precipitation about 4 inches.
Parent Material: Alluvium from mixed rocks but mainly from granitic formations.
Physiography: Old stream terrace - nearly level; ground water about 50 feet below surface. Elevation: About 2,200 feet. Drainage: Little or no runoff; moderate to slow permeability.

HORIZON	DESCRIPTION
S55Calif-36-8-1 A 0- 4"	Reddish brown (5YR 5/3 dry, 5/4 moist) coarse sand; single grain; soft when dry, loose when moist; noncalcareous, slightly micaceous; very rapidly permeable; pH on dilution of 1 to 5 with thymol blue 8.1; few very fine roots; lower boundary abrupt and smooth.
S55Calif-36-8-2 B_2 4-11"	Reddish brown (5YR 5/ 4 dry, 4/4 moist) sandy clay loam; moderately defined coarse prismatic; very hard when dry, friable when moist; noncalcareous; micaceous; slowly permeable; pH on dilution 9.0; few fine plant roots; lower boundary abrupt and smooth.
S55Calif-36-8-3 B_{2ca} 11-18"	Light reddish brown (5YR 6/4 dry, reddish brown 4.5/4 moist; sandy clay loam; weak coarse prismatic; very hard when dry, friable when moist; mildly to moderately calcareous, strongly calcareous on lime veins which range from few to common; moderately to slowly permeable; slightly micaceous; very few fine plant roots; pH on dilution 9.0; lower boundary abrupt and smooth.
S55Calif-36-8-4 B_3 18-24"	Brown (7.5YR 5/4 dry and moist) coarse sandy loam; weakly medium to fine subangular blocky; very hard when dry, friable when moist; strongly calcareous with many fine distinct white flecks of lime; slightly micaceous; moderately permeable; practically devoid of plant roots; pH on dilution 8.8; lower boundary abrupt and smooth.
S55Calif-36-8-5 C 24-32"	Light brown (7.5YR 6/4 dry, brown 5/4 moist) gravelly loamy sand; massive; hard when dry, very friable when moist; mildly calcareous in spots where occasional white lime flecks occur; slightly micaceous; moderately to rapidly permeable; devoid of plant roots; pH on dilution 8.2; lower boundary abrupt and smooth.
S55Calif-36-9-6 B_{2cab} 32-44"	Light yellowish brown (10YR 6/4 dry, dark grayish brown 4/2 moist) loam; moderately defined medium subangular blocky; very hard when dry, very friable when moist; very strongly calcareous with many lime flecks, streaks and splotches; moderate to moderately slow permeability; pH on dilution 9.3; devoid of plant roots; lower boundary abrupt and smooth.
S55Calif-36-9-7	Pale brown (10YR 6/3 dry, brown 5/2.5 moist) loamy sand;

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aquic Natrargid, fine-loamy, mixed, mesic

SOIL Loyalton fine sandy loam SOIL Nos. S61Calif-32-11 LOCATION Sierra Valley, Plumas County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 61659 - 61667

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.2)			Coarse fragments 2A2 ≥ 2	2 - 19	19 - 76
		Pct. of < 2 mm														Pct.	Pct. of < 76 mm	
0-1/4	A11	78.7	18.5	2.8	5.5	16.1	13.5	28.9	14.7	11.8	6.7	43.6	46.9			2		
1/4-5	A12	72.2	23.1	4.7	4.5	11.4	12.7	28.1	15.5	10.1	13.0	42.1	40.2			2		
5-8	A2	71.8	23.8	4.4	4.8	13.4	12.7	27.4	13.5	11.3	12.5	40.7	42.4			2		
8-12	B21t	58.7	18.9	22.4	5.0	10.2	10.8	21.7	11.0	8.0	10.9	31.2	35.5			4		
12-17	B22tca	51.6	18.7	29.7	5.3	9.4	9.3	18.6	9.0	8.1	10.6	27.7	32.0			3		
17-23	B31ca	43.5	20.6	35.9	3.3	7.8	8.6	16.1	7.7	8.1	12.5	24.6	27.0			5		
23-33	B32ca	53.0	21.4	25.6	7.7	10.8	10.7	16.1	7.7	8.6	12.8	24.4	37.2			3		
33-53	IIC1	83.0	8.2	8.8	20.5	24.6	19.7	15.9	2.3	3.4	4.8	11.3	75.1			39		
53-63+	IIC2	88.7	6.2	5.1	39.6	24.1	9.5	13.4	2.1	3.3	3.9	10.0	81.0			38		

Depth (In.)	6A1a Organic carbon	6B1a Nitrogen	C/N		6E1b Carbonate as CaCO3		Bulk density: 4A1f 1/3 bar	4A1h Oven-Dry			Water content: 4B1c 1/3 bar		4B2 15 bar	4D1 Vert. Extensibility	4A C*		pH: 8C1b Sat. Paste	8C1a 1:10
	Pct.	Pct.			Pct.		g/cc	g/cc	g/cc		Pct.	Pct.	Pct.	in./in.				
0-1/4	1.23	0.073	16.8		-								5.0				7.1	7.3
1/4-5	0.66	0.056	11.8		-		1.54	1.57			12.1		4.7	0.006	0.99		8.0	8.3
5-8	0.38	0.034	11.2		-								4.8				8.6	8.7
8-12	0.37	0.035	10.6		1		1.14	1.65			45.9		20.3	0.115	0.98		9.4	10.1
12-17	0.31				6		1.20	1.47			38.1		23.1	0.065	0.99		9.8	10.2
17-23	0.19				9		1.17	1.41			39.7		24.4	0.061	0.98		9.9	10.3
23-33	0.17				18		1.30	1.46			32.1		20.9	0.039	0.99		9.5	10.2
33-53	0.02						1.32	1.44			25.8		8.9	0.028	0.76		8.9	9.6
53-63+	0.02												7.1				8.5	9.0

Depth (In.)	Extractable bases 5B1a: 6N2c Ca	6O2c Mg	6P2a Na	6Q2a K	6H2a Ext. Acidity	Cat. Exch. Cap. NaOAc 5A2a	5B1b Exch. Na	Water extract from saturated paste 8A1: 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g							meq/liter									mmho/cm
0-1/4	4.9	2.4	1.0	1.5	1.1	11.2	0.7	3.4	3.5	9.5	1.5		0.4	9.7	5.1		1.52
1/4-5	5.2	2.6	1.9	1.5	0.6	12.4	1.7	0.5	0.4	7.6	0.3		0.4	5.8	2.4		0.88
5-8	4.4	2.6	3.8	0.8	0.4	13.5	3.5	0.6	0.4	12.3	0.1		0.7	6.1	6.4		1.44
8-12	10.1	2.2	30.9	0.4	-	32.4	27.3	0.8	0.6	66.5	0.1		4.6	14.8	30.9		5.54
12-17	10.5	2.4	33.3	0.3	-	32.9	27.0	0.6	0.6	77.3	0.1		14.6	19.8	21.8		5.21
17-23	13.1	3.6	38.3	0.3	-	35.3	31.2	0.8	0.5	68.5	0.1		16.3	18.1	17.0		6.25
23-33	12.8	4.4	29.0	0.2	-	31.0	26.0	0.7	0.6	47.0	0.1		6.1	12.7	15.8		4.63
33-53	4.0	3.5	11.2	0.1	0.6	20.0	10.5	0.7	0.4	26.4	-		1.6	6.1	9.8		2.70
53-63+	4.6	4.3	8.4	0.1	0.8	18.4	7.8	1.1	0.9	22.1	0.1		1.1	3.7	11.0		2.50

Depth (In.)	8A Water at Saturation	5D1 Exchangeable Na	Clay Mineralogy (<0.002 mm.): Montmorillonite	Vermiculite	Mica	Kaolinite	Kaolinite % DTA
	Pct.	Pct.	X-ray				
0-1/4	31.0	6					
1/4-5	24.0	14	x	x	-	t	t
5-8	25.5	26					
8-12	53.4	84					
12-17	81.7	82	xxx	x	-	t	3
17-23	108.4	88					
23-33	65.4	84	xxx	-	-	t	4
33-53	28.1	52					
53-63+	27.1	42					

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

* Fractional volume of the horizon occupied by <2 mm. soil material.

LOYALTON FINE SANDY LOAM

S61Calif-32-11

Location: Sierra Valley, Plumas County, California. Lucky Hereford property - 200 feet west of Plumas County Highway A-24. 200 feet west and 200 feet north of S1/4 corner section 24, T. 22 N., R. 15 E. Date of Sampling: September 12, 1961.
Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, James McLaughlin, Eugene Begg.
Vegetation: Rabbit brush, silver sage, big sage and salt grass. Climate: Estimated annual precipitation 12 inches. Nearest weather station (Sierraville): Mean annual temperature 46° F.; January average 28.2° F.; July average 63.6° F.; mean annual precipitation 25.78 inches.
Parent Material: Medium textured alluvium from mixed sources. Relief: Low terrace near valley floor with slightly hummocky topography. Slope less than 1 percent. Elevation: 4889 feet.
Drainage: Moderately well drained; runoff is slow; permeability very slow.
Remarks: This is a Solodized Solonetz soil. The surface (A11) has a winnowed appearance. The A11 may be absent or in the deeper spots may be several inches thick. The B21t and B22t horizons

have organic staining on the ped faces. The IIC2 horizon has one inch thick, intermittently cemented bands which are dark red (2.5YR 3/6) when moist and yellowish red (5YR 4/6 and 5/6) when dry. Classification: Aquic Natrargid, fine-loamy, mixed, mesic.

HORIZON	DESCRIPTION
A11	0 to 1/4 inch, light brownish gray (10YR 6/2) loamy sand, very dark grayish brown
RSL No.	(10YR 3/2) when moist; single grain; soft, very friable, nonsticky and nonplastic;
61659	neutral (pH 7.0); abrupt, smooth boundary.
A12	1/4 to 5 inches, gray (10YR 6/1) fine sandy loam, very dark grayish brown (10YR 3/2)
RSL No.	when moist; moderate coarse platy structure; soft, very friable, nonsticky and nonplastic;
61660	abundant very fine, plentiful fine to coarse roots; many very fine interstitial, common very fine, few fine and medium tubular pores; neutral (pH 6.8); clear, smooth boundary.
A2	5 to 8 inches, light gray (10YR 7/1) sandy loam, dark grayish brown (10YR 4/2) when moist;
RSL No.	weak very fine platy structure; slightly hard, friable, nonsticky and nonplastic;
61661	plentiful very fine, fine, and medium roots; common very fine tubular and interstitial pores; few thin clay films in pores; moderately alkaline (pH 8.0)(pH 8.6 with thymol blue); abrupt, smooth boundary.
B21t	8 to 12 inches, variegated grayish brown (10YR 5/2) and light gray (10YR 7/2) sandy clay,
RSL No.	very dark grayish brown (10YR 3/2) when moist; strong medium columnar structure;
61662	extremely hard, very firm, sticky and very plastic; abundant very fine, few fine to medium, mainly exped roots; common very fine and few fine tubular pores; continuous thick clay films on ped faces; moderately alkaline (pH 8.4)(pH 9.6 with thymol blue); very slightly effervescent, lime is concentrated in areas; clear, smooth boundary.
B22tca	12 to 17 inches, variegated grayish brown (10YR 5/2) and light gray (10YR 7/2) sandy
RSL No.	clay, variegated very dark grayish brown (10YR 3/2) and brown (10YR 5/3) when moist;
61663	strong medium prismatic structure; extremely hard, very firm, sticky and very plastic; few very fine inped, and few fine exped roots; common very fine tubular pores; continuous thick clay films on ped faces; moderately alkaline (pH 8.4)(pH 9.6 with thymol blue); strongly effervescent, lime occurs in soft masses; clear, slightly wavy boundary.
B31ca	17 to 23 inches, variegated light gray (10YR 7/2) and pale brown (10YR 6/3) sandy clay,
RSL No.	variegated dark grayish brown (10YR 4/2), brown (10YR 5/3), and light gray (10YR 7/2)

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aquic Natrargid, fine-loamy, mixed, mesic
SOIL Lovalton fine sandy loam **SOIL Nos.** S61Calif-32-10 **LOCATION** Sierra Valley, Plumas County, California
SOIL SURVEY LABORATORY Riverside, California **LAB. Nos.** 61649 - 61658

Depth (in.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	3A1 Int. II (0.2-0.02)	(2-0.2)			Coarse fragments 2A2: > 2 Pct.	2 - 19	19 - 76
		Size class and particle diameter (mm) — Pct. of < 2 mm															Pct. of < 76 mm	
0-2	A1	89.1	7.7	3.2	21.2	19.4	14.5	25.1	8.9	4.5	3.2	27.1	66.5			6		
2-7	A2	63.7	31.3	5.0	7.0	9.7	9.9	24.7	12.4	12.1	19.2	39.6	36.2			3		
7-10	A & B	71.8	21.5	6.7	8.6	12.0	11.8	27.5	11.9	8.4	13.1	36.7	43.5			5		
10-16	B21t	60.5	19.7	19.8	6.7	9.8	10.1	23.0	10.9	7.3	12.4	31.8	36.0			4		
16-21	B22t	42.2	23.7	34.1	3.4	7.3	6.9	16.4	8.2	8.2	15.5	26.6	23.8			1		
21-30	B31t	59.9	16.7	23.4	7.0	11.5	9.1	22.7	9.6	7.4	9.3	31.4	35.9			7		
30-39	B32t	74.5	11.3	14.2	14.9	14.0	8.1	26.8	10.7	5.7	5.6	33.2	47.0			29		
39-49	IIC1	91.8	5.5	2.7	20.6	19.8	14.6	29.9	6.9	3.4	2.1	26.9	68.3			31		
49-60	IIC2m	89.8	5.2	5.0	29.7	35.0	11.8	11.5	1.8	2.7	2.5	10.1	82.4			39		
60-65+	IIC3	83.7	6.3	10.0	25.3	22.0	15.3	18.1	3.0	3.1	3.2	14.2	72.6			33		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		6E1b Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	4B2 15 bar Pct.			pH 8C1b Sat. Paste	8C1a 1:10
0-2	0.85	0.057	15		-								3.2			6.4	6.5
2-7	0.31	0.030	10		-								3.0			7.0	6.8
7-10	0.28	0.027	10		-								4.2			7.8	7.7
10-16	0.26				-								15.7			8.4	9.3
16-21	0.19				-								24.3			8.7	9.5
21-30	0.09				1								15.6			8.8	9.8
30-39	0.04				1								10.0			8.6	9.6
39-49	0.02				-								3.1			8.0	7.5
49-60	0.02				-								5.1			7.8	7.8
60-65+	0.02				-								8.8			7.5	8.0

Depth (in.)	Extractable bases 6N2c Ca	6O2c Mg	6P2a Na	5B1a 6Q2a K	6H2a Ext. Acidity	Cat. Exch. Cap. 5A2a NaOAc		5B1b Exch. Na	Water extract from saturated paste 8A1: 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-2	2.7	1.3	0.1	0.5	2.2	7.2		0.1	1.5	1.2	1.3	0.9		-	3.0	1.0		0.52
2-7	2.6	1.5	0.6	0.3	1.6	7.2		0.6	0.4	0.6	3.0	0.1		-	1.7	1.6		0.40
7-10	1.8	1.3	3.7	0.1	0.9	8.6		3.4	2.4	0.6		0.1		-	3.4	9.6		1.59
10-16	2.8	3.9	16.3	0.2	0.7	24.0		15.1	0.8	0.7	26.5	0.1		0.8	6.2	14.7		2.91
16-21	3.7	6.2	26.1	0.3	-	32.9		24.2	0.9	0.5	20.9	0.1		0.3	4.0	9.9		2.20
21-30	6.2	4.6	19.4	0.3	-	24.6		17.2	0.6	0.6	32.2	-		0.9	5.6	13.7		3.37
30-39	5.7	3.4	13.9	0.1	0.3	20.5		12.5	0.5	0.6	41.8	0.1		1.3	6.4	23.4		4.59
39-49	2.0	1.7	2.3	-	1.5	7.3		2.2	0.4	0.1	2.4	-		-	1.2	0.8		0.28
49-60	1.2	3.2	1.9	0.1	0.9	11.3		1.8	0.3	0.2	2.6	-		-	1.4	0.9		0.32
60-65+	6.7	5.5	2.8	0.1	1.4	17.1		2.8	0.2	0.3	2.5	-		0.1	2.7	0.9		0.32

Depth (in.)	8A Water at Saturation Pct.	5D1 Exchangeable Na Pct.		Gypsum Pct.		
0-2	25.5	1				
2-7	18.6	8				
7-10	19.7	39				
10-16	48.6	63				
16-21	91.2	74				
21-30	69.0	70				
30-39	34.2	61				
39-49	20.9	30				
49-60	22.7	16				
60-65+	25.2	16				

LOYALTON FINE SANDY LOAM

S61Calif-32-10

Location: Sierra Valley, Plumas County, California. 100 yards east of County Highway A-24, 100 yards east of center section 24, T. 22 N., R. 15 E. Lucky Hereford property, approximately 4½ miles north of Loyalton along Beckworth Lane. Date of Sampling: September 11, 1961.
Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, James McLaughlin, Eugene Begg.
Vegetation: Silver sage, rabbit brush, salt grass, big sage. Climate: Estimated annual precipitation 12 inches. Nearest weather station (Sierraville): Mean annual temperature 46° F.; January average 28.2° F.; July average 63.6° F.; mean annual precipitation 25.78 inches.
Parent Material: Medium textured alluvium from mixed sources. Relief: Low terrace near valley floor, slightly hummocky. Slope less than 1 percent. Elevation: 4890 feet.
Drainage: Moderately well drained, runoff slow, very slowly permeable.
Soil Moisture: Moist below 10 inches when sampled.
Remarks: This is a Solodized Solonetz soil. The upper part of the A2 horizon has somewhat more pronounced coarse platy structure than the lower portion. The IIC2m horizon is made up of coarse sand which is somewhat intermittently cemented in bands. The profile appears to be degrading. Evidences for degradation are the gray coatings on peds of the B21t horizon and "ghost columns" in the A & B horizon. Classification: Aquic Natrargid, fine-loamy, mixed, mesic.

HORIZON	DESCRIPTION
A1 RSL No. 61649	0 to 2 inches, light brownish gray (10YR 6/2) loamy coarse sand, very dark grayish brown (10YR 3/2) when moist; single grain; loose, very friable, nonsticky and nonplastic; slightly acid (pH 6.2); noncalcareous.
A2 RSL No. 61650	2 to 7 inches, gray (10YR 6/1) fine sandy loam, dark brown (10YR 3/3) when moist; strong very coarse platy structure; soft, very friable, nonsticky and nonplastic; plentiful very fine and fine, few medium and coarse roots; abundant very fine interstitial, few very fine tubular pores; medium acid (pH 6.0); noncalcareous; abrupt, smooth boundary.
A & B RSL No. 61651	7 to 10 inches, variegated gray (10YR 6/1) and dark brown (10YR 4/3) sandy loam (near loam), dark gray (10YR 4/1) when moist; moderate coarse columnar structure; very hard, firm, slightly sticky and slightly plastic; plentiful very fine and fine, few medium and coarse mainly exped roots; common very fine tubular and interstitial, and few fine and medium tubular pores; common thin clay films in pores and on ped faces; moderately alkaline (pH 8.0)(pH 8.4 with thymol blue); noncalcareous; abrupt, smooth boundary.
B21t RSL No. 61652	10 to 16 inches, dark gray (10YR 4/1) sandy clay, very dark brown (10YR 2/2) when moist; strong coarse columnar structure; extremely hard, firm, sticky and very plastic; plentiful very fine, few fine and medium mainly exped roots; few very fine and fine tubular pores; continuous thick clay films in pores and on ped faces; strongly alkaline (pH 8.5) (pH 9.0 with thymol blue); noncalcareous; clear, smooth boundary.
B22t RSL No. 61653	16 to 21 inches, grayish brown (10YR 5/2) sandy clay, dark brown (10YR 3/3) when moist; moderate coarse prismatic structure; extremely hard, firm, sticky and very plastic; few very fine roots; few very fine pores; many moderately thick clay films in pores and on ped faces; strongly alkaline (pH 8.5)(pH 9.4 with thymol blue); slightly effervescent; lime is segregated; clear, smooth boundary.
B31t RSL No. 61654	21 to 30 inches, variegated pale brown and light gray (10YR 6/3) and (10YR 7/2) sandy clay loam (near sandy clay), dark brown (10YR 3/3) when moist; weak coarse prismatic structure; extremely hard, very firm, sticky, and plastic; very few very fine to medium roots; common very fine interstitial, few fine tubular pores; many thin clay films on ped faces; strongly alkaline (pH 8.5)(pH 9.6 with thymol blue); slightly effervescent; lime is segregated; clear, smooth boundary.
B32t RSL No. 61655	30 to 39 inches, pale brown (10YR 6/3) coarse sandy clay loam, brown (10YR 4/3) when moist; massive; very hard, very firm, slightly plastic and slightly sticky; very few very fine to medium roots; few very fine tubular and interstitial pores; common thin clay films in pores; strongly alkaline (pH 8.5)(pH 9.4 with thymol blue); slightly effervescent; lime is segregated; abrupt, slightly wavy boundary.
IIC1 RSL No. 61656	39 to 49 inches, light gray (10YR 7/1) coarse sand, dark grayish brown (10YR 4/2) when moist; single grain; loose, nonsticky and nonplastic; very few very fine to medium roots; many very fine and common fine interstitial pores; slightly acid (pH 6.3); noncalcareous; abrupt, slightly wavy boundary.
IIC2m RSL No. 61657	49 to 60 inches, pale brown (10YR 6/3) coarse sand, brown (10YR 4/3) when moist; massive; very hard, very firm, nonsticky and nonplastic; very few very fine to medium roots; many very fine interstitial pores; slightly acid (pH 6.5); noncalcareous; abrupt, slightly wavy boundary.
IIC3 RSL No. 61658	60 to 65 inches +, light brownish gray (2.5Y 6/2) loamy sand, brown (10YR 4/3) when moist; massive; friable, nonsticky and nonplastic; very few very fine to medium roots; common very fine interstitial pores; neutral (pH 7.0); noncalcareous.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Churchill silt loam (taxadjunct) **LOCATION** Lassen County, California.

SOIL FAMILY: Duric Natrargid, fine, montmorillonitic, mesic

SOIL NOS. S55-Cal-18-1 **LAB. NOS.** 55479-487

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1	2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2	
0-1½	A11	0.3	1.0	1.4	11.6	20.7	60.3	4.7	57.1	32.6	0	sil
1½-3	A12	0.4	0.6	1.0	8.8	17.7	62.7	8.8	50.2	37.1	0	sil
3-10	B2	0.1	0.2	0.4	3.2	5.6	45.0	45.5	19.9	33.1	2	sic
10-16	B3ca	0.3	0.6	0.4	2.2	4.9	60.2	31.4	20.2	46.6	3	sicl
16-28	Cca	0.3	0.6	1.0	8.1	10.4	61.1	18.5	33.4	44.5	0	sil
28-33	C1	0.3	2.0	2.7	8.9	6.9	62.1	17.1	26.8	48.0	0	sil
33-43	C2	1.5	1.4	1.1	4.2	6.8	69.1	15.9	24.4	54.4	2	sil
43-55	C3	0.4	1.1	1.6	7.6	10.9	63.8	14.6	33.5	46.7	6	sil
55-64	C4	1.2	1.5	0.9	2.1	4.5	79.2	10.6	22.6	62.5	9	sil

8C1b pH SATURATED PASTE 1:1	8C1a pH 1:5	8C1a pH 1:10	6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	4B2 MOISTURE TENSIONS 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
7.3		8.2	0.68	.054	12.6		6.2		-			7.4
7.4		8.6	0.30	.035	8.6		22.2		-			6.0
8.7		9.6	0.39				18.1	2	-			28.5
9.1		9.8	0.17				21.3	17	-			36.8
8.9		9.8	0.13				16.0	16	-			27.1
8.9		9.8	0.11				12.3	10	-			27.4
8.8		9.6	0.11				8.2	8	-			28.3
8.4		9.4	0.07				6.6	4	-			27.3
8.2		9.1	0.07				8.4	3	-			22.5
5A2a	EXTRACTABLE CATIONS 5B					5D1	SATURATION EXTRACT SOLUBLE 8A1					8A

CHURCHILL SILT LOAM (taxadjunct) 1/
S55Calif-18-1

Location: About 10 miles east of Susanville, 56/100 mile west of the east quarter-corner of section 1, T 29 N, R 13 E, Honey Lake Valley SCD, Lassen County, California. 80 feet south of an east-west gravelled road.
Description by: W. M. Johnson. Collectors: S. B. Johnson, Ralph E. Nelson, Stuart Porter, Reuben E. Nelson, John Cady, E. A. Naphan and W. M. Johnson. Date of Sampling:
Classification: Duric Natrargid, fine, montmorillonitic, mesic.
Vegetation: Sparse (probably about 15 percent density) cover of shrubs and grasses. Big sage, rabbitbrush, greasewood, spiny hopsage, salt grass are dominant. Minor components are giant wild rye and squirreltail. Climate: At Susanville the January average temperature is 30.7 °F, the July average is 69.5° F, and the frost-free period averages 142 days. Summers are hot and dry, winters are cold and snowy. Mean annual rainfall at Susanville is 17.8 inches, most of which falls during the period October to May. At the site where samples were taken it is estimated that the mean annual rainfall is 8 to 10 inches. Parent Material: Mainly alluvial sediments deposited by Willow Creek, including a large proportion of reworked Lake Lahontan sediments. Calcareous silty and clayey materials. The lower horizon may be undisturbed Lake Lahontan sediments. Relief: Smooth, gently sloping basin rim or old basin floor of ancient Lake Lahontan. There is a fairly regular microrelief of 3 to 6 inches due to wind reworking of the surface. High spots are under shrubs; low spots are barren or sparsely covered with grass and herbs. Slope about ½ percent, slightly convex, towards the east. Elevation: About 4080 feet. Use: Pasture in poor condition. Drainage: Runoff, medium. Permeability moderately slow. There is probably a high water table during part of the year. Other Features: The soil surface is characterized by many barren patches, and many small patches with a white salt crust. The surface in more than one-half of the area has a fine polygonal pattern of cracks from 1/4 to 1/2 inch wide, 3 to 4 inches deep. These cracked areas are assumed to have higher salt content in the upper part of the profile than the site sampled. The soil sampled occurs in complex association with a friable, highly saline soil having a granular rather than columnar-blocky B horizon. Reaction: Measure of free carbonates obtained by observing reaction to dilute HCl. pH's determined with thymol blue in about 1:5 soil-indicator dilution.

HORIZON	DESCRIPTION
A11 RSL No. 55479	0 to 1½ inches, light brownish gray (10YR 6/2, dry; 4/2, moist) silt loam. Very weak thick and very thick platy. Soft, friable, slightly plastic, slightly sticky. Roots numerous, but do not branch. Very porous--strongly vesicular. Noncalcareous. pH 8.0. Abrupt, smooth lower boundary.
A12 RSL No. 55480	1½ to 3 inches, light gray (10YR 7/2, dry; 4/2, moist) silt loam. Moderate medium and thin platy. Slightly hard, friable, slightly plastic, slightly sticky. Slightly porous--moderate number of very fine vesicles. Roots numerous, but do not branch. Noncalcareous. pH 8.3. Abrupt, slightly wavy lower boundary.
B2t RSL No. 55481	3 to 10 inches, brown (10YR 5/3, dry; 3/3, moist) silty clay loam. Weak medium columns break to strong fine and very fine blocks with thick, continuous clay skins. Very hard, firm, very plastic, very sticky. Roots numerous and permeate the soil mass; they "spread out" in this horizon. Soil mass as a whole is porous, but individual peds are nonporous. Noncalcareous. pH 9.2. Clear, wavy lower boundary.
B3tca RSL No. 55482	10 to 16 inches, light brownish gray (10YR 6.5/2.5, dry; 4/3, moist) silty clay loam with many coarse, distinct mottles of very pale brown (dry). Moderate fine and very fine subangular blocky. Friable, very plastic, very sticky. Peds have thin, continuous clay skins and are nonporous. Roots numerous. Strongly calcareous. pH 9.4. Gradual, wavy lower boundary.
Cca RSL No. 55483	16 to 28 inches, light brownish gray (10YR 6.5/2.5, dry; 4.5/3, moist) silty clay loam with common, fine, thread-like mottles of white and a few coarse mottles of very pale brown (dry). Moderate medium and fine subangular blocky. Very firm, plastic, very sticky. There are friable masses in this horizon, but most of the material appears to be slightly cemented. Moderate number of roots. Many thin patches of clay skins on vertical and horizontal surfaces and in root channels. Strongly calcareous. pH 9.3. Gradual, wavy lower boundary.
C1 RSL No. 55484	28 to 33 inches, light gray (10YR 7/2.5, dry; 5.5/3, moist) heavy silt loam with common, fine, faint mottles of white (dry). Weak medium subangular blocky. Varies from friable to very firm in different parts of the same horizon. Peds have thin, scattered patches of clay skins. Contains a few snail shells. Few roots. Some places are noncalcareous, but most are strongly calcareous. pH 9.3. Gradual, wavy lower boundary.
C2 RSL No. 55485	33 to 43 inches, very pale brown (10YR 7/3, dry; 5.5/3, moist) heavy silt loam with many fine, faint mottles of light yellowish brown, yellowish brown, and white. Weak very thick platy (stratified). Consistence varies from friable to very firm in different parts of the horizon. Has thin lenses of very firm material that appear to be slightly cemented; these are strongly calcareous. Friable spots are noncalcareous. Very few roots. A few thin patches of clay skins along lines of cleavage. Porous. pH 9.0. Gradual, wavy lower boundary.
C3 RSL No. 55486	43 to 55 inches, light gray (10YR 7/2.5, dry; 5/3, moist) heavy silt loam with common, fine mottles of white and light brown (dry). Massive-stratified. Friable matrix, with thin and thick cemented lamina or lenses. The friable parts are noncalcareous. Contains many fine and medium hard carbonate concretions that look like root channel fillings. Very few thin patches of clay skins. Porous. Very few roots. pH 8.9. Gradual, slightly wavy lower boundary.
C4 RSL No. 55487	55 to 64 inches+, white (2.5Y 8/2, dry; 5.5/2, moist) silt loam or silt with many prominent, coarse mottles of strong brown (dry). Massive-stratified. Friable matrix, with many thin and thick cemented lenses that are strongly calcareous. Most of the matrix material is noncalcareous. A few thin patches of thin clay skins, especially in tubules. Very few roots. pH 8.4.

1/ The Churchill series is in a fine, montmorillonitic, mesic family of Typic Natrargids. This pedon has a C horizon that is weakly cemented by silica.

U S DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Xerollic Natrargid, fine-silty, mixed, mesic
SOIL Skumpah silt loam (taxadjunct) SOIL Nos. 854Calif-18-2 LOCATION Lassen County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 54297 - 54304

Depth (In.)	Horizon	1B1b Total Sand (2–0.05)	Total Silt (0.05–0.002)	Total Clay (< 0.002)	Sand Very coarse (2–1)	Sand Coarse (1–0.5)	Sand Medium (0.5–0.25)	Sand Fine (0.25–0.1)	Sand Very fine (0.1–0.05)	Silt 0.05–0.02	Silt Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)			Coarse fragments 2A2 > 2 Pct.	2 – 19	19 – 76
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm															Pct of < 76 mm	
0-2	A11	29.6	53.9	16.5	0.4	1.5	2.7	12.9	12.1	19.2	34.7		17.5			0		
2-4	A12	26.5	52.9	20.6	0.3	1.4	2.7	12.4	9.7	17.5	35.4		16.8			0		
4-12	B2t	15.2	42.8	42.0	0.0	0.6	1.5	6.4	6.7	12.5	30.3		8.5			0		
12-20	B3	23.1	48.7	28.2	0.0	3.2	4.3	9.2	6.4	14.4	34.3		16.7			0		
20-38	C1		Incomplete dispersion - Gypsum													0		
38-52	C2		Possible incomplete dispersion													0		
52-60	C3	11.9	67.7	20.4	0.0	0.7	1.0	4.9	5.3	18.5	49.2		6.6			0		
60-80	C4		Incomplete dispersion - Gypsum													0		

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		6E1b Carbonate as $CaCO_3$ Pct		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	4B2 Pct.				pH 8C1b Sat. Paste	8C1a 1:10 H_2O
0-2	1.12	0.072	15.6		tr.								9.2				7.1	8.0
2-4	1.10	0.064	17.2		-								11.1				7.2	8.1
4-12	0.79	0.066	12.0		-								22.2				7.5	8.6
12-20	0.72	0.062	11.6		-								28.2				7.7	8.2
20-38	0.49				tr.								24.1				8.3	8.7
38-52	0.24				10								22.5				8.2	9.1
52-60	0.14				7								25.3				8.2	8.9
60-80	0.11				3								24.7				8.3	8.8
					4													

Depth (In.)	Extractable bases 6N1a Ca 5B1a	6O1a Mg 5B1a	6P2a Na 5B1b	6Q2a K 5B1b	Sum of bases	5A2a CEC NaOAc			Water extract from saturated paste 6N1a 6O1a Ca+Mg		6P1a Na	6Q1a K		CO_3	HCO_3	Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-2	9.1	9.2	1.0	3.4	22.7	20.9			3.4		5.9	1.6						0.98
2-4	9.6	7.5	2.7	3.9	23.7	24.9			1.3		8.6	0.9						1.01
4-12	11.1	12.1	9.7	7.5	40.4	38.4			3.7		25.0	1.6						2.80
12-20	19.5	17.1	15.9	4.3	56.8	49.3			60.4		216.0	12.0						18.70
20-38	86.8	18.0	17.0	3.3	125.1	42.7			76.2		371.9	15.9						30.20
38-52	46.3	15.8	15.2	3.2	80.5	43.3			57.9		400.0	11.6						31.60
52-60	39.7	16.8	23.2	7.1	86.8	44.9			57.3		390.6	11.6						30.40
60-80	78.1	17.9	28.6	5.1	129.7	39.8			46.1		390.6	11.6						30.50

Depth (In.)	8A Water at Saturation Pct.	5D1 Exchangeable Na Pct.		6F1a Gypsum Pct.		
0-2	36.3	5		-		
2-4	35.3	11		-		
4-12	72.4	25		-		
12-20	79.2	32		2		
20-38	67.9	40		10		
38-52	87.5	35		tr.		
52-60	82.5	52		tr.		
60-80	68.6	72		10		

SKUMPAH SILT LOAM (taxadjunct) 1/
S54Calif-18-2

SOIL SURVEY LABORATORY Riverside, California

Calneva silt

SOIL TYPE [illegible] **LOCATION** [illegible]

SOIL NOS. S55-Cal. 18-3 **LAB. NOS.** 55498-505

DEPTH INCHES	HORIZON	PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	3A1 0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-1	Av-1	0.9	5.1	6.2	17.6	15.6	45.1	9.5	42.8	29.4	2	l
1-2	Av-2	0.5	4.0	5.3	13.2	10.9	46.6	19.5	31.7	34.3	1	l
2-5	B21	0.4	2.6	3.6	7.6	5.5	36.4	43.9	16.7	29.8	0	c
5-10	B22	0.5	2.1	2.1	6.4	8.4	56.0	24.5	24.2	44.6	1	sil
10-22	C1	7.0	8.8	4.6	9.0	7.6	43.4	19.6	22.3	34.4	0	l
22-32	C2	10.8	10.9	4.6	8.8	7.7	39.7	17.5	27.9	25.0	0	l
32-33	D1	4.1	11.9	13.1	22.5	8.9	28.4	11.1	28.0	22.2	2	fsl
33-41	D2	0.5	7.6	14.3	46.6	19.1	6.6	5.3	52.6	3.7	0	fs

pH 8C1b SATURATED PASTE	pH 8C1a 1:5	pH 8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON	ORGANIC MATTER 6B1a NITROGEN	ORGANIC MATTER C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS	6E1b $CaCO_3$ equivalent	6F1a GYPSUM mo./100g. SOIL	MOISTURE TENSIONS 4B2 1/10 ATMOS.	MOISTURE TENSIONS 4B2 1/3 ATMOS.	MOISTURE TENSIONS 4B2 15 ATMOS.

CALNEVA SILT LOAM (taxadjunct) 1/
S55Calif-18-3

Location: About 8 miles east of Susanville. 4/10 mile east of the west quarter-corner of Section 3, T. 29 N., R. 13 E.; 165 feet north of east-west graveled road. Honey Lake Valley SCD, Lassen County, California.
Date of Sampling: August, 1955.
Description by: S. B. Johnson. Collectors: S. B. Johnson, Ralph E. Nelson, Stuart Porter, Reuben E. Nelson, John Cady, E. A. Naphan, and W. M. Johnson.
Classification: Typic Paleargid, fine-loamy, mixed, mesic.
Vegetation: Shrub-grass, with many barren spots. Total density about 17 percent. Composition: Greasewood, 40 percent; big sage, 40 percent; rabbitbrush, 16 percent; annual weeds, 2 percent; giant and blue wild rye, 1 percent; salt grass, 1 percent; trace of Indian rice grass and squirreltail. Forage production, 100 to 150 pounds per acre. Climate: At Susanville the January average temperature is 30.7° F., the July average is 69.5° F., and the frost-free period averages 142 days. Summers are hot and dry, winters are cold and snowy. Mean annual rainfall at Susanville is 17.8 inches, most of which falls during the period October to May. At the site where samples were taken it is estimated that the mean annual rainfall is 8 to 10 inches.
Parent Material: Loamy calcareous alluvium, probably mainly reworked Lake Lahontan sediments. The sandy substratum may be an ancient bar or beach. This site is a few feet higher than that of S55Calif-18-4. Use: Pasture in poor condition. Relief: Smooth, very gently sloping basin rim or old basin floor of ancient Lake Lahontan. There is 3 to 6 inch microrelief, probably due to wind erosion, with the high spots occurring under shrubs. Slope is about ½ percent, plane, toward the southwest. Elevation: About 4,100 feet.
Drainage: Runoff, slow. Permeability, moderate or moderately slow. Soil is probably imperfectly drained. It probably has a high water table during part of the year. Other Features: Profile was slightly moist below the Av. This soil exists in complex with a Canby-like soil and with a darker colored, more strongly saline soil. The sandy substratum is intermittent. Would probably correlate as sandy substratum phase of the soils whose samples are numbered S55Calif-18-2 and S55Calif-18-4. Reaction: Free carbonates estimated by observing reaction to dilute HCl. pH's determined with thymol blue in about 1:5 soil-indicator dilution.

HORIZON	DESCRIPTION
Av1 RSL No. 55498	0 to 1 inch, grayish brown (10YR 5.5/2, dry; 3.5/2, moist) silt loam or very fine sandy loam. Weak medium platy. Very soft, very slightly plastic. Strongly vesicular. Noncalcareous. pH 8.5. Abrupt, wavy lower boundary.
Av2 RSL No. 55499	1 to 2 inches, light brownish gray (10YR 6.5/2, dry; 4/2, moist) silt loam, with common, prominent medium mottles of brown (dry) in the lower part. Moderate fine and medium platy. Soft, slightly plastic. pH 8.8. Calcareous. Abrupt, wavy lower boundary.
B21 RSL No. 55500	2 to 5 inches, dark grayish brown (10YR 4/2, dry; 3/2, moist) light silty clay loam. Moderate fine and very fine granular. Hard, plastic. Calcareous. Lime is segregated, so that some spots are highly calcareous. Clay skins along root channels. pH 9.6. Abrupt lower boundary.
B22 RSL No. 55501	5 to 10 inches, light yellowish brown (2.5Y 6/4, dry; 4.5/4, moist) silty clay loam, with common, faint, medium mottles of light brownish gray (dry). Medium and coarse granular. Hard, plastic. Thin clay skins along root channels. Strongly calcareous. pH 9.8 plus. Clear lower boundary.
C1 RSL No. 55502	10 to 22 inches, light yellowish brown (10YR 6/4, dry; 5/4, moist) silt loam with common, faint fine mottles of light brownish gray and a few distinct medium mottles of light gray (dry). Massive or very weak, very coarse platy. Crushes to weak medium and coarse granular. Firm, slightly plastic. Thin clay skins along root channels. Strongly calcareous. pH 9.8 plus. Probably an "alkali pan." Clear, wavy lower boundary.
C2 RSL No. 55503	22 to 32 inches, light brownish gray (2.5Y 6.5/2, dry; 4/4, moist) silt loam with common faint medium mottles of light gray (dry). Weak very coarse platy, crushing to weak coarse granular. Extremely firm; slightly cemented. Few thin clay skins. Strongly calcareous. pH 9.8 plus. Abrupt, wavy lower boundary.
D1 RSL No. 55504	32 to 33 inches, pale yellow (2.5Y 7/4, dry; 5/4, moist) fine sandy loam with a few prominent fine mottles of strong brown (dry). Massive. Soft, very friable. Strongly calcareous. pH 9.8 plus. Abrupt, wavy lower boundary.
D2 RSL No. 55505	33 to 41 inches, light brownish gray (2.5Y 6/2, dry; 4/2, moist) loamy sand. Massive. Very soft, nearly loose. One rodent burrow observed, filled with roots. Strongly calcareous. There is a thin (2-4 mm.) strongly cemented layer at the top of the horizon; this layer is also strongly calcareous. Contains a few fine gravels. pH 9.8 plus. Clear lower boundary.
D3 RSL No.	41 to 47 inches, grayish brown (2.5Y 5/2, dry; 3.5/2 moist) loamy fine sand with a few prominent fine mottles of strong brown (dry). Massive. Nearly loose. Slightly micaceous. Noncalcareous. pH 9.6. Abrupt lower boundary.
D4	47 to 51 inches, grayish brown (2.5Y 5/2, dry; 3.5/2, moist) fine gravelly sand. Massive. Slightly micaceous. Very soft. Noncalcareous. pH 8.0. Abrupt lower boundary.
D5 RSL No.	51 to 61 inches +, grayish brown (2.5Y 5/2, dry; 4/2, moist) fine sand. Massive. Very soft, nearly loose. Slightly micaceous. Noncalcareous. pH 8.8.

Roots go through Av horizon without branching, spread out in the B2 horizon, then decrease in the C and spread out again in the sandy substratum.

1/ The Calneva series is in a fine, montmorillonitic, mesic family of Xeralfic Natrargids. This pedon is dry in all parts of the moisture control section more than 3/4 of the time the soil temperature at 50 cm. is 5° C or more.

SOIL SURVEY LABORATORY Riverside, California

MORALES CLAY LOAM
S55Calif-36-2

Location: 1120 feet N and 70 feet E of SW Corner of section 29, T 5 N, R 1 E ~~about 4 miles NE~~ of Lucerne Valley, San Bernardino County, California.
Collectors: Fred Eggers, W. G. Harper, R. E. Nelson, January, 1955.
Elevation: About 3000 ft. Annual precipitation about 4 inches
Classification: Typic Calciorthid, fine, montmorillonitic, thermic.
California Grouping: III. Has had one crop this year
Vegetation: Barley probably was cut for hay - fair to poor stubble
Parent Material: Alluvium from mixed rocks but mainly from granitic formations
Physiography: Broad, nearly level plain a few feet above basin of playa
Drainage: Rapid runoff where slope is sufficient and no vegetation occurs. Permeability is slow
Description by: W. G. Harper

HORIZON		DESCRIPTION
S55Calif-36-2-1 A_p	0- 4"	Very pale brown (10YR 7/3, pale brown 5.5/3 moist) clay loam; moderately defined very coarse prismatic; very hard when dry, firm when moist; very strongly calcareous; slowly permeable; pH on 1 to 5 dilution with thymol blue 8.6; plant roots common; diffused lower boundary.
S55Calif-36-2-2 C_1	4- 9"	Very pale brown (10YR 7/3 dry, pale brown 6/3 moist) clay; weakly defined very coarse prismatic; very hard when dry; firm when moist; very strongly calcareous; slowly permeable; pH on dilution 9.0; few plant roots; abrupt and smooth lower boundary.
S55Calif-36-2-3 C_2	9-16"	Very pale brown (10YR 7/3 dry, 7/2 crushed, light grayish brown 6/2 moist) clay; weak fine subangular blocky; slightly hard when dry, friable when moist; very strongly calcareous with about 25 percent of surface mottled by lime or gypsum; few soft lime nodules; moderately slow to moderate permeability; few fine plant roots; pH on dilution 8.0; clear and smooth lower boundary.
S55Calif-36-2-4 C_{ca1}	16-23"	A mixture of white and light grayish brown (10YR 8/2 and 6/2 dry, light gray 7/2 moist) fine nodular clay; weak fine to medium subangular blocky; very hard when dry, friable when moist; very strongly calcareous with many lime nodules ranging up to 1/2 inch in diameter; moderate to moderately slow permeability; pH on dilution 8.0; practically devoid of plant roots; clear and smooth lower boundary.
S55Calif-36-2-5 C_{ca2}	23-30"	A mixture of white and light grayish brown (10YR 8/2 and 6/2 dry, pale brown 6.5/3 moist) gravelly or nodular clay; moderately medium subangular blocky and nodular; very hard when dry, friable when moist; moderate to moderately slow permeability; very strongly calcareous with common to many lime nodules ranging up to 1/2 inch in diameter, together with some gypsum; pH on dilution 8.0; practically no plant roots; clear and smooth lower boundary.
S55Calif-36-2-6 C_{ca3}	30-46"	A mixture of white and light brownish gray (10YR 8/2 and 6/2 dry, 6/2 moist) clay; moderately defined medium subangular blocky; very hard when dry, friable when moist;

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Morales silty clay loam (taxadjunct) **LOCATION** San Bernardino County, California

SOIL FAMILY: Typic Calciorthid, fine, montmorillonitic, thermic.

SOIL NOS. 855Calif-36-4 **LAB. NOS.** 5526 thru 5531

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE SAND	COARSE SAND	MEDIUM SAND	FINE SAND	VERY FINE SAND	SILT	CLAY			> 2	

MORALES SILTY CLAY LOAM (taxadjunct) 1/
S55Calif-36-4

Sampled by: Fred Eggers, W. G. Harper, Reuben Nelson, January, 1955.
Description by: W. G. Harper

Location: 250 ft. E and 250 ft. N of W1/4 Cor., Sec. 30, T5N, R1E, or about 300 ft. NE of Schieldmyer's pump, about 3 miles NE of Lucerne Valley, San Bernardino County, California. Elevation: About 3000 feet.
Classification: Typic Calciorthid, fine, montmorillonitic, thermic.
California Group: III
Vegetation: Shadscale 10 to 15 inches high. 98 percent of surface barren.
Parent Material: Alluvium from mixed rocks but mainly from granitic formations
Physiography: Playa - nearly level
Drainage: Little or no runoff, very slow permeability. 50 feet to water table

HORIZON	DESCRIPTION
S55Calif-36-4-1 C1sa 0- 3"	Very pale brown (10YR 7/3, pale brown 6/3 moist) weakly vesicular and moderately fine platy; hard when dry, friable when moist; very strongly calcareous; slowly permeable; pH on 1 to 5 dilution with thymol blue 8.2; few fine plant roots; lower boundary abrupt and smooth.
S55Calif-36-4-2 C2sa 3- 6"	Pale brown (10YR 6/3 dry, brown 5/3 moist) silty clay loam; weakly defined medium to fine granular; hard when dry, friable when moist; very strongly calcareous and with many very fine white gypsum and salt crystals; slowly permeable; pH on dilution 8.0; very few fine plant roots; lower boundary abrupt and smooth.
S55Calif-36-4-3 C3sa 6-16"	Same color as layer above; silty clay; moderately medium subangular blocky; very hard but easily crumbled when dry, firm when moist; slowly permeable; very strongly calcareous with many white lime mottlings and salt or gypsum crystals; pH on dilution 8.2; almost devoid of plant roots; lower boundary clear and smooth.
S55Calif-36-4-4 C4sa 16-27"	Brown (10YR 5/3 moist) silty clay; moderately defined coarse to medium subangular blocky; extremely hard when dry, firm when moist; very strongly calcareous with many medium and fine lime mottlings and/or salt and gypsum crystals; pH on dilution 8.3; slowly permeable; devoid of plant roots; lower boundary clear and smooth.
S55Calif-36-4-5 C5 27-45"	Pale brown (10YR 6/3 moist) clay; massive; extremely hard when dry, very firm when moist; very strongly calcareous with a few medium white lime mottlings; very slowly permeable; pH on dilution 9.6; devoid of plant roots; lower boundary clear and smooth.
S55Calif-36-4-6 C6 45-60"	Slightly finer in texture. Otherwise similar to layer above.

Remarks: This is a strongly saline Alluvial soil. In nearby locations on Mr. Schieldmyer's farm, the distribution of gypsum is erratic ranging in

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aquic Calciorthid, coarse-loamy, mixed, mesic

SOIL Balman loam (taxadjunct) SOIL Nos. S61Calif-46-4 LOCATION Sierra Valley, Sierra County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 61668 - 61675

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt (0.05-0.02)	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.2)			Coarse fragments 2A2 ≥ 2 Pct.	2 - 19	19 - 76
		Pct. of < 2 mm															Pct. of < 76 mm	
0-2	Ap1ca	30.2	42.7	27.1	0.8	3.7	3.6	12.5	9.6	11.8	30.9	29.8	12.2			2		
2-11	Ap2ca	27.5	43.4	29.1	0.9	2.9	3.0	11.1	9.6	13.4	30.0	30.5	10.4			0		
11-18	ACca	35.7	37.9	26.4	1.1	4.3	4.2	14.7	11.4	11.4	26.5	32.6	14.5			0		
18-27	IIC1	60.6	29.8	9.6	1.3	9.8	8.6	24.6	16.3	12.7	17.1	44.8	28.5			2		
27-37	IIC2	67.6	18.0	14.4	3.0	11.0	9.8	28.3	15.5	8.8	9.2	42.4	34.0			6		
37-48	IIC3	73.1	16.2	10.7	4.8	11.9	10.6	29.5	16.3	7.8	8.4	42.6	38.3			7		
48-63	IIC4	74.8	14.1	11.1	9.1	18.5	12.7	24.3	10.2	6.9	7.2	30.8	50.9			16		
63-75+	IIIC5	28.8	38.5	32.7	0.6	2.1	2.5	9.2	14.4	14.5	24.0	35.6	7.7			0		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		6E1b Carbonate as $CaCO_3$ Pct.	6C2a Ext. Iron as Fe Pct.	Bulk density g/cc	Bulk density g/cc	Bulk density g/cc		Water content Pct.	Water content Pct.	4B2 15 bar Pct.			pH 8C1b Sat. Paste	pH 8C1a 1:10
0-2	3.26	0.319	10		27	0.1							20.1			8.3	8.7
2-11	3.00	0.293	10		29	0.1							21.0			8.6	9.2
11-18	1.35	0.136	10		35	0.1							22.2			8.0	8.7
18-27	0.25				5	0.5							9.5			7.9	8.7
27-37	0.11				2	0.6							9.0			7.7	8.7
37-48	0.05				1	0.7							7.8			7.8	8.5
48-63	0.04				2	0.7							8.7			7.9	8.6
63-75+	0.04				tr.	0.3							15.7			7.4	8.4

Depth (in.)	Extractable bases 5B1a: 6N2c Ca (meq/100 g)	6O2c Mg	6P2a Na	6Q2a K	6H2a Ext. Acidity	Cat. Exch. Cap. 5A2a NaOAc	5B1b Exch. Na	Water extract from saturated paste 8A1: 6N1a Ca (meq/liter)	6O1a Mg	6P1a Na	6Q1a K	6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity mmho/cm
0-2	31.0	9.2	1.0	2.1	-	32.0	0.8	2.6	1.8	3.1	2.5	1.2	6.3	0.4		0.72
2-11	23.1	10.7	4.7	0.8	-	31.1	3.9	1.7	1.8	13.7	0.2	1.6	8.1	2.4		1.53
11-18	22.2	14.1	3.5	0.1	-	28.4	2.3	8.8	15.4	20.3	-	0.8	2.7	23.4		4.15
18-27	22.5	10.0	2.0	0.2	-	22.6	1.6	12.9	19.3	13.0	0.1	0.5	1.6	27.5		4.08
27-37	18.2	8.3	1.1	0.2	-	20.6	0.8	16.4	17.2	9.3	0.1	-	1.6	34.3		3.64
37-48	14.1	6.3	0.7	0.2	0.1	17.5	0.5	5.5	5.6	4.3	0.1	-	1.8	10.6		1.62
48-63	16.9	5.9	0.5	0.2	0.2	18.5	0.4	2.2	1.7	1.9	0.1	-	2.5	1.9		0.60
63-75+	20.2	8.5	0.5	0.4	0.3	26.6	0.4	1.6	1.2	1.4	0.1	-	2.8	0.4		0.35

Depth (in.)	8A Water at Saturation Pct.	5D1 Exchangeable Na Pct.		Gypsum Pct.			
0-2	59.1	2					
2-11	60.4	12					
11-18	59.8	8					
18-27	32.4	7					
27-37	31.9	4					
37-48	28.0	3					
48-63	28.6	2					
63-75+	43.5	1					

BALMAN LOAM (taxadjunct) 1/

S61Calif-46-4

Location: Sierra Valley, Sierra County, California; 0.25 mile south of N1/4 corner, 100 yards east of Hariot Lane, section 4, T. 21 N., R. 15 E. Date of Sampling: September 13, 1961.
Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, James McLaughlin, Eugene Begg.
Vegetation: Cropped; now planted to dryland seeding of tall wheat grass. Climate: Estimated annual precipitation 14 inches. Nearest weather station (Sierraville): Mean annual temperature 46° F.; January average 28.2° F.; July average 63.6° F.; mean annual precipitation 25.78 inches.
Parent Material: Somewhat stratified medium textured alluvium from mixed sources. Relief: Old lake terrace mound with slightly hummocky topography and less than 1 percent slope. Elevation: 4885 feet.
Drainage: Moderately well drained; slow runoff, moderately slowly permeable. Soil Moisture: Slightly moist below 27 inches.
Remarks: This is a Calcium Carbonate Solonchak in the Chestnut soil zone. The surface 1/4 inch tends to be crusted. Surface appears to seal. **Classification: Aquic Calciorthid, coarse-loamy, mixed, mesic.**

HORIZON	DESCRIPTION
Aplca RSL No. 61668	0 to 2 inches, light gray (10YR 7/1) loam, dark gray (10YR 4/1) when moist; weak coarse platy structure; soft, very friable, slightly sticky and slightly plastic; abundant very fine roots below surface crust; common very fine tubular and interstitial pores; moderately alkaline (pH 8.3)(8.2 with thymol blue); violently effervescent, lime disseminated; abrupt, smooth boundary.
Ap2ca RSL No. 61669	2 to 11 inches, gray (10YR 6/1) loam, dark gray (10YR 4/1) when moist; weak medium platy structure; soft, very friable, slightly sticky and slightly plastic; abundant very fine, many fine, few medium roots; many very fine tubular and common very fine interstitial pores; moderately alkaline (pH 8.3)(8.8 with thymol blue); violently effervescent, lime disseminated; clear, smooth boundary.
ACca RSL No. 61670	11 to 18 inches, light gray (10YR 7/1) loam, dark gray (10YR 4/1) when moist; weak coarse platy breaking to moderate medium subangular blocky structure; slightly hard, friable, slightly sticky and slightly plastic; plentiful very fine and fine, few medium roots; common very fine tubular and interstitial, few fine and medium tubular pores; moderately alkaline (pH 8.3)(pH 8.0 with thymol blue); violently (?) effervescent, lime disseminated; clear, smooth boundary.
IIC1 RSL No. 61671	18 to 27 inches, light gray (2.5Y 7/2) sandy loam, brown (10YR 4/3) when moist; massive; slightly hard, slightly firm, slightly sticky and slightly plastic; plentiful very fine and fine, few medium roots; common very fine tubular and interstitial, few fine and medium tubular pores; moderately alkaline (pH 8.2)(pH 8.0 with thymol blue); strongly effervescent, lime disseminated; clear, smooth boundary.
IIC2 RSL No. 61672	27 to 37 inches, light brownish gray (10YR 6/2) sandy clay loam (near sandy loam) with white segregated lime (10YR 8/2), dark grayish brown (10YR 4/2) when moist; massive; hard, firm, slightly sticky and slightly plastic; few very fine, plentiful fine and medium roots; common very fine tubular and interstitial, few fine to medium tubular pores; few thin clay films in pores; few dark concretions; moderately alkaline (pH 8.2)(pH 8.0 with thymol blue); strongly effervescent, lime occurs in seams; gradual, smooth boundary.
IIC3 RSL No. 61673	37 to 48 inches, light gray (10YR 7/2) sandy loam with white (10YR 8/2) lime in seams, dark brown (10YR 4/3) when moist; massive; slightly hard, slightly firm, nonsticky and slightly plastic; few very fine, plentiful fine to medium roots; common very fine tubular and interstitial, few fine and medium tubular pores; mildly alkaline (pH 8.2)(pH 8.0 with thymol blue); strongly effervescent, lime occurs in seams; clear, irregular boundary.
IIC4 RSL No. 61674	48 to 63 inches, very pale brown (10YR 7/3) sandy loam, dark yellowish brown (10YR 4/4) when moist; massive; soft, friable, nonsticky and nonplastic; few very fine, plentiful fine to medium roots, many very fine interstitial, common very fine and few fine to medium tubular pores; moderately alkaline (pH 8.2)(pH 8.2 with thymol blue); strongly effervescent, lime disseminated; abrupt, slightly wavy boundary.
IIIC5 RSL No. 61675	63 to 73 inches, pale yellow (5Y 7/3) and white (5Y 8/2) silty clay lake sediments, dark grayish brown (2.5Y 4/2) with common fine distinct dark brown (10YR 3/3) mottles when moist; massive; very hard, very firm, sticky and plastic; no roots, pores or clay films observed; moderately alkaline (pH 8.0)(pH 8.4 with thymol blue); strongly effervescent, lime occurs in soft masses.

1/ This pedon has less clay in the 10 to 40 inch control section than required for the Balman

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aquic Calciorthid, fine-loamy, mixed, mesic

SOIL Balman loam SOIL Nos. S61Calif-32-12 LOCATION Sierra Valley, Plumas County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 61676 - 61685

Depth	Horizon	1B1b Total Sand	Total Silt	Total Clay	Sand Very	Sand Coarse	Sand Medium	Sand Fine	Sand Very fine	Silt	Silt Int. III	Size class and particle diameter (mm) 3A1 Int. II (0.2-0.02)	(2-0.2)			Coarse fragments 2A2 > 2	2 - 19	19 -76

BALMAN LOAM
S61Calif-32-12

Location: Sierra Valley, Plumas County, California, Southeast corner SW1/4NW1/4 section 35, T. 22 N., R. 15 E. Approximately 2-3/4 miles north and 1-3/4 miles west of Loyalton.
Date of Sampling: September 13, 1961.
Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, James McLaughlin, Eugene Begg.
Vegetation: Cropland, now planted to dryland seeding of tall wheat grass. Climate: Estimated annual precipitation 13 inches. Nearest weather station (Sierraville): Mean annual temperature 46° F.; January average 28.2° F.; July average 63.6° F.; mean annual precipitation 25.78 inches.
Parent Material: Stratified medium textured alluvium from mixed sources. Relief: Old lake terrace mound, slightly hummocky, slope less than 1 percent. Elevation: 4890 feet. Drainage: Moderately well drained, runoff slow, permeability moderately slow. Soil Moisture: Slightly moist below 24 inches, moist at 43 inches.
Remarks: This soil is a Calcium Carbonate Solonchak in the Chestnut soil zone. It often occurs in complex with Ramelli silty clay loam, the Ramelli occupying the swales and lower topography and the Hovey the mounds on higher ground. The C2ca has dark stains in pores.
Classification: Aquic Calciorthid, fine-loamy, mixed, mesic.

HORIZON	DESCRIPTION
Ap1ca RSL No. 61676	0 to 2½ inches, light brownish gray (10YR 6/2) loam, very dark grayish brown (10YR 3/2) when moist; moderate medium platy structure; soft, very friable, slightly sticky and slightly plastic; abundant very fine, few fine and medium roots; many very fine mostly horizontal tubular and common fine interstitial pores; moderately alkaline (pH 8.4)(pH 9.2 with thymol blue); violently effervescent, lime disseminated; clear, smooth boundary.
Ap2ca RSL No. 61677	2½ to 10 inches, gray (10YR 6/1) loam, very dark grayish brown (10YR 3/2) when moist; moderately fine platy structure; soft, very friable, slightly sticky and slightly plastic; abundant, very fine, plentiful fine and medium roots; abundant very fine, few fine and medium tubular and common fine interstitial pores, moderately alkaline (pH 8.4)(pH 9.0 with thymol blue); violently (?) effervescent, lime disseminated; clear, smooth boundary.
ACca RSL No. 61678	10 to 17 inches, gray (10YR 6/1) loam, very dark grayish brown (10YR 3/2) when moist; massive breaking to weak medium and coarse subangular blocky structure; soft, friable to firm, slightly sticky and slightly plastic; abundant very fine, common fine, few medium roots; many very fine and fine, few medium tubular and few very fine interstitial pores; few thin clay films; moderately alkaline (pH 8.0)(pH 8.6 with thymol blue); violently effervescent, lime disseminated; clear, smooth boundary.
C1ca RSL No. 61679	17 to 22 inches, gray (10YR 6/1) loam, very dark grayish brown (10YR 3/2) when moist; massive breaking to weak medium and coarse subangular blocky structure; slightly hard, firm, slightly sticky and slightly plastic; plentiful very fine, few fine roots; common very fine, few fine and medium tubular pores, few very fine interstitial pores; few thin clay films in pores; moderately alkaline (pH 8.0)(pH 8.2 with thymol blue); violently effervescent, lime disseminated; clear, smooth boundary.
C2ca RSL No. 61680	22 to 30 inches, gray (10YR 6/1) loam, very dark brown (10YR 2/2) when moist; massive breaking to weak medium and coarse subangular blocky structure; hard, very firm, slightly sticky and slightly plastic; few very fine and fine roots; common very fine to medium tubular, common very fine, few fine interstitial pores; common thin clay films in pores; moderately alkaline (pH 8.0) (pH 8.2 with thymol blue); violently effervescent, lime disseminated; gradual, smooth boundary.
C3ca RSL No. 61681	30 to 42 inches, light gray (10YR 7/2) sandy loam, dark grayish brown (10YR 4/2) when moist; massive breaking to weak medium and coarse subangular blocky structure; hard, firm, slightly sticky and slightly plastic; very few very fine and fine roots; many very fine, common fine and medium tubular and common very fine and few fine interstitial pores; common moderately thick

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aquic Duric Calciorthid, coarse-loamy, mixed, mesic
SOIL Kisring silt loam (taxadjunct) SOIL Nos. S56Calif-18-4 LOCATION Lassen County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 55506 - 55515

Depth (In)	Horizon	1B1b Total Sand (2-0 05)	Total Silt (0 05-0.002)	Total Clay (< 0.002)	Sand Very coarse (2-1)	Sand Coarse (1-0 5)	Sand Medium (0 5-0.25)	Sand Fine (0.25-0.1)	Sand Very fine (0 1-0.05)	Silt 0.05-0.02	Silt Int. III (0.02-0.002)	Int. II (0.2-0.02)	3A1 (2-0.1)			Coarse fragments 2A2 > 2	2 - 19	19 -76
		Pct. of < 2 mm														Pct	Pct. of < 76 mm	
0-1	Av1	33.4	56.9	9.7	0.4	1.9	2.3	12.0	16.8	22.1	34.8	47.8	16.6			1		
1-2	Av2	26.4	59.6	14.0	0.3	1.3	1.9	9.4	13.5	19.7	39.9	40.0	12.9			0		
2-4	B21	38.3	51.3	10.4	0.3	1.2	1.3	13.0	22.5	19.0	32.3	52.7	15.8			0		
4-6	B22	42.0	46.2	11.8	0.4	0.9	1.1	17.5	22.1	18.0	28.2	54.8	19.9			0		
6-11	B3	32.2	57.0	10.8	1.1	1.6	0.9	8.9	19.7	18.9	38.1	46.1	12.5			0		
11-20	C1	30.4	50.5	19.1	5.9	6.2	2.9	6.9	8.5	13.0	37.5	26.1	21.9			0		
20-31	C2	33.0	47.7	19.3	5.0	8.7	4.3	8.6	6.4	11.0	36.7	22.8	26.6			0		
31-37	C3	30.1	57.2	12.7	3.0	5.6	3.4	9.1	9.0	16.1	41.1	31.0	21.1			0		
37-49	C4	25.6	56.5	17.9	6.2	4.6	2.6	5.3	6.9	14.5	42.0	24.8	18.7			0		
49-60+	Cu	8.2	72.7	19.1	1.4	1.5	0.8	2.0	2.5	10.6	62.1	14.4	5.7			2		

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		6E1b Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct	Pct	4B2 15 bar Pct			pH 8C1b Sat. Paste	8C1a 1:10
0-1	0.40	0.043	9.3		3								8.2			8.8	9.7
1-2	0.35	0.038	9.2		5								12.9			9.1	9.9
2-4	0.33	0.040	8.3		7								26.4			9.9	10.0
4-6	0.35				6								32.2			10.1	10.1
6-11	0.19				8								33.1			10.2	10.2
11-20	0.16				18								29.3			10.2	10.2
20-31	0.15				17								29.6			9.7	10.1
31-37	0.12				20								26.3			9.1	9.7
37-49	0.09				11								27.1			8.8	9.6
49-60+	0.12				3								27.5			8.8	9.4

Depth (In)	Extractable bases Ca	Mg	5B1b Exch. Na 6P2a	5B1b Exch. K 6Q2a		5A2a CEC NaOAc			Water extract from saturated paste 6N1a 6O1a Ca + Mg	Mg	6P1a Na	6Q1a K		CO_3	HCO_3	Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-1			12.9	3.6		21.2			1.1		26.0	0.5						2.4
1-2			23.6	3.0		29.6			0.5		43.5	0.7						4.5
2-4			44.1	7.6		53.6			0.4		97.0	1.0						8.5
4-6			49.2	7.2		61.0			0.7		142.0	1.3						11.9
6-11			56.1	6.5		63.0			1.7		156.0	1.2						12.3
11-20			53.0	5.0		59.4			0.5		109.0	0.8						9.4
20-31			42.1	4.4		54.2			0.4		45.5	0.3						4.4
31-37			32.9	3.6		48.3			0.5		18.8	0.2						1.7
37-49			30.1	3.1		49.2			0.5		13.1	0.1						1.3
49-60+			25.0	2.4		47.4			0.5		11.0	0.1						1.1

Depth (In)	8A Water at Saturation Pct.	5D1 Exchangeable Na Pct		6F1a Gypsum Pct.			
0-1	25.6	61		-			
1-2	32.9	80		-			
2-4	65.5	82		-			
4-6	68.4	81		-			
6-11	73.6	89		-			
11-20	68.3	89		-			
20-31	73.7	78		-			
31-37	72.1	68		-			
37-49	71.8	61		-			
49-60+	116.4	53		-			

KISRING SILT LOAM (taxadjunct) 1/
S55Calif-18-4

Location: About 8 miles east of Susanville. 41/100 mile east of west quarter-corner of Section 3, T. 29 N., R. 13 E., and 400 feet north of east-west graveled road; Honey Lake Valley SCD, Lassen County, California. Date of Sampling: August, 1955.
Description by: W. M. Johnson. Collectors: S. B. Johnson, Ralph E. Nelson, Stuart Porter, Reuben E. Nelson, John Cady, E. A. Naphan, and W. M. Johnson.
Classification: Aquic Duric Calciorthid, coarse-loamy, mixed, mesic.
Vegetation: Shrub-grass with many barren patches. Total density, about 15 percent. Composition: Big sage, 30 percent; greasewood, 30 percent; rabbitbrush, 37 percent; giant and blue wild rye, 1 percent; salt grass, 1 percent; lupine and poverty weed, 1 percent; squirreltail and Indian rice grass, trace. Forage production about 100 to 200 pounds per acre. Climate: At Susanville the January average temperature is 30.7° F., the July average is 69.5° F., and the frost-free period averages 142 days. Summers are hot and dry, winters are cold and snowy. Mean annual rainfall at Susanville is 17.8 inches, most of which falls during the period October to May. At the site where samples were taken it is estimated that the mean annual rainfall is 8 to 10 inches. Parent Material: Calcareous silty alluvium, probably derived from the reworking of Lake Lahontan sediments. The lowest horizon may be the bedded lacustrine sediments. Use: Pasture in poor condition. Relief: Smooth, gently sloping basin rim or old basin floor of ancient Lake Lahontan. There is microrelief of 3-6 inches, probably due to wind erosion and deposition. High spots are around shrubs. Slope is about 1 percent, plane, toward the northeast. Elevation: About 4,100 feet. Drainage: Runoff is slow and permeability is moderate or moderately slow. Soil is probably imperfectly drained. It gives the appearance of being influenced by relatively high water table during a part of the year. At the time the profile was examined, there was free water at a depth of 84 inches. Other Features: The profile was dry in the Av, moist below to a depth of 49 inches, and very moist below 49 inches. The surface of the soil is characterized by many barren "slick spots", some of which have small patches of salt crust. This soil occurs in complex with a similar soil having sandy substratum, with a Canby-like soil, and with a dark-colored, more strongly saline soil. Reaction: Free carbonates were estimated by observing reaction to dilute HCl. pH's were determined with thymol blue in about 1:5 soil-indicator dilution.

HORIZON	DESCRIPTION
Av1 RSL No. 55506	0 to 1 inch, light brownish gray (10YR 6.5/2, dry; 4/2, moist) silt loam. Very weak thick platy, nearly massive. Soft, friable, slightly plastic, slightly sticky. Roots are numerous, but do not branch in this horizon. Very porous--strongly vesicular. Moderately calcareous. pH 9.0. Abrupt, smooth lower boundary.
Av2 RSL No. 55507	1 to 2 inches, light brownish gray (10YR 6.5/2, dry; 4/2, moist) silt loam. Moderate medium platy; soft, friable, slightly plastic, slightly sticky. Moderately porous--vesicular. Roots are numerous but do not branch in this horizon. Moderately calcareous. pH 9.4. Abrupt, smooth lower boundary.
B21 RSL No. 55508	2 to 4 inches, pale brown (10YR 6/3, dry; 3.5/3, moist) silty clay loam. Weak medium prismatic, breaking to strong fine granular. Firm, very plastic, very sticky. Moderate continuous clay skins. Roots profuse. Peds are nonporous. Moderately to strongly calcareous, the lime being disseminated and in tiny soft segregations. pH 9.8. Clear, slightly wavy lower boundary.
B22 RSL No. 55509	4 to 6 inches, pale brown (10YR 5.5/3, dry; 4/3.5, moist) silty clay loam. Weak medium prismatic, breaking to strong medium and fine granular. Firm, very plastic, very sticky. Moderate continuous clay skins. Roots profuse. Peds are nonporous. Moderately to strongly calcareous, the carbonates being disseminated and in tiny soft segregations. pH 9.6. Clear, slightly wavy lower boundary.
B3 RSL No. 55510	6 to 11 inches, pale brown (1Y 6/3, dry; 4.5/3, moist) light silty clay loam. Weak medium subangular blocky, breaking to moderate medium granular. Friable, plastic, very sticky. Moderate, nearly continuous clay skins. Roots are numerous. Peds are nonporous. Strongly calcareous, the lime being mainly disseminated. pH 9.6. Clear, wavy lower boundary.
C1 RSL no. 55511	11 to 20 inches, pale brown (1Y 6/3, dry; 4.5/3, moist) light silty clay loam with common medium and fine mottles of white. Moderate medium subangular blocky. Friable, with many firm peds; plastic, sticky. Few roots. Peds are slightly porous. Moderate, nearly continuous clay skins. Strongly calcareous, the lime existing in disseminated form and in common fine soft segregations. pH 9.6. Gradual, wavy lower boundary.
C2 RSL No. 55512	20 to 31 inches, light gray (10YR 7.5/2, dry; 5/3, moist) light silty clay loam with common medium mottles of white and a few brown specks (dry). Weak medium subangular blocky. Firm, with a few friable patches; plastic, sticky. Numerous distinct patches of clay skins. Very few roots. Peds are slightly porous. Very strongly calcareous; lime is disseminated and in common fine soft segregations. pH 9.6. Gradual, wavy lower boundary.
C3 RSL No. 55513	31 to 37 inches, white (10YR 8/2, dry; 5/2, moist) silt loam with many fine mottles of strong brown (dry). Very weak medium subangular blocky. Friable, with many very firm peds or lumps of lacustrine sediment (cemented ?). A very few thin patches of clay skins. Very few roots. Reaction to HCl varies from none to very strong within short distances in this horizon. pH 9.4. Gradual, broken lower boundary.
C4 RSL No. 55514	37 to 49 inches, distinctly mottled pale yellow and dark grayish brown (2.5Y 7/3 and 10YR 4.5/2, dry; 2.5Y 5/3 and 10YR 2/2, moist) silt loam. Massive. Firm, with many irregular lumps of material that appear to be very firm "alkali pan". No roots observed. No clay skins observed. Porous. Moderately to strongly calcareous. pH 9.1. Clear, wavy lower boundary.
Cu RSL No. 55515	49 to 60 inches +, white (2.5Y 8/2, dry; 6/2, moist) silt loam with many thin horizontal bands of reddish yellow (dry). Massive--laminated. Friable, plastic, sticky. This appears to be a substratum of lacustrine sediments in place. No roots observed. Slightly porous. Slightly calcareous. pH 8.9.

1/ The Kisring series is in a fine-loamy, mixed, mesic family of Aquic Duric Calciorthids. This pedon has less than 18 percent clay in the control section.

SOIL SURVEY LABORATORY Riverside, California

OIL TYPE Laveen sandy clay loam (taxadjunct) **LOCATION** San Bernardino County, California

SOIL FAMILY: Typic Camborthid, coarse-loamy, mixed, thermic.

SOIL NOS. 855Calif-36-9 **LAB. NOS.** 5560 thru 5567

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 2	A	25.9	25.2	12.5	21.1	6.9	6.1	2.3	24.5	0.6	9	cos
2- 8	B2	15.0	20.1	11.5	20.6	7.1	10.1	15.6	24.7	4.6	15	sl
8-17	B2ca	4.4	12.6	12.6	32.6	11.8	13.4	12.6	39.8	4.9	6	fsl
17-25	C1	6.5	14.0	9.8	16.3	8.8	25.0	19.6	30.2	13.1	5	sl
25-31	C2	5.5	16.0	14.9	29.6	11.9	13.9	8.2	36.4	6.2	1	sl
31-38	C3	6.3	20.2	14.6	20.9	6.8	19.5	11.7	29.1	9.1	1	cosl
38-51	C4	15.3	34.4	19.5	17.8	3.9	4.7	4.4	13.0	3.6	5	cos
51-60	C5	35.5	25.6	6.1	7.6	3.9	10.2	11.1	11.4	6.7	17	cosl

8C1b pH SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	6C1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM meq./100g. SOIL	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
8.6		9.0	0.05	0.007			0.85	0	0		3.8	1.2
8.7		9.4	0.04	0.006			19.8	2.0	0			5.7
9.3		9.7	0.02				15.7	4.6	0			5.7
9.3		9.6	0.01				17.5	1.6	0			9.5
9.3		9.4	0.00				13.9	0	0			4.7
9.4		9.5	0.01				15.5	0.6	0			6.5
9.5		9.4	0.10				11.1	0	0			3.3
9.5		9.7	0.02				16.9	0.9	0			6.5

5A2a CATION EXCHANGE CAPACITY (+ Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	5B Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	5D1 Exch. Na %	SATURATION EXTRACT SOLUBLE 6P1a Na	6Q1a K	6N1a Ca+Mg 6O1a	8A1	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
3.7				0.7	0.3	19	9.5	0.2	1.3			18.4
15.3				8.8	0.5	58	238	1.3	7.6			25.1
13.8				8.4	0.2	61	193	1.3	2.4			30.2
21.9				15.5	0.2	71	218	1.3	2.0			51.5
12.1				9.8	0.1	81	158	1.3	1.0			38.4
17.5				13.6	0.1	78	180	1.3	1.3			42.3
4.5				3.1	0.2	69	173	1.3	1.0			28.2
14.0				9.6	Tr	69	203	1.3	0.6			26.4

1/ Percent of the whole soil

LAVEEN SANDY CLAY LOAM (taxadjunct) 1/
S55Calif-36-9

Sampled by: Fred Eggers, W. G. Harper, Reuben E. Nelson.
Location: 300 ft. S and 40 ft. W of the NE cor., Sec. 1, T8N, R3E, about 4 mi. E of Newberry, San Bernardino County, California.
Elevation: About 1800 ft. Annual precipitation about 4 inches
Description by: W. G. Harper. Sampling date: January, 1955.
Classification: Typic Camborthid, coarse-loamy, mixed, thermic.
Vegetation: Shadscale, creosote, mesquite bushes on nearby sand dunes, and desert sage. About 70 percent of surface barren
Parent Material: Alluvium from mixed rocks but mainly from granitic formations.
Physiography: Valley plain - less than 1 percent slope
Drainage: Water table is about 16 ft. below surface at present. Runoff is nil. Permeability of profile moderate to moderately slow

HORIZON		DESCRIPTION
S55Calif-36-9-1 A	0-2"	Pale brown (10YR 6/3 dry, dark grayish brown 4/2 moist) coarse sand; single grain; soft when dry, loose when moist; noncalcareous; rapidly permeable; pH at 1 to 5 dilution 8.4; no plant roots apparent; lower boundary abrupt and smooth.
S55Calif-36-9-2 B_2	2-8"	Reddish brown (5YR 5/4 dry, 4/4 moist) with a few fine faint iron mottlings; sandy loam; moderately defined coarse prismatic; hard when dry, friable when moist; micaceous; mildly calcareous and strongly calcareous where occasional medium flecks of lime occur; pH on dilution 9.6; few fine plant roots; moderately permeable.
S55Calif-36-9-3 B_{2ca}	8-17"	Light brown (7.5YR 6/4 dry, brown 5/4 moist) fine sandy loam with common medium faint (5YR 4/4) iron mottlings and common medium (10YR 8/4) lime flecks; very weak [illegible]

OIL SURVEY LABORATORY Riverside, California

OIL TYPE Milham sandy clay loam (taxadjunct) **LOCATION** San Bernardino County, California

SOIL FAMILY: Typic Camborthid, fine-loamy, mixed, thermic.

SOIL NOS. S55Calif-36-10 **LAB. NOS.** 5568 thru 5572

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 6	A	20.0	23.6	13.1	27.3	9.2	3.8	3.0	27.7	2.4	5	cos
6-11	B2	5.3	16.4	18.9	38.8	5.7	3.3	11.6	26.1	3.5	4	lfs
11-16	B2ca	6.2	11.0	17.2	47.5	6.5	3.5	8.1	32.8	2.0	15	lfs
16-36	D1	9.1	14.9	7.6	12.9	7.5	22.7	25.3	20.7	17.0	3	scl
36-57	D2	1.9	6.9	8.3	23.6	20.1	27.1	12.1	48.4	13.8	1	fsl

8C1b pH SATURATED PASTE 1:1	8C1a pH 1:5	8C1a pH 1:10	6A1a ORGANIC MATTER ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8F1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	4B2 MOISTURE TENSIONS 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
8.4		8.5	0.08	0.006			1.3-	0	0		3.5	1.6
8.4		9.4	0.07	0.007			11.7	3.4	0		13.9	4.8
8.8		9.5	0.04				12.8	1.4	0		10.8	4.0
8.5		9.1	0.03				20.0	0.9	2.3			12.4
8.7		9.4	0.02				17.4	0	0			6.7

MILHAM SANDY CLAY LOAM (taxadjunct) 1/
S55Calif-36-10

Sampled by: Fred Eggers, W. G. Harper, Reuben E. Nelson.
Location: 200 ft. S and 250 ft. W of NE cor., Sec. 36, T9N, R3E - about 4 mi. E of Newberry, San Bernardino County, California.
Elevation: About 1800 ft. Annual precipitation about 4 inches
Description by: W. G. Harper. Sampling date: January, 1955.
Classification: Typic Camborthid, fine-loamy, mixed, thermic.
California Group: III
Vegetation: Desert sage, shadscale, creosote bushes and rabbitbrush occur on occasional dunes that are 5 to 6 feet high and about 250 ft. long - 80 percent of surface is barren.
Parent Material: Alluvium from mixed rocks but mainly from granitic formations.
Physiography: Valley plain less than 1 percent slope, with occasional dunes 6 to 8 feet high
Drainage: No runoff, no surface drainage channels, moderate to slow internal permeability

HORIZON	DESCRIPTION
S55Calif-36-10-1 A 0- 6"	Light yellowish brown (10YR 6/4 dry, grayish brown 4.5/2 moist) coarse sand; single grain; soft when dry, loose when moist; noncalcareous; rapidly permeable; pH at 1 to 5 dilution with thymol blue 8.2; almost devoid of plant roots; lower boundary abrupt and smooth.
S55Calif-36-10-2 B_2 6-11"	Light reddish brown (5YR 6/4 dry, reddish brown 5/4 moist) loamy fine sand with common, medium to coarse prominent white lime mottling; hard when dry, friable when moist; mildly calcareous, but strongly calcareous on the white lime flecks and spots; moderately permeable; pH on dilution 9.4; almost devoid of plant roots; lower boundary clear and smooth.
S55Calif-36-10-3 B_{2ca} 11-16"	Pinkish gray (7.5YR 6/2 dry, brown 5/4 moist) fine gravelly loamy sand with common, fine to medium prominent white lime veins and flecks; hard when dry, friable when moist; strongly calcareous; moderately permeable and almost devoid of plant roots; pH on dilution 9.6; lower boundary abrupt and smooth.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Salorthid, very-fine, montmorillonitic, thermic.
SOIL Morales clay (taxadjunct) SOIL Nos. S55Calif-36-3 LOCATION San Bernardino County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 5517 - 5523

Depth (cm)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: (0.05-0.02)	Int. III (0.02-0.002)	Int. II (0.2-0.02)	2-0.1			Coarse fragments 1A2a: > 2 Pct. of whole soil	2-19	19-76
		Size class and particle diameter (mm) 3A1																
		Pct. of <2 mm															Pct. of <76mm	
0-5	C1	12.4	25.4	62.2	0.2	0.5	0.5	3.9	7.3	1.3	24.1	11.9	5.1			1		
5-15	C2sa	10.2	24.7	65.1	0.1	0.4	0.6	3.3	5.8	1.8	22.9	10.2	4.4			0		
15-25	C3sa	8.9	45.4	45.7	0.3	0.5	0.5	2.9	4.7	13.2	32.2	20.1	4.2			0		
25-58	C4sa	6.8	29.5	63.7	0.1	0.2	0.3	1.8	4.4	2.2	27.3	8.0	2.4			0		
58-86	C5sa	4.4	30.5	65.1	0.1	0.3	0.2	1.1	2.7	2.1	28.4	5.6	1.7			0		
86-105	C6sa	5.4	24.0	70.6	0.2	0.4	0.4	1.5	2.9	2.4	21.6	6.5	2.5			0		
105-135	C7	1.9	34.1	64.0	0.0	0.1	0.0	0.5	1.3	2.6	31.5	4.3	0.6			0		

Depth (cm)	6A1a Organic carbon a/ Pct.	6B1a Nitrogen Pct.	C/N	6C2b Ext. iron as Fe Pct.	Carbonate as $CaCO_3$: 6E1b <2 mm Pct.	3A1a <0.002 mm Pct.	Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content: 4B1c 1/3 bar Pct.		4B2 15 bar Pct.	4D1 Extensibility COLEf cm/cm	4D1 Extensibility COLE cm/cm	pH: 8C1b Saturated paste	8C1a H_2O 1:10	8C1c IN KCl 1:1
0-5	0.10	0.013			13.8								14.6			8.6	9.8	
5-15	0.17	0.012			13.6								17.0			8.5	9.1	
15-25	0.20	0.013			13.9								17.7			8.4	9.0	
25-58	0.16	0.015			16.7								18.6			8.4	9.4	
58-86	0.18				21.9								18.9			8.5	9.7	
86-105	0.19				24.0								18.6			8.6	9.7	
105-135	0.13				28.5								18.1			8.6	9.8	

Depth (cm)	Extractable bases: 6N4c Ca 5B3a	6O4c Mg 5B3a	6P2a Exch. Na 5B1b	6Q2a Exch. K 5B1b	6H2a Ext. acidity	5A2a CEC Na OAc	5D1 Exchangeable Sodium Pct.	5E Sodium adsorption ratio	8A Water at saturation Pct.	Water extract from saturated paste 8A1: 6O1a 6N1a Ca + Mg		6P1a Na	6Q1a K	6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1b SO_4	8A1a Electrical conductivity mmho/cm
			meq/100 g							meq/liter								
0-5			20.2	6.2		33.0	61	107	47.7	2.1		110	5.0					10.8
5-15			24.5	6.2		32.8	75	352	43.9	34.4		1460	45.0					92.6
15-25			28.6	5.3		31.3	91	399	44.1	30.6		1560	30.0					99.0
25-58			24.5	6.2		33.7	73	181	53.4	34.2		750	27.0					63.8
58-86			22.6	4.3		35.4	64	154	64.4	13.3		396	10.0					36.4
86-105			26.1	4.0		36.2	72	153	56.4	8.2		310	10.0					30.2
105-135			25.8	3.1		33.3	77	153	59.1	3.9		213	6.8					22.0

Depth (cm)	Ratios to clay 8D1: CEC Na OAc	Ext. iron	15 bar water	Salt Pct.	6F1a Gypsum Pct.
0-5	0.53		0.23	0.3	-
5-15	0.50		0.26	3.9	0.73
15-25	0.68		0.39	4.1	2.06
25-58	0.53		0.29	2.5	0.09
58-86	0.54		0.29	1.6	-
86-105	0.51		0.26	1.1	-
105-135	0.52		0.28	0.8	-

a/ 1.7 kg of organic carbon per square meter to a depth of 1 meter

MORALES CLAY (taxadjunct) 1/
S55Calif-36-3

Location: San Bernardino County, California. About 5 miles northeast of Lucerne Valley, 400 feet west and 200 feet south of center of sec. 24, T. 5 N., R. 1 W. Date of Sampling: January, 1955.
Description by: W. G. Harper. Collectors: Fred Eggers, W. G. Harper, and Reuben E. Nelson.
Classification: Typic Salorthid, very-fine, montmorillonitic, thermic.
Vegetation: Occasional clumps of shad scale (Atriplex canescens), about 24 inches high. 99 percent of surface barren. Climate: Annual precipitation about 4 inches. Parent Material: Alluvium, mainly from granitic rocks. Elevation: About 3000 feet. Physiography: Playa - nearly level. Drainage: Well drained; no runoff, no surface outlet, very slow permeability through soil. Water table about 50 feet. Remarks: Weakly defined very coarse prisms are present in the C2sa and C3sa horizons.

HORIZON	DESCRIPTION
	Colors are for the dry soil unless indicated otherwise.
C1 RSL No. 5517	0-2 inches (0-5 cm.), pale brown (10YR 6/3) clay, yellowish brown (10YR 5/4) moist; weak coarse prisms breaking to strong very fine subangular blocks; very hard (dry), firm (moist); violently effervescent; few pebbles; abrupt smooth boundary.
C2sa RSL No. 5518	2-6 inches (5-15 cm.), light yellowish brown (10YR 6/4) clay, yellowish brown (10YR 5/4) moist; strong very fine granules; soft (dry), very friable (moist); very few roots; 5 to 10 percent by volume of salt crystals up to 3 mm. in diameter; abrupt wavy boundary.
C3sa RSL No. 5519	6-10 inches (15-25 cm.), yellowish brown (10YR 5/4) silty clay, dark yellowish brown (10YR 4/4) moist; weak very coarse prisms breaking to moderate very fine to medium granules; slightly hard (dry), friable (moist); violently effervescent; few shad scale roots; abrupt smooth boundary.
C4sa RSL No. 5520	10-23 inches (25-58 cm.), brown (10YR 5/3) clay, brown (10YR 5/3) moist; very weak medium to coarse prisms breaking to weak fine subangular blocks; very hard (dry), firm (moist); violently effervescent; few shad scale roots; clear smooth boundary.
C5sa RSL No. 5521	23-34 inches (58-86 cm.), brown (10YR 5/3) clay moist; weak medium subangular blocks; hard (dry), firm (moist); violently effervescent; very few fine shad scale roots; abrupt smooth boundary.
C6sa RSL No. 5522	34-41 inches (86-105 cm.), brown (10YR 5/3) clay moist; massive; very hard (dry), firm (moist); violently effervescent, common fine to medium white (10YR 8/1) carbonate mottles; very few roots; abrupt smooth boundary.
C7 RSL No. 5523	41-53 inches (105-135 cm.), pale brown (10YR 6/3) clay moist; massive; very hard (dry), firm (moist); violently effervescent, many white (10YR 8/2) carbonate mottles.

1/ The Morales series is in a fine, montmorillonitic, thermic family of Typic Salorthids. This pedon has a salic horizon and more clay than is typical of the series.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Rosamond silt loam (taxadjunct) **LOCATION** San Bernardino County, California

SOIL FAMILY: Typic Torrifluvent, fine, montmorillonitic, thermic.

SOIL NOS. S55Calif-36-1 **LAB. NOS.** 553 thru 559

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0- 4	C1	0.1	0.4	0.4	7.4	16.5	55.3	19.9	37.9	40.0	0	sil
4-12	C2	0.2	0.9	0.8	13.4	19.9	38.5	26.3	45.5	24.4	0	l
12-22	C3	0.2	1.3	1.4	7.5	12.4	42.9	34.3	30.7	30.4	0	cl
22-28	C4	0.2	1.1	1.4	4.8	7.6	43.0	41.9	22.9	31.0	0	sic
28-41	C5	0.4	1.8	1.9	5.4	7.1	33.3	50.1	20.1	24.0	1	c
41-51	C6	1.7	4.8	2.9	7.4	7.0	24.3	51.9	18.0	18.3	14	c
51-63	C7	6.6	17.0	9.2	13.5	7.0	13.4	33.3	19.3	8.7	2	scl

8C1b pH SATURATED PASTE 1:1	8C1a pH 1:5	8C1a pH 1:10	6A1a ORGANIC MATTER ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	4B2 MOISTURE TENSIONS 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
7.7		8.6	0.56	0.064	8.8		26.9	38.6	0			9.0
7.7		8.4	0.48	0.065	7.4		38.5	17.3	1.2			10.8
7.8		8.8	0.27	0.029	9.3		22.9	25.2	0			11.7
7.9		8.4	0.25				19.9	38.5	20.0			13.5
8.2		8.6	0.27				19.1	49.1	13.6			15.8
8.3		8.8	0.21				16.7	43.2	2.3			18.6
8.1		8.7	0.10				11.8	16.5	2.9			11.2

5A2a CATION EXCHANGE	5B1a EXTRACTABLE CATIONS 5B1a	6H2a	Exch. 5B1b	Exch. 5B1h	5D1 Exch.	SATURATION EXTRACT SOLUBLE 6P1a	6Q1a	6N1a	8A1	6K1a	8A MOISTURE AT

ROSAMOND SILT LOAM (taxadjunct) 1/
S55Calif-36-1

Sampled by: Fred Eggers, W. G. Harper, Reuben Nelson, January, 1955.
Description by: W. G. Harper
Location: On Stringer farm - 1/4 mi. E and 1120 ft. S of NW Cor., Sec. 6, T4N, R1E, about 2 miles NE of Lucerne Valley, San Bernardino County, Calif.
Elevation: About 3000 ft. Annual precipitation about 4 inches
Classification: Typic Torrifluvent, fine, montmorillonitic, thermic.
California Group: III
Vegetation: Shadscale 12 to 14 inches high, and some Mormon tea. About 85 percent of surface is barren
Parent Material: Alluvium deposited in playa - derived largely from granitic rocks but with a mixture of other rock forms
Physiography: Playa - nearly level
Drainage: No surface outlet. Water table at about 50 ft.

HORIZON	DESCRIPTION
S55Calif-36-1-1 C1 0- 4"	Very pale brown (10YR 7/3 dry, brown 5/3 moist) silt loam; vesicular in topmost 1/2 inch; moderately defined fine to medium platy; slightly hard when dry, friable when moist; moderately permeable; low in organic carbon; very strongly calcareous and almost devoid of plant roots; pH on 1 to 5 dilution with thymol blue 8.5.
S55Calif-36-1-2 C2 4-12"	Same color as surface soil; loam; weakly defined medium subangular blocky; hard when dry, friable when moist; very strongly calcareous; moderately permeable; pH on 1 to 5 dilution 8.5. This contains a few shadscale roots.
S55Calif-36-1-3 C3 12-22"	Pale brown (10YR 6/3 dry, brown 5/3 moist) clay loam; moderately defined medium subangular blocky; very hard when dry, very firm when moist; very strongly calcareous; slowly to very slowly permeable; pH on 1 to 5 dilution 8.6.
S55Calif-36-1-4 C4 22-28"	Same color as layer above; silty clay; moderately defined medium to coarse subangular blocky; very hard when dry; very firm when moist; strongly calcareous; very slowly permeable; pH on 1 to 5 dilution 8.5. This contains a few shadscale roots and a few fine distinct white flecks of lime and gypsum.
S55Calif-36-1-5 C5 28-41"	Pale brown (10YR 6/3 dry and moist) clay; weak medium subangular blocky; very hard when dry, very firm when moist; very slowly permeable; very strongly calcareous with common, fine to medium flecks of white (10YR 8/2) lime and gypsum; very few shadscale roots; pH on dilution 8.6.
S55Calif-36-1-6 C6 41-51"	This contains a few small lime carbonate concretions, but otherwise appears similar to the layer above but contains no plant roots; pH on 1 to 5 dilution 8.8.
S55Calif-36-1-7 C7 51-63"	White (10YR 8/2 dry and moist) sandy clay loam; moderately defined medium to coarse platy (laminated) extremely hard when dry, friable when moist; slowly permeable; very strongly calcareous; pH on dilution 8.7; no plant roots.

Remarks: This is a stratified soil laid down in slowly moving water. The lime concentrated in the three lower layers may have been accumulated from a high water table. The gypsum probably precipitated from ground water.

1/ The Rosamond series is in a fine-loamy, mixed (calcareous), thermic family of Typic Torriorthents. This pedon has more clay and an organic carbon content that decreases irregularly with depth.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE San Emigdio loam **LOCATION** Riverside County, California

SOIL FAMILY: Typic Xerofluvent, coarse-loamy, mixed, calcareous, thermic

SOIL NOS. S55Calif-33-11 **LAB. NOS.** 55578-583

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1	2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	
0- 8	Ap1	0.3	1.4	2.6	15.3	13.7	49.3	17.4	43.7	30.1	0	l
8-14	Ap2	0.7	2.3	4.5	19.3	13.9	40.6	18.7	41.5	25.9	1	l
14-28	C1	1.7	3.5	5.0	20.0	14.8	37.1	17.9	43.7	21.7	2	l
28-39	C2	1.2	3.0	4.7	23.4	18.5	36.0	13.2	51.5	19.0	2	l
39-54	C3	1.6	4.5	7.6	30.7	16.2	28.5	10.9	47.4	16.1	2	fsl
54-64	C4	0.1	0.7	1.6	11.7	14.6	51.2	20.1	42.2	32.4	1	sil

SAN EMIGDIO LOAM
S55Calif-33-11

Sampled by: Fred Eggers, Arnold Knecht, F. G. Stephens, W. G. Harper, R. E. Nelson and F. F. Harradine. **Description by:** W. G. Harper.

Location: 1700' W of Wolfskill-Moreno ranch headquarters, 1000' E of the S 1/4 corner Sec. 18, T3S, R2W, near Moreno, Riverside County, California. **Sampling date:** December 6, 1955.

Elevation: 1490 feet

Precipitation: 13 inches

Classification: Typic Xerofluvent, coarse-loamy, mixed, calcareous, thermic.

California Group: I

Vegetation: Used for dry farm grain. Fallow this season.

Parent Material: Alluvium from San Timoteo Badlands.

Physiography: Near lower end broad alluvial fan.

Drainage: Well drained.

HORIZON		DESCRIPTION
Ap1	0- 8"	Pale brown (10YR 6/3 dry, dark brown 3/3 or 3.5/3 moist) loam; weakly defined; medium to fine crumb structure; very hard, friable; moderately permeable; many roots; very slightly calcareous; pH 8.2; abrupt smooth boundary.
Ap2	8-14"	Pale brown (10YR 6/3 dry, dark brown 4/3 moist) loam; massive but breaks up into various sized angular blocky aggregates; very hard, friable; moderately permeable; many roots; effervesces strongly; white lime veins; pH 8.2; abrupt smooth boundary.
C1	14-28"	Pale brown (10YR 6/3 dry, dark brown 4/3 moist) loam; massive; very hard; friable; moderately permeable; strong effervescense; many pores and many worm holes; moderate amount of roots; pH 8.2; abrupt smooth boundary.
C2	28-39"	Brown (10YR 5.5/3 dry, dark brown 4.5/3 moist) fine sandy loam; massive; slightly hard, very friable; moderately permeable; few roots; strong effervescense; pH 8.2; clear smooth boundary.
C3	39-54"	Brown (10YR 5.5/3 dry, dark brown 4.5/3 moist) sandy loam; slightly hard, very friable; moderately permeable; very few roots; strong effervescense; pH 8.2; clear smooth boundary.
C4	54-64"	Light yellowish brown (10YR 6/4 dry, dark brown 4.5/3 moist) silt loam; massive; may have some clay films; very hard, friable; moderately permeable; very few roots; effervesces strongly; pH 8.2; many pores

SOIL SURVEY LABORATORY Riverside, California

OIL TYPE San Emigdio fine sandy loam **LOCATION** Riverside County, California

SOIL FAMILY: Typic Xerofluvent, coarse-loamy, mixed, calcareous, thermic.

SOIL NOS. S57Calif-33-1 **LAB. NOS.** 57468 - 57475

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0- 7	Ap1	0.8	5.3	10.5	35.5	14.8	26.3	6.8	51.8	12.7	0	fsl
7-11	Ap2	0.5	4.5	11.9	40.7	14.4	22.0	6.0	50.5	9.9	0	fsl
11-15	C1	0.4	10.3	23.1	45.5	9.7	8.5	2.5	36.3	3.9	0	s
15-18	C2	0.4	3.3	7.8	29.0	14.2	38.5	6.8	49.6	20.9	0	fsl
18-34	C3	0.6	4.5	9.7	40.8	19.8	19.9	4.7	58.7	7.3	0	lfs
34-45	C4	1.7	9.0	12.3	35.4	15.4	20.1	6.1	49.2	7.3	0	fsl
45-47	C5	0.9	4.1	4.7	16.9	9.8	49.6	14.0	39.6	30.8	0	l
47-60+	C6	3.7	11.8	12.9	35.3	12.7	17.9	5.7	43.6	7.4	0	ls

pH 8C1b Saturated Paste 1:1	pH 8C1a 1:10	4A3a Bulk Density g/cc	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	6I1a CO_3 meq/l	8A1a Elec. Conduc. mmhos/cm. at 25° C.	6E1b $CaCO_3$ equivalent %	6J1a HCO_3 meq/l	MOISTURE TENSIONS 4B1a 1/10 ATMOS. %	4B1a 1/3 ATMOS. %	4B2 15 ATMOS. %
7.5	7.6		0.60	0.058	10.3	1.2	1.1	<1.0	7.4	21.4	13.4	4.3
7.8	8.4		0.25	0.017	14.7	0.0	1.1	1.0	4.7	18.4	11.4	3.5
8.0	8.3		0.05	0.004		0.0	0.5	<1.0	2.9	7.3	3.9	1.5
7.5	8.2		0.22			0.0	1.2	1.4	3.4	30.2	17.7	4.7
7.8	8.5	1.35	0.13			0.0	0.8	1.2	2.8	19.3	8.5	2.7
7.8	8.5	1.37	0.13			0.0	0.8	1.3	2.9	20.1	9.2	3.0
7.7	8.5		0.30			0.0	1.8	3.4	3.7			7.8
8.0	8.7	1.37	0.15			0.0	0.8	<1.0	2.9	18.7	8.6	2.8

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	5B1b Exch. Na 6P2a	5B1b Exch. K 6Q2a	5D1 Exch. Na %	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a Ca	6O1a Mg	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
11.9	15.7	3.0		0.7	0.4	6	4.6	0.4	5.6	2.6	1.9	32.6
12.3	13.7	1.8		0.6	0.3	5	4.0	0.5	5.4	1.0	1.4	29.8
8.0	6.7	0.8		0.5	0.2	6	2.1	0.3	2.1	1.2	0.4	27.3
17.5	17.9	1.6		0.4	0.5	2	7.8	0.8	5.9	1.0	2.8	34.5
13.3	15.5	1.2		0.5	0.3	4	3.5	0.3	3.3	0.6	1.2	29.8
12.7	16.9	1.2		0.6	0.2	5	3.7	0.3	3.2	0.8	1.2	30.8
24.1	25.3	2.2		1.1	0.6	5	9.2	0.4	9.5	1.4	4.7	44.2
9.5	15.3	1.2		0.9	0.2	9	4.7	0.3	2.6	0.7	1.2	28.1

SAN EMIGDIO FINE SANDY LOAM
S57Calif-33-1

Location: Riverside County, California. 3,050 feet west and 150 feet north of southeast corner of Sec. 8, T. 5 S., R. 1 E. On the Lovell Ranch, 150 feet north of Acacia, about 3 miles east of Hemet. Date of sampling: June 3, 1957.
Description by: Fred Eggers. Collectors: George Borst, R. E. Nelson, Cliff Henry, and Arnold Knecht.
Classification: Typic Xerofluvent, coarse-loamy, mixed, calcareous, thermic.
Vegetation: Walnut grove. Precipitation: 10-14 inches. Parent material: Alluvium from the Bautista Beds. Physiography: Flood plain. Elevation: 1,765 feet. Drainage: Well drained.
Remarks: Very thin silt lenses in all horizons. These were not sampled except in C2 and C5. The entire profile is highly micaceous. There are grass roots in Ap1, Ap2 and C1 and walnut tree roots throughout the profile.

HORIZON	DESCRIPTION
Ap1 RSL No. 57468	0-7 inches, grayish brown (10YR 5/2 dry, very dark grayish brown 3/2 moist) fine sandy loam; weakly defined medium to fine subangular blocky structure; slightly hard when dry, firm when moist; nonsticky and nonplastic when wet; moderately rapid permeability; many plant roots; pH 7.6; strong effervescence; clear smooth lower boundary.
Ap2 RSL No. 57469	7-11 inches, grayish brown (10YR 5/2 dry, very dark grayish brown 3/2 moist) fine sandy loam; moderately defined medium angular blocky structure; hard when dry, very firm when moist; nonsticky and nonplastic when wet; moderately permeable; many plant roots; pH 8.0; strong effervescence; clear wavy lower boundary. This is a moderately compacted cultivation pan.
C1 RSL No. 57470	11-15 inches, gray (10YR 6/1 dry, dark gray 4/1 moist) loamy sand, single grain; loose when dry, loose when moist; nonsticky and nonplastic when wet; rapid permeability; many plant roots; pH 8.0; strong effervescence; clear smooth lower boundary.
C2 RSL No. 57471	15-18 inches, gray (10YR 6/1 dry, dark gray 4/1 moist) very fine sandy loam; moderately defined fine platy; slightly hard when dry; very firm when moist; slightly sticky and nonplastic when wet; moderate permeability; many plant roots; pH 8.0; strong effervescence; clear smooth lower boundary. This is a mixture of very thin silt lenses and loamy fine sand.
C3 RSL No. 57472	18-34 inches, gray (10YR 6/1 dry, dark gray 4/1 moist) loamy fine sand, single grain; loose when dry, loose when moist, nonsticky and nonplastic when wet; moderately rapid permeability; many plant roots; pH 8.0; strong effervescence; clear smooth lower boundary.
C4 RSL No. 57473	34-45 inches, gray to light brownish gray (10YR 6/1-6/2 dry, dark gray to dark grayish brown 4/1-4/2 moist), loamy sand, single grain; loose when dry, loose when moist; nonsticky and nonplastic when wet; rapid permeability; clear smooth lower boundary.
C5 RSL No. 57474	45-47 inches, gray (10YR 6/1 dry, dark gray 5/1 moist) silt loam; moderately defined fine platy; slightly hard when dry, firm when moist; slightly sticky and plastic when wet; moderately permeable; common plant roots; pH 8.0; clear smooth lower boundary.
C6 RSL No. 57475	47-60 inches +, gray (10YR 5/1 dry, dark gray 4/1 moist) loamy sand, single grain; loose when dry, loose when moist; nonsticky and nonplastic when wet; rapid permeability; common plant roots; pH 8.2; strong effervescence.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE San Emigdio fine sandy loam **LOCATION** Riverside County, California

SOIL FAMILY: Typic Xerofluvent, coarse-loamy, mixed, calcareous, thermic.

SOIL NOS. S57Calif-33-2 **LAB. NOS.** 57476 - 57482

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0- 8	Ap1	1.1	7.3	8.4	28.7	18.2	26.8	9.5	50.9	12.8	0	fsl
8-13	Ap2	0.9	6.8	8.2	29.5	18.1	28.3	8.2	52.1	13.5	0	fsl
13-23	C1	1.0	6.2	7.9	28.4	17.3	30.2	9.0	50.7	14.8	0	fsl
23-30	C2	0.9	6.5	8.0	27.3	15.4	31.6	10.3	48.0	16.0	0	fsl
30-35	C3	0.7	5.5	8.2	26.4	15.2	32.9	11.1	47.6	16.8	0	fsl
35-42	C4	0.6	5.4	8.1	28.6	18.1	30.9	8.3	52.9	14.5	0	fsl
42-60+	C5	0.7	8.1	10.1	35.2	18.1	19.7	8.1	52.3	7.9	0	fsl

8C1b pH SATURATED PASTE 1:1	8C1a pH 1:10	4A3a Bulk Density g/cc	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	6I1a CO_3 meq/L	6A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C	6E1b $CaCO_3$ equivalent %	6J1a HCO_3 meq./l.	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
7.6	7.8		0.52	0.055	9.5	0.5	1.7	Tr	6.2	23.7	14.0	4.8
7.8	8.3	1.47	0.24	0.026		0.0	0.7	<1.0	4.0	22.7	13.3	4.6
7.8	8.6	1.36	0.19	0.021		0.0	0.9	1.9	3.0	24.0	14.5	4.9
7.7	8.5	1.32	0.17	0.022		0.0	1.7	2.3	3.0			5.3
7.9	8.6	1.16	0.26	0.021		0.0	1.8	2.1	2.5			5.7
7.8	8.4	1.42	0.13			0.0	1.7	1.1	2.5			5.0
7.9	8.7	1.50	0.12			0.0	1.9	0.0	15.0			3.5

5A2a CATION EXCHANGE CAPACITY (-Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	5D1 EXCH. Na %	SATURATION EXTRACT SOLUBLE 6P1a Na	6Q1a K	6N1a Ca	6O1a Mg	6K1a Cl	8A1 8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
12.0	9.4	2.7	-	0.7	0.9	6	4.0	1.7	9.0	2.1	0.5	30.5
12.2	17.6	1.9	-	0.7	0.5	6	3.8	0.4	3.0	0.8	0.1	31.2
14.6	22.1	1.5	-	0.8	0.3	5	3.4	0.3	5.1	0.7	1.0	32.2
14.4	23.0	1.4	-	0.9	0.3	6	6.5	0.2	9.0	1.2	4.8	33.7
15.7	24.1	1.5	-	0.8	0.2	5	7.2	0.2	9.3	1.2	5.3	34.2
13.6	21.4	1.6	-	0.9	0.2	7	6.9	0.2	7.5	1.3	5.1	32.3
9.5	17.5	1.4	-	0.7	0.2	7	7.8	0.3	8.6	1.7	6.7	29.6

SAN EMIGDIO FINE SANDY LOAM
S57Calif-33-2

Location: Riverside County, California. 900 feet east and 800 feet south of the center of Sec. 14, T. 5 S., R. 1 W., 450 feet west and 550 feet north of the intersection of Yale and Johnston Streets, on the J. Rite Ranch, Hemet. Date of sampling: June 3, 1957.
Description by: Fred Eggers. Collectors: George Borst, R. E. Nelson, Cliff Henry, and Arnold Knecht.
Classification: Typic Xerofluvent, coarse-loamy, mixed, calcareous, thermic.
Vegetation: Apricot grove. Precipitation: 10-14 inches. Parent material: Alluvium from Bautista Beds. Physiography: Alluvial fan. Elevation: 1,648 feet. Drainage: Well drained.
Remarks: Lime in seams and pores in all C horizons.

HORIZON	DESCRIPTION
Ap1 RSL No. 57476	0-8 inches, grayish brown (10YR 5/2 dry, dark brown 3/3 moist) fine sandy loam; weakly defined medium subangular blocky structure; hard when dry, friable when moist; nonsticky and nonplastic when wet; moderately rapid permeability; many plant roots; pH 8.0; slight effervescence; abrupt smooth lower boundary.
Ap2 RSL No. 57477	8-13 inches, grayish brown (10YR 5/2 dry, dark brown 3/3 moist) fine sandy loam; moderately defined medium angular blocky structure; very hard when dry, friable when moist; nonsticky and nonplastic when wet; moderately rapid permeability; many plant roots; pH 8.0; strongly effervescent; diffuse smooth lower boundary.
C1 RSL No. 57478	13-23 inches, grayish brown (2.5Y 5/2 dry, dark brown 10YR 3/3 moist) fine sandy loam; massive; breaks to weakly defined fine angular blocky structure; slightly hard when dry, friable when moist; nonsticky and plastic when wet; moderately rapid permeability; many plant roots; pH 8.0; strong effervescence; gradual smooth lower boundary.
C2 RSL No. 57479	23-30 inches, grayish brown (2.5Y 5/2 dry, dark brown 10YR 3/3 moist) very fine sandy loam; massive; breaks to weakly defined fine angular blocky structure; slightly hard when dry, firm when moist; nonsticky and plastic when wet; moderately permeable; many plant roots; pH 8.0; strong effervescence; clear smooth lower boundary.
C3 RSL No. 57480	30-35 inches, light brownish gray to grayish brown (2.5Y 6/2-5/2 dry, dark brown 10YR 3/3-3/4 moist) fine sandy loam; massive; slightly hard when dry, firm when moist; nonsticky and plastic when wet; moderately rapid permeability; many plant roots; pH 8.0; strong effervescence; clear smooth lower boundary.
C4 RSL No. 57481	35-42 inches, light brownish gray (2.5Y 6/2 dry, dark yellowish brown 10YR 3/4 moist) fine sandy loam; massive; slightly hard when dry, firm when moist; nonsticky and plastic when wet; moderately rapid permeability; many plant roots; pH 8.0; strong effervescence; clear smooth lower boundary.
C5 RSL No. 57482	42-60 inches +, light brownish gray (2.5Y 6/2 dry, dark yellowish brown 10YR 3/4 moist) loamy fine sand; massive; loose when dry, loose when moist; nonsticky and nonplastic when wet; rapid permeability; common plant roots; pH 8.0; strong effervescence.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Campbell clay loam (taxadjunct) **LOCATION** Contra Costa County, California

SOIL FAMILY: Aquic Xerofluvent, fine-silty, mixed, nonacid, thermic

SOIL NOS. S57Calif-07-2 **LAB. NOS.** 5824 - 5829

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0-5	A1	0.1	0.2	0.2	2.3	3.1	52.7	41.4	18.8	38.8	0	sic
5-13	B1	0.1	0.2	0.4	8.4	9.9	46.4	34.6	33.9	29.2	0	sicl
13-20	B21	0.1	0.2	0.5	10.1	10.9	47.7	30.5	38.4	28.5	0	cl
20-28	B22	0.1	0.5	0.9	12.0	11.3	47.6	27.6	40.0	28.4	0	cl
28-44	C11	0.4	1.2	1.0	8.7	7.2	50.4	31.1	30.5	33.6	0	sicl
44+	C12	0.1	0.2	0.7	8.8	11.1	47.7	31.4	36.1	29.7	0	cl

pH 8C1b SATURATED PASTE 1:1	pH 1:5	pH 8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	ORGANIC MATTER 6B1a NITROGEN %	ORGANIC MATTER C/N	6I1a CO_3 meq/l	8A1a Elec. Conduc. mmhos/cm. at 25° C.	6E1b $CaCO_3$ equivalent %	6J1a HCO_3 meq/l	MOISTURE TENSIONS 1/10 ATMOS. %	MOISTURE TENSIONS 1/3 ATMOS. %	4B2 15 ATMOS. %
8.2		8.5	0.37	0.032	11.6	0.0	38.4	Tr	6.5			20.0
7.6		8.3	0.82	0.062	13.2	0.0	35.8	Tr	4.5			15.6
7.8		8.4	0.50	0.041	12.2	0.0	40.0	Tr	4.0			15.1
8.3		8.7	0.39	0.031	12.6	5.0	40.2	Tr	0.5			14.6
8.3		9.0	0.39	0.032	12.2	0.0	44.9	1	11.4			15.7
7.2		8.0	1.20	0.102	11.8	0.0	13.1	Tr	3.5			12.9

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	5B Exch. 5B1b Na 6P2a	5B Exch. 5B1b K 6Q2a	5D1 EXCH. Na %	SATURATION EXTRACT SOLUBLE 6P1a Na	6Q1a K	8A1 6N1a	6O1a	6K1a	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
34.4	43.7	7.3	--	13.0	0.1	37.8	387.5	0.5	32.1	78.9	321	62.1
29.8	9.8	13.6	--	14.9	0.5	50.0	380.0	0.2	25.5	84.8	283	56.9
29.1	18.3	6.5	--	9.3	0.2	32.0	377.5	0.2	31.1	98.3	364	54.4
28.1	25.0	6.0	--	10.8	0.1	38.4	410.0	0.3	32.8	87.4	357	53.5
30.7	19.3	15.6	--	17.7	0.4	57.7	372.0	0.3	33.7	84.2	388	56.8
29.0	11.9	9.8	--	4.0	0.7	13.8	105.0	1.0	40.7	45.9	68	48.3

CAMPBELL CLAY LOAM (taxadjunct) 1/
S57Calif-07-2

Collector: R. C. Huff, May 16, 1957

[redacted]

Native Vegetation: Salt bush and meadow barley.
Salt or alkali: Appears to be strongly saline-alkali.
Climate: Rainfall 18". Cool, moist winters and dry summers.
Parent Material: Mixed alluvium - dominately sedimentary.
Relief: Flat to very gently sloping on the basin rim.
Elevation: 20 feet.
Drainage: Surface and internal drainage is slow. Free water at 5 feet at time of sampling.
Root Distribution: Many 0 - 5". Some large salt bush roots extend to 5' or more.
Classification: Aquic Xerofluvent, fine-silty, mixed, nonacid, thermic.

HORIZON		DESCRIPTION
57Calif-07-2-1		Gray brown (10YR 5/2 dry), very dark grayish brown
A_1	0- 5"	(10YR 3/2 moist) clay loam; strong, medium angular blocky;

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Oxalis clay loam (taxadjunct) **LOCATION** Contra Costa County, California

SOIL FAMILY: Aquic Xerofluvent, fine, montmorillonitic, nonacid, thermic

SOIL NOS. S57Calif-07-1 **LAB. NOS.** 5818 - 5823

DEPTH INCHES	HORIZON	1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
		PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1		
0-6	A1	0.3	1.7	3.0	13.1	4.1	36.5	41.3	19.6	29.4	0	c
6-13	B1	0.0	1.0	2.0	9.8	3.0	28.0	56.2	13.4	24.1	0	c
13-24	B21	0.1	1.2	2.1	10.9	3.2	28.8	53.7	13.6	25.7	0	c
24-31	B22	0.1	1.8	3.3	18.6	4.3	25.7	46.2	20.0	22.5	0	c
31-47	C11	0.2	2.8	5.3	27.7	5.2	22.4	36.4	27.0	18.7	0	cl
47+	C12	1.1	5.7	10.3	28.6	6.3	17.6	30.4	26.3	13.3	0	scl

8C1b SATURATED PASTE 1:1	pH 1:5	8C1a 1:10	6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	6I1a CO_3 meq/l	8A1a Elec. Conduc. mmhos/cm. at 25°C.	6E1b $CaCO_3$ equivalent %	6J1a HCO_3 meq/l	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
			ORGANIC MATTER									
6.7		7.5	1.85	0.163	11.3	0.0	4.8	--	3.2			15.3
7.8		8.7	1.00	0.075	13.3	0.0	20.6	--	4.0			18.9
8.0		8.5	0.34	0.034	10.0	0.0	22.4	Tr	5.0			18.7
8.3		9.1	0.40	0.033	12.1	0.0	18.3	Tr	8.9			16.4
8.2		9.0	0.25	0.045	5.5	0.0	22.0	Tr	9.4			14.2
8.0		8.5	0.18	0.018		0.0	15.3	3	4.0			11.6

5A2a CATION EXCHANGE CAPACITY (+ Na)	5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	5D1 EXCH. Na %	6P1a Na	6Q1a K	6N1a	6O1a	6K1a	8A MOISTURE AT SATURATION %
	EXTRACTABLE CATIONS						SATURATION EXTRACT SOLUBLES 8A1					
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
35.5	11.5	14.4	--	5.0	1.2	14.1	33.8	0.2	4.1	7.9	9	56.0
44.0	12.4	17.3	--	16.1	0.8	36.6	192.5	0.2	19.0	29.5	109	68.7
42.2	11.5	16.9	--	15.6	0.7	37.0	250.0	0.2	26.9	48.5	138	69.9
37.0	10.2	16.6	--	15.1	0.5	40.8	218.8	0.2	10.8	35.1	145	63.7
27.6	8.2	15.2	--	8.8	0.4	31.9	220.0	0.3	11.5	39.6	132	54.4
23.8	24.0	7.3	--	7.5	0.1	31.5	3.8	0.2	23.3	28.2	63	47.4

OXALIS CLAY LOAM (taxadjunct) 1/
S57Calif-07-1

Collector: R. C. Huff, May 16, 1957
Location: Contra Costa County, California. Eastern Contra Costa, SCD, SW corner Delta and Byron Roads. 200 feet west and 300 feet south of power pole in corner.
Vegetation: Meadow barley and weeds.
Salt or alkali: Appears to be strongly saline-alkali.
Climate: Rainfall 18". Cool moist winters and hot dry summers.
Parent Material: Mixed alluvium-dominately sedimentary.
Relief: Flat to very gently sloping on the basin rim.
Elevation: 10 feet.
Drainage: Surface and internal drainage is slow. Free water at 5 feet at time of sampling. Root Distribution: Many 0 to 5 inches, few below.
Classification: Aquic Xerofluvent, fine, montmorillonitic, nonacid, thermic.

HORIZON	DESCRIPTION
57Calif-07-1-1 A_1 0- 6"	Grayish brown (10YR 5.5/2 dry), very dark grayish brown (10YR 3/2 moist) clay loam; moderate, medium, angular blocky, plastic, sticky and extremely hard. The pH is 6.5. Clear, wavy lower boundary.
57Calif-07-1-2 B_1 6-13"	Very dark gray (10YR 3/1) moist clay; massive, very sticky, very plastic, and firm. The soil does not react to dilute HCl but salt crystals (probably gypsum) occur as seams. The pH is 8+. Clear, wavy lower boundary.
57Calif-07-1-3 B_{21} 13-24"	Very dark grayish brown (10YR 3/2 moist) clay; massive, very sticky and very plastic; clayskins are thin and discontinuous. The soil effervesces with dilute HCl and lime and gypsum are very prominent as seams or soft masses. The pH is 8+. Clear, wavy lower boundary.
57Calif-07-1-4 B_{22} 24-31"	Very dark grayish brown to dark grayish brown (10YR 3.5/2 moist) clay; massive, firm, plastic and sticky; clayskins prominent and continuous. Soil reacts strongly to dilute HCl. Salts (probably gypsum) and lime are very prominent as seams and as masses. The pH is 8+. Diffuse, wavy lower boundary.
57Calif-07-1-5 C_{11} 31-47"	Dark grayish brown (10YR 4.5/2) clay loam; massive, plastic, sticky, and firm; thin, patchy clayskins. Soil reacts strongly to dilute HCl. Salts (gypsum?) and lime are disseminated and in seams; there are also a few soft iron or manganese concretions. The pH is 8+. There are common, fine and faint mottles.
57Calif-07-1-6 C_{12} 47"+	Dark yellowish brown (10YR 4/4 moist) clay loam; massive, sticky, plastic, and firm to very firm. Salts (probably gypsum) and lime occur as seams and masses. Soil reacts violently to dilute HCl. The pH is 8+. Mottles, which are dark olive gray (5Y 3/2) are common, medium, and distinct. Soft iron or manganese concretions are common. This horizon appears to have the firmness and brittleness of a fragipan.

1/ The Oxalis series is in a fine, montmorillonitic (calcareous), thermic family of Aquic Xerorthents. This pedon has an organic carbon content that decreases irregularly with depth.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Rosamond sandy clay I am **LOCATION** San Bernardino County, California

SOIL FAMILY: Typic Torriorthent, fine-loamy, mixed, calcareous, thermic

SOIL NOS. 855Calif-36-6 **LAB. NOS.** 5538 thru 5546

ROSAMOND SANDY CLAY LOAM
S55Calif-36-6

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE San Emigdio loam (taxadjunct) **LOCATION** Riverside County, California

SOIL FAMILY: Typic Xerorthent, coarse-loamy, mixed, calcareous, thermic.

SOIL NOS. S55Calif-33-13 **LAB. NOS.** 55589-592

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-8	Ap	2.1	4.1	5.7	22.3	20.2	37.1	8.5	54.2	18.3	0	vfsl
8-22	C1	1.5	4.6	6.1	23.2	19.9	36.5	8.2	53.0	18.9	7	vfsl
22-42	C2	1.6	4.3	6.3	24.1	19.7	36.2	7.8	55.1	16.9	7	vfsl
42-60	C3	1.5	4.3	5.7	21.6	19.6	38.2	9.1	52.5	19.9	6	vfsl

pH 8C1b SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B1a 1/10 ATMOS. %	4B1a 1/3 ATMOS. %	4B2 15 ATMOS. %
7.6		8.2	0.61	0.060	10.2		0.84	<1		25.9	14.2	5.0
7.8		8.9	0.24	0.028			0.62	1		22.1	13.1	4.9
8.0		9.0	0.16				0.42	2		21.5	13.3	4.8
8.0		8.9	0.12				0.46	3				5.4

5A2a CATION EXCHANGE CAPACITY (+ Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	5D1 EXCH. Na %	SATURATION EXTRACT SOLUBLE 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg	8A1	6K1a	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
13.8	13.1	1.0		0.2	0.6	1	0.5	0.4	7.9			28.0
14.7	19.4	4.2		0.2	0.3	1	0.6	0.1	5.1			26.2
14.7	19.1	6.0		0.2	0.2	1	0.7	0.1	3.5			26.2
16.3	19.4	6.1		0.3	0.2	2	1.0	0.1	3.5			26.6

1/ Percent of the whole soil

SAN EMIGDIO LOAM (taxadjunct) 1/
S55Calif-33-13

Sampled by: Fred Eggers, Arnold Knecht, F. G. Stephens, W. G. Harper, R. E. Nelson and F. F. Harradine. Description by: W. G. Harper

Location: 100' NE of State Hwy. 79, 800' E and 400' S of NW corner Sec. 16, T3S, R2W, about 3 miles E of Moreno, Riverside County, California.

Sampling date: December 6, 1955.

Elevation: 1600 feet

Precipitation: 14 inches

Classification: Typic Xerorthent, coarse-loamy, mixed, calcareous, thermic.

California Group: I

Vegetation: Rabbitbrush, cheat grass and much alfileria.

Parent Material: Alluvium from San Timoteo Badlands.

Physiography: Gently sloping, alluvial fan; slightly convex.

Drainage: Good.

HORIZON		DESCRIPTION
Ap	0- 8"	Pale brown (10YR 6/3 dry, very dark grayish brown 3.5/2.5 moist) fine sandy loam; weakly defined medium to fine crumb structure; hard, very friable; moderately permeable; many roots; pH 8.2; effervesces slightly with weak acid; abrupt smooth boundary.
C1	8-22"	Pale brown (10YR 6/3 dry, grayish brown 5/2 moist) fine sandy loam; massive; hard, very friable; moderately permeable; few roots; effervesces strongly; contains mycelium-like lime; pH 8.2; clear smooth boundary.
C2	22-42"	Pale brown (10YR 6/3 dry, dark grayish brown 4/2 moist) fine sandy loam; massive; slightly hard, very friable; moderately permeable; few roots; effervesces strongly; contains mycelium-like lime; pH 8.2; clear smooth boundary.
C3	42-60"	Pale brown (10YR 6/3 dry, dark grayish brown 4.5/2.5 moist) fine sandy loam; massive; slightly hard, very friable; moderately permeable; effervesces strongly with acid; contains mycelium-like lime; pH 8.2

Remarks: This profile contains some mica throughout. It has a slightly compacted tillage pan between 8 and 22 inches.

1/ The San Emigdio series is in a coarse-loamy, mixed, calcareous, thermic family of Typic Xerofluvents. This pedon has a regular decrease in organic carbon.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Xerorthent, coarse-loamy, mixed, nonacid, thermic

SOIL Hanford gravelly sand loam (taxadjunct) SOIL Nos. S4557-2 LOCATION Tehama County, California

SOIL SURVEY LABORATORY _____ LAB. Nos. _____

Depth (In.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Size class and particle diameter (mm) Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	3A1 (2-0.1)			Coarse fragments > 2 Pct.	2-19	19-76
		Pct. of < 2 mm															Pct. of < 76mm	
0-8	Ap	57.4	35.2	7.4	8.4	11.1	7.6	14.8	15.5	17.1	18.1		41.9					
8-13	A11	55.4	35.9	8.7	6.7	11.0	7.9	14.9	14.9	16.5	19.4		40.5					
13-18	B21	55.6	35.1	9.3	7.6	11.2	8.1	15.0	13.7	16.2	18.9		41.9					
18-22	B22	56.5	34.1	9.4	5.8	10.5	8.4	17.2	14.6	15.3	18.8		41.9					
22-27	B23	59.6	30.5	9.9	6.3	11.1	9.7	19.1	13.4	12.4	18.1		59.6					
27-36	IIC1	72.6	18.3	9.1	16.0	23.8	12.2	13.5	7.1	6.3	12.0		65.5					
36-44	IIC2	75.5	16.4	8.1	14.2	28.4	14.0	12.7	6.2	6.6	9.8		69.3					

Depth (In.)	6A2a Organic carbon Pct	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	Pct			pH		8C1a (1:1)
0-8	0.44	0.053																5.5
8-13	0.35	0.048																5.8
13-18	0.22	0.038																6.3
18-22	0.13	0.032																6.8
22-27	0.13	0.030																6.9
27-36	0.09	0.023																7.1
36-44	0.10	0.020																7.1

Depth (In.)	Extractable bases: Ca 6N2b	Mg 602b	Na 1/	K 2/	5B1a Sum of bases	6H2a Ext. Acidity	5A3a CEC Sum									Base saturation: 5C3 Sum Pct	Pct.
	meq/100 g																
0-8	2.2	0.6	tr	0.21	3.0	3.1	6.1									49	
8-13	2.9	0.8	0.01	0.19	3.9	2.7	6.6									59	
13-18	4.4	1.3	<0.01	0.13	5.8	2.1	7.9									74	
18-22	5.0	1.8	0.01	0.08	6.9	2.0	8.9									77	
22-27	5.1	2.7	0.01	0.07	7.9	2.3	10.2									77	
27-36	4.6	3.2	0.02	0.02	7.8	1.8	9.6									82	
36-44	4.2	3.2	0.02	0.01	7.4	1.8	9.2									81	

Depth (In.)					

1/ Gravimetrically as sodium uranyl magnesium acetate

2/ Precipitated as potassium sodium cobaltinitrite and titrated as potassium permanganate

HANFORD GRAVELLY SANDY LOAM (taxadjunct) 1/
S4557-2

Location: Tehama County, California. Pit located 0.1 mile south of the N¼ corner, Sec. 32, T25N, R3W, MDB&M. Date of Sampling: April 1, 1945.
Description by: R. C. Roberts and R. A. Gardner. Collectors: R. C. Roberts, F. O. Youngs, R. A. Gardner, L. T. Alexander, H. Jenny and R. E. Storie.
Classification: Typic Xerorthent, coarse-loamy, mixed, nonacid, thermic.

Vegetation: Cultivated - grain. Climate: Average annual precipitation is about 19 inches. Average annual temperature is about 62°F., January 45°F., and July 82°F. Parent Material: Alluvium from sedimentary rocks. Topography: Nearly level fans adjacent to stream channels on general slope of about 1 or 2 percent. Elevation: 310 feet. Drainage: Well drained, moderate permeability, slow

Ap 0 to 8 inches, brown (10YR 5/3) gravelly sandy loam, grayish brown (10YR 4/2) moist; very weak blocky structure; slightly hard, very friable, nonplastic; moderate number of roots; strongly acid (pH 5.5); abrupt smooth boundary.

A11 8 to 13 inches, brown (10YR 5/3) gravelly sandy loam, grayish brown (10YR 4/2) moist; weak blocky structure; slightly more compact than layer above; slightly hard, very friable, nonplastic; moderate number of roots; medium acid (pH 5.8); very gradual boundary.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Xerorthent, coarse-loamy, mixed, nonacid, thermic

SOIL Hanford sandy loam SOIL Nos. 865Calif-27-12 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65449 - 65455

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand Very coarse (2-1)	Sand Coarse (1-0.5)	Sand Medium (0.5-0.25)	Sand Fine (0.25-0.1)	Sand Very fine (0.1-0.05)	Silt 0.05-0.02	Silt Int. III (0.02-0.002)	Silt Int. II (0.2-0.02)	Silt (2-0.1)	3A1a Carbonate clay	3A1a Noncarbonate clay	* Coarse fragments 1A2a ≥ 2 Pct.	2-19	19-76	
		Size class and particle diameter (mm) 3A1															Pct. of < 76mm		
		Pct. of < 2 mm																	
0-3	Ap1	76.6	15.4	8.0	16.6	21.8	12.6	17.7	7.9	8.4	7.0	25.2	68.7	0	8	16	16	0	**
3-12	Ap2	73.4	17.4	9.2	15.9	19.0	12.4	18.0	8.1	9.5	7.9	26.7	65.3	0	9	17	17	0	**
12-26	A13	72.9	17.0	10.1	14.5	23.5	13.0	15.4	6.5	9.0	8.0	22.4	66.4	0	10	16	16	0	**
26-38	A14	73.3	18.2	8.5	21.4	22.8	10.3	12.9	5.9	10.2	8.0	22.4	67.4	0	9	20	20	0	**
38-52	C1	79.0	14.2	6.8	24.1	25.0	10.7	14.0	5.2	7.5	6.7	19.1	73.8	0	7	24	24	0	**
52-73	C2	87.9	5.8	6.3	20.6	33.1	15.2	15.6	3.4	3.1	2.7	12.6	84.5	0	6	25	25	0	**
73-91+	C3	90.3	3.0	6.7	20.1	34.5	17.4	15.9	2.4	1.5	1.5	9.5	87.9	0	7	36	36	0	**

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1		
0-3	0.55	0.047	12	0.4	-								3.4			7.3		
3-12	0.51	0.044	12	0.4	-		1.67	1.67		0.89		8.7	3.9	0.001	0.001	7.2		
12-26	0.41				-		1.50	1.51		0.90		9.5	4.1	0.003	0.003	7.8		
26-38	0.22			0.4	-		1.70	1.71		0.81		9.2	3.7	0.002	0.002	8.0		
38-52	0.06			0.4	-		1.75	1.76		0.83		7.5	2.7	0.002	0.001	7.6		
52-73	0.03			0.3	-								2.4			7.7		
73-91+	0.03			0.4	-								2.5			7.6		

Depth (in.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A1a NH_4OAc	5A3a Sum		6G1d KCl Ext. Al^{+++}		Atterberg limits Plastic limit	Liquid limit	Plastic index	5D2 Exch. Na		Base saturation 5C1 NH_4OAc	
	meq/100 g											Pct.	of <0.4 mm		Pct.		Pct.	Pct.
0-3	4.2	1.3	0.2	0.3	6.0		5.4										100+	
3-12	4.5	1.3	0.2	0.3	6.3		5.8										100+	
12-26	5.2	1.7	0.2	0.3	7.4		6.5					16	17	1			100+	
26-38	3.6	1.3	0.2	0.2	5.3		4.9										100+	
38-52	2.7	1.1	0.2	0.2	4.2		4.0								5		100+	
52-73	2.4	1.0	0.2	0.2	3.8		3.4						-	N.P.	6		100+	
73-91+	2.7	1.3	0.2	0.0	4.2		4.1								5		100+	

Depth (in.)	Clay mineralogy (<0.002 mm) 7A Mont.	Verm.	Mica	Kaolin	
	X-ray				
0-3					
3-12			xxxx	x	
12-26					
26-38					
38-52					
52-73					
73-91+		x	xxxx	x	

* From characterization sample (5Kg) (except as noted).
** 2-19 mm. from characterization sample, 19-76 mm. from field volume estimate.

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

HANFORD SANDY LOAM
S65Calif-27-12

Location: Monterey County, California, adjacent to south edge at dirt road, 0.7 mile from Old Stage Road, 0.33 mile from intersection Old Stage Road and Chualar Road, Salinas Valley. Date of Sampling: May 20, 1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton and B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich. Site selection by T. D. Cook.
Classification: Typic Xerorthent, coarse-loamy, mixed, nonacid, thermic. (brunizemic Alluvial).
Vegetation: Irrigated cropland. Sugar beets. Climate: Precipitation is 12 inches; mean annual temperature 58°F., mean January 50°F., mean July 65°F. Parent Material: Alluvium from phaneritic, intermediate igneous rocks. Topography: Nearly flat, slightly convex ridge on alluvial fan surface having a west-facing slope of one percent. Elevation: 160 feet. Drainage: Well to somewhat excessively drained, moderately rapid permeability, medium runoff. Soil Moisture: Moist, near field capacity throughout.
Remarks: Field pH determinations by Truog Kit. Reaction much higher than typical for series. No satisfactory explanation. No paired profile. Classification assumes that this soil series as a whole has structure or is only slightly hard when dry. Only a few areas have had structure destroyed by excessive tillage. Asterisks indicate horizons sampled for California Division of Highways.

HORIZON	DESCRIPTION
Ap1 RSL 65449	0 to 3 inches, dark grayish brown (10YR 4/2) coarse sandy loam, very dark grayish brown (10YR 3/2) moist; massive; slightly hard, very friable, nonsticky and nonplastic; plentiful very fine roots; many very fine interstitial and few fine tubular pores; neutral (pH 7.0); clear smooth boundary.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Honcut fine sandy loam **LOCATION** Riverside County, California

SOIL FAMILY: Typic Xerorthent, coarse-loamy, mixed, nonacid, thermic.

SOIL NOS. S55Calif. 33-12 **LAB. NOS.** 55584-588

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)							3A1		2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	
0- 8	Ap	6.2	10.5	8.4	20.7	20.0	27.2	7.0	52.1	8.6	6	fsl
8-22	C1	6.6	10.5	8.7	23.4	19.1	25.2	6.5	50.3	9.1	11	fsl
22-34	C2	5.2	9.6	9.3	25.2	17.7	25.1	7.9	50.3	8.8	11	fsl
34-58	C3	7.1	11.9	9.3	24.1	17.5	23.9	6.2	46.6	10.1	15	fsl
58-64+	C4	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	18	fsl

HONCUT FINE SANDY LOAM
S55Calif-33-12

Sampled by: Fred Eggers, Arnold Knecht, F. G. Stephens, W. G. Harper, R. E. Nelson and F. F. Harradine. Description by: W. G. Harper

Location: On N. side of Indian Avenue 1320' N and 500' E of the SW corner of

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Xerorthent, fine-loamy, mixed, calcareous, thermic

SOIL Shedd silty clay loam (taxadjunct) SOIL Nos. S65Calif-27-9 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65546 - 65551

Depth (In.)	Horizon	1B1b Total Sand (2-0 05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0 25-0.1)	Sand: Very fine (0 1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	1/ Coarse fragments 1A2a > 2 Pct.	2 - 19 (Pct. of < 76 mm)	19 -76 (Pct. of < 76 mm)
		Pct. of < 2 mm																
0-5	A11	22.0	47.9	30.1	1.3	4.1	3.3	6.5	6.8	19.0	28.9	29.3	15.2	4	26			
5-15	A12	21.2	47.8	31.0	1.2	3.2	2.9	6.2	7.7	18.6	29.2	29.9	13.5	2	29			
15-26	A13	23.3	42.2	34.5	1.2	3.7	3.2	6.4	8.8	17.9	24.3	30.5	14.5	4	31			
26-50	C1ca	35.4	40.9	23.7	0.9	8.0	5.7	9.0	11.8	16.4	24.5	33.2	23.6	2	22			
50-60+	C2	70.4	20.2	9.4	0.2	2.1	2.1	13.2	52.8	7.8	12.4	71.9	17.6	0	9			
50-60+	C2 2/																	

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	Bulk density: 4A1f 1/3 bar g/cc	Bulk density: 4A1h Oven dry g/cc	Bulk density: g/cc	Water content: Pct	Water content: 4B1c 1/3 bar Pct.	Water content: 4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1b Saturated Paste	pH: 8C1a H_2O 1:1	pH: 8C1a H_2O 1:10
0-5	2.34	0.230	10		17						25.8				7.8	
5-15	1.32	0.071	19		18						24.6				8.0	
15-26	1.14				19						25.2				8.0	
26-50	0.22				15						34.2				7.9	
50-60+	0.03				1						16.5				8.1	
50-60+											17.0					

Depth (In.)	Extractable bases: 6N2a Ca	6O2a Mg	6P2a Na	5B1a 6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 8A1: 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-5	40.2	6.3	0.6	0.7	47.8			32.5	8.8	2.2	2.1	0.3		0.8	7.4	3.0		1.15
5-15	40.7	7.2	0.4	0.3	48.6			32.2	4.6	1.4	1.8	0.1		-	4.3	2.3		0.72
15-26	39.6	8.1	0.8	0.3	48.8			33.6	10.9	2.1	3.5	0.1		-	3.7	7.1		1.19
26-50	36.0	10.5	3.3	0.4	50.2			32.4	18.2	8.5	27.9	0.2		-	4.4	42.0	1.9	2.85
50-60+	19.4	7.3	3.2	0.2	30.1			18.2	6.4	3.2	20.2	0.1		-	1.8	21.8	2.1	3.02
50-60+								18.6	5.4	2.7	18.9	0.1		-	2.0	19.3		2.68

Depth (In.)	8A Water at Saturation Pct.	5D2 Exchangeable Na Pct	5B1b Exch. Na meq./100g.	Gypsum Pct.	5C1 Base Sat. NH_4OAc Pct.	Clay mineralogy (<0.002 mm.) 7A: Mont.	Verm.	Mica	Kaolin.
						X-ray			
0-5	64.6		0.5		100+				
5-15	66.3		0.3		100+				
15-26	65.6		0.6		100+				
26-50	83.0		1.0		100+				
50-60+	56.3	11	2.0		100+				
50-60+	57.4								

1/ Apparent coarse fragments dispersed readily. Hence the characterization sample was ground whole and all chemical and physical data are on a whole soil basis.

2/ C2 horizon material (shale) from which lime coatings were removed.

SHEDD SILTY CLAY LOAM (taxadjunct) 1/
S65Calif-27-9

Location: Monterey County, California, center NE¼ SE¼ NW¼ Sec. 24, T20S, R9E at the top of the watershed at Hamilton Canyon about 10 miles SE of King City. Date of

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Xerorthent, fine-silty, mixed, calcareous, thermic
SOIL Shedd silty clay loam SOIL Nos. S65Calif-27-8 LOCATION Monterey County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65538 - 65545

Depth (in.)	Horizon	1B1b Total Sand (2–0.05)	Total Silt (0.05–0.002)	Total Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	3A1 Int. II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	* Coarse fragments ≥ 2 Pct.	1A2a 2 – 19 (Pct. of < 76 mm)	1A2a 19 – 76 (Pct. of < 76 mm)
		Pct. of < 2 mm																
0-5	A11	10.0	58.5	31.5	0.5	0.9	1.0	3.7	3.4	24.2	34.3	30.2	6.1	0	32			
5-12	A12	10.0	56.4	33.6	0.3	0.6	0.8	3.7	4.6	23.1	33.3	30.0	5.4	1	33			
12-23	A13	11.9	54.9	33.2	0.4	0.9	1.1	4.3	5.2	21.8	33.1	29.6	6.7	2	31			
23-30	A14	13.0	54.7	32.3	0.4	1.0	1.6	5.1	4.9	20.8	33.9	28.6	8.1	2	30			
30-50	C1ca	7.9	55.7	36.4	0.1	0.3	0.6	4.9	2.0	17.3	38.4	22.2	5.9	0	36			
50-67+	C2	9.5	56.1	34.4	0.2	1.0	1.1	4.8	2.4	15.9	40.2	21.1	7.1	0	34			
50-67+	C2w**																	

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6E1b Carbonate as $CaCO_3$ Pct.	Atterberg limits: Plastic limit Pct.	Liquid limit of <0.4 mm	Plastic index	Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	3B2 Cm	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-5	2.10	0.200	11	4				1.18	1.30	1.00		33.6	27.5	0.033	0.033		7.8	
5-12	1.18	0.118	10	6				1.04	1.19	1.00		37.0	23.2	0.045	0.045		8.0	
12-23	1.00			8	37	52	15	0.98	1.12	1.00		38.4	24.4	0.044	0.044		8.0	
23-30	0.71			8				1.00	1.13	1.00		40.0	28.2	0.043	0.043		8.0	
30-50	0.10			2				1.15	1.30	1.00		48.8	38.4	0.042	0.042		7.9	
50-67+	0.06			tr.	48	81	33	1.17	1.33	1.00		47.0	36.2	0.043	0.043		7.8	
50-67+													30.6				7.7	

Depth (in.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	5B1a Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 8A1 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-5	46.5	6.3	0.3	0.6	53.7			43.1	6.7	1.3	1.5	0.2		0.8	5.5	1.5		0.81
5-12	46.8	7.1	0.4	0.4	54.7			42.4	4.0	0.9	1.6	0.1		-	4.0	1.2		0.56
12-23	44.9	7.9	0.4	0.4	53.6			38.0	3.6	1.3	1.6	0.1		-	3.6	1.1		0.57
23-30	44.8	9.3	0.8	0.3	55.2			39.9	3.8	1.2	2.7	0.1		-	2.9	3.4		0.76
30-50	38.6	12.6	4.0	0.6	55.8			43.0	6.7	2.9	12.7	0.2		-	2.4	17.5	0.9	2.33
50-67+	32.8	13.4	4.6	0.7	51.5			48.2	7.3	3.9	18.4	0.2		-	1.6	24.0	1.1	3.03
50-67+								43.6										

Depth (in.)	8A Water at Saturation Pct.	5D2 Exchangeable Na Pct.	5B1b Exch. Na meq./100g.	Gypsum Pct.	5C1 Base Sat. NH_4OAc Pct.	Clay mineralogy (<0.002 mm.) 7A: Mont.	Verm.	Mica	Kaolin.	
						X-ray				
0-5	66.0		0.2		100+	xxxx			t	
5-12	65.6		0.3		100+					
12-23	68.8		0.3		100+					
23-30	71.8		0.6		100+					
30-50	102	6	2.7		100+					
50-67+	100	6	2.8		100+	xxxx			t	
50-67+										

* Apparent coarse fragments dispersed readily. Hence the characterization sample was ground whole and all of the chemical and physical data are on a whole soil basis.

** C2 horizon material (shale) from which lime coatings were removed.

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

SHEDD SILTY CLAY LOAM
S65Calif-27-8

Location: Monterey County, California, center of SE¼ of NW¼ Sec. 11, T20S, R9E, about 9 miles east of King City up Wild Horse Canyon. Coordinate Grid (C65 J10) about 30 feet down from ridge top. Date of Sampling: May 19, 1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich; site selection by T. D. Cook.
Classification: Typic Xerorthent, fine-silty, mixed, calcareous, thermic. (brunizemic Regosol).
Vegetation: Range - wild oats, soft chess, bur clover, ripgut brome, small thistles. Climate: Precipitation 10 inches; mean annual temperature 62°F., January mean 59°F., July mean 68°F. Parent Material: Soft calcareous sandstone and shale, Plio-Pleistocene, non-marine. Topography: Upland. Moderately steep slightly convex slope of 20 percent on south to southwest exposure. Slope becomes much steeper 100 feet below sample site. Elevation: 1250 feet. Drainage: Well drained, permeability is moderately slow, runoff is medium. Soil Moisture: Very slightly moist, near wilting point or below.
Remarks: The paired samples for Shedd are samples #S65Calif-27-9-1 through 5. Slight erosion at sample site but steeper slopes are moderately eroded in the general vicinity. About 3 percent shale fragments throughout. Asterisks indicate horizons sampled for California Division of Highways.

HORIZON	DESCRIPTION
A11 RSL 65538	0 to 5 inches, gray (5Y 6/1) light silty clay loam, dark grayish brown (2.5Y 4/2) moist; moderate medium angular blocky structure with some tendency to weak coarse platy structure due to cattle trampling; hard, very friable, sticky and plastic; plentiful very fine roots, moist on exterior of peds; common very fine interstitial and common very fine tubular pores; moderately alkaline (pH 8.0), strongly effervescent with disseminated lime; about 3 percent fine shale fragments in an A11 and A12; clear smooth boundary.
A12	5 to 12 inches, gray (5Y 6/1) toward (5Y 5/1) light silty clay loam, very dark

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Daulton gravelly loam **LOCATION** Riverside County, California

SOIL FAMILY: Lithic Xerorthent, loamy, mixed, nonacid, thermic

SOIL NOS. S57Calif-33-4 **LAB. NOS.** 57488 - 57490

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 3	A11	5.4	9.1	5.7	14.4	13.9	43.5	8.0	45.5	21.5	38	1
3- 8	A12	4.0	8.4	6.0	15.0	14.2	41.8	10.6	45.4	20.4	40	1
8-19+	C1	Parent Material										

pH 8C1b SATURATED PASTE 1:1	pH 1:5	pH 8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	ORGANIC MATTER 6B1a NITROGEN %	ORGANIC MATTER C/N	6I1a CO_3 meq/l	8A1a Elec. Conduc. mmhos/cm. at 25°C.	6E1b $CaCO_3$ equivalent %	6J1a HCO_3 meq/l	MOISTURE TENSIONS 4B1a 1/10 ATMOS. %	4B1a 1/3 ATMOS. %	4B2 15 ATMOS. %
6.4		6.3	1.07	0.096	11.1	0.0	0.6	0	3.8			5.1
6.2		6.3	0.52	0.048	10.7	0.0	0.4	0	2.0	21.3	15.7	4.9
Parent Material												

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a Ca	6O1a Mg	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						milliequivalents per liter →					
12.4	7.5	2.3	2.0	0.5	0.7	85	1.2	0.6	3.5	1.4	1.3	32.2
13.6	7.1	2.8	2.0	0.5	0.3	84	1.3	0.2	1.8	1.1	0.9	24.5
Parent Material												

1/ Percent of the whole soil

DAULTON GRAVELLY LOAM
S57Calif-33-4

Sampled by: George Borst, R. E. Nelson, Clifford Henry, Arnold Knecht and Fred Eggers. Date: June 4, 1957.

Description by: Fred Eggers.

Location: 150' E of W 1/4 corner of Sec. 27,T5S,R4W, about 4 miles NE of Elsinore, Riverside County, California.

Elevation: 1640'

Precipitation: 10"

Classification: Lithic Xerorthent, loamy, mixed, nonacid, thermic.

California Group: VIII

Vegetation: California sage, wild buckwheat, annual grasses and weeds, and scattered cactus.

Parent Material: Bedford Canyon formation (known locally as Elsinore metamorphic) metamorphic fine grain sandstone.

Physiography: Low rolling hills.

Drainage: Well drained.

HORIZON	DESCRIPTION
S57Calif-33-4-1 A11 0-3"	Brown to dark brown (10YR 5/3-4/3 dry, dark brown 3/3 moist) gravelly loam; moderately defined medium angular blocky with slight indication of platy structure; breaks to weakly defined fine granular; slightly hard when dry, friable when moist; nonsticky and slightly plastic when wet; moderately permeable; many roots; pH 6.0; non-calcareous; many fine pores; clear smooth lower boundary.
S57Calif-33-4-2 A12 3-8"	Brown (7.5YR 5/4 dry, dark yellowish brown 10YR 3/4 moist) gravelly loam; moderately medium angular blocky; to weakly defined fine granular; hard when dry, friable when moist; nonsticky and slightly plastic when wet; moderate permeability; many plant roots; pH 6.2; non-calcareous; many fine pores; diffused irregular lower boundary.
S57Calif-33-4-3 C1 8-19"+	Decomposing metamorphic fine grain sandstone containing a small amount of soil. There are occasional fragmentary weak clay skins; pH 6.2.

CONSECUTIVE PEDON NUMBER C 68 72 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES CAGWIN SOIL NO. S68 CALI. 9-21 SAMPLE NO. 8 576-8 583 TAHOE BASIN AREA

SOIL FAMILY TYPIC CRYOPSAMMENT MIXED RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1-	MS .5-	FS .25-	VFS .1-	SILTS CSI .05-	FSI .02-	CLAYS CO. 2-.2U	FINE LT .2U	CARBO NATE 3A1A	1A2A INT. II .2-.02	COARSE FRAGMENTS PCT. OF WHOLE SOIL	I 75-	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														--A	--	--Z	--
8 576	5- 0	01																		
8 577	0- 8	A11	81.1	14.7	4.2	11.9	23.5	14.2	21.4	10.1	8.2	6.5				28.8	17A	--	17Z	N
8 578	8- 20	A12	79.2	16.3	4.5	10.2	22.7	13.9	21.3	11.1	8.9	7.4				30.5	14A	--	14Z	N
8 579	20- 46	AC	85.4	10.3	4.3	15.6	28.0	14.7	19.5	7.6	5.6	4.7				22.5	26*	--	19Z	N
8 580	46- 69	C 1	83.6	11.6	4.8	11.7	25.3	15.3	21.8	9.5	6.4	5.2				26.3	30*	--	23Z	N
8 581	69- 91	C 2	83.2	12.2	4.6	16.3	26.9	13.9	17.9	8.2	6.8	5.4				23.8	38*	--	32Z	N
8 582	91-120	C 3	88.9	8.6	2.5	28.0	28.8	12.5	13.2	6.4	5.1	3.5				18.0	43A	--	43Z	N
8 583	0- 20	A															19A	--	19Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		-PERCENT-				PCT.NO.40		GRAMS PER CC				-PERCENT-			-PERCENT-	
8 576	5- 0											0.40		1.00						
8 577	0- 8	1.92	0.068							NP	NP		1.24	1.24	0.91		14.5	3.6		
8 578	8- 20	1.16	0.046										1.05	1.05	0.94		13.5	2.7		
8 579	20- 46	0.80								NP	NP		1.35	1.35	0.85		7.1	2.8		
8 580	46- 69	0.37											1.37	1.37	0.82		6.6	2.5		
8 581	69- 91	0.20											1.34	1.34	0.76		6.8	2.3		
8 582	91-120												1.81	1.81	0.66		8.8	1.3		
8 583	0- 20		0.058									1.15			0.91			3.3		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	CEC SUM 5A3A	CEC NH4-OAC 5A6A	BASE SATURATION SUM+ ACI-DITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 576	5- 0															
8 577	0- 8	3.6	0.3	TR	0.3	4.2	3.2		7.4	8.0	57		53		6.2	5.5
8 578	8- 20						6.5		6.5						6.1	5.4
8 579	20- 46						4.5		4.5						6.0	5.2
8 580	46- 69	1.3	0.2	TR	0.2	1.7	2.8		4.5	4.2	38		41		5.8	5.0
8 581	69- 91	1.1	0.1	0.1	0.2	1.5	2.2		3.7	3.5	41		43		5.8	5.0
8 582	91-120	0.2	0.1	0.1	0.1	0.5			0.5	1.7	100+		29		5.9	5.2
8 583	0- 20														6.2	5.4

SAMPLE NO	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY I 7A2E	II 7A2E	III 7A2E	IV 7A2E	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY I 7A2E	II 7A2E	III 7A2E	MINERAL CODE	RELATIVE AMOUNT
8 576	5- 0											1 TRACE
8 577	0- 8	VR	3	KK	2						VR VERMICLT	2 SMALL
8 578	8- 20										KK KAOLINIT	3 MODERATE
8 579	20- 46	VR	3	KK	2							4 ABUNDANT
8 580	46- 69	VR	3	KK	3							5 DOMINANT
8 581	69- 91											6 INDETER.
8 582	91-120											
8 583	0- 20											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.
N=NOT DETERMINED.
*=THE 75-250MM FRACTION IS INCLUDED.

CAGWIN LOAMY SAND
S68Calif-9-21

Location: Tahoe Basin, El Dorado County, California; northwest corner of SW¼ of sec. 8, T. 12 N., R. 18 E., ½ mile north of Tahoe Blvd., 50 feet east of old logging road on Tahoe Mountain. Date of Sampling: August 5, 1968.
Description by: G. M. Kennedy. Collectors: L. N. Langan, G. L. Anderson, L. I. Larsen, D. Palmquist, B. R. Brasher, W. D. Nettleton, J. H. Rogers.
Classification: Typic Cryopsamment, mixed.
Vegetation: Not cultivated; Jeffrey pine, incense cedar, white fir, squaw carpet, greenleaf manzanita, mountain whitethorn, chinquapin, lupine. Climate: Precipitation is 35 inches. Mean annual temperature is 42°F., to 45°F., January 26°F., July 61°F. Parent Material: Granodiorite residuum. Topography: Mountainous uplands, 35 percent east slope. Elevation: 6550 feet. Drainage: Somewhat excessively drained; runoff is rapid; permeability is rapid. Soil Moisture: Dry throughout.
Remarks: Field determinations for pH by Hellige-Truog triplex indicator. Composite A taken from 20 satellite samples.

HORIZON	DESCRIPTION
O1 RSL 68576	(5-0 cm.) 2 to 0 inches, squaw carpet and pine needle litter.
A11 RSL 68577	(0-8 cm.) 0 to 3 inches, brown (10YR 5/3) loamy sand, dark brown (10YR 3/3) moist; moderate fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine and fine roots; many very fine, few fine interstitial pores; 10 percent ½ inch minus gravel; slightly acid (pH 6.3); clear wavy boundary.
A12 RSL 68578	(8-20 cm.) 3 to 8 inches, brown (10YR 5/3) loamy sand, dark brown (10YR 3/3) moist; weak fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine, common coarse roots; many very fine, few fine interstitial pores; 10 percent ½ inch minus gravel; medium acid (pH 6.0); clear wavy boundary.
AC RSL 68579	(20-46 cm.) 8 to 18 inches, yellowish brown (10YR 5/4) light loamy sand, dark yellowish brown (10YR 3/4) moist; weak fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine, fine, medium and coarse roots; many very fine, few fine interstitial pores; 10 percent 1½ inch minus gravel; 5 percent fragments greater than 3 inches; medium acid (pH 6.0); gradual smooth boundary.
C1 RSL 68580	(46-69 cm.) 18 to 27 inches, light yellowish brown (10YR 6/4) sand, yellowish brown (10YR 5/4) moist; massive; soft, very friable, nonsticky and nonplastic; many very fine, fine, medium and coarse roots; many very fine, few fine interstitial pores; 10 percent ½ inch minus gravel; 5 percent fragments greater than 3 inches; medium acid (pH 6.0); clear wavy boundary.
C2 RSL 68581	(69-91 cm.) 27 to 36 inches, light yellowish brown (10YR 6/4) gravelly coarse sand, yellowish brown (10YR 5/4) moist; massive; soft, very friable, nonsticky and nonplastic; many very fine and fine, common medium and coarse roots; many very fine, few fine interstitial pores; 30 percent gravel; 5 percent fragments greater than 3 inches; strongly acid (pH 5.5); abrupt wavy boundary.
C3 RSL 68582	(91-120 cm.) 36 to 48 inches, white (10YR 8/1, 8/2) grus; massive; few very fine, fine and medium roots in cracks; medium acid (pH 6.0).

CONSECUTIVE PEDON NUMBER C 68 73 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES CAGWIN SOIL NO. S68 CALI. 9-24 SAMPLE NO. 8 584-8 592 TAHOE BASIN AREA

SOIL FAMILY TYPIC CRYOPSAMMENT MIXED RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1-.05	MS .5-.25	FS .25-.10	VFS .1-.05	SILTS CSI .05-.02	FSI .02-.002	CLAYS CO. 2-.2U	FINE LT .2U	CARBONATE 3A1A.02	INT. II .2-	1A2A COARSE FRAGMENTS PCT. OF WHOLE 20 I 75-	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														SOIL	PCT. OF LT 75	
																	--A	--	--
8 584	3- 0	O1+O2																	
8 585	0- 10	A11	83.7	13.4	2.9	24.1	22.7	9.9	16.9	10.1	7.2	6.2				26.2	23A	—	23Z N
8 586	10- 20	A12	84.8	12.4	2.8	25.4	23.5	10.0	17.3	8.6	6.3	6.1				24.3	39A	—	39Z N
8 587	20- 30	AC	84.9	12.1	3.0	23.3	23.1	10.3	18.5	9.7	6.4	5.7				26.1	33A	--	33Z N
8 588	30- 51	C 1	85.1	12.3	2.6	28.6	24.0	9.4	15.7	7.4	6.4	5.9				22.1	39A	--	39Z N
8 589	51- 66	C 2	88.4	8.9	2.7	34.9	25.1	8.8	13.5	6.1	4.4	4.5				17.3	44A	--	44Z N
8 590	66- 94	C 3															43A	—	43Z N
8 591	94-110	C 4															56A	--	56Z N
8 592	0- 25	A															32A	--	32Z N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	4A1H	DRY 3B2	CM WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		-PERCENT-				PCT.NO.40		GRAMS PER CC			-PERCENT-			-PERCENT-	
8 584	3- 0																		
8 585	0- 10	3.39	0.109									0.92	1.03	0.90	20.1	5.2	3.8	3.4	
8 586	10- 20	1.65	0.054									1.28	1.28	0.77	10.6	3.2			
8 587	20- 30	1.17								NP	NP	1.29	1.31	0.80	9.6	3.0	0.5	0.4	
8 588	30- 51	0.66										1.32	1.32	0.76	6.5	2.4			
8 589	51- 66	0.37										1.39	1.39	0.71	4.6	2.2			
8 590	66- 94											1.53	1.53	0.70	2.8	1.6			
8 591	94-110											1.74	1.74	0.54	4.8	1.3			
8 592	0- 25	2.74									1.25			0.82		4.3			

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACI- DITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
8 585	0- 10	4.0	0.7	0.1	0.4	5.2	9.0		14.2	12.5	37		42		5.8	4.9
8 586	10- 20	2.5	0.3	TR	0.3	3.1	8.5		11.6	9.4	27		33		5.8	4.9
8 587	20- 30	1.6	0.1	TR	0.2	1.9	7.1		9.0	7.2	22		27		5.7	5.0
8 588	30- 51	1.2	0.2	TR	0.2	1.6	6.3		7.9	5.9	21		28		5.7	5.0
8 589	51- 66	0.7	0.1	0.1	0.2	1.1	4.1		5.2	4.0	21		27		5.8	5.1
8 590	66- 94	0.1	0.1	0.1	0.2	0.5			0.5	2.5	100+		20		5.9	5.2
8 591	94-110	0.2	0.1	0.1	0.2	0.6	2.8		3.4	2.7	18		22		5.8	5.0
8 592	0- 25														5.6	4.8

SAMPLE NO	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM MINERAL X RAY 7A2E I	7A2E II	7A2E III	CODE	RELATIVE AMOUNT
8 584	3- 0											1 TRACE
8 585	0- 10										VR VERMICLT	2 SMALL
8 586	10- 20										KK KAOLINIT	3 MODERATE
8 587	20- 30	VR 3	KK 3									4 ABUNDANT
8 588	30- 51											5 DOMINANT
8 589	51- 66											6 INDETER.
8 590	66- 94											
8 591	94-110											
8 592	0- 25											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

CAGWIN LOAMY COARSE SAND
S68Calif-9-24

Location: Tahoe Basin, El Dorado County, California; 0.7 miles east and 0.2 miles north of the southwest corner, sec. 12, T. 12 N., R. 18 E., 80 feet north of High Meadow Road, 4 miles south of Stateline. Date of Sampling: August 8, 1968.
Description by: L. N. Langan. Collectors: L. I. Larsen, B. R. Brasher, W. D. Nettleton, J. H. Rogers.
Classification: Typic Cryopsamment, mixed.
Vegetation: Not cultivated. Jeffrey pine, white fir, greenleaf manzanita. Climate: Precipitation is 35 inches. Mean annual temperature is 42°F., January 26°F., July is 60°F. Parent Material: Granodiorite residuum. Topography: Mountainous upland, 40 percent southwest slope. Elevation: 6500 feet. Drainage: Somewhat excessively drained; runoff is rapid; permeability is rapid. Soil Moisture: Dry.
Remarks: Field determinations for pH by Hellige-Truog triplex indicator. Composite A taken from 10 satellite samples.

HORIZON	DESCRIPTION
01 & 02 RSL 68584	(3-0 cm.) 1 to 0 inches, pine litter and duff, abrupt smooth boundary.
A11 RSL 68585	(0-10 cm.) 0 to 4 inches, dark grayish brown (10YR 4/2) loamy coarse sand, very dark brown (10YR 2/2) moist; massive; soft, very friable, nonsticky and nonplastic; common very fine and fine roots; many very fine and fine interstitial pores; strongly acid (pH 5.5); clear smooth boundary.
A12 RSL 68586	(10-20 cm.) 4 to 8 inches, grayish brown (10YR 5/2) loamy coarse sand, very dark brown (10YR 2/2) moist; massive; soft, very friable, nonsticky and nonplastic; common very fine and fine, few medium and coarse roots; many very fine and fine interstitial pores; medium acid (pH 5.7); clear wavy boundary.
AC RSL 68587	(20-30 cm.) 8 to 12 inches, grayish brown (10YR 5/2) loamy coarse sand, dark brown (10YR 3/3) moist; massive; soft, very friable, nonsticky and nonplastic; common very fine and fine, few medium and coarse roots; many very fine and fine interstitial pores; medium acid (pH 5.8); clear wavy boundary.
C1 RSL 68588	(30-51 cm.) 12 to 20 inches, pale brown (10YR 6/3) coarse sand, brown (10YR 4/3) moist; massive; soft, very friable, nonsticky and nonplastic; common very fine, few fine and medium roots; many very fine and fine interstitial pores; medium acid (pH 5.9); clear smooth boundary.
C2 RSL 68589	(51-66 cm.) 20 to 26 inches, pale brown (10YR 6/3) gravelly coarse sand, brown (10YR 4/3) moist; massive; soft, very friable, nonsticky and nonplastic; common very fine, few fine and medium roots; many very fine and fine interstitial pores; medium acid (pH 5.8); abrupt wavy boundary.
C3 RSL 68590	(66-94 cm.) 26 to 37 inches, white and gray (10YR 8/2, 5/1) grus; very few very fine roots; strongly acid (pH 5.5).
C4 RSL 68591	(94-110 cm.) 37 to 44 inches, white and gray (10YR 8/2, 6/1) less weathered grus.

CONSECUTIVE PEDON NUMBER C 68 70 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES CAGWIN DEEP VARIANT SOIL NO. S68 CALI. 9-27 SAMPLE NO. 8 553-8 564 TAHOE BASIN AREA

SOIL FAMILY TYPIC CRYOPSAMMENT MIXED RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	SIZE, CLASS AND PARTICLE DIAMETER (MM) 181B,3A1 TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SANDS VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	CSI .05- .02	FSI .02- .002	SILTS CO. 2- .2U	CLAYS FINE LT .2U	CARBONATE 3A1A	1A2A INT. II .2- .02	COARSE FRAGMENTS PCT. OF WHOLE SOIL	I 75- 20 PCT. OF LT 75	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														--A	--	--	--
8 553	3- 0	01																		
8 554	0- 5	A11	86.5	9.3	4.2	28.6	30.8	11.2	11.9	4.0	4.1	5.2				13.1	4A	--	4Z	N
8 555	5- 15	A12	80.8	13.5	5.7	14.7	24.2	13.7	19.2	9.0	6.9	6.6				25.1	23A	5	18Z	N
8 556	15- 20	A13	79.3	13.7	7.0	20.0	26.0	12.2	14.7	6.4	6.3	7.4				19.4	22A	--	22Z	TR
8 557	20- 33	C 1	77.3	14.4	8.3	14.8	26.7	13.8	16.5	5.5	6.5	7.9				19.5	22A	--	22Z	N
8 558	33- 51	C 2	78.7	13.1	8.2	19.1	26.3	12.4	14.5	6.4	5.6	7.5				18.8	29A	6	23Z	N
8 559	51- 89	C 3	78.8	12.7	8.5	19.3	26.3	12.9	14.5	5.8	5.4	7.3				17.8	54A	32	22Z	N
8 560	89-140	C 4	89.9	5.6	4.5	18.3	45.5	13.4	10.2	2.5	2.3	3.3				8.7	47A	17	30Z	N
8 561	140-185	2C 5	89.1	7.4	3.5	18.4	30.9	17.5	17.1	5.2	3.2	4.2				16.4	28A	13	15Z	N
8 562	185-200	3C 6	91.7	5.7	2.6	27.2	36.5	11.3	12.6	4.1	2.8	2.9			--	13.2	43A	7	36Z	N
8 563	0- 20	A															27A	--	27Z	N
8 564	0- 02	C 5															--A	--	--	--

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		PERCENT				PCT.	NO.40	GRAMS PER CC			PERCENT			-PERCENT-	
8 553	3- 0											0.40							
8 554	0- 5	5.85	0.190				TR			NP	NP	0.45	0.60	0.99	60.7	8.2	10.1	10.0	
8 555	5- 15	1.42	0.063									1.47	1.47	0.86	11.5	3.9			
8 556	15- 20	1.25	0.052							NP	NP	1.40	1.40	0.87	10.0	4.7			
8 557	20- 33	0.48										1.45	1.45	0.86	10.9	4.7			
8 558	33- 51	0.47										1.35	1.36	0.82	11.0	4.6	0.2	0.2	
8 559	51- 89	0.27										1.35	1.35	0.63	12.1	5.0			
8 560	89-140	0.12								NP	NP	1.45	1.47	0.67	10.5	3.3	0.5	0.3	
8 561	140-185	0.07										1.45		0.82		3.0			
8 562	185-200											1.45		0.71		2.4			
8 563	0- 20	2.49										1.45		0.83		5.2			
8 564	0- 0																		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A	MG 5B4A	NA 5B4A	K 5B4A	SUM 6N2E	EXT. ACIDITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 6O2D	NH4-OAC 5A3A	BASE SATURATION SUM+ ACIDITY 5A6A	SUM+ AL	NH4-OAC 5C1	NAF 2 MIN 8C1D	PH H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 553	3- 0															
8 554	0- 5	4.5	0.8	0.1	0.3	5.7	8.8	0.5	14.5	15.6	39	92	37		5.0	4.4
8 555	5- 15	2.1	0.3	TR	0.3	2.7	6.3		9.0	8.4	30		33		5.8	4.9
8 556	15- 20	1.9	0.2	TR	0.3	2.4	7.7		10.1	8.6	24		28		5.8	5.0
8 557	20- 33	1.4	0.1	0.1	0.2	1.8	5.3		7.1	6.1	25		30		5.8	4.9
8 558	33- 51	1.6	0.1	0.1	0.2	2.0	5.1		7.1	6.7	28		30			
8 559	51- 89	2.3	0.2	0.1	0.2	2.8	4.5		7.3	6.9	38		41		5.8	5.1
8 560	89-140	1.6	0.2	0.1	0.3	2.2	2.6		4.8	5.0	46		44		6.2	5.3
8 561	140-185	1.4	0.2	0.1	0.3	2.0	1.0		3.0	3.9	67		51		6.0	5.3
8 562	185-200	1.1	0.1	0.1	0.2	1.5	1.0		2.5	3.9	60		38		5.9	5.1
8 563	0- 20														6.0	5.2
8 564	0- 0															

SAMPLE NO	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 553	3- 0											1 TRACE
8 554	0- 5	VR	3	KK	2						VR VERMICLT	2 SMALL
8 555	5- 15										KK KAOLINIT	3 MODERATE
8 556	15- 20	VR	3	KK	2						MI MICA	4 ABUNDANT
8 557	20- 33										VM VM/MICA	5 DOMINANT
8 558	33- 51											6 INDETER.
8 559	51- 89											
8 560	89-140	MI	2	KK	2	VM	2					
8 561	140-185											
8 562	185-200											
8 563	0- 20											
8 564	0- 0											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

CAGWIN LOAMY COARSE SAND, DEEP VARIANT
S68Calif-9-27

Location: Tahoe Basin, El Dorado County, California; 900 feet west and 500 feet north of southwest corner, sec. 21, T. 12 N., R. 18 E., 1 mile northeast of Meyers, north of Pioneer Trail Road. Date of Sampling: August 9, 1968.
Description by: L. N. Langan. Collectors: J. H. Rogers, L. I. Larsen, G. M. Kennedy, G. L. Anderson, D. Palmquist, B. R. Brasher, W. D. Nettleton.
Classification: Typic Cryopsamment, mixed. (Pedon lacks a paralithic contact.)
Vegetation: Not cultivated. Jeffrey pine, white fir, lodgepole pine, big sagebrush, mules ear, mountain whitethorn, greenleaf manzanita, bitterbrush, gooseberry, squirrel tail. Climate: Precipitation is 30 to 35 inches. Mean annual temperature is 43°F., January is 27°F., July 61°F. Parent Material: Mixed alluvium. Topography: Crest of lateral moraine, pre-Tahoe?, 1 percent slope. Elevation: 6528 feet. Drainage: Somewhat excessively drained; runoff is very slow; permeability is rapid. Soil Moisture: Dry.
Remarks: Field determinations for pH by Hellige-Truog triplex indicator. Pit dug 40 feet west of Elmira S67-Calif-9-12. Composite A taken from 10 satellite samples. Bottom three horizons highly weathered, bottom horizon the most.

HORIZON	DESCRIPTION
	(3-0 cm.)
01	1 to 0 inches, pine needle litter.
RSL 68553	(0-5 cm.) [illegible]
RSL 68554	moist; moderate very fine and fine granular structure; soft, very friable, nonsticky, and nonplastic; many very fine roots; many very fine and fine interstitial pores; horizon contains partially decomposed pine litter; strongly acid (pH 5.5); clear wavy boundary.
	(5-15 cm.)
A12 RSL 68555	2 to 6 inches, grayish brown (10YR 5/2) loamy coarse sand, very dark grayish brown (10YR 3/2) moist; massive; soft, very friable, nonsticky and nonplastic; many very fine and fine roots; many very fine and fine interstitial pores; 10 percent gravel; medium acid (pH 5.8); abrupt smooth boundary.
	(15-20 cm.)
A13 RSL 68556	6 to 8 inches, brown (10YR 5/3) loamy coarse sand, dark brown (10YR 3/3) moist; weak fine and medium platy structure; soft, very friable, nonsticky and nonplastic; many very fine, few fine and medium roots; many very fine and fine interstitial pores; 10 percent gravel; medium acid (pH 5.8); abrupt wavy boundary.
	(20-33 cm.)
C1 RSL 68557	8 to 13 inches, pale brown (10YR 6/3) loamy coarse sand, brown (10YR 4/3) moist; massive; soft, very friable, nonsticky and nonplastic; many very fine and fine, common medium and coarse roots; many very fine and fine interstitial pores; 10 percent gravel; medium acid (pH 5.8); clear wavy boundary.

CONSECUTIVE PEDON NUMBER C 68 74 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES CAGWIN SOIL NO. S68 CALI. 9-32 SAMPLE NO. 8 593-8 601 TAHOE BASIN AREA

SOIL FAMILY TYPIC CRYOPSAMMENT MIXED RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SANDS VCS 2-1	CS 1- .5	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBONATE 3A1A	1A2A INT. II .2- .02	COARSE FRAGMENTS PCT. OF WHOLE SOIL	I 75- 20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75			
8 593	0- 5	A11	86.7	10.6	2.7	26.4	21.9	10.7	18.9	8.8	5.2	5.4				24.0	19A	--	19Z	N
8 594	5- 13	A12	81.2	15.4	3.4	23.3	19.0	9.9	18.3	10.7	7.5	7.9				28.4	14A	--	14Z	N
8 595	13- 33	C 1	82.4	14.3	3.3	20.4	21.4	10.7	19.9	10.0	6.8	7.5				27.2	17A	--	17Z	N
8 596	33- 51	C 2	82.9	13.7	3.4	22.6	21.8	10.4	19.0	9.1	1.5	12.2				20.1	19A	--	19Z	N
8 597	51- 66	C 3	81.7	15.0	3.3	18.4	21.3	10.7	21.4	9.9	7.3	7.7				28.0	20A	--	20Z	N
8 598	66- 76	C 4															15A	--	15Z	N
8 599	76- 97	C 5													--	--	21A	--	21Z	N
8 600	97-115	C 6															16A	--	16Z	N
8 601	0- 13	A															13A	--	13Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 4A1F	1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		PERCENT				PCT.NO.40		GRAMS PER CC				PERCENT			-PERCENT-	
8 593	0- 5	2.25	0.074									1.00	1.10	0.92		12.0	3.7		3.2	3.0
8 594	5- 13	1.18	0.042									1.17	1.18	0.93		8.2	3.2		0.3	0.3
8 595	13- 33	0.52								NP	NP	1.35	1.35	0.90		4.9	2.4			
8 596	33- 51	0.24										1.39	1.39	0.89		5.5	2.4			
8 597	51- 66	0.27										1.44	1.44	0.88		5.8	2.4			
8 598	66- 76	0.23					0.1					1.39	1.39	0.91		5.4	2.9			
8 599	76- 97						TR					1.49	1.49	0.87		4.3	2.1			
8 600	97-115											2.15	2.15	0.86		1.7	1.8			
8 601	0- 13	1.92									1.10			0.94			3.5			

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACI-DITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 593	0- 5	3.6	0.3	0.1	0.2	4.2	7.7		11.9	9.8	35		43		5.8	5.0
8 594	5- 13	2.1	0.1	TR	0.2	2.4	6.5		8.9	8.3	27		29		5.8	5.1
8 595	13- 33	0.7	0.1	0.1	0.1	1.0	5.5		6.5	5.6	15		18		5.7	4.9
8 596	33- 51	0.8	0.1	TR	0.1	1.0	3.9		4.9	4.0	21		26		5.7	5.0
8 597	51- 66	0.5	0.1	TR	0.2	0.8	3.7		4.5	4.3	19		20		5.7	5.0
8 598	66- 76	0.5	0.1	0.1	0.2	0.9	2.6	0.6	3.5	4.7	26	60	19		5.5	4.8
8 599	76- 97	0.5	0.1	TR	0.2	0.8	2.8	0.4	3.6	3.6	23	68	24		5.6	4.8
8 600	97-115	0.1	TR	0.1	0.1	0.3	2.0	0.3	2.3	2.9	14	52	11		5.5	4.8
8 601	0- 13														5.9	5.1

SAMPLE NO	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 593	0- 5											1 TRACE
8 594	5- 13										VR VERMICLT	2 SMALL
8 595	13- 33	VR 3	KK 3								KK KAOLINIT	3 MODERATE
8 596	33- 51											4 ABUNDANT
8 597	51- 66											5 DOMINANT
8 598	66- 76											6 INDETER.
8 599	76- 97											
8 600	97-115											
8 601	0- 13											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

CAGWIN LOAMY SAND
S68Calif-9-32

Location: Tahoe Basin, El Dorado County, California; 1.3 miles south and 0.8 miles east of the northwest corner sec. 1, T.12N., R.18E., 125 feet south of apex of access road to Heavenly Valley Ski Lodge. Date of Sampling: August 13, 1968.
Description by: J. H. Rogers. Collectors: L. I. Larsen, L. C. Leifer, B. R. Brasher, W. D. Nettleton, J. H. Rogers.
Classification: Typic Cryopsamment, mixed.
Vegetation: Red fir, western white pine, Jeffrey pine, prostrate manzanita. Climate: Precipitation is 40 inches. Mean annual temperature is 41°F., January 25°F., July 60°F.
Parent Material: Granodiorite residuum. Topography: Mountainous uplands, 25 percent

CONSECUTIVE PEDON NUMBER C 68 85 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES TOEM SOIL NO. 568 CALI. 9-20 SAMPLE NO. 8 697-8 703 TAHOE BASIN AREA

SOIL FAMILY TYPIC CRYOPSAMMENT MIXED,SHALLOW RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1- .5	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A.02	INT. II .2- PCT. OF WHOLE 20	1A2A COARSE FRAGMENTS I 75-	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														SOIL PCT. OF LT 75		
																	--A	--	--	--
8 697	2- 0	01																	
8 698	0- 13	A11	88.2	9.1	2.7	18.1	29.5	15.1	19.0	6.5	5.3	3.8				20.5	33A	--	33Z	N
8 699	13- 20	A12	87.8	9.6	2.6	13.6	24.2	15.7	22.9	9.4	5.4	4.2				25.9	45A	--	45Z	N
8 700	20- 48	C 1	82.2	15.0	2.8	11.9	24.0	14.5	20.9	10.9	7.9	7.1				29.3	55A	--	55Z	N
8 701	48- 76	C 2	80.7	15.9	3.4	13.4	23.8	14.4	19.0	10.1	8.6	7.3				28.7	56A	--	56Z	N
8 702	76- 84	C 3	82.5	15.4	2.1	24.2	24.3	11.9	14.2	7.9	8.4	7.0				23.7	65A	--	65Z	N
8 703	0- 20	A															33A	--	33Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR-BON 6A1A	NITRO-GEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT PYROP AL 6G7A	FE 6C5A	AL 6G5A	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY 1/3 EST. 4A1F	1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		PERCENT				PCT.NO.40		GRAMS PER CC				PERCENT			-PERCENT-	
8 697	2- 0														1.00					
8 698	0- 13	2.43	0.091			0.5				NP	NP		1.33	1.34	0.80		8.9	4.0	0.2	0.2
8 699	13- 20	2.12	0.047										1.38	1.38	0.70		9.7	3.2		
8 700	20- 48	0.62				0.5				NP	NP		1.28	1.30	0.63		9.8	2.6	0.5	0.3
8 701	48- 76	0.33										1.50			0.58			2.1		
8 702	76- 84	0.13											1.70	1.70	0.46		13.3	2.1		
8 703	0- 20	1.89	0.081									1.35			0.80			3.9		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6Q2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACI-DITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 697	2- 0															
8 698	0- 13	3.6	0.5	0.1	0.3	4.5	9.1		13.6	11.0	33		41		5.6	4.9
8 699	13- 20	1.9	0.3	0.1	0.2	2.5	6.5		9.0	7.3	28		34		5.6	4.9
8 700	20- 48	1.6	0.3	0.1	0.2	2.2	4.9		7.1	5.6	31		39		5.7	5.0
8 701	48- 76	1.6	0.3	0.1	0.2	2.2	2.6	0.6	4.8	5.3	46	79	42		5.5	4.7
8 702	76- 84	0.8	0.2	0.1	0.1	1.2	3.0	1.1	4.2	4.2	29	52	29		5.4	4.6
8 703	0- 20														5.7	5.0

CLAY MINERALOGY

SAMPLE NO	DEPTH CM.	LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE		RELATIVE AMOUNT
8 697	2- 0												1 TRACE
8 698	0- 13	VR 3	MI 2	KK 2							VR	VERMICLT	2 SMALL
8 699	13- 20										MI	MICA	3 MODERATE
8 700	20- 48	VR 2	MI 2	KK 2							KK	KAOLINIT	4 ABUNDANT
8 701	48- 76												5 DOMINANT
8 702	76- 84												6 INDETER.
8 703	0- 20												

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

TOEM GRAVELLY LOAMY COARSE SAND
S68Calif-9-20

Location: Tahoe Basin, El Dorado County, California. 300 feet south and 50 feet east of west ¼ corner sec. 17, T.12N., R.18E., 2 miles north of Meyers. Compass bearing S 89°W on Mt. Tallac, S 48°W on Angora Peak. Date of Sampling: August 5, 1968.
Description by: G. M. Kennedy. Collectors: J. H. Rogers, G. L. Anderson, B. R. Brasher, W. D. Nettleton.
Classification: Typic Cryopsamment, mixed, shallow.
Vegetation: Not cultivated. Jeffrey pine, white fir, greenleaf manzanita, bitterbrush, mountain whitethorn, squaw carpet, huckleberry,oak and some perennial grasses.
Climate: Precipitation is 35 inches. Mean annual temperature is 42°F. to 45°F., January 27°F., July 61°F. Parent Material: Granodiorite residuum. Topography: Mountainous uplands, 20 percent southwest slope. Elevation: 6500 feet. Drainage: Excessively drained; runoff is medium; permeability is rapid. Soil Moisture: Dry.
Remarks: Field determinations for pH by Hellige-Truog indicator. Composite A was sampled on 20 satellites. 10 percent rock outcrop.

HORIZON	DESCRIPTION
01 RSL 68697	(2-0 cm.) 3/4 to 0 inches, pine litter.
	(0-13 cm.) [illegible]
RSL 68698	brown (10YR 2/2) moist; moderate fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine and fine, common medium roots; many very fine interstitial pores; 20 percent gravel; strongly acid (pH 5.5); clear smooth boundary.
A12 RSL 68699	(13-20 cm.) 5 to 8 inches, grayish brown (10YR 5/2) gravelly loamy coarse sand, dark brown (10YR 3/3) moist; weak fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine and fine, common medium roots; many very fine interstitial pores; 20 percent gravel; strongly acid (pH 5.5); clear smooth boundary.
C1 RSL 68700	(20-48 cm.) 8 to 19 inches, yellowish brown (10YR 5/4) gravelly loamy coarse sand, dark yellowish brown (10YR 3/4) moist; weak fine granular structure; soft, very friable, nonsticky and nonplastic; many very fine, fine and medium, common coarse roots; many very fine interstitial pores; 25 percent gravel; strongly acid (pH 5.5); gradual wavy boundary.
C2 RSL 68701	(48-76 cm.) 19 to 30 inches, white and brownish yellow (10YR 8/2, 6/6) grus with black micro crystals; common very fine, fine, medium and coarse roots; strongly acid (pH 5.5); clear wavy boundary.
	(76-84 cm.)

CONSECUTIVE PEDON NUMBER C 68 86 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES TOEM SOIL NO. S68 CALI. 9-23 SAMPLE NO. 8 704-8 712 TAMDE BASIN AREA

SOIL FAMILY TYPIC CRYOPSAMMENT MIXED,SHALLOW RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 TOTAL FRACTION 2- .05	.05- .002	LT .002	VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	CSI .05- .02	FSI .02- .002	CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A.02	1A2A INT. .2- II	COARSE FRAGMENTS PCT. OF WHOLE SOIL	I 75- 20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														--A	--	--	--
8 704	3- 0	O1																		
8 705	0- 5	A11	88.6	8.8	2.6	41.2	25.2	8.1	10.4	3.7	4.4	4.4				13.0	35A	--	35Z	N
8 706	5- 23	A12	85.5	10.9	3.6	26.0	25.0	10.9	16.5	7.1	5.7	5.2				21.2	33A	--	33Z	N
8 707	23- 36	C 1	85.6	11.4	3.0	27.0	24.6	9.8	16.1	8.1	5.9	5.5				22.7	41A	--	41Z	N
8 708	36- 43	C 2	88.9	8.7	2.4	29.6	25.7	11.3	15.7	6.6	5.0	3.7				19.7	48A	--	48Z	N
8 709	43- 69	C 3															43A	--	43Z	N
8 710	69- 79	C 4															27A	--	27Z	N
8 711	0- 23	A															44A	--	44Z	N
8 712	0- 0																--A	--	--	--

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	AL 6G7A	EXT PYROP FE 6C5A	AL 6G5A	ATTERBERG LIMIT EST. LL 4F2	PL 4F1	BULK DENSITY 1/3 BAR 4A1F	DRY 4A1F	CM 4A1H	3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		PERCENT				PCT.NO.40		GRAMS PER CC				PERCENT			-PERCENT-	
8 704	3- 0											0.40								
8 705	0- 5	4.63	0.168			0.4				NP	NP		1.04	1.09	0.82		12.6	7.1	1.6	1.3
8 706	5- 23	1.40	0.520										1.22	1.22	0.82		10.1	4.0		
8 707	23- 36	0.58				0.3				NP	NP		1.49	1.50	0.72		5.8	2.9	0.2	0.2
8 708	36- 43	0.42										1.56			0.65			2.7		
8 709	43- 69	0.08											1.65	1.65	0.68		5.2	1.6		
8 710	69- 79												1.64	1.64	0.82		5.8	1.6		
8 711	0- 23	1.97	0.067									1.18			0.74			3.7		
8 712	0- 0																			

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6Q2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACI-DITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 704	3- 0															
8 705	0- 5	8.3	0.9	TR	0.4	9.6	8.0		17.6	17.4	55		55		5.8	5.1
8 706	5- 23	6.1	0.8	TR	0.5	7.4	5.3		12.7	11.2	58		66		6.5	5.6
8 707	23- 36	2.4	0.4	TR	0.3	3.1	4.3		7.4	6.6	42		48		6.1	5.3
8 708	36- 43						3.2		3.2						6.1	5.2
8 709	43- 69														6.2	5.6
8 710	69- 79	0.6	0.1	0.1	0.2	1.0			1.0	2.7	100+		37		6.1	5.4
8 711	0- 23														5.8	5.1
8 712	0- 0															

CLAY MINERALOGY

SAMPLE NO	DEPTH CM.	LESS THAN 0.002 MM X RAY I 7A2E	II 7A2E	III 7A2E	IV 7A2E	DTA 7A3 -PERCENT-	L.T. 0.0002 MM X RAY I 7A3	II 7A2E	III 7A2E	MINERAL CODE	RELATIVE AMOUNT
8 704	3- 0										1 TRACE
8 705	0- 5	VR	3	MI	2	KK	2			VR VERMICLT	2 SMALL
8 706	5- 23									MI MICA	3 MODERATE
8 707	23- 36	VR	2	KK	2					KK KAOLINIT	4 ABUNDANT
8 708	36- 43										5 DOMINANT
8 709	43- 69										6 INDETER.
8 710	69- 79										
8 711	0- 23										
8 712	0- 0										

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

TOEM GRAVELLY LOAMY COARSE SAND
S68Calif-9-23

Location: Tahoe Basin, El Dorado County, California, 0.7 miles east and 0.2 miles north of southwest corner sec. 12, T. 12 N., R. 18 E., 100 feet north of High Meadow Road, 4 miles south of Stateline. Date of Sampling: August 7, 1968.
Description by: L. N. Langan. Collectors: L. I. Larsen, B. R. Brasher, W. D. Nettleton.
Classification: Typic Cryopsamment, mixed, shallow.
Vegetation: Not cultivated. Jeffrey pine, white fir, greenleaf manzanita. Climate: Precipitation is 35 inches. Mean annual temperature 42°F., January 26°F., July 60°F. Parent Material: Granodiorite residuum. Topography: Mountainous uplands, 35 percent south slope. Elevation: 7000 feet. Drainage: Excessively drained; runoff is rapid; permeability is rapid. Soil Moisture: Dry.
Remarks: Field determinations for pH by Hellige-Truog triplex indicator. Composite A was sampled on 10 satellites.

HORIZON	DESCRIPTION
01 RSL 68704	(3-0 cm.) 1 to 0 inches, pine litter-undecomposed.
A11 RSL 68705	(0-5 cm.) 0 to 2 inches, grayish brown (10YR 5/2) gravelly coarse sand, very dark brown (10YR 2/2) moist; very dark grayish brown pockets; weak very fine and fine granular structure; soft, very friable, nonsticky and nonplastic; many very fine roots; many very fine and fine interstitial pores; 20 percent ½ inch minus gravel; medium acid (pH 6.0); clear wavy boundary.
A12 RSL 68706	(5-23 cm.) 2 to 9 inches, grayish brown (10YR 5/2) gravelly loamy coarse sand, very dark grayish brown (10YR 3/2) moist; dark gray and very dark grayish brown pockets and seams adjacent to coarse roots or old root channels; weak fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine, few coarse roots; many very fine and fine interstitial pores; 30 percent 3/4 inch minus gravel; slightly acid (pH 6.5); abrupt wavy boundary.
C1 RSL 68707	(23-36 cm.) 9 to 14 inches, pale brown (10YR 6/3) gravelly loamy coarse sand, dark grayish brown (10YR 4/2) moist; massive; soft, very friable, nonsticky and nonplastic; many very fine, few fine, medium and coarse roots; many very fine and fine interstitial pores; 40 percent ½ inch minus gravel; medium acid (pH 6.0); clear wavy boundary.
C2 RSL 68708	(36-43 cm.) 14 to 17 inches, pale brown (10YR 6/3) very gravelly coarse sand, dark grayish brown (10YR 4/2) moist; single grained; soft, very friable, nonsticky and nonplastic; many very fine, fine and medium roots; many very fine and fine interstitial pores; 55 percent ½ inch minus gravel; medium acid (pH 6.0); abrupt wavy boundary.
C3 RSL 68709	(43-69 cm.) 17 to 27 inches, weathering granodiorite grus; medium acid (pH 6.0).
C4 RSL 68710	(69-79 cm.) 27 to 31 inches, grus, less weathered than C3.

CONSECUTIVE PEDON NUMBER C 68 87 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES TOEM SOIL NO. S68 CALI. 9-31 SAMPLE NO. 8 713-8 720 TAHOE BASIN AREA

SOIL FAMILY TYPIC CRYOPSAMMENT MIXED,SHALLOW RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 181B,3A1 SANDS VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBONATE 3A1A	1A2A INT. II .2- .02	COARSE FRAGMENTS PCT. OF WHOLE SOIL	I 75- 20 PCT.	II 20-5 OF LT 75	III 5-2
			PERCENT OF LESS THAN 2MM																	
8 713	0- 8	A11	89.1	8.4	2.5	32.1	27.8	10.4	13.6	5.2	4.4	4.0				16.2	19A	--	19Z	N
8 714	8- 15	A12	91.2	6.9	1.9	44.6	25.7	8.3	9.4	3.2	3.6	3.3				11.2	20A	--	20Z	N
8 715	15- 25	AC	88.8	9.2	2.0	32.1	26.7	10.6	14.5	4.9	4.8	4.4				16.5	24A	--	24Z	N
8 716	25- 33	C 1	84.0	13.4	2.6	24.7	25.7	9.0	16.8	7.8	7.6	5.8				24.1	25A	--	25Z	N
8 717	33- 41	C 2	86.2	11.1	2.7	27.7	24.7	10.1	16.2	7.5	5.8	5.3				21.3	30A	--	30Z	N
8 718	41- 61	C 3												--	--		35A	--	35Z	N
8 719	61- 99	C 4															37A	--	37Z	N
8 720	3- 25	A															18A	--	18Z	N

SAMPLE NO.	DEPTH CM.	ORG. MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LL 4F2	PL 4F1	LI 4A1F	MIT EST. 1/3 BAR 4A1F	BULK DENSITY 1/3 BAR 4A1H	DRY 3B2	CM	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		PERCENT		PERCENT		PERCENT				PCT. NO.40		GRAMS PER CC					PERCENT			PERCENT	
8 713	0- 8	2.29	0.077			0.3				NP	NP		1.12	1.14	0.91		10.1	3.9		0.6	0.5
8 714	8- 15	0.94	0.034										1.19	1.21	0.90		6.5	3.0		0.6	0.5
8 715	15- 25	0.82	0.032			0.3				NP	NP		1.28	1.28	0.87		7.6	2.9			
8 716	25- 33	0.66				0.3				NP	NP		1.31	1.31	0.86		7.1	2.5			
8 717	33- 41	0.34											1.43	1.43	0.81		4.9	1.7			
8 718	41- 61	0.15											1.37	1.37	0.78		5.6	1.6			
8 719	61- 99												1.49	1.49	0.75		6.0	1.3			
8 720	3- 25	1.92	0.070									1.20			0.91			3.4			

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 584A 6N2E	MG 584A 602D	NA 584A 6P2A	K 584A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	CEC SUM 5A3A	CEC NH4-OAC 5A6A	BASE SATURATION SUM+ ACI-DITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 713	0- 8	3.1	0.3	0.1	0.3	3.8	5.0		8.8	11.4	43		33		5.6	5.0
8 714	8- 15	1.4	0.2	0.1	0.3	2.0	3.2		5.2	7.0	38		29		5.8	5.1
8 715	15- 25	0.9	0.2	TR	0.2	1.3	5.9		7.2	5.2	19		26		5.7	5.1
8 716	25- 33	0.6	0.1	0.1	0.2	1.0	4.5		5.5	4.8	18		21		5.6	5.0
8 717	33- 41	0.2	0.1	0.1	0.2	0.6			0.6	2.4	100+		25		5.8	5.2
8 718	41- 61	0.1	TR	0.1	0.1	0.3	1.6		1.9	2.4	18		14		5.6	5.0
8 719	61- 99	0.1	0.3	0.1	0.1	0.6			0.6	2.2	100+		27		5.5	4.9
8 720	3- 25														5.7	5.1

SAMPLE NO	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
		PERCENT										
8 713	0- 8	VR 2	KK 2								VR VERMICLT	1 TRACE
8 714	8- 15										KK KAOLINIT	2 SMALL
8 715	15- 25	VR 2	MI 2	KK 2								3 MODERATE
8 716	25- 33	VR 2	MI 2	KK 2							MI MICA	4 ABUNDANT
8 717	33- 41											5 DOMINANT
8 718	41- 61											6 INDETER.
8 719	61- 99											
8 720	3- 25											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

TOEM LOAMY COARSE SAND
S68Calif-9-31

Location: Tahoe Basin, El Dorado County, California, 1.3 miles south and 0.8 mile east of northwest corner, sec. 1, T. 12 N., R. 18 E., 210 feet upslope from apex of switchback on access road to Heavenly Valley Ski Lodge. Date of Sampling: August 13, 1968. Description by: L. I. Larsen. Collectors: L. C. Leifer, B. R. Brasher, W. D. Nettleton. Classification: Typic Cryopsamment, mixed, shallow.
Vegetation: Red fir, western white pine, prostrate manzanita. Climate: Precipitation is 40 inches. Mean annual temperature is 41°F., January 25°F., July 60°F. Parent Material: Granodiorite residuum. Topography: Mountainous uplands, 14 percent northeast slope. Elevation: 8000 feet. Drainage: Excessively drained; runoff is

medium; permeability is rapid. Soil Moisture: Top 2 inches moist, dry below.
Remarks: Field determinations for pH by Hellige-Truog triplex indicator. Composite A was taken from 10 satellite samples. 15 to 18 percent rock outcrops.

HORIZON	DESCRIPTION
	(1-0 cm.)
01	½ to 0 inches, nearly continuous 01 horizon from manzanita litter at immediate site; in vicinity more than half of surface lacks a thin 0 horizon but has a thin coat of "white" fine gravel and coarse sand grains.
	(0-8 cm.)
A11 RSL 68713	0 to 3 inches, dark grayish brown (10YR 4/2) loamy coarse sand, very dark grayish brown (10YR 3/2) moist; weak fine granular structure; soft, very friable, nonsticky and nonplastic; common very fine roots; many very fine and fine interstitial pores; slightly acid (pH 6.3); clear smooth boundary.
	(8-15 cm.)
A12 RSL 68714	3 to 6 inches, brown (10YR 5/3) loamy coarse sand, very dark grayish brown (10YR 3/2) moist; weak fine and medium granular structure; soft, very friable, nonsticky and nonplastic; common very fine and fine roots; many very fine and fine interstitial pores; slightly acid (pH 6.3); clear smooth boundary.
	(15-25 cm.)
AC RSL 68715	6 to 10 inches, light yellowish brown (10YR 6/4) light loamy coarse sand, dark grayish brown (10YR 4/2) moist; massive; soft with slightly hard lumps making up about 15 percent of the horizon, lumps have some accommodation and some pores, particularly tubular, very friable, nonsticky and nonplastic; common very fine and fine roots; many very fine and fine interstitial pores; medium acid (pH 6.0); clear smooth boundary.
	(25-33 cm.)
C1 RSL 68716	10 to 13 inches, light gray (2.5Y 7/2) coarse sand, dark grayish brown (2.5Y 4/2) moist; massive; soft, very friable, nonsticky and nonplastic; common very fine, few fine roots; many very fine and fine interstitial pores; slightly acid (pH 6.2); clear wavy boundary.
	(33-41 cm.)
C2 RSL 68717	13 to 16 inches, light gray (5Y 7/2) coarse sand, with black and white sand grains, grayish brown (2.5Y 5/2) moist; single grained; slightly hard, very friable, nonsticky and nonplastic; common very fine roots; common very fine interstitial pores; 25 percent gravel; has wick-rock structure; medium acid pH

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Xeric Torripsamment, mixed, mesic

SOIL Ormsby coarse sandy loam **variant** SOIL Nos. S61Calif-32-17 LOCATION Sierra Valley, Plumas County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 61718 - 61723

Depth (in.)	Horizon	1B1b Size class and particle diameter (mm) Total: Sand (2-0.05)	Total: Silt (0.05-0.002)	Total: Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	3A1 Int. II (0.2-0.02)	(2-0.2)			Coarse fragments 2A2 ≥ 2 Pct.	2 - 19 Pct. of < 76 mm	19-76 Pct. of < 76 mm
		Pct. of < 2 mm																
0-2	A11	88.1	7.2	4.7	8.5	22.0	16.1	32.4	9.1	3.8	3.4	30.0	61.9			8		
2-14	A12	84.4	9.8	5.8	6.3	19.0	15.5	33.3	10.3	6.0	3.8	35.4	55.0			9		
14-22	AC	85.2	7.8	7.0	6.1	13.2	16.2	37.7	12.0	3.9	3.9	37.0	52.1			7		
22-33	C1	85.7	7.8	6.5	5.3	13.4	15.6	39.4	12.0	5.6	3.2	39.0	51.3			7		
33-45	C2	84.8	8.2	7.0	10.2	15.9	14.5	31.5	12.7	4.8	3.4	36.1	53.5			12		
45-58+	C3	92.0	4.2	3.8	18.2	21.3	18.0	28.5	6.0	2.4	1.8	22.5	71.9			21		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		6E1b Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	4B2 15 bar Pct.			pH 8C1b Sat. Paste	8C1a 1:10
0-2	1.02	0.081	13		-								3.8			5.7	6.3
2-14	0.57	0.049	12		-								4.2			6.2	6.6
14-22	0.43	0.036	12		-								4.2			6.7	6.7
22-33	0.16				-								3.9			7.2	7.4
33-45	0.09				tr.								4.3			8.2	8.8
45-58+	0.05				tr.								2.7			8.5	9.0

Depth (in.)	Extractable bases 5B1a 6N2c Ca	6O2c Mg	6P2a Na	6Q2a K	6H2a Ext. Acidity	Cat. Exch. Cap. NaOAc 5A2a		5B1b Exch. Na	Water extract from saturated paste 8A1 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-2	3.7	1.9	0.1	0.7	2.9	8.8		0.1	4.4	3.2	0.3	1.9		-	1.7	0.1		1.16
2-14	4.9	1.6	0.2	0.6	2.4	10.4		0.2	1.0	0.6	0.4	0.4		-	0.8	0.1		0.26
14-22	4.6	1.9	0.1	0.6	1.7	10.0		0.1	1.0	0.6	0.7	0.3		-	1.0	-		0.24
22-33	4.0	4.7	0.2	0.4	0.8	8.6		0.2	1.0	1.1	0.6	0.2		-	1.2	-		0.16
33-45	8.4	4.7	0.2	0.3	-	9.1		0.2	0.9	2.2	1.5	0.1		0.2	3.1	0.1		0.10
45-58+	5.2	3.2	0.6	0.1	-	6.1		0.5	0.6	1.1	4.6	-		0.5	3.1	0.5		0.60

Depth (in.)	8A Water at Saturation Pct.	Exchangeable Na Pct.		Gypsum Pct.	5C3 Base Sat. (Sum) Pct.		
0-2	25.6				69		
2-14	22.4				75		
14-22	21.9				81		
22-33	21.8				92		
33-45	20.9				100		
45-58+	19.4				100		

ORMSBY LOAMY COARSE SAND variant 1/
S61Calif-32-17

Location: Sierra Valley, Plumas County, California. 0.6 mile north and 0.3 mile west of southeast corner of section 21, T. 22 N., R. 15 E. 0.1 mile north of road. About 4.7 miles north and 3.3 miles west of Loyalton. Date of Sampling: September 15, 1961.
Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, James McLaughlin, Eugene Begg.
Vegetation: The vegetation was silver sage and dryland sedge. This area has been sprayed resulting in almost 100 percent kill of the existing vegetation. Climate: Estimated annual precipitation 15 inches. Nearest weather station (Portola): Mean annual temperature 45.1° F.; January average 28.7° F.; July average 62.6° F.; mean annual precipitation 20.62 inches. Parent Material: Coarse textured alluvium dominantly from granitic sources. Relief: Flood plain with relatively smooth topography and a slope

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE La Posta loamy coarse sand (taxadjunct) **LOCATION** San Diego County, California

SOIL FAMILY: Typic Xerorpsamment mixed mesic

SOIL NOS. S60Calif-37-3 **LAB. NOS.** 60406 - 60410

DEPTH INCHES	HORIZON	1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
		PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)										
0-1½	A11	12.4	30.7	12.1	22.3	7.1	10.6	4.8	24.8	4.4	23	lcos
1½-6	A12	9.6	20.2	13.4	27.3	11.1	11.6	6.8	33.3	5.2	14	lcos
6-19	C1	12.3	26.7	11.4	21.2	7.9	11.9	8.6	25.8	5.3	27	lcos-cosl
19-30	C2	12.6	26.9	11.9	21.3	8.8	11.8	6.7	26.8	5.2	33	lcos
30-48 +	C3	17.4	34.0	11.6	18.9	6.2	8.5	3.4	21.3	3.4	40	cos

pH 8C1a H_2O 1:1	pH 8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	Atterberg limits Plastic limit Pct.	Liquid limit of 0.4	Plastic index mm.	Free Fe_2O_3 6C1b %	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.8	6.2		1.55	0.133	11.7				0.5			4.[illegible]
6.5	5.3		0.73	0.062	11.8				0.6			3.8
6.7	4.7		0.33	0.028		N.P.	-	N.P.	0.7			5.4
6.9	4.3		0.14	0.015					0.6			5.3
6.8	4.3		0.06			N.P.	-	N.P.	0.5			3.8

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 6N2b Ca	602c Mg	6H2a H	6P2a Na	6Q2a K	5B1c BASE SAT. % 5C3	Sum of Cations me./100 g. soil 5A3a	K				MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →							milliequivalents per liter →				
9.1	7.6	1.0	1.5	0.1	0.5	86	10.7					
9.1	6.2	0.8	2.1	0.1	0.2	78	9.4					

LA POSTA LOAMY COARSE SAND (taxadjunct) 1/

S60Calif-37-3

Description: George Borst
Date of Sampling: November 9, 1962
Collectors: Reuben Nelson, Lothair Grant, Fred C. Eggers, John Wachtell, and George Borst
Location: Approximately 1000 feet west of southeast corner, sec. 18, T. 11 S., R. 5 E., 75 feet north of Montezuma Valley Road at entrance to Chimney Rock Ranch, **San Diego County, California.**
Elevation: 4,150 feet
Relief: Strongly sloping, 10 percent
Use: Range
Vegetation: Native chaparral
Drainage: Somewhat excessively drained
Parent Material: Deeply weathered granitic rock
Climate: This soil occurs under about 17" annual rainfall. The mean annual temperature is about 57°, the mean January temperature about 44°, and the mean July temperature about 73°.
Remarks: A layer of angular quartz fragments, 2 to 6 mm. in size, occurs on the surface. This soil is somewhat shallower and has less prominent clay films than is considered representative of the series.
Reaction: pH is determined in the field with a Truog (Hellige) soil reaction kit
Classification: **Typic Xeropsamment, mixed, mesic.**

HORIZON		DESCRIPTION
A11	0-1½"	Grayish brown (10YR 5/2) loamy coarse sand, containing light gray (10YR 7/2) angular quartz fragments, very dark grayish brown (10YR 3/2) when moist; weak very fine granular structure; soft, very friable, nonsticky and nonplastic; micaceous; abundant fine random roots; neutral (pH 7.0); clear wavy boundary.
A12	1½-6"	Dark grayish brown (10YR 4/2) loamy coarse sand, very dark grayish brown (10YR 3/2) when moist; weak fine crumb structure; soft, friable, nonsticky and nonplastic; abundant fine random pores; micaceous; slightly acid (pH 6.2); abrupt wavy boundary.
C1	6-19"	Brown (10YR 5/3) loamy coarse sand, containing very pale brown (10YR 7/3) and dark gray (10YR 4/1) mineral particles, brown (10YR 4/3), pale brown (10YR 6/3), and black (10YR 2/1) when moist; massive; hard, friable, nonsticky and nonplastic; plentiful medium random roots; common fine random tubular pores; colloid in bridges between mineral grains; micaceous; medium acid (pH 6.0); gradual smooth boundary.
C2	19-30"	Brown (10YR 5/3), very pale brown (10YR 7/3), and dark gray (10YR 4/1) disintegrated granite with loamy coarse sand texture, dark brown (10YR 4/3) pale brown (10YR 6/3), and very dark gray (10YR 3/1) when moist; massive; hard, friable, nonsticky and nonplastic; few medium random roots; medium acid (pH 6.0); gradual smooth boundary.
C3	30-48"+	Very pale brown (10YR 7/3) brown (10YR 5/3), and dark gray (10YR 4/1) disintegrated granite with coarse sand texture, pale brown (10YR 6/3) dark brown (10YR 4/3), and very dark gray (10YR 3/1) when moist; massive; hard, friable, nonsticky and nonplastic; very few coarse random roots; medium acid (pH 6.0).

1/ This pedon has slightly less organic carbon and slightly paler moist colors than the La Posta series.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Cieneba very fine sandy loam variant **LOCATION** Riverside County, California

SOIL FAMILY: Typic Xeropsamment, mixed, shall w, thermic

SOIL NOS. S55Calif-33-8 **LAB. NOS.** 55565-568

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-1	A11	4.6	9.9	9.8	35.0	19.0	16.7	5.0	55.2	3.0	9	lfs
1-13	A12	3.8	10.5	12.5	39.6	16.3	11.9	5.4	49.4	2.3	39	ls
13-32	C	3.0	14.8	19.3	43.4	10.5	6.9	2.1	40.2	0.2	35	s
32-50	Dr	6.2	15.6	14.7	34.3	14.2	13.0	2.0	43.7	3.4	22	ls

pH 8C1b SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Condc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B1a 1/10 ATMOS. %	4B1a 1/3 ATMOS. %	4B2 15 ATMOS. %
5.9		6.0	0.80	0.074	10.8		0.32			19.0	9.4	3.1
6.5		6.5	0.30	0.029	10.3		0.26			14.6	7.3	3.3
6.9		6.3	0.06				0.12			7.5	4.4	2.2
6.8		6.4	0.02				0.11			16.2	8.5	2.8

CIENEBA VERY FINE SANDY LOAM variant 1/
S55Calif-33-8

Collectors: George Borst, Arnold Knecht, and R. E. Nelson.

Date of Sampling: December 5, 1955. Description by: W. G. Harper

Location: Searl Bros. Ranch in Diamond Valley on rocky knob, N. side of Newport Road near N 1/4 corner of Sec. 5, T6S, R1W, 2 miles W of State St., about 5 miles SW of Hemet, Riverside County, California.

Elevation: 1650 feet

Precipitation: 13 inches

Classification: Typic Xeropsamment, mixed, shallow, thermic.

Vegetation: California sage, wild buckwheat, cheat grass, wild oats and some squirrel tail.

Parent Material: Mica schist, triassic marine meta-sediments.

Physiography: On top of narrow ridge in rolling hills with moderate amount of rock outcrop 3 to 4 feet above surface and 3 to 10 feet across.

Drainage: Well drained Remarks: This soil is inusually high in mica.

HORIZON		DESCRIPTION
A11	0- 1"	Brown (10YR 5/3 dry, very dark grayish brown, 3/2 moist) very fine sandy loam; weakly defined very fine crumb structure; soft, loose; moderately permeable; many plant roots and much organic matter; pH 7.0; abrupt smooth boundary.
A12	1-13"	Brown (10YR 5/3 dry, dark brown 3/3 moist) very fine sandy loam; weakly defined very fine crumb structure to nearly massive; soft, very friable; moderately permeable; few roots and medium to low content of organic matter; pH 6.5; clear smooth boundary.
C	13-32"	Soft decomposing rock containing salt and pepper-like material composed of many gray (10YR 5/1) nearly colorless quartz fragments; massive; extremely hard, extremely firm; slowly permeable; very few

CONSECUTIVE PEDON NUMBER C 68 80 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES ELMIRA SOIL NO. S68 CAL1. 9-22 SAMPLE NO. 8 664-8 673 TAHOE BASIN AREA

SOIL FAMILY ALFIC XEROPSAMMENT MIXED,FRIGID RIVERSIDE SOIL SURVEY LABORATORY

SIZE, CLASS AND PARTICLE DIAMETER (MM) 181B,3A1 1A2A

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SANDS VCS 2-1	CS 1-.05	MS .5-.25	FS .25-.10	VFS .1-.05	CSI .05-.02	FSI .02-.002	SILTS CO. 2-.2U	FINE .2U	CLAYS CARBO LT .2U	NATE 3A1A	INT. II .2-.02	COARSE FRAGMENTS PCT. I OF 75- WHOLE SOIL	II 20-5 20 PCT.	III 5-2 OF LT 75
			PERCENT OF LESS THAN 2MM																	
8 664	0- 3	A11	90.7	7.3	2.0	26.9	33.6	13.4	12.5	4.3	3.7	3.6					13.7	11A	--	11Z N
8 665	3- 10	A12	88.9	8.0	3.1	31.9	24.5	11.6	14.7	6.2	4.2	3.8					18.0	15A	--	15Z N
8 666	10- 23	A3	86.5	9.9	3.6	20.6	25.6	13.6	18.0	8.7	5.0	4.9					23.4	15A	--	15Z N
8 667	23- 36	B2	87.3	9.0	3.7	16.4	27.7	16.3	19.3	7.6	3.9	5.1					21.3	14A	--	14Z N
8 668	36- 63	B3	86.0	10.3	3.7	20.8	26.5	13.8	17.4	7.5	5.5	4.8					22.1	14A	--	14Z N
8 669	63- 99	C 1	86.0	10.6	3.4	12.2	26.7	16.7	21.1	9.3	4.9	5.7					25.2	13A	--	13Z N
8 670	99-135	C 2	89.1	8.1	2.8	18.0	29.6	14.0	19.8	7.7	3.9	4.2					22.3	16A	--	16Z N
8 671	135-155	2C 3	78.1	17.9	4.0	9.6	19.1	11.5	24.8	13.1	8.4	9.5					36.3	4A	--	4Z N
8 672	155-175	2C 4	75.9	19.4	4.7	4.1	11.1	9.0	31.9	19.8	10.5	8.9					49.1	1A	--	1Z N
8 673	0- 23	A																18A	--	18Z N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6B5A	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		PERCENT				PCT.NO.40		GRAMS PER CC			PERCENT			-PERCENT-	
8 664	0- 3	2.65	0.091			0.5						1.30		0.94			4.2		
8 665	3- 10	1.32	0.044			0.5				NP	NP	1.39	1.41	0.91	8.1	2.7	0.5	0.4	
8 666	10- 23	0.78	0.028			0.6						1.49	1.49	0.91	8.0	2.9			
8 667	23- 36	0.42				0.6				NP	NP	1.53	1.53	0.91	6.4	2.7			
8 668	36- 63	0.29				0.6						0.78	1.25	0.91	78.0	2.4	17.0	0.2	
8 669	63- 99	0.20				0.6						1.54	1.54	0.92	6.4	2.3			
8 670	99-135	0.06										1.55	1.55	0.90	5.6	1.7			
8 671	135-155											1.53	1.53	0.98	9.7	2.7			
8 672	155-175									NP	NP	1.50	1.50	0.99	9.7	3.6			
8 673	0- 23	1.40	0.054									1.45		0.89		2.7			

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACIDITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACIDITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CACL2 8C1E
		MEQ/100G									PERCENT					
8 664	0- 3	2.2	0.3	0.1	0.2	2.8		0.4	2.8	8.8	100+	88	32		5.4	4.5
8 665	3- 10	1.8	0.2	0.1	0.2	2.3	6.5	0.5	8.8	7.1	26	82	32		5.7	4.8
8 666	10- 23	1.8	0.2	0.1	0.3	2.4	5.5		7.9	6.0	30		40		6.0	5.1
8 667	23- 36	1.5	0.1	TR	0.2	1.8	6.5		8.3	5.1	22		36		6.0	5.2
8 668	36- 63	1.2	0.1	0.1	0.2	1.6	3.8		5.4	4.3	30		37		5.9	5.2
8 669	63- 99	0.5	0.3	0.1	0.2	1.1	2.4		3.5	3.5	31		31		5.9	5.1
8 670	99-135	1.3	0.3	0.1	0.1	1.8			1.8	1.6	100+		100+		6.0	5.2
8 671	135-155	3.4	0.5	0.1	0.1	4.1			4.1	5.3	100+		77		5.9	5.2
8 672	155-175	4.6	0.7	0.1	0.1	5.5			5.5	7.2	100+		76		5.9	5.2
8 673	0- 23	2.5	0.3	0.1	0.3	3.2	6.5		9.7	8.6	33		37		5.9	5.1

CLAY MINERALOGY

SAMPLE NO	DEPTH CM.	LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 664	0- 3											1 TRACE
8 665	3- 10	VR 3	KK 2			MI 1					VR VERMICLT	2 SMALL
8 666	10- 23										KK KAOLINIT	3 MODERATE
8 667	23- 36	VR 3	KK 3								MI MICA	4 ABUNDANT
8 668	36- 63											5 DOMINANT
8 669	63- 99											6 INDETER.
8 670	99-135											
8 671	135-155											
8 672	155-175	KK 3	MI 2	VR 1								
8 673	0- 23											

- SAMPLE PREPARATION CODES
  A=STANDARD PREPARATION.
  N=NOT DETERMINED.
  Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

ELMIRA LOAMY COARSE SAND
S68Calif-9-22

Location: Tahoe Basin, El Dorado County, California. 0.7 miles east and 0.4 miles south of northwest corner, sec. 3, T. 12 N., R. 18 E., 300 feet from intersection of Happy Home Cemetery Road, 70 feet northeast of Al Tahoe Blvd., 1 mile south of Bijou. Date of Sampling: August 7, 1968.

Description by: L. N. Langan. Collectors: J. H. Rogers, G. M. Kennedy, R. C. Huff, B. R. Brasher, W. D. Nettleton.

Classification: Alfic Xeropsamment, mixed, frigid.

Vegetation: Not cultivated; Jeffrey pine, big sagebrush, bitterbrush, squirrel tail, and other scattered perennials, lupine. Climate: Precipitation is 25 to 30 inches. Mean annual temperature is 43°F., January 27°F., July 61°F. Parent Material: Mixed, predominantly granitic, alluvium. Topography: Nearly level glacial outwash, 1 percent slope. Elevation: 6265 feet. Drainage: Somewhat excessively drained; runoff is slow; permeability is very rapid. Soil Moisture: Essentially dry.

Remarks: Field determination for pH by Hellige-Truog triplex indicator. Composite A taken from 10 satellite samples. Thin mat of pine needles on surface.

HORIZON	DESCRIPTION
A11 RSL 68664	(0-3 cm.) 0 to 1 inch, dark grayish brown (10YR 4/2) loamy coarse sand, very dark brown (10YR 2/2) moist; single grained; loose, very friable, nonsticky and nonplastic; common very fine roots; many very fine and fine interstitial pores; strongly acid (pH 5.2); abrupt smooth boundary.
A12 RSL 68665	(3-10 cm.) 1 to 4 inches, dark grayish brown (10YR 4/2) loamy coarse sand, very dark grayish brown (10YR 3/2) moist; weak fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine, common fine and medium roots; many very fine and fine interstitial pores; strongly acid (pH 5.5); clear wavy boundary.
A3 RSL 68666	(10-23 cm.) 4 to 9 inches, brown (10YR 5/3) loamy coarse sand, dark brown (10YR 3/3) moist; massive; soft, very friable, nonsticky and nonplastic; many very fine, few fine and medium roots; many very fine interstitial pores; medium acid (pH 6.0); clear wavy boundary.
B2 RSL 68667	(23-36 cm.) 9 to 14 inches, light yellowish brown (10YR 6/4) loamy coarse sand, dark yellowish brown (10YR 4/4) moist; massive; soft, very friable, nonsticky and nonplastic; many very fine, few fine and medium roots; many very fine interstitial pores; slightly acid (pH 6.2); clear wavy boundary.
B3 RSL 68668	(36-63 cm.) 14 to 25 inches, light yellowish brown (10YR 6/4) loamy coarse sand, strong brown (7.5YR 5/6) moist; massive; soft, very friable, nonsticky and nonplastic; many fine, medium and coarse, common very fine roots; many very fine interstitial pores; 10 percent gravel; slightly acid (pH 6.2); gradual smooth boundary.
C1 RSL 68669	(63-99 cm.) 25 to 39 inches, pale brown (10YR 6/3) light loamy coarse sand, strong brown (7.5YR 5/6) moist; massive; soft, very friable, nonsticky and nonplastic; common very fine, fine, medium and coarse roots; many very fine interstitial pores; 10 percent gravel; slightly acid (pH 6.2); gradual smooth boundary.
C2 RSL 68670	(99-135 cm.) 39 to 53 inches, very pale brown (10YR 8/3) coarse sand with bands of coarse sandy loam, very pale brown (10YR 8/3) moist; massive; soft, very friable, nonsticky and nonplastic; coarse sandy loam bands slightly plastic; common very fine and fine roots; many very fine interstitial and few tubular pores; few 1/8 inch minus clay bands, wavy, discontinuous, and horizontal, other bands unoriented; bands have few moderately thick and many thin clay films in bridges; 5 percent gravel; slightly acid (pH 6.2); abrupt smooth boundary.
IIC3 RSL 68671	(135-155 cm.) 53 to 61 inches, very pale brown (10YR 8/3) sand, very pale brown (10YR 8/3) moist; massive; soft and slightly hard, very friable; few very fine and fine roots; many very fine interstitial and few tubular pores; clay pockets and 3/4 inch minus discontinuous clay bands; pockets and bands have many moderately thick clay films in bridges; strongly acid (pH 5.5); clear smooth boundary.
IIC4 RSL 68672	(155-175 cm.) 61 to 69 inches, very pale brown (10YR 8/3) fine sandy loam, very pale brown (10YR 8/3) moist; massive; slightly hard and very hard, very friable and friable; few very fine and fine roots; many very fine interstitial and few very fine tubular pores; common ½ inch minus clay bands and pockets of clay; common very coarse distinct strong brown (7.5YR 5/6) iron stains; many moderately thick clay films in bridges of bands and pockets; strongly acid (pH 5.5).

CONSECUTIVE PEDON NUMBER C 68 75 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES ELMIRA TAXADJUNCT SOIL NO. S68 CALI. 9-26 SAMPLE NO. 8 602-8 613 TAHOE BASIN AREA

SOIL FAMILY ALFIC XEROPSAMMENT MIXED,FRIGID RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 TOTAL FRACTION 2- .05	SILT .05- .002	CLAY LT .002	SANDS VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBONATE 3A1A.02	1A2A INT. II .2-	COARSE FRAGMENTS PCT. OF WHOLE SOIL	I 75- 20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75			
8 602	0- 0	01															--A	--	--Z	--
8 603	0- 4	A11	84.2	10.6	5.2	28.5	26.1	12.5	13.1	4.0	4.6	6.0				14.2	20A	8	12Z	N
8 604	4- 13	A12	82.8	11.9	5.3	23.2	24.8	13.3	16.2	5.3	5.1	6.8				17.5	22A	6	16Z	N
8 605	13- 28	A3	81.3	12.7	6.0	16.1	23.3	15.2	20.1	6.6	5.5	7.2				21.2	24A	7	17Z	N
8 606	28- 48	B1 T	80.8	12.5	6.7	13.7	23.2	16.0	20.8	7.1	4.9	7.6				21.5	21A	12	9Z	N
8 607	48- 74	2B21T	80.0	12.6	7.4	12.5	22.8	16.4	21.3	7.0	4.6	8.0	--	--		20.9	43A	36	7Z	N
8 608	74- 89	3B22T	83.0	8.6	8.4	18.7	28.7	13.9	16.5	5.2	3.6	5.0				16.5	26A	17	9Z	N
8 609	89-100	3B31T	86.3	7.8	5.9	17.5	34.8	14.4	14.9	4.7	2.9	4.9				14.3	27A	8	19Z	N
8 610	100-130	3B32T	87.1	8.5	4.4	11.2	30.5	17.5	21.5	6.4	4.0	4.5				20.4	26A	15	11Z	N
8 611	130-160	4C 1SI	86.6	8.5	4.9	7.9	21.9	20.7	27.7	8.4	4.4	4.1				26.2	36A	26	10Z	N
8 612	160-175	4C 2	88.7	7.0	4.3	20.4	26.3	17.5	20.3	4.2	4.1	2.9				17.1	35A	12	23Z	N
8 613	0- 28	A															15A	--	15Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR-BON	NITRO-GEN	CARBONATE AS CACO3 LT 2	2-20	SESQUIOXIDES DI-CIT EXT FE	AL	PYROP EXT FE	AL	ATTERBERG L I M I T LL	PL	EST. 1/3 BAR	BULK DENSITY DRY 1/3 BAR		CM WATER CONTENT 1/10 BAR	1/3 BAR	15 BAR	EXTNSBLTY LEF	LE
		6A1A	6B1A	6E1B	6E1B	6C2B	6G7A	6C5A	6G5A	4F2	4F1	4A1F	4A1F	4A1H	3B2	4B1C	4B1C	4B2	4D1	4D1
		PERCENT		PERCENT		PERCENT				PCT	NO 40	GRAMS	PER	CC				PERCENT		
8 603	0- 4	3.48	0.119		2.1					NP	NP		0.52	0.65	0.95		31.9	5.0	7.7	7.4
8 605	13- 28	1.01	0.042			2.8				NP	NP		1.35	1.35	0.86		7.6	4.0		
8 606	28- 48	0.59				2.9				NP	NP		1.46	1.46	0.87		8.3	3.8		
8 607	48- 74	0.21				3.0							1.61	1.61	0.69		7.5	3.8		
8 608	74- 89	0.11				2.4				NP	NP		1.48	1.48	0.84		8.4	4.2		
8 609	89-100	0.11				1.7				NP	NP		1.47	1.47	0.83		8.0	3.3		
8 610	100-130	0.13				1.8							1.36	1.36	0.85		11.7	2.9		
8 611	130-160					1.9							1.53	1.53	0.75		11.5	3.2		
8 612	160-175					2.2				NP	NP		1.63	1.63	0.75		9.9	3.3		
8 613	0- 28	1.52	0.066									1.20			0.93			4.1		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA	MG	NA	K	SUM	EXT. ACI-DITY	KCL EXT. AL	C E C SUM	NH4-OAC	BASE SATURATION SUM+ ACI-DITY	SUM+ AL	NH4-OAC	PH NAF 2 MIN	H2O 1/1	CA-CL2
		5B4A	5B4A	5B4A	5B4A											
		6N2E	6D2D	6P2A	6Q2A		6H2A	6G1E	5A3A	5A6A			5C1	8C1D	8C1A	8C1E
		MEQ/100G									PERCENT					
8 602	0- 0															
8 603	0- 4	6.6	0.5	0.1	0.2	7.4	7.7		15.1	14.4	49		51		5.7	5.0
8 604	4- 13	5.2	0.4	0.1	0.2	5.9	6.9		12.8	10.3	46		57		6.0	5.3
8 605	13- 28	3.7	0.3	TR	0.2	4.2	4.9		9.1	7.6	46		56		6.3	5.6
8 606	28- 48	2.7	0.3	TR	0.2	3.2	4.5		7.7	5.7	42		57		6.2	5.4
8 607	48- 74	2.0	0.3	0.1	0.2	2.6			2.6	4.3	100+		60		6.2	5.5
8 608	74- 89	2.1	0.3	0.1	0.3	2.8			2.8	3.9	100+		72		6.2	5.6
8 609	89-100	1.7	0.3	0.1	0.2	2.3			2.3	3.5	100+		66		6.3	5.6
8 610	100-130	1.8	0.3	TR	0.2	2.3			2.3	3.7	100+		63		6.2	5.6
8 611	130-160	1.7	0.4	0.1	0.2	2.4			2.4	3.6	100+		67		6.2	5.5
8 612	160-175	1.6	0.6	0.1	0.1	2.4			2.4	3.7	100+		65		6.2	5.6
8 613	0- 28															

CLAY MINERALOGY

SAMPLE NO	DEPTH CM.	LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 602	0- 0											1 TRACE
8 603	0- 4	VR 3	KK 3								VR VERMICLT	2 SMALL
8 604	4- 13										KK KAOLINIT	3 MODERATE
8 605	13- 28	VR 3	KK 3								MI MICA	4 ABUNDANT
8 606	28- 48	VR 3	KK 3	MI 2								5 DOMINANT
8 607	48- 74	KK 3	VR 2	MI 1								6 INDETER.
8 608	74- 89	KK 3	VR 1									
8 609	89-100											
8 610	100-130											
8 611	130-160											
8 612	160-175	KK 3										
8 613	0- 28											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.
N=NOT DETERMINED

ELMIRA LOAMY COARSE SAND (TAXADJUNCT)*
S68Calif-9-26

Location: Tahoe Basin, El Dorado County, California. 1600 feet east and 400 feet north of center sec. 11, T. 12 N., R. 18 E., east of Cold Creek Trail Road, 500 feet east of water tank, 3 miles south of Bijou. Date of Sampling: August 8, 1968.
Description by: L. N. Langan. Collectors: J. H. Rogers, G. M. Kennedy, D. Palmquist, B. R. Brasher, W. D. Nettleton, G. L. Anderson.
Classification: Alfic Xeropsamment, mixed, frigid.
Vegetation: Not cultivated; Jeffrey pine, white fir, red fir, greenleaf manzanita, mountain whitethorn, squaw carpet, mules ear. Climate: Precipitation is 30 inches. Mean annual temperature is 42°F. to 45°F., January 27°F., July 61°F. Parent Material: Mixed gravelly alluvium. Topography: Old glacial outwash terrace, 3 percent slope. Elevation: 6600 feet. Drainage: Well drained; runoff is slow; permeability is moderate over slow. Soil Moisture: Essentially dry.
Remarks: Field determinations for pH by Hellige-Truog triplex indicator. Composite A taken from 10 satellite samples. *Has weak cementation in lower part not included in current concept of series.

HORIZON	DESCRIPTION
	(1-0 cm.)
01	1/8 to 0 inches, pine needles - intermittent, about 1 inch near trees.
RSL 68602	(0-4 cm.)
A11 RSL 68603	0 to 1½ inches, brown (7.5YR 4/4) loamy coarse sand, dark brown (7.5YR 3/2) moist; weak fine and medium granular structure; soft, very friable, nonsticky and nonplastic; common very fine roots; many very fine and fine interstitial pores; 10 percent 1 inch minus gravel; medium acid (pH 5.7); abrupt smooth boundary.
A12 RSL 68604 (4-13 cm.)	1½ to 5 inches, brown (7.5YR 4/4) loamy coarse sand, dark brown (7.5YR 3/4) moist; moderate fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine, few fine roots; many very fine and fine interstitial pores; 10 percent 1 inch minus gravel; medium acid (pH 6.0); clear wavy boundary.
A3 RSL 68605 (13-28 cm.)	5 to 11 inches, reddish brown (5YR 5/4) loamy coarse sand, dark reddish brown (5YR 3/4) moist; weak fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine, common fine and medium roots; many very fine and fine interstitial pores; 10 percent 1 inch minus gravel; medium acid (pH 6.0); clear wavy boundary.
B1t RSL 68606 (28-48 cm.)	11 to 19 inches, reddish brown (5YR 5/4) loamy coarse sand, reddish brown (5YR 4/4) moist; weak fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine, fine, medium and coarse roots; many very fine and fine interstitial pores; few thin clay films in bridges; 10 percent 1 inch minus gravel; medium acid (pH 6.0); abrupt wavy boundary.
IIB21t RSL 68607 (48-74 cm.)	19 to 29 inches, light reddish brown (5YR 6/4) very gravelly coarse sandy loam, yellowish red (5YR 4/6, 5/6 crushed) moist; massive; hard, very friable, nonsticky and nonplastic; common very fine, few fine, medium and coarse roots; common very fine and fine interstitial, few fine tubular pores; common thin clay films in bridges, many thin clay films coating grains; 45 percent gravel and 10 percent cobbles; clear wavy boundary.
IIIB22t RSL 68608 (74-89 cm.)	29 to 35 inches, strong brown (7.5YR 5/6) dry and moist, gravelly coarse sandy loam; massive; hard, very friable, nonsticky and slightly plastic; few very fine, fine and medium roots; common very fine and fine interstitial pores; many moderately thick and few thick yellowish red (5YR 5/6) clay films in bridges; 25 percent gravel; medium acid (pH 6.0); clear wavy boundary.
IIIB31t RSL 68609 (89-100 cm.)	35 to 40 inches, strong brown (7.5YR 5/6) gravelly loamy coarse sand, strong brown (7.5YR 5/6) moist; massive; slightly hard, very friable, nonsticky and nonplastic; few very fine, fine and medium roots; many very fine and fine interstitial pores; thin ¼ inch minus reddish brown (5YR 4/4) clay bands with few thick and common thin clay films in bridges; 30 percent gravel; medium acid (pH 6.0); clear wavy boundary.
IIIB32t RSL 68610 (100-130 cm.)	40 to 51 inches, reddish yellow (7.5YR 6/6) gravelly light loamy coarse sand, strong brown (7.5YR 5/6) moist; massive; soft, very friable, nonsticky and nonplastic; few very fine, fine and medium roots; many very fine and fine interstitial pores; thin reddish brown (5YR 4/4) discontinuous clay bands with very few thin and moderately thick clay films in bridges and coating sand grains; 15 percent 1 inch minus gravel; medium acid (pH 6.0); abrupt wavy boundary.
IVC1si RSL 68611 (130-160 cm.)	51 to 63 inches, reddish yellow (7.5YR 6/6) very gravelly coarse sand, reddish yellow (7.5YR 6/6) moist; massive; very hard with hard and slightly hard areas, very friable to firm, nonsticky and nonplastic; roots matted on top horizon; many very fine and fine interstitial pores; few moderately thick strong brown (7.5YR 5/6) clay films in bridges and few thin and moderately thick clay films coating sand grains; 70 percent gravel; few pockets appear to be cemented by silica and clay; some clods do not slake in water, but most do; medium acid (pH 6.0); gradual wavy boundary.
IVC2 RSL 68612 (160-175 cm.)	63 to 69 inches, reddish yellow (7.5YR 6/6) very gravelly coarse sand, strong brown (7.5YR 5/6) moist; massive; slightly hard, very friable, nonsticky and nonplastic; few very fine, fine and medium roots; many very fine and fine interstitial pores; 50 percent gravel; medium acid (pH 6.0).

CONSECUTIVE PEDON NUMBER C 68 71 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES ELMIRA SOIL NO. S68 CALI. 9-28 SAMPLE NO. 8 565-8 575 TAHOE BASIN AREA

SOIL FAMILY ALFIC XEROPSAMMENT MIXED,FRIGID RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A.02	INT. II .2- WHOLE 20	1A2A COARSE FRAGMENTS PCT. OF SOIL	I 75-	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														--A	--	--	--
8 565	4- 0	01																		
8 566	0- 5	A11	83.3	11.1	5.6	31.7	27.6	10.1	10.3	3.6	3.6	7.5				12.0	24A	--	24Z	N
8 567	5- 17	A12	81.0	12.0	7.0	28.4	27.4	10.4	10.6	4.2	4.0	8.0				13.2	25A	6	19Z	N
8 568	17- 48	C 1	79.7	12.4	7.9	19.2	29.2	13.0	13.2	5.1	4.4	8.0				15.8	18A	--	18Z	N
8 569	48- 69	C 2	78.2	12.7	9.1	18.2	26.6	12.7	14.5	6.2	4.6	8.1				17.9	31A	12	19Z	N
8 570	69- 79	2B1	79.8	11.0	9.2	17.2	31.7	13.7	12.9	4.3	3.2	7.8				13.8	47A	30	17Z	N
8 571	79- 89	2B2	79.8	11.1	9.1	19.6	29.5	12.4	13.5	4.8	3.7	7.4				15.0	45A	17	28Z	N
8 572	89-110	2B3	83.9	8.6	7.5	27.1	32.1	10.6	10.5	3.6	2.9	5.7				11.2	42A	26	16Z	N
8 573	110-150	2C 1															28A	7	21Z	N
8 574	150-185	3C2	87.1	8.7	4.2	23.9	33.0	13.9	13.0	3.3	4.3	4.4				13.2	28A	16	12Z	N
8 575	0- 17	A															20A	--	20Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		PERCENT		PERCENT		PERCENT				PCT.NO.40		GRAMS PER CC				PERCENT			PERCENT	
8 565	4- 0											0.40								
8 566	0- 5	3.77	0.137				0.1					0.45			0.95			5.7		
8 567	5- 17	1.94	0.070							NP	NP		1.27	1.28	0.86		11.1	4.6	0.3	0.2
8 568	17- 48	0.52											1.34	1.34	0.90		9.3	4.6		
8 569	48- 69	0.34											1.38	1.39	0.81		11.4	5.1	0.2	0.2
8 570	69- 79	0.15											1.36	1.37	0.69		14.6	4.8	0.2	0.2
8 571	79- 89	0.14								NP	NP		1.36	1.36	0.71		13.8	6.2		
8 572	89-110												1.23	1.24	0.74		14.3	4.4	0.3	0.2
8 573	110-150											1.45			0.82			2.7		
8 574	150-185									NP	NP	1.45			0.83			2.5		
8 575	0- 17	1.87	0.071									1.20			0.90			4.5		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6Q2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACI-DITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 565	4- 0															
8 566	0- 5	2.8	0.4	0.1	0.4	3.7	7.9	0.8	11.6	13.3	32	82	28		5.3	4.5
8 567	5- 17	2.7	0.3	TR	0.3	3.3	7.1		10.4	10.1	32		33		5.9	5.0
8 568	17- 48	1.6	0.2	TR	0.2	2.0	5.1		7.1	6.8	29		30		5.9	5.1
8 569	48- 69	1.9	0.3	0.1	0.3	2.6	4.3		6.9	6.8	38		38		5.8	5.0
8 570	69- 79	2.0	0.3	TR	0.4	2.7	3.9		6.6	6.1	41		45		5.8	5.1
8 571	79- 89	2.2	0.4	0.1	0.5	3.2	3.3		6.5	6.6	49		48		5.8	5.2
8 572	89-110	1.6	0.3	0.1	0.5	2.5	3.2		5.7	5.1	44		49		5.7	5.0
8 573	110-150	1.4	0.3	0.1	0.2	2.0	1.8		3.8	4.6	53		43		5.8	4.9
8 574	150-185	1.3	0.3	0.1	0.2	1.9	2.2		4.1	4.2	46		45		5.8	5.0
8 575	0- 17															

SAMPLE NO	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 565	4- 0											1 TRACE
8 566	0- 5										VR VERMICLT	2 SMALL
8 567	5- 17	VR 3	KK 3								KK KAOLINIT	3 MODERATE
8 568	17- 48										VM VM/MICA	4 ABUNDANT
8 569	48- 69											5 DOMINANT
8 570	69- 79											6 INDETER.
8 571	79- 89	VR 3	KK 3	VM 2								
8 572	89-110											
8 573	110-150											
8 574	150-185	KK 3	VM 3	VR 2								
8 575	0- 17											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

ELMIRA GRAVELLY LOAMY COARSE SAND
S68Calif-9-28

Location: Tahoe Basin, El Dorado County, California, 2800 feet east and 1700 feet south of the northwest corner of sec. 21, T. 12 N., R. 18 E., 3 miles south of Tahoe Valley, north of Pioneer Trail Road. Date of Sampling: August 8, 1968.
Description by: L. N. Langan. Collectors: J. H. Rogers, G. M. Kennedy, L. I. Larsen, D. Palmquist, B. R. Brasher, W. D. Nettleton.
Classification: Alfic Xeropsamments, mixed, frigid.
Vegetation: Not cultivated; Jeffrey pine, white fir, lodgepole pine, squaw carpet, greenleaf manzanita, mountain whitethorn. Climate: Precipitation is 30 to 35 inches. Mean annual temperature is 43°F., January 27°F., July 61°F. Parent Material: Mixed alluvium. Topography: Crest of lateral moraine, pre-Tahoe?, 1 percent slope. Elevation: 6,502 feet. Drainage: Somewhat excessively drained; runoff is slow; permeability is rapid. Soil Moisture: Dry.
Remarks: Field determinations for pH by Hellige-Truog triplex indicator. Composite A taken from 10 satellite samples.

HORIZON	DESCRIPTION
O1	(1-0 cm.) 1/2 to 0 inches, pine litter.
RSL 68565 A11	(0-5 cm.) 0 to 2 inches, grayish brown (10YR 5/2) gravelly loamy coarse sand, very dark grayish brown (10YR 3/2) moist; seams of dark gray (10YR 4/1) dry; moderate fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine roots; many very fine and fine interstitial pores; 17 percent gravel; medium acid (pH 5.8); abrupt wavy boundary.
RSL 68566 A12	(5-17 cm.) 2 to 6½ inches, brown (10YR 5/3) gravelly loamy coarse sand, very dark grayish brown (10YR 3/2) moist; seams of grayish brown and dark grayish brown (10YR 5/2, 4/2) dry; weak fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine, few fine roots; many very fine and fine interstitial pores; 17 percent gravel; medium acid (pH 6.0); clear wavy boundary.
RSL 68567 C1	(17-48 cm.) 6½ to 19 inches, pale brown (10YR 6/3) gravelly loamy coarse sand, brown (10YR 4/3) moist; weak fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine, fine, medium and coarse roots; many very fine and fine interstitial pores; 13 percent gravel; medium acid (pH 6.0); clear wavy boundary.
RSL 68568 C2	(48-69 cm.) 19 to 27 inches, pale brown (10YR 6/3) gravelly loamy coarse sand, brown (10YR 4/3) moist; massive; soft, very friable, nonsticky and nonplastic; many fine, medium and coarse roots; many very fine and fine interstitial pores; 22 percent gravel; medium acid (pH 6.0); clear wavy boundary.
RSL 68569 IIB1	(69-79 cm.) 27 to 31 inches, pale brown (10YR 6/3) gravelly loamy coarse sand, brown (10YR 4/3) moist; massive; soft, very friable, nonsticky and nonplastic; many fine and common very fine, medium and coarse roots; many very fine and fine interstitial pores; few very thin clay films in bridges and coating sand grains, occurring in pockets and thin bands; 32 percent gravel; medium acid (pH 6.0); abrupt wavy boundary.
RSL 68570 IIB2	(79-89 cm.) 31 to 35 inches, light yellowish brown (10YR 6/4) gravelly loamy coarse sand, dark yellowish brown (10YR 4/4) moist; massive; hard, very friable, nonsticky and nonplastic; many fine, common very fine and medium roots; common very fine and fine interstitial pores; common very thin clay films as bridges and coatings on sand grains in pockets and thin discontinuous bands; 31 percent gravel; medium acid (pH 5.8); abrupt wavy boundary.
RSL 68571 IIB3	(89-110 cm.) 35 to 44 inches, pale brown (10YR 6/3) gravelly loamy coarse sand, brown (10YR 4/3) moist; massive; soft, very friable, nonsticky and nonplastic; many fine, common very fine and medium roots; many very fine and fine interstitial pores; few very thin clay films as bridges, common very thin clay coatings on sand grains in pockets and thin discontinuous bands; 31 percent gravel; medium acid (pH 5.8); clear wavy boundary.
RSL 68572 IIC1	(110-150 cm.) 44 to 59 inches, pale brown (10YR 6/3) gravelly coarse sand, brown (10YR 4/3) moist; massive; soft, very friable, nonsticky and nonplastic; many fine, common very fine and medium roots; many very fine and fine interstitial pores; 20 percent gravel; strongly acid (pH 5.5); abrupt smooth boundary.
RSL 68573 IIIC2	(150-185 cm.) 59 to 72 inches, light gray (10YR 7/2) very gravelly coarse sand, grayish brown (10YR 5/2) moist; massive; soft, very friable, nonsticky and nonplastic; many very fine, few fine roots; many fine and few very fine interstitial pores; some silica and clay coatings on bottom side of gravel; gravel strongly weathered; 19 percent gravel; medium acid (pH 5.8).
RSL 68574	

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Alfic Xeropsamment, mixed, thermic
SOIL Oceano sand SOIL Nos. S62Calif-42-1 LOCATION Santa Barbara County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 6269 - 6276

Depth (in.)	Horizon	1B1b Total: Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Size class and particle diameter (mm) 3A1 — Sand: Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt: 0.05–0.02	Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)			Coarse fragments 2A2: > 2	2 - 19	19 - 76
		Pct. of < 2 mm														Pct.	Pct. of < 76mm	
0-1½	A11	84.9	11.9	3.2	0.2	29.1	39.7	13.5	2.4	8.3	3.6	14.2	82.5			0		
1½-4	A12	79.3	17.0	3.7	0.1	24.8	36.5	14.2	3.7	13.2	3.8	21.1	75.6			0		
4-15	A13	87.2	9.8	3.0	0.3	29.3	39.9	15.0	2.7	6.8	3.0	13.9	84.5			0		
15-20	AC	85.7	11.4	2.9	0.0	27.3	40.4	15.2	2.8	7.9	3.5	15.5	82.9			0		
20-39	C1	83.5	13.0	3.5	0.2	24.0	38.5	16.1	4.7	8.8	4.2	19.0	78.8			0		
39-55	C2	80.9	15.0	4.1	0.2	22.6	37.0	17.0	4.1	10.1	4.9	20.1	76.8			0		
55-75	C3	86.7	9.4	3.9	0.3	25.7	40.3	17.2	3.2	6.3	3.1	14.9	83.5			0		
75-90+	C4	93.8	4.0	2.2	0.3	27.4	45.3	18.5	2.3	2.1	1.9	10.0	91.5			0		
Depth	6A1a Organic	6B1a Nitrogen	C/N			6C2a Ext	Bulk density 4A3a				Water content 4B1d	4B1d	4B2			pH 8C1a	8C1c	

OCEANO SAND
S62Calif-42-1

Location: Santa Barbara County, California. Central SE¼ of NW¼ Sec. 36, T.10N., R.34W.; 3.7 airline miles south, southeast of center of Santa Maria. Date of Sampling: March 13, 1962.
Description by: L. C. Leifer. Collectors: R. E. Nelson, B. R. Brasher, G. E. Shipman, C. J. Hilliker, L. C. Leifer.
Classification: Alfic Xeropsamment, mixed, thermic.
Vegetation: Annuals used for range; alfilaria, annual lupine, ripgut brome, other small weeds. Climate: Precipitation 14 inches. Mean annual temperature 57°F., January 50°F., July 63°F. Parent Material: Aeolian sands. Topography: Low rolling dune-like hills, locally plane 5 percent southwest slope. Elevation: 400 feet. Drainage: Excessively drained. Runoff is very slow. Permeability is rapid. Soil Moisture: Moist throughout.
Remarks: Field determinations of pH by Hellige-Truog triplex indicator. Paired site sample numbers are Oceano S62Calif-42-2-1 through 9 and 3A horizons only S62Calif-42-5-1 through 3.

HORIZON	DESCRIPTION
A11 RSL 6269	0 to 1½ inches, grayish brown (10YR 5/2) sand, dark grayish brown (10YR 4/2) when moist; single grain; soft, very friable, nonsticky and nonplastic; abundant very fine roots; many very fine interstitial pores; slightly acid (pH 6.1); clear smooth boundary.
A12 RSL 6270	1½ to 4 inches, grayish brown (10YR 5/2) sand, dark grayish brown (10YR 4/2) when moist; massive; soft, very friable, nonsticky and nonplastic; plentiful very fine roots; many very fine interstitial pores; medium acid (pH 5.9); gradual smooth boundary.
A13 RSL 6271	4 to 15 inches, light brownish gray (10YR 6/2) sand, brown (10YR 4/3) when moist; massive; weakly coherent, very friable, nonsticky, nonplastic; few very fine roots; many very fine interstitial pores, very few fine tubular pores;

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Alfic Xeropsamment, mixed, thermic

SOIL: Oceano sand — SOIL Nos. S62Calif-42-2 — LOCATION: Santa Barbara County, California

SOIL SURVEY LABORATORY: Riverside, California — LAB. Nos. 6277 - 6285

Depth (in.)	Horizon	1B1b Total: Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Sand: Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt: 0.05–0.02	Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)			Coarse fragments 2A2: > 2 Pct.	2 - 19	19 - 76
		Pct. of < 2 mm															Pct. of < 76mm	
0-2½	A11	79.9	16.9	3.2	0.1	18.7	40.0	16.8	4.3	13.8	3.1	23.9	75.6			0		
2½-14	A12	89.7	7.8	2.5	0.1	21.4	46.8	17.4	4.0	4.9	2.9	14.8	85.7			0		
14-21	A13	87.8	9.8	2.4	0.0	21.0	45.0	17.9	3.9	7.4	2.4	17.5	83.9			0		
21-27	AC	87.8	10.1	2.1	0.0	22.2	44.0	17.2	4.4	7.4	2.7	17.9	83.4			0		
27-40	C1	87.8	10.6	1.6	0.0	24.7	42.5	16.8	3.8	7.1	3.5	16.9	84.0			0		
40-53	C2	87.0	11.3	1.7	0.0	24.5	42.1	16.4	4.0	7.1	4.2	16.9	83.0			0		
53-65	C3	84.9	13.0	2.1	0.0	26.9	40.0	14.2	3.8	8.5	4.5	17.3	81.1			0		
65-80	C4	82.5	12.5	5.0	0.0	26.4	37.7	14.0	4.4	8.0	4.5	17.5	78.1			0		
80-94+	C5	92.3	3.7	4.0	0.0	25.4	44.7	19.5	2.7	2.2	1.5	11.2	89.6			0		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.		Bulk density: 4A3a Field moist cores g/cc	4A1h Oven-dry clods g/cc	g/cc		Water content: 4B1d 1/10 bar cores Pct.	4B1d 1/3 bar cores Pct.	4B2 15 bar Pct.			pH: 8C1a H_2O 1:1	8C1c 1 N KCl 1:1	(1:1)
0-2½	0.87	0.078	11				1.41				8.3	6.0	2.7			6.2	5.0	
2½-14	0.36						1.56				6.8	4.2	1.6			6.0	3.4	
14-21	0.18						1.56				5.0	3.1	1.4			6.2	4.8	
21-27	0.11						1.58				4.7	3.0	1.4			6.2	4.6	
27-40	0.05						1.60				4.8	3.0	1.0			6.5	4.7	
40-53	0.03						1.60				5.3	3.4	1.1			6.7	4.7	
53-65	0.03						1.61				6.2	3.9	1.1			6.8	4.8	
65-80	0.02						1.64	1.86*			7.4	4.3	2.1			6.7	4.6	
80-94+	0.01						1.58				4.7	3.2	1.7			6.7	4.8	

Depth (in.)	Extractable bases 5B1a: 6N2b Ca	6O2c Mg	6P2a Na	6Q2a K		6H2a Ext. Acidity	Cation Exch. Capacity: 5A3a Sum								Base saturation: 5C3 Sum	
	meq/100 g														Pct.	Pct.

OCEANO SAND
S62Calif-42-2

Location: Santa Barbara County, California. SE¼ of NW¼ Sec. 31, T9N, R33W; 1½ miles south of Betteravia Road on Telephone Road, 3400 feet west and northwest on oil well road, 100 feet up slope; 4.2 miles south southeast of center of Santa Maria. Date of Sampling: March 15, 1962.

Description by: L. C. Leifer. Collectors: R. E. Nelson, B. R. Brasher, G. E. Shipman C. J. Hilliker, L. C. Leifer.

Classification: Alfic Xeropsamment, mixed, thermic.

Vegetation: Annuals used for range; annual lupine, red brome, white stem alfilaria, fiddle neck, small weeds. Climate: Precipitation 14 inches. Mean annual temperature 57°F., January 50°F., July 63°F. Parent Material: Aeolian sands. Topography: Low rolling [illegible] Elevation: 400 feet. Drainage: Excessively drained. Runoff is very slow. Permeability is rapid. Soil Moisture: Moist throughout.

Remarks: Field determinations of pH by Hellige-Truog triplex indicator. Paired site sample numbers are Oceano S62Calif-42-1-1 through 8 and 3 A horizons only, S62Calif-42-5-1 through 3.

HORIZON	DESCRIPTION
A11 RSL 6277	0 to 2½ inches, brown (10YR 5/3) sand, dark grayish brown (10YR 4/2) when moist; 0-1 inch single grain, 1-2½ inches very weak, coarse platy structure; soft, very friable, nonsticky and nonplastic; abundant very fine roots; many very fine interstitial pores; medium acid (pH 6.0); clear smooth boundary.
A12 RSL 6278	2½ to 14 inches, brown (10YR 5/3) sand, dark grayish brown (10YR 4/2) when moist; massive; soft, very friable, nonsticky and nonplastic; plentiful very fine roots; many very fine interstitial and very few fine tubular pores; medium acid (pH 5.6); gradual smooth boundary.
A13 RSL 6279	14 to 21 inches, brown (10YR 5/3) sand, darker brown (10YR 4/3) when moist; massive; soft, very friable, nonsticky and nonplastic; very few fine roots; many very fine interstitial and very few fine tubular pores; medium acid (pH 5.8); gradual smooth boundary.
AC RSL 6280	21 to 27 inches, pale brown (10YR 6/3) sand, brown (7.5YR 5/4) when moist; massive; soft, very friable, nonsticky and nonplastic; very few fine roots; many very fine interstitial pores; medium acid (pH 5.8); gradual smooth boundary.
C1 RSL 6281	27 to 40 inches, pinkish gray (7.5YR 7/3) sand; brown (7.5YR 5/4) when moist; massive; soft, very friable, nonsticky and nonplastic; very few very fine roots; many very fine interstitial pores; medium acid (pH 5.9); lower boundary of C horizons coincide with upper bands in each following horizon; clear smooth boundary.
C2 RSL 6282	40 to 53 inches, pinkish gray (7.5YR 7/3) sand, brown (7.5YR 5/4) when moist; massive; slightly hard, very friable, nonsticky and nonplastic; no roots; many very fine interstitial pores; one very thin weak lamella top of horizon, several short discontinuous lamellae, lower lamella 1/4 inch thick; medium acid (pH 6.0); clear wavy boundary.
C3 RSL 6283	53 to 65 inches, pinkish gray (7.5YR 7/2) sand, brown (7.5YR 5/4) when moist; massive; soft, very friable, nonsticky and nonplastic; no roots; many very fine interstitial pores; several continuous weak lamellae 1/8 inch to 1/4 inch thick near top, several very weak lamellae 4 to 6 inches long lower part; medium acid (pH 6.0); clear wavy boundary.
C4	65 to 80 inches, very pale brown (10YR 7/3) sand, light yellowish brown (10YR 6/4)

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aquic Duorthidic Xeropsamment, mixed, mesic

SOIL Tangair sand SOIL Nos. S62Calif-42-7 LOCATION Santa Barbara County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 62111 - 62115

1B1b Size class and particle diameter (mm) 3A1

TANGAIR SAND
S62Calif-42-7

Location: Santa Barbara County, California. 120° 34' 10" W., 34° 44' 50" N; Vandenberg Air Force Base; 1/4 mi. west of missile control center, 4.5 miles north northeast of Surf. Field Sheet SA23. Date of Sampling: April 18, 1962.
Description by: L. C. Leifer. Collectors: R. E. Nelson, Lothair Grant, G. E. Shipman, C. J. Hilliker, L. C. Leifer.
Classification: **Aquic Durorthidic Xeropsamment, mixed, mesic.**
Vegetation: Prostrate manzanita, sticky monkey flower, few small weeds, considerable bare ground and most shrubs only 2 to 3 feet high. Climate: Precipitation 14 inches. Mean annual temperature 55°F., July 61°F., January 49°F. Parent Material: Marine sands. Topography: Nearly level terrace, locally a little less than 1 percent slope. Elevation: 350 feet. Drainage: Imperfectly drained; in unusually rainy springs the soils are saturated to within a few inches of the surface for several months. Runoff is very slow. Permeability is rapid to a depth of 6 to 12 feet with slow or very slowly permeable material below. Soil Moisture: Moist, free water at 44 inches.
Remarks: Field pH determinations by Wallace-Truog triplex indicator. Pair site sample

HORIZON	DESCRIPTION
A1 RSL 62111	0 to 4 inches, light gray (10YR 6/1) sand, very dark grayish brown (10YR 3/2) when moist; weak coarse granular structure; loose, nonsticky and nonplastic; plentiful very fine and few fine horizontal roots; many very fine interstitial pores; medium acid (pH 6.0); clear wavy boundary.
A2 RSL 62112	4 to 24 inches, light gray (10YR 7/2d and 10YR 7/2m) sand; massive; loose, nonsticky and nonplastic; few fine and very few medium roots; many fine interstitial pores; incipient concretions 1/16 to 1/8" across make up less than 1/2 percent of mass, uniformly distributed, reddish brown (5YR 5/4 to 6/4m); medium acid (pH 6.0); gradual wavy boundary.
B21ir RSL 62113	24 to 36 inches, very pale brown (10YR 7/3) sand, light gray (10YR 7/2) when moist; massive; loose, nonsticky and nonplastic; very few fine roots; many very fine interstitial pores; concretions make up from 5 to 15 percent of

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aquic Durorthidic Xeropsamment, mixed, mesic
SOIL Tangair sand SOIL Nos. S62Calif-42-8 LOCATION Santa Barbara County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 62116 - 62121

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	3A1 Int. II (0.2-0.02)	(2-0.3)			Coarse fragments 2A2: > 2 Pct.	2-19	19-76
		Pct. of < 2 mm															Pct. of < 76mm	
0-4	A11	85.4	12.4	2.2	2.0	24.2	36.0	20.5	2.7	7.6	4.8	15.9	82.7			5		
4-11	A12	88.1	10.2	1.7	1.4	22.0	37.7	23.7	3.3	6.0	4.2	16.1	84.8			0		
11-29	A2	88.0	10.6	1.4	1.3	22.0	37.5	24.0	3.2	6.3	4.3	16.1	84.8			0		
29-44	B211r	87.6	8.5	3.9	2.0	25.3	35.4	22.2	2.7	4.1	4.4	13.3	84.9			0		
29-44	B211rm																	
44-72	B221r	89.4	7.4	3.2	0.9	20.5	39.4	25.6	3.0	4.1	3.3	14.3	86.4			0		
72-78	C1g	84.2	14.3	1.5	0.7	8.7	49.0	23.7	2.1	10.8	3.5	17.4	82.1			0		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.	6C2a Ext. Iron as Fe Pct.	Bulk density 4A1f 1/3 bar clods g/cc	4A1h Oven-dry clods g/cc	g/cc		Water content Pct.	4B1c 1/3 bar clods Pct.	4B2 15 bar Pct.	4D1 Extensibility COLE In./in.		pH 8C1a H_2O 1:1	8C1c 1 N KCl 1:1	(1:1)
0-4	2.49	0.156	16			0.1							6.4			5.9	5.0	
4-11	0.33					0.1							1.4			6.6	5.5	
11-29	0.10					0.1							1.0			6.7	5.5	
29-44	0.11					0.8	1.85	1.85				5.9	2.4	0.000		5.8	4.4	
29-44						0.7												
44-72	0.05					0.5							1.9			5.8	4.3	
72-78	0.03					0.1							1.0			5.5	4.4	

Depth (in.)	Extractable bases 5B1a: 6N2b Ca	5O2c Mg	6P2a Na	6Q2a K		6H2a Ext. Acidity	Cation Exch. Capacity 5A3a Sum									Base saturation 5C3 Sum Pct.	Pct.
	meq/100 g																
0-4	5.9	2.1	0.1	0.2		5.0	13.3									62	
4-11	1.4	0.6	0.1	0.1		0.6	2.8									79	
11-29	0.7	0.5	0.1	0.1		0.2	1.6									88	
29-44	0.5	0.9	0.2	0.2		1.5	3.3									54	
29-44	0.6	0.8	0.2	0.2		1.7	3.5									51	
44-72	0.5	0.8	0.3	0.1		0.8	2.5									68	
72-78	0.3	0.5	0.3	Tr.		0.2	1.3									85	

Depth (in.)

* Concretion

TANGAIR SAND
S62Calif-42-8

Location: Santa Barbara County, California. 120° 34' 10" W. and 34° 43' 10" N; Vandenberg Air Force Base, 1000 feet southwest of south end of air strip; 200 feet west and 50 feet south of railroad siding intersection with old road to Lompoc Landing; 3 miles northeast of Surf. Field sheet SD13. Date of Sampling: April 18, 1962.
Description by: L. C. Leifer. Collectors: R. E. Nelson, Lothair Grant, G. E. Shipman, C. J. Hilliker, L. C. Leifer.
Classification: Aquic Durorthidic Xeropsamment, mixed, mesic.
Vegetation: Chamise, black sage (Salvia mellifera), low manzanita and low Ceanothus; brush fairly dense and much about 5 feet high. Climate: Precipitation 14 inches. Mean annual temperature 55°F., July 61°F., January 49°F. Parent Material: Marine sands. Topography: Nearly level terrace, locally a little less than 1 percent slope. Elevation: 360 feet. Drainage: Imperfectly drained; in unusually rainy springs the soils are saturated to within a few inches of the surface for several months. Runoff is very slow. Permeability is rapid to a depth of 6 to 12 feet with slow or very slowly permeable material below. Soil Moisture: Moist, free water at 75 inches.
Remarks: Field pH determination by Hellige-Truog triplex indicator. Pair site sample numbers are Tangair S62Calif-42-7-1 through 5.

HORIZON	DESCRIPTION
A11 RSL 62116	0 to 4 inches, gray (10YR 5/1) light loamy sand, very dark gray (10YR 3/1) when moist; weak medium platy and weak coarse granular structure; soft, very friable, nonsticky and nonplastic; plentiful very fine and very few medium roots; many very fine interstitial pores; medium acid (pH 5.8); clear smooth boundary.
A12 RSL 62117	4 to 11 inches, gray (10YR 6/1) sand, pale brown (10YR 6/3) when moist; massive; loose, nonsticky and nonplastic; few fine and medium roots; many very fine interstitial pores; 10 to 15 percent of soil is diffuse to clear, old root channels 1/8 to 1" across; filled with A1 soil material; medium acid (pH 6.0); diffuse smooth boundary.
A2 RSL 62118	11 to 29 inches, light gray (10YR 7/2d, 7/2m) sand; massive; loose, nonsticky and nonplastic; few fine and medium roots; many very fine interstitial pores; 1 percent of mass distinct to faint very dark gray organic filled old root channels; medium acid (pH 6.0); clear wavy boundary.
B21ir RSL 62119	29 to 44 inches, light gray (10YR 7/2d, 7/2m) sand; massive; loose, nonsticky and nonplastic; very few fine and medium roots; many very fine interstitial pores; 15 to 35 percent of horizon made up of round to oblong concretions 1 to 10" in diameter, tend to occur in pockets 2 to 3' across, reddish yellow to yellowish red (7.5YR 6/6m to 5YR 4/6m and 5YR 4/6d) in harder shell; interior of large concretions light yellowish brown (10YR 6/4d); concretions friable to very firm, usually firm; medium acid (pH 5.8); gradual wavy boundary.
62119a	Concretions in the B21ir analyzed separately.
B22ir RSL 62120	44 to 72 inches, light gray (10YR 7/2d, 7/2m) sand; massive; loose, nonsticky and nonplastic; very few fine and medium roots; many very fine interstitial pores; 35 percent of horizon made up of firm concretions 1/2 to 3" across, yellowish red (5YR 5/6d) with blotches of light brownish gray (2.5Y 6/2m) few concretions dark gray on outside and reddish on inside; medium acid (pH 5.6); gradual wavy boundary.
Cg? RSL 62121	72 to 78 inches +, white (10YR 8/1) sand, light gray (10YR 7/2) when moist; massive; loose, nonsticky and nonplastic; no roots; many very fine interstitial pores; medium acid (pH 5.6).

CONSECUTIVE PEDON NUMBER C 68 89 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES WACA SOIL NO. S68 CALI. 31- 1 SAMPLE NO. 8 727-8 734 TAHOE BASIN AREA

SOIL FAMILY TYPIC CRYANDEPT CINDERY RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A	1A2A INT. II .2- .02	COARSE FRAGMENTS PCT. OF WHOLE 20 I 75-	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														SOIL	PCT. OF LT 75	
8 727	5- 0	D1+D2															--A	--	--
8 728	0- 23	A11	66.4	28.9	4.7	19.2	14.5	7.8	14.1	10.8	11.6	17.3				30.3	63*	20	23Z
8 729	23- 36	A12	66.1	29.1	4.8	14.7	14.1	9.1	15.8	12.4	11.6	17.5				33.1	52*	24	25Z
8 730	36- 53	AC	66.6	28.7	4.7	14.1	14.0	9.1	16.2	13.2	12.1	16.6				34.9	53*	24	27Z
8 731	53- 79	C 1	67.6	28.4	4.0	12.7	15.1	9.9	17.2	12.7	11.2	17.2				33.8	55*	21	29Z
8 732	79- 91	C 2	68.2	27.6	4.2	11.1	17.4	10.7	17.6	11.4	10.7	16.9			--	32.1	73*	37	31Z
8 733	91- 0	R															94*	--	--
8 734	0- 53	A															29A	--	29Z

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT EXT FE 6C2B	AL 6G7A	PYROP EXT FE 6C5A	AL 6G9A	ATTERBERG LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		PERCENT		PERCENT		PERCENT				PCT.NO.40		GRAMS PER CC				PERCENT			PERCENT	
8 727	5- 0														1.00					
8 728	0- 23	4.85	0.178			1.0				NP	NP		0.90	0.91	0.64		27.8	13.8	0.4	0.2
8 729	23- 36	1.97	0.109									0.90			0.73			11.8		
8 730	36- 53	1.63	0.089			1.1				NP	NP		0.88	0.88	0.73		21.9	12.7		
8 731	53- 79	1.60				1.1				NP	NP	0.76			0.74			14.4		
8 732	79- 91	1.44				1.0				NP	NP	0.71			0.58			13.9		
8 733	91- 0											1.63			0.10					
8 734	0- 53	4.04	0.175									0.75			0.90			13.1		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 584A 6N2E	MG 5B4A 6D2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	CEC SUM 5A3A	CEC NH4-OAC 5A6A	BASE SATURATION SUM+ ACI-DITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 727	5- 0															
8 728	0- 23	5.0	0.6	0.1	0.9	6.6	23.7		30.3	23.6	22		28		6.2	5.5
8 729	23- 36	3.2	0.4	0.1	0.9	4.6	21.2		25.8	21.0	18		22		6.2	5.5
8 730	36- 53	4.2	0.5	0.1	0.7	5.5	20.4		25.9	21.8	21		25		6.3	5.4
8 731	53- 79	5.5	0.4	0.1	0.7	6.7	21.7		28.4	24.1	24		28		6.2	5.3
8 732	79- 91	5.9	0.5	0.1	0.7	7.2	20.0		27.2	25.5	26		28		5.9	5.1
8 733	91- 0															
8 734	0- 53														6.1	5.5

SAMPLE NO	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 727	5- 0											1 TRACE
8 728	0- 23	AM 6									AM AMORPHOS	2 SMALL
8 729	23- 36											3 MODERATE
8 730	36- 53	AM 6										4 ABUNDANT
8 731	53- 79	AM 6										5 DOMINANT
8 732	79- 91											6 INDETER.
8 733	91- 0											
8 734	0- 53											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
*=THE 75-250MM FRACTION IS INCLUDED.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

WACA COBBLY COARSE SANDY LOAM
S68Calif-31-1

Location: Tahoe Basin, Placer County, California; 200 yards north of S¼ corner, sec. 34, T. 15 N., R. 16 E. Compass bearing: S. 37° W to Barker Peak, N 85° W to Stanford Rock, 100 feet south of Blackwood Canyon Road. Date of Sampling: August 6, 1968.
Description by: R. C. Huff. Collectors: J. H. Rogers, L. N. Langan, G. M. Kennedy, G. L. Anderson, B. R. Brasher, W. D. Nettleton.
Classification: Typic Cryandept, cindery.
Vegetation: Not cultivated; white fir, ribies sp., snow berry. Climate: Precipitation is 40 to 45 inches. Mean annual temperature is 40°F., January 24°F., July 60°F. Parent Material: Andesitic tuff breccia. Topography: Mountainous uplands, 35 percent northwest-facing slope. Elevation: 6800 feet. Drainage: Well drained; runoff is rapid; permeability is moderately rapid. Soil Moisture: Near wilting point.
Remarks: Field determinations for pH by Hellige-Truog triplex indicator. Composite A taken from 10 satellite samples.

HORIZON	DESCRIPTION
	(5-0 cm.)
01 & 02 RSL 68727	2 to 0 inches, fir litter and duff.
	(0-23 cm.)
A11 RSL 68728	0 to 9 inches, brown (10YR 5/3) cobbly coarse sandy loam, very dark grayish brown (10YR 3/2) moist; massive; soft, very friable, nonsticky and nonplastic; very few fine roots; 30 percent gravel and 14 percent cobbles and stones; slightly acid (pH 6.2); abrupt smooth boundary.
	(23-36 cm.)
A12 RSL 68729	9 to 14 inches, brown (10YR 5/3) gravelly coarse sandy loam, dark brown (10YR 3/3) moist; massive; soft, very friable, nonsticky and nonplastic; common very fine and fine, many medium roots; many very fine and fine interstitial pores; 35 percent gravel and 3 percent cobbles and stones; medium acid (pH 6.0); diffuse wavy boundary.
	(36-53 cm.)
AC RSL 68730	14 to 21 inches, brown (10YR 5/3) gravelly coarse sandy loam, dark brown (10YR 3/3) moist; weak medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine, fine, medium and coarse roots; many very fine and fine interstitial pores; 45 percent gravel and 3 percent cobbles and stones; medium acid (pH 5.8); diffuse wavy boundary.
	(53-79 cm.)
C1 RSL 68731	21 to 31 inches, pale brown (10YR 6/3) very gravelly coarse sandy loam, brown (10YR 4/3) moist; weak medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine, fine, medium and coarse roots; many very fine and fine interstitial pores; 35 percent gravel and 3 percent cobbles and stones; medium acid (pH 5.6); diffuse wavy boundary.
	(79-91 cm.)
C2 RSL 68732	31 to 36 inches, pale brown (10YR 6/3) very gravelly coarse sandy loam, dark yellowish brown (10YR 4/6) moist; weak medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine, fine, medium and coarse roots; many very fine and fine interstitial pores; 60 percent gravel; very strongly acid (pH 5.0); abrupt irregular boundary.
	(91 cm.)
R	36 inches, slightly weathered andesitic tuff, fractured with some roots and soil

CONSECUTIVE PEDON NUMBER C 68 90 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES WACA SOIL NO. S68 CALI. 31- 2 SAMPLE NO. 8 735-8 745 TAHOE BASIN AREA

SOIL FAMILY TYPIC CRYANDEPT CINDERY RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	CSI .05- .02	FSI .02- .002	CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A	1A2A INT. II .2- .02	-COARSE FRAGMENTS- PCT. OF WHOLE SOIL	I 75- 20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														--A	--	--	--
8 735	5- 0	01																		
8 736	0- 3	A11	59.3	32.8	7.9	16.5	10.6	6.6	13.7	11.9	11.4	21.4				31.5	49A	--	49Z	N
8 737	3- 8	A12	59.9	35.4	4.7	12.5	14.4	8.3	14.3	10.4	15.0	20.4				33.1	50A	23	27Z	N
8 738	8- 20	A13	61.7	32.4	5.9	16.1	14.0	7.7	13.1	10.8	10.5	21.9				29.2	46A	20	26Z	N
8 739	20- 33	AC	61.8	32.7	5.5	14.8	13.8	7.7	13.4	12.1	11.7	21.0				31.8	49A	25	24Z	N
8 740	33- 53	C 1	60.5	34.8	4.7	12.4	14.1	7.7	14.1	12.2	12.5	22.3				32.4	64A	52	12Z	N
8 741	53- 74	C 2	62.9	33.4	3.7	6.7	14.9	10.1	17.9	13.3	13.3	20.1				36.7	63A	44	19Z	N
8 742	74- 89	C 3	64.2	32.2	3.6	11.4	15.3	9.7	15.8	12.0	12.7	19.5		--	--	32.8	58A	31	27Z	N
8 743	89- 99	C 4															60A	25	35Z	N
8 744	99-100	R															67A	30	37Z	N
8 745	0- 33	A															29A	--	29Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR- BON 6A1A -PERCENT-	NITRO GEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT EXT FE 6C2B	AL 6G7A	PYROP EXT FE 6C5A	AL 6G5A	ATTERBERG L I M I T LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 4A1F	1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
				-PERCENT-		-------PERCENT-------				PCT.NO.40		GRAMS PER CC				----PERCENT---			-PERCENT-	
8 735	5- 0														1.00					
8 736	0- 3	19.18	0.682									0.24	0.32	0.92		102.9	37.0	10.1	9.1	
8 737	3- 8	9.19	0.329			1.3				NP	NP	0.68	0.70	0.79		38.6	28.6	1.0	0.8	
8 738	8- 20	7.66	0.293									0.68	0.70	0.82		37.7	20.0	1.0	0.8	
8 739	20- 33	5.18	0.224			1.4				NP	NP	0.72	0.73	0.80		40.4	18.8	0.5	0.4	
8 740	33- 53	3.76				1.3				NP	NP	0.76	0.76	0.66		40.7	20.1			
8 741	53- 74	3.06										0.71	0.73	0.69		41.0	20.5	0.9	0.6	
8 742	74- 89	2.88										0.74	0.75	0.72		39.6	19.2	0.4	0.3	
8 743	89- 99	3.12										0.94	0.98	0.65		46.3	22.2	1.4	0.9	
8 744	99-100											1.63	1.67	0.44		20.6	14.2	0.8	0.4	
8 745	0- 33	9.07	0.352								0.70			0.90			25.8			

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6Q2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI- DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4- OAC 5A6A	BASE SATURATION SUM+ ACI- DITY	SUM+ AL	NH4- OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA- CL2 8C1E
		MEQ/100G									PERCENT					
8 735	5- 0															
8 736	0- 3	10.1	5.0	0.1	0.7	15.9	39.1	1.3	55.0	59.0	29	92	27	5.3		4.7
8 737	3- 8	1.9	0.2	0.1	0.5	2.7	23.2	1.5	25.9	40.0	10	64	7		5.4	4.9
8 738	8- 20	2.0	0.4	0.1	0.5	3.0	36.7	0.7	39.7	34.7	8	81	9		5.5	4.9
8 739	20- 33	0.8	0.1	0.1	0.4	1.4	36.9	0.3	38.3	32.5	4	82	4		5.5	5.0
8 740	33- 53	0.8	0.2	0.1	0.2	1.3	35.2	0.1	36.5	32.3	4	93	4		5.5	5.1
8 741	53- 74	0.7	0.1	0.1	0.1	1.0	34.1	0.3	35.1	34.7	3	77	3		5.5	5.0
8 742	74- 89	0.7	0.2	0.1	0.1	1.1	35.7	0.3	36.8	33.7	3	79	3		5.4	5.0
8 743	89- 99	0.6	0.2	0.2	0.1	1.1	33.2	0.6	34.3	37.6	3	65	3		5.4	4.9
8 744	99-100						27.2	0.9	27.2	27.9					5.3	4.8
8 745	0- 33														5.3	4.9

SAMPLE NO	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 735	5- 0											1 TRACE
8 736	0- 3										AM AMORPHOS	2 SMALL
8 737	3- 8	AM 6									KH HALOYSIT	3 MODERATE
8 738	8- 20											4 ABUNDANT
8 739	20- 33	AM 6										5 DOMINANT
8 740	33- 53	KH 1										6 INDETER.
8 741	53- 74											
8 742	74- 89											
8 743	89- 99											
8 744	99-100											
8 745	0- 33											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

WACA GRAVELLY COARSE SANDY LOAM
S68Calif-31-2

Location: Tahoe Basin, Placer County, California. SE¼ of sec. 8, T. 14 N., R. 16 E., 0.4 mile east of Barker Pass, 70 feet south of Blackwood Canyon Road. Date of Sampling: August 6, 1968.
Description by: G. M. Kennedy. Collectors: R. C. Huff, L. N. Langan, J. H. Rogers, B. R. Brasher, W. D. Nettleton.
Classification: Typic Cryandept, cindery.
Vegetation: Not cultivated; Red fir, mountain hemlock. Climate: Precipitation is 45 to 50 inches. Mean annual temperature is 39°F., January 20°F., July 59°F. Parent Material: Andesitic breccia. Topography: Mountainous uplands, 30 percent, north-facing slope. Elevation: 7800 feet. Drainage: Well drained; runoff is medium; permeability is moderately rapid. Soil Moisture: Slightly moist.
Remarks: Field determinations for pH by Hellige-Truog triplex indicator. Composite A taken from 10 satellite samples. Samples taken of C1 and C2 horizons for Division of Highways. 51°F. at 20 inches.

HORIZON	DESCRIPTION
01 RSL 68735	(5-0 cm.) 2 to 0 inches, needle duff.
A11 RSL 68736	(0-3 cm.) 0 to 1 inch, brown (10YR 4/3) gravelly coarse sandy loam, very dark brown (10YR 2/2) moist; massive; soft, very friable, nonsticky and nonplastic; common very fine, few fine roots; many very fine interstitial and few fine tubular pores; strongly acid (pH 5.2); abrupt smooth boundary.
A12 RSL 68737	(3-8 cm.) 1 to 3 inches, brown (10YR 4/3) cobbly coarse sandy loam, very dark brown (10YR 2/2) moist; moderate fine and medium granular structure; soft, very friable, nonsticky and nonplastic; common very fine and few fine roots; many very fine interstitial and few fine tubular pores; 30 percent gravel and 5 percent cobbles; medium acid (pH 6.0); clear wavy boundary.
A13 RSL 68738	(8-20 cm.) 3 to 8 inches, brown (10YR 4/3) gravelly coarse sandy loam, very dark grayish brown (10YR 3/2) moist; moderate fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine and fine, common medium and coarse roots; many very fine interstitial, few fine tubular pores; 30 percent gravel; medium acid (pH 6.0); clear wavy boundary.
AC RSL 68739	(20-33 cm.) 8 to 13 inches, brown (10YR 5/3) gravelly coarse sandy loam, dark brown (7.5YR 3/2) moist; weak fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine and fine, common medium and few coarse roots; many very fine interstitial and few very fine and fine tubular pores; 35 percent gravel and 5 percent cobbles; medium acid (pH 6.0); clear wavy boundary.
C1 RSL 68740	(33-53 cm.) 13 to 21 inches, yellowish brown (10YR 5/4) very gravelly coarse sandy loam, dark brown (7.5YR 3/2) moist; massive; soft, very friable, nonsticky and slightly plastic; many very fine, fine, medium and coarse roots; many very fine and few fine tubular pores; 50 percent gravel and 5 percent cobbles and stones; medium acid (pH 6.0); gradual wavy boundary.
C2 RSL 68741	(53-74 cm.) 21 to 29 inches, light yellowish brown (10YR 6/4) very gravelly coarse sandy loam, brown (7.5YR 4/4) moist; massive; soft, very friable, nonsticky and slightly plastic; many very fine, fine, medium and coarse roots; common very fine tubular and many very fine, few fine interstitial pores; 50 percent gravel and 5 percent cobbles; medium acid (pH 6.0); gradual smooth boundary.
C3 RSL 68742	(74-89 cm.) 29 to 35 inches, light yellowish brown (10YR 6/4) very gravelly coarse sandy loam, brown (7.5YR 4/4) moist; massive; soft, very friable, nonsticky and slightly plastic; many very fine and fine, common medium and few coarse roots; many very fine interstitial and few very fine and fine tubular pores; 45 percent gravel; medium acid (pH 6.0); abrupt wavy boundary.
C4 RSL 68743	(89-99 cm.) 35 to 39 inches, very pale brown (10YR 7/4) very gravelly coarse sandy loam, brown (7.5YR 4/4) with reddish brown (5YR 4/3) stains, moist; massive; soft, very friable, nonsticky and nonplastic; common medium and coarse, few fine roots; 45 percent gravel; medium acid (pH 6.0); gradual wavy boundary.
R RSL 68744	39 to 40 inches, andesitic tuff breccia. (99-100 cm.)

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Windy sandy loam **LOCATION** Amador County, California

SOIL FAMILY: Typic Dystrandept, cindery, frigid

SOIL NOS. S61Calif-3-30 **LAB. NOS.** 61121 - 61127

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	2/ CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 8	A_{11}	13.0	18.6	6.1	12.3	7.7	33.7	8.6	25.7	23.0	18	
8-16	A_{12}	12.9	18.7	6.7	13.3	7.8	31.2	9.4	25.0	21.9	25	
16-21	AB	15.8	19.4	6.4	12.2	7.4	29.3	9.5	24.0	19.6	9	
21-35	B	11.1	20.7	8.3	14.6	7.5	28.7	9.1	25.5	19.3	7	
35-47	C_1	11.9	22.1	7.6	13.9	7.5	29.4	7.6	24.9	20.1	6	
47-57	C_2	17.5	26.7	8.5	13.6	6.7	22.6	4.4	20.0	16.7	46	
57+	R	14.9	25.0	7.3	11.2	5.8	31.9	3.9	19.5	24.3	13	

pH 8C1b Saturated Paste 1:1	8C1c KCl 1:10	Organic Matter 6A1a Organic Carbon Pct.	6B1a Nitrogen Pct.	C/N	6C1b Free iron as % Fe_2O_3	4B2 S 11 water 15 atmos. Pct.
5.9	4.5	6.12	0.214	28.7	1.4	18.6
5.9	4.6	3.87	0.174	22.2	1.1	18.3
6.0	4.6	2.18	0.091	24.0	0.8	15.8
6.0	4.5	0.97	0.055	17.6	0.4	16.0
5.9	4.4	0.90	0.051	17.6	0.4	16.2
6.0	4.2	0.21			0.8	14.5
6.0	4.0	0.11			0.5	15.1

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 6N2b Ca	6O2c Mg	6H2a H	6P2a Na	6Q2a K	5B1a BASE SAT. % 5C3	5A3a Sum of Cations					
	milliequivalents per 100g. soil						milliequivalents per 100 g soil					
44.1	5.7	1.2	25.0	0.2	1.0	25	33.1					
40.1	4.1	0.8	25.0	0.2	0.8	19	30.9					
29.6	4.3	1.1	18.7	0.2	0.8	26	25.1					
26.9	4.6	2.3	14.8	0.2	1.0	35	22.9					
24.5	2.8	2.2	15.3	0.2	1.0	29	21.5					
24.3	2.7	1.7	12.8	0.2	1.0	30	18.4					
28.6	6.2	3.0	12.9	0.2	0.8	44	23.1					

1/ Percent of the whole soil

2/ Incompletely dispersed

WINDY SANDY LOAM
S61Calif-3-30

Collectors: Grant Kennedy, Jack McClelland, John Rogers, Paul Nazar, David Estrada. **Description by: Grant Kennedy.**
Date of Sampling: May 8, 1961
Location: Amador County, California, 0.5 mile southwest of Hams Station; 1/4 mile east of southwest corner of sec. 13, T. 8 N., R. 14 E., on old Highway 88
Elevation: 5,445 feet
Relief: Hilly uplands with a 25 percent slope to the south.
Drainage: Well drained; slow runoff; rapid permeability
Parent Material: Andesitic tuff breccia
Climate: Mean annual precipitation 47 inches, some as snow; mean annual temperature 50°
Vegetation: Ponderosa pine, incense cedar, red fir, white fir
Classification: **Typic Dystrandept, cindery, frigid.**
Remarks: The C1 horizon is 90 percent rotten rock, with soil in crevices in tongues. The C2 horizon is fractured rotten rock with a little soil in cracks. The A11 horizon contains about 10 percent gravels, A12 contains 15 percent gravels, A13 contains 40 percent gravels and cobbles, and the B contains 50 percent gravels, cobbles and fractured stones.

HORIZON		DESCRIPTION
O1	2- ½"	Partially decomposed needles
O2	½- 0"	Well decomposed needles; very strongly acid, pH 5.0.
A11	0- 8"	Dark grayish brown (10YR 4/2) sandy loam, very dark brown (10YR 2/2) when moist; moderate very fine granular; slightly hard when dry; friable when moist; nonsticky and nonplastic when wet; plentiful very fine to medium roots; many very fine interstitial pores; medium
RSL No.		

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Windy sandy loam **LOCATION** Amador County, California

SOIL FAMILY: Typic Dystrandept, cindery, frigid

SOIL NOS. S61Calif-3-31 **LAB. NOS.** 61128 - 61133

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	2/ CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 7	A_{11}	20.1	15.1	6.8	10.9	6.6	32.4	8.1	22.1	23.3	27	
7-14	A_{12}	16.4	15.9	7.4	12.3	6.7	33.9	7.4	25.2	22.8	24	
14-21	A3	18.8	16.0	7.0	11.8	7.7	30.4	8.3	24.4	20.8	26	
21-34	B	18.8	18.4	7.8	13.4	7.1	28.1	6.4	24.7	18.5	10	
34-66	$C_{1&R}$	15.9	18.7	9.4	15.2	7.6	25.9	7.3	25.9	16.7	20	
66+	R	12.7	18.3	8.6	14.2	7.8	34.7	3.7	34.5	16.3	4	

pH 8C1b Saturated Paste 1:1	pH 8C1c KCl 1:10	Organic Matter 6A1a Organic Carbon Pct.	6B1a Nitrogen Pct.	C/N					6C1b Free iron as % Fe_2O_3			4B2 Soil water 15 atmos. Pct.
6.2	5.1	7.63	0.273	27.9					1.9			21.3
6.2	4.9	5.02	0.182	27.6					2.6			20.4
6.2	5.0	2.43	0.123	19.8					2.6			17.9
6.3	5.0	1.28	0.084	15.2					2.5			17.4
6.1	4.6	0.47	0.032	14.7					2.0			21.5

WINDY SANDY LOAM

S61Calif-3-31

Collectors: Grant Kennedy, Jack McClelland, John Rogers, Paul Nazar, David Estrada **Description by: Grant Kennedy.**
Date of Sampling: May 8, 1961
Location: Amador County, California; 3/4 mile south southwest of Cooks Station; NW1/4SW1/4.sec. 28, T. 8 N., R. 14 E.
Elevation: 4800 feet
Relief: Hilly uplands with a 25 percent slope to the south.
Drainage: Well drained; medium runoff; moderately rapid permeability
Parent Material: Andesitic tuff breccia
Climate: Mean annual precipitation 46 inches - some is snow. Mean annual temperature 50°
Vegetation: Ponderosa pine, incense cedar, bear clover, black oak; sparse stand, burned and cutover
Classification: **Typic Dystrandept, cindery, frigid.**
Remarks: The C1&R horizon is fractured rotten rock, breaking to angular cobbles and gravels, with less than 5 percent soil in cracks, except for the upper 10 inches. The B horizon contains 35 percent cobbles and gravels by volume. The A3 horizon contains 30 percent cobbles and gravels by volume; 20 percent of the volume of the A12 and the A3 horizons are roots. There are 2 inches of loose pine needles and 1 to 1½ inches of partially decomposed pine needles and organic matter on the surface.

HORIZON		DESCRIPTION
A11	0- 7"	Dark grayish brown (10YR 4/2) sandy loam, very dark grayish brown (10YR 3/2) when moist; strong very fine granular; soft when dry, very friable when moist, and nonsticky and nonplastic when wet; plentiful very fine to medium roots; many very fine interstitial pores; medium acid, pH 6.0; gradual smooth boundary.
RSL No. 61128		
A12	7-14"	Dark grayish brown (10YR 4/2) sandy loam, very dark grayish brown (10YR 3/2) when moist; moderate very fine granular; soft when dry;

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Vitrandept, medial-skeletal, mesic

SOIL McCarthy loam SOIL Nos. S62Calif-3-1 LOCATION Amador County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 62435 - 62440

Depth (In.)	Horizon	1B1b Total Sand (2–0.05)	Total Silt (0.05–0.002)	Total Clay (< 0.002)	Sand Very coarse (2–1)	Sand Coarse (1–0.5)	Sand Medium (0.5–0.25)	Sand Fine (0.25–0.1)	Sand Very fine (0.1–0.05)	Silt 0.05–0.02	Silt Int III (0.02–0.002)	Silt Int. II (0.2–0.02)	Silt (2–0.1)			Coarse fragments 1A2a > 2 1/ Pct	2 – 19 (Pct. of < 76mm)	19 – 76 (Pct. of < 76mm)
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm																
0–4	A11	56.2	35.9	7.9	8.7	13.3	8.5	16.4	9.3	12.4	23.5	29.9	46.9			25		
4–7	A12	58.9	34.7	6.4	10.5	14.4	8.6	15.3	10.1	13.3	21.4	31.6	48.8			22		
7–16	A3	59.2	33.5	7.3	8.3	13.8	9.1	16.8	11.2	12.2	21.3	32.6	48.0			15		
16–24	B2	50.1	40.1	9.8	5.0	11.6	8.8	16.0	8.7	17.8	22.3	34.8	41.4			47		
24–34	B3	53.9	34.0	12.1	4.5	12.0	9.4	17.3	10.7	11.5	22.5	31.2	43.2			46		
34+	R	52.7	35.1	12.2	3.5	10.4	9.0	18.1	11.7	12.3	22.8	33.8	41.0			56		

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	Bulk density 4A1h Oven dry g/cc	Bulk density g/cc	4A C'	Water content Pct	Water content 4B1c 1/3 bar Pct.	Water content 4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1		
0–4	9.30	0.337	28										22.8			6.4		
4–7	3.11	0.124	25										15.8			6.6		
7–16	1.70	0.093	18										15.4			6.4		
16–24	0.75	0.050	15				0.93	0.99		0.74		33.7	15.8	0.021	0.016	6.1		
24–34	0.52	0.040	13				1.09	1.13		0.74		35.5	17.0	0.015	0.009	5.7		
34+	0.53	0.044	12				0.99	1.05		0.68		42.3	17.7	0.020	0.013	5.4		

[redacted]

McCARTHY SANDY LOAM
S62Calif-3-1

Location: 0.3 mile N and 0.2 mile W of the southeast corner of section 26, T. 8 N., R. 14 E., Amador County, California; 2.3 miles southwest of Ham's station, 100 feet S of the intersection of Winton Lumber Company field roads Nos. 5 and 6. Date of Sampling: December 11, 1962.
Description by: John E. McClelland. Collectors: Klaus W. Flach, Grant M. Kennedy, John Rogers, and John E. McClelland.
Classification: Typic Vitrandept, medial-skeletal, mesic.
Vegetation: Dominantly white fir, with ponderosa pine, incense cedar, sugar pine and an understory of mountain misery. Climate: Precipitation 47 inches (estimated), some as snow; mean annual temperature 48° F. (estimated). Parent Material: Tuff breccia. Relief: Simple north facing slope of 14 percent towards a saddle in the mountainous uplands of the Sierra Nevada. Elevation: 4740 feet. Drainage: Well drained; moderately rapid permeability; very slow runoff. Biological Activity: None observed. Remarks: The site appeared to be uneroded and no surface stones were observed. The soil was moist when sampled.

Horizon	DESCRIPTION
O1	2 to 0 inches, fresh pine needles, leaves, and twigs that are quite well decomposed in the lower portion.
A11 RSL No. 62435	0 to 4 inches, dark brown (7.5YR 3/2) sandy loam, very dark brown (7.5YR 2/2) when moist, strong very fine granular structure; soft, very friable, nonsticky and nonplastic; abundant very fine, few fine roots; many very fine interstitial pores; slightly acid, pH 6.3; clear, smooth boundary.
A12 RSL No. 62436	4 to 7 inches, dark brown (7.5YR 4/3) sandy loam, dark brown (7.5YR 3/2) when moist; strong very fine granular structure; soft, very friable, nonsticky and nonplastic; plentiful very fine to coarse roots; many very fine interstitial pores; about 10 percent of volume consists of rotten cobbles; soil is difficult to wet and appears slightly grayish because of mycelia; medium acid, pH 6.0; clear, smooth boundary.
A13 62437	7 to 16 inches, similar to above except lacks mycelia; gradual, smooth boundary.
B2 RSL No. 62438	16 to 24 inches, brown (7.5YR 5/3) sandy loam, dark brown (7.5YR 3/3) when moist; massive but crumbles readily to moderate very fine granules; soft, friable, nonsticky and nonplastic; roots and pores as above; about 25 percent rotten cobbles containing common very fine pores with yellowish brown colloidal stains in them and in adjacent areas; medium acid, pH 5.8; gradual smooth boundary.
B3 62439	24 to 34 inches, similar to above but with 40 percent rotten cobbles and strongly acid, pH 5.5; clear, slightly irregular boundary.
R RSL No. 62440	34 inches +, gray, light brownish gray, yellowish brown, with a few flecks of very dark gray (10YR 6/1, 6/2, 5/6, and 3/1) decomposing coarse sandy tuffaceous conglomerate of sandy loam texture when crushed, brown to grayish brown and black (7.5YR 4/2 to 2.5Y 5/2 and 2/2) when moist; massive; hard, very firm; few fine to coarse roots along fracture planes; many very fine irregular pores; some with thin red, others light brownish gray

Additional notes: Sands are uniformly but thinly coated dark brown (7.5YR hue) and all but the coarser grains are well rounded. Most of the sand grains appear

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Piper sandy loam (taxadjunct) **LOCATION** Contra Costa County, California

SOIL FAMILY: Aeric Haplaquept, sandy, mixed, calcareous, thermic.

SOIL NOS. 57Calif-7-3 **LAB. NOS.** 5830 - 5834

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0-2	A1	0.4	4.6	16.2	51.3	7.2	10.9	9.4	37.7	7.4	0	lfs
2-8	AC	0.2	6.5	18.5	45.4	6.4	12.7	10.3	32.6	8.6	0	fsl
8-26	C11	0.1	5.4	18.3	54.4	5.8	5.8	10.2	35.9	4.0	0	lfs
26-44	C12	0.0	7.2	20.5	54.0	6.0	5.3	7.0	34.9	3.7	0	lfs
44+	C13	0.0	7.4	22.7	55.1	6.6	3.7	4.5	35.8	2.3	0	fs

pH 8C1b SATURATED PASTE 1:1	pH 1:5	pH 8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	ORGANIC MATTER 6B1a NITROGEN %	ORGANIC MATTER C/N	6I1a CO_3 meq/l	8A1a Elec. Conduc. mmhos/cm at 25°C.	6E1b $CaCO_3$ equivalent %	6J1a HCO_3 meq/l	MOISTURE TENSIONS 1/10 ATMOS. %	MOISTURE TENSIONS 1/3 ATMOS. %	4B2 15 ATMOS. %
7.7		8.1	5.69	0.433	13.1	2.0	2.0	2	18.3			14.4
8.2		8.7	1.77	0.185	9.6	0.0	1.2	7	7.3			9.2
8.4		9.1	0.30	0.023	13.0	0.0	0.7	9	10.9	13.8	11.8	4.6
8.3		9.0	0.21			0.0	0.9	5	3.7	11.6	7.8	3.5
8.1		8.7	0.15			0.0	1.4	Tr	2.4	8.8	6.4	2.8

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	5D1 EXCH. Na %	SATURATION EXTRACT SOLUBLE 6P1a Na	6Q1a K	6N1a	6O1a	6K1a	8A1 8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
24.1	19.3	11.7	--	0.2	1.3	0.8	2.2	2.6	8.1	9.3	2.3	65.3
21.9	12.4	13.9	--	0.3	1.7	1.4	3.1	1.8	2.6	7.1	3.3	44.9
7.5	13.1	5.5	--	0.1	0.4	1.3	2.2	0.5	2.1	3.9	0.7	27.2
6.3	16.5	4.4	--	0.2	0.6	3.1	3.2	0.1	2.5	3.9	1.0	32.0
5.2	4.8	2.9	--	0.3	0.9	5.8	2.1	0.1	3.0	5.1	4.0	30.7

PIPER SANDY LOAM (taxadjunct) 1/
S57Calif-7-3

Collector: R. C. Huff. **Date of sampling:** May 14, 1957.
Location: Contra Costa County, California. 100 feet south NE corner of NW1/4NE1/4 sec. 26, T. 2. N., R. 3 E., projected. Eastern Contra Costa SCD. Holland Tract-Dutra Ranch pit silo. 1 mile due north of barn.
Classification: Aeric Haplaquept, sandy, mixed, calcareous, thermic.
Native Vegetation: Annual weeds and grasses, dominately ripgut (Bromus ridigus) red brome (Bromus rubens).
Climate: Rainfall 18". Cool moist winters, dry summers with occasional fog.
Parent Material: Mixed alluvium.
Relief: Undulating with mounds or long low ridges surrounded by peat or muck which has subsided from around these areas.
Elevation: 0 feet.
Drainage: Excessive surface and good internal.
Root Distribution: Numerous 0-2", some 2-10", few to 60".

HORIZON		DESCRIPTION
57Calif-07-3-1 A_1	0- 2"	Gray (10YR 5/1 dry); very dark gray (10YR 3/1 moist) sandy loam. Very friable, loose, nonsticky and nonplastic. Soil reacts violently to dilute HCl. The pH is 8.0. Clear and smooth lower boundary.
57Calif-07-3-2 AC	2- 8"	Light grayish brown (10YR 5.5/2 dry); dark grayish brown (10YR 4/2 moist) sandy loam. Massive, soft, friable, nonsticky and nonplastic. Soil reacts violently to dilute HCl. The pH is 8.0. Lime disseminated and as soft light gray blotches (10YR 7/2 dry). Root holes have rust colored stainings. Abrupt, smooth lower boundary.
57Calif-07-3-3 C_{11}	8-26"	Light gray (10YR 7/2 dry), light grayish brown (10YR 6/2 moist) sandy loam. Massive, slightly hard, friable, nonsticky, and nonplastic. Soil reacts violently to dilute HCl. The pH is 8.0. Diffuse, wavy lower boundary.
57Calif-07-3-4 C_{12}	26-44"	Pale brown (10YR 6/3 dry), brown to yellowish brown (10YR 5/3.5 moist) sandy loam. Massive, soft, firm, nonsticky and nonplastic. Soil reacts violently to dilute HCl. The pH is 8.0. Strong brown mottlings are common, medium, and prominent. Root holes have rust colored stainings. Diffuse, wavy lower boundary.
57Calif-07-3-5 C_{13}	44"+	Light yellowish brown (10YR 6/4 dry), yellowish brown (10YR 5/4 moist) sandy loam. Massive, friable, soft, nonsticky and nonplastic. Soil reacts violently to dilute HCl. The pH is 8.0. Strong brown mottlings - many, coarse, and prominent increasing in amounts with depth. Root holes have rust colored coatings.

1/ The Piper series is in a coarse-loamy, mixed, calcareous, thermic family of Aeric Haplaquepts. This pedon has less clay and more than 50 percent fine sand.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Reyes silty clay loam **LOCATION** Sonoma County, California

SOIL FAMILY: Sulfic Haplaquept, fine, mixed, acid, thermic.

SOIL NOS. S61Calif-49-13 **LAB. NOS.** 61478 - 61483

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	3A1 0.02-0.002	2A2 > 2	TEXTURAL CLASS
0- 7	Ap1	0.1	0.3	0.2	0.5	0.3	39.7	58.9	3.8	36.5	0	c-sic
7-14	[illegible]	0.1	0.2	0.1	0.5	0.4	33.9	60.7	2.3	36.2	0	[illegible]

REYES SILTY CLAY LOAM
S61Calif-49-13

Location: Sonoma County, California. Leveroni Ranch - 57° and 3800 feet from railroad bridge at Wingo. SW1/4, SW1/4, section 4, T. 4 N., R. 5 W. Date of Sampling: June 28, 1961.
Collectors: Reuben Nelson, Leo C. Klameth, V. C. Miller, G. Lambert, J. H. Kashiwagi, F. Graham.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Reyes silty clay loam **LOCATION** Sonoma County, California

SOIL FAMILY: Sulfic Haplaquept, fine, mixed, acid, thermic.

SOIL NOS. S61Calif-49-14 **LAB. NOS.** 61484 - 61489

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1	2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2	
0- 7	Ap1	0.0	0.3	0.2	0.7	0.5	45.2	53.1	4.9	41.2	0	sic
7-12	Ap2	0.0	0.2	0.1	0.7	0.5	44.7	53.8	3.9	41.7	0	sic
12-23	C3g	0.0	0.2	0.1	0.5	0.4	46.0	52.8	4.5	42.2	0	sic
23-39	C4g	0.1	0.2	0.3	1.0	0.7	41.6	56.1	2.6	40.3	0	sic
39-49	C5g	0.3	0.3	0.1	0.3	0.2	41.5	57.3	1.8	40.1	0	sic
49-59	C6g	0.2	0.3	0.1	0.2	0.3	46.5	52.4	4.6	42.3	0	sic

pH 8C1a H_2O 1:1	pH 8C1c KCl		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	SATURATION EXTRACT SOLUBLE 6P1a Na^+	6Q1a K^+	6N1a Ca^{++}	6O1a Mg^{++}	6K1a Cl^-	6I1a $CO_3^=$	8A1 6J1a HCO_3^-
						← milliequivalents per liter →						
4.1			3.41	0.299	11	11.2	0.6	2.7	8.7	19.1	-	-
4.1			3.15	0.290	11	15.3	0.6	3.8	11.9	28.4	-	-
3.7			2.01	0.198	10	35.9	1.5	3.8	14.4	40.2	-	-
3.5			2.52	0.235	11	64.3	1.5	6.0	25.1	62.6	-	-
3.1			5.44	0.341	16	228.	3.6	22.4	200.	225	-	-
3.7			4.11	0.261	16	77.8	5.4	25.4	242	271	-	-

5A2a CATION EXCHANGE CAPACITY (-Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 1/	EXCH. Na % 1/	8A1a Elec. Conduc. mmhos/cm. at 25° C.	Total Sulfur S 7C1 %	6C1b Ext. Iron % Fe	4B2 Soil water 15 atmos. Pct.	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →											
42.7	4.5	11.1	27.3	1.1	1.0			2.6	0.28	1.4	23.8	70.3
43.6	4.9	11.2	27.4	1.4	0.8			3.4	0.20	1.3	24.1	72.5
41.1	3.3	8.5	32.8	2.7	0.9			5.2	0.37	1.6	34.1	88.3
51.4	8.9	27.6	39.8	3.5	1.2			8.2	0.49	2.0	39.2	96.5
45.1	8.5	28.4	41.3	5.6	1.2			28.3	2.06	0.9	40.4	99.8
45.5	10.0	30.3	31.3	26.2	2.0			25.5	3.24	0.6	38.6	94.7

1/ Because of oxidizeable S in the soil base saturation and ESP have no meaning.

MEYERS SILTY CLAY LOAM
S61Calif-49-14

Location: Sonoma County, California. Reading Ranch - 278° and 4500 feet from railroad bridge at Wingo. NW1/4, NW1/4, section 8, T. 4 N., R. 5 W. Date of Sampling: June 29, 1961.
Collectors: Reuben Nelson, Leo C. Klameth, V. C. Miller, G. Lambert, J. H. Kashiwagi, F. Graham.
Classification: Sulfic Haplaquept, fine, mixed, acid, thermic.
Vegetation: Natural before cultivation - tules, saltgrass, hydrophytic plants. Use: Oat hay production. Climate: Subhumid mesothermal - annual precipitation 24.7 inches, mean annual January temperature 46.7° F., mean July temperature 66.3° F., mean annual temperature 57.5° F. Parent Material: San Pablo Bay sediments. Elevation: Minus 1 foot. Relief: Gently undulating on 0-1 percent slope. Drainage: Poor. Biological Activity: Anaerobic bacteria in waterlogged subsoil.
Remarks: Soil conditions in this profile have been influenced by reclamation of tidal flats which have been diked and drained. Apparently deep plowing in past years has brought about a mixing of colors in the surface horizons and also reaction may vary from one aggregate to another in most of the horizons. This soil resists rewetting after it has dried beyond the "moist to the feel stage".

HORIZON	DESCRIPTION
Ap1 * RSL No. 61484	0 to 7 inches, mixed light brownish gray (10YR 6/2) and light gray (10YR 7/2) silty clay loam, olive gray (5Y 4/2) when moist; fine granular structure; slightly hard when dry, friable when moist, sticky and plastic when wet; abundant very fine and fine roots; reaction ranges from very strongly acid to slightly acid, pH 5.0 - 6.1; gradual, wavy boundary.
Ap2 RSL No. 61485	7 to 12 inches, grayish brown (10YR 5/2) silty clay loam, olive gray (5Y 4/2) when moist; granular structure; hard when dry, friable when moist, sticky and plastic when wet; abundant very fine and fine roots; very strongly acid, pH 5.0; abrupt, wavy boundary.
C3g RSL No. 61486	12 to 23 inches, gray (10YR 6/1) clay with common fine distinct yellowish brown (10YR 5/6) mottles when dry; dark grayish brown (2.5Y 4/2) with common, fine, distinct yellowish brown (10YR 5/4) mottles when moist; mottling occurs from sulfur residues, root channel deposits and fine organic matter; massive; extremely hard when dry;

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Marcuse silty clay (taxadjunct) **LOCATION** Riverside County, California

SOIL FAMILY: Vertic Haplaquept, fine, montmorillonitic, calcareous, thermic.

SOIL NOS. S57Calif-33-7 **LAB. NOS.** 57501 - 57506

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0- 7	Ap	0.1	0.3	0.2	1.5	3.3	48.6	46.0	12.7	40.3	0	sic
7-18	B1	0.1	0.2	0.1	0.8	2.6	43.8	52.4	10.9	36.1	0	sic
18-28	B21	0.0	0.1	0.1	0.8	2.6	42.4	54.0	11.0	34.6	0	sic
28-44	B22	0.1	0.1	0.1	0.6	2.2	42.2	54.7	9.9	35.0	0	sic
44-56	Cca1	0.2	0.1	0.1	0.8	2.2	44.2	52.4	11.1	35.9	0	sic
56-64+	Cca2	0.2	0.2	0.2	1.0	2.8	43.0	52.6	12.0	34.6	0	sic

8C1b pH SATURATED PASTE 1:1	8C1a pH 1:10	6J1a HCO_3* meq/l	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	6I1a CO_3* meq/l	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL		4A3a Bulk Density g/cc	4B2 Soil Water 15 atmos. %
7.8	8.2	5.1	1.03	0.097	10.6	0.0	5.0	2.3				19.2
7.7	8.0	3.0	0.77	0.062	12.4	0.0	9.8	3.5			1.49	20.3
7.9	8.4	3.3	0.70	0.051	13.7	0.0	9.8	4.7			1.54	21.6
8.0	8.7	3.8	0.55	0.037	14.9	0.0	9.8	4.0			1.60	21.5
8.0	8.2	2.9	0.40	0.027	14.8	0.0	14.7	3.9	8.0		1.53	22.3
8.0	8.1	2.9	0.32			0.0	17.7	3.6	11.5		1.49	23.2

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	5D1 EXCH. Na %	SATURATION EXTRACT SOLUBLES 8A1 6P1a Na	6Q1a K	6N1a	6O1a	6K1a	8A MOISTURE AT SATURATION %
		milliequivalents per 100g. soil					milliequivalents per liter					
40.3	38.4	10.4	-	5.2	1.6	13	30.3	0.5	14.5	6.2	27.4	71.6
41.0	39.3	13.1	-	7.7	1.4	19	63.0	0.5	29.0	8.0	82.9	74.3
41.0	34.9	12.7	-	14.6	1.4	36	78.0	0.4	12.2	3.4	83.5	78.4
41.1	27.8	9.9	-	19.8	1.3	48	89.0	0.2	5.0	3.9	81.6	88.3
44.1	59.8	9.9	-	18.4	1.1	42	141.0	0.4	26.5	3.9	91.7	83.8
44.8	59.7	24.9	-	21.2	1.0	47	166.0	0.5	28.4	5.6	107.3	86.5

MARCUSE SILTY CLAY (taxadjunct) 1/
S57Calif-33-7

Location: Riverside County, California. 1100 feet west and 1000 feet south of the N1/4 corner of Sec. 26, T. 4 S., R. 3 W. In barley field east side of San Jacinto River. Floyd Bonge Ranch, about 4 miles east-northeast of Perris.
Date of Sampling: June 7, 1957.
Description by: Fred C. Eggers. Collectors: George Borst, R. E. Nelson, Cliff Henry, and Arnold Knecht.
Classification: Vertic Haplaquept, fine, montmorillonitic, calcareous, thermic.
Vegetation: Irrigated barley--an excellent crop. Precipitation: 12 inches.
Parent Material: Mixed alluvium, chiefly granitic. Physiography: Flood plain.
Elevation: 1,418 feet. Drainage: Slow.
Remarks: Colorado River water used on this field three years.

HORIZON	DESCRIPTION
S57Calif-33-7-1 Ap 0-7"	Light brownish gray (10YR 6/2 dry, dark gray 4/1 moist) silty clay; strongly defined medium subangular blocky breaking to strongly defined medium granular; hard when dry, firm when moist; sticky and very plastic when wet; moderately slow permeability; many plant roots; many fine pores; pH 8.4; strong effervescence; clear smooth lower boundary.
S57Calif-33-7-2 B1 7-18"	Gray (10YR 5/1 dry, very dark gray 3/1 moist) silty clay; strongly defined medium subangular blocky with slight evidence of prismatic structure; very hard when dry, very firm when moist; sticky and very plastic when wet; moderately slow permeability; many plant roots; many pores; pH 8.2; strong effervescence; gradual smooth lower boundary.
S57Calif-33-7-3 B21 18-28"	Gray (10YR 5/1 dry, dark gray to very dark gray 4/1-3/1 moist) silty clay; strongly defined medium subangular blocky with slight evidence of prismatic structure; very hard when dry, very firm when moist; sticky and very plastic when wet; slow permeability; occasional plant roots; many fine pores; pH 8.2; strong effervescence; gradual smooth lower boundary.
S57Calif-33-7-4 B22 28-44"	Gray (10YR 5/1 dry, dark gray to very dark gray 4/1-3/1 moist) silty clay; strongly defined medium subangular blocky with slight evidence of prismatic structure; very hard when dry, very firm when moist; sticky and very plastic when wet; slow permeability; occasional plant roots; many fine pores; pH 8.2; strong effervescence; clear wavy lower boundary.
S57Calif-33-7-5 Ccal 44-56"	Gray (10YR 6/1-5/1 dry, dark gray 4/1 moist) silty clay; massive; very hard when dry, very firm when moist; sticky and very plastic when wet; [illegible]

SOIL SURVEY LABORATORY Riverside, California

SOIL FAMILY: Histic Humaquept, coarse-loamy, mixed, calcareous, thermic

SOIL NOS. S57Calif-7-4 **LAB. NOS.** 5835 - 5839

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2	
0-4	A1	Organic matter										
4-10	AC	0.2	6.2	14.8	46.2	5.9	8.9	17.8	31.6	6.7	0	fsl

PIPER SANDY LOAM (taxadjunct) 1/
S57Calif-7-4

Collector: R. C. Huff. **Date of Sampling:** May 14, 1957.
Location: Contra Costa County, California. NE corner of SE1/4SW1/4 sec. 10, T. 2 N., R. 3 E. MDB&M, projected. Eastern Contra Costa SCD. Bethel Island, Honegger Ranch at Pit Silo 1/2 mile due east of house.
Classification: Histic Humaquept, coarse-loamy, mixed, calcareous, thermic.
Native Vegetation: Annual weeds and grasses - mainly ripgut brome (Bromus ridigus) and Red Brome (Bromus rubens).
Climate: Rainfall 18". Cool moist winters. Dry summers with occasional fog.
Parent Material: Mixed alluvium.
Relief: Undulating with mounds or long low ridges surrounded by peat and muck which has subsided from around these areas.
Elevation: 0 feet.
Drainage: Excessive surface and good internal.
Root Distribution: Numerous 0-4", some 4-10", few to 60".

HORIZON	DESCRIPTION
57Calif-07-4-1 A_1 0- 4"	Very dark gray (10YR 3/1 dry); black (10YR 2/1 moist) sandy loam. Crumb structure, loose, nonstick y and nonplastic. The soil mass is noncalcareous but lime occurs as small nodules which appear to have been brought up from the underlying horizon. The pH is 8.0. Clear, wavy lower boundary.
57Calif-07-4-2 AC 4-10"	Light gray (10YR 7/1 dry); grayish brown (10YR 5/2 moist) sandy loam. Massive, soft, very friable, nonsticky and nonplastic. The soil reacts violently to dilute HCl and lime is disseminated. The pH is 8.0. Root holes have iron coatings. Krotovinas are common. Clear, smooth lower boundary.
57Calif-07-4-3 C_{11} 10-24"	Light gray (10YR 7/1 dry); grayish brown (10YR 5/2, moist) sandy loam. Massive, firm, friable, nonsticky and nonplastic. The soil reacts violently to dilute HCl, the pH is 8.0. Rust-colored iron stains follow root holes into horizon below. Diffuse and wavy boundary.
57Calif-07-4-4 C_{12} 24-38"	Light gray (10YR 7/2, dry); light grayish brown (10YR 6/2 moist) sandy loam. Massive, hard, firm, nonsticky and nonplastic. Brownish colored mottlings (10YR 6/4-7/4 dry) are common, medium and distinct. Root holes have rust colored coatings. Soil reacts violently to dilute HCl. The pH is 8.0. Diffuse and wavy boundary.
57Calif-07-4-5 C_{13} 38"+	Pale brown (10YR 6.5/3 dry); yellowish brown (10YR 5/4 moist) sandy loam. Massive, hard, firm, nonsticky and nonplastic. Soil reacts violently to dilute HCl. The pH is 8.0. Rust-colored mottlings (7.5YR 5/6) many, coarse, and prominent.

NOTE: The A_1 horizon of this soil is higher in organic matter (admixture of peat) than the A_1 of 57Calif-07-3-1.

1/ The Piper series is in a coarse-loamy, mixed, calcareous, thermic family of Aeric Haplaquepts. This pedon has a histic epipedon.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Dystric Cryochrept, loamy-skeletal, mixed

SOIL Umpa gravelly sandy loam (taxadjunct) SOIL Nos. S68Calif-31-7 LOCATION Placer County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. R7021 - 7025

Depth cm.	Horizon	1B1b Size class and particle diameter (mm) 3A1 Total Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Sand Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt 0.05–0.02	Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	1/ Coarse fragments 1A2a > 2	2 - 19	19 - 76
		Pct. of < 2 mm														Pct.	Pct of < 76mm	
0-8	A1	53.6	37.0	9.4	11.4	10.8	8.4	14.0	9.0	11.8	25.2	28.5	44.6					
8-28	A3	52.4	33.2	14.4	10.8	11.2	8.1	14.0	8.3	9.7	23.5	25.5	44.1					
28-51	B1	52.9	32.9	14.2	6.3	11.3	9.7	16.7	8.9	9.7	23.2	27.5	44.0					
51-89	B2	46.1	36.8	17.1	4.2	11.1	8.4	13.9	8.5	8.9	27.9	24.9	37.6					
89-125	R	42.9	38.1	19.0	3.5	10.3	8.2	13.3	7.6	7.3	30.8	22.3	35.3					

Depth cm.	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2b Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility LEf Pct	4D1 Extensibility LE Pct	pH 8C1a H_2O 1:1	8C1e $CaCl_2$
0-8	3.20														6.3	5.8
8-28	0.48														6.1	5.2
28-51	0.45														5.9	5.0
51-89	0.57														5.7	5.0
89-125	0.23														5.5	4.9

Depth cm.	Extractable bases 5B4a 6N2e Ca	6O2d Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A6a NH_4OAc	5A3a Sum	6G1e KCl Ext. Al^{+++}	Base saturation 5C1 NH_4OAc Pct.	Pct
	meq/100 g										
0-8	3.9	0.5	0.2	0.7	5.3		18.9			28	
8-28	2.9	0.9	0.2	0.8	4.8		13.3			36	
28-51	2.5	1.0	0.2	0.9	4.6		14.4			31	
51-89	3.9	0.8	0.2	1.0	5.9		15.6			37	
89-125	5.3	1.2	0.3	0.8	7.6		15.0			50	

Depth cm.	Clay mineralogy (<0.002 mm.) 3A1 Mont.	Verm.	Mica 7A2 X-ray	Kaolin.	
0-8					
8-28					
28-51					
51-89					
89-125					

UMPA GRAVELLY SANDY LOAM (taxadjunct) 1/
S68Calif-31-7

Location: Tahoe Basin, Placer County, California; 0.4 mile north of SE corner, sec. 23, T. 16 N., R. 16 E., 75 feet SW of logging road. Date of Sampling: September 19, 1968.
Description by: G. M. Kennedy and G. L. Anderson. Collectors: J. H. Rogers, L. I. Larsen, W. E. Dollarhide.
Classification: Dystric Cryochrept, loamy-skeletal, mixed.
Vegetation: Red fir, white fir, western white pine, pinemat manzanita. Climate: Precipitation is 35 to 40 inches. Mean annual temperature 40°F., January 23°F., July 57°F. Parent Material: Andesite. Topography: Uplands, 8 percent slope, east aspect. Elevation: 7700 feet. Drainage: Well drained; runoff is medium; permeability is moderate. Soil Moisture: Dry throughout profile.
Remarks: This is not the type location for Survey Area. Field determination of pH by Hellige Truog trialex indicator. Five percent of surface area covered with stones and

an occasional boulder. Soil temperature at 20 inches 52°F. Fine to coarse roots found in fracture planes at 35 to 49 inches. B2 horizon appeared marginal to an argillic horizon. Coarse fragments estimated as percent by volume.

HORIZON	DESCRIPTION
	(3-0 cm.)
01	1 to 0 inch, fresh and decomposed conifer needles. Not sampled.
	(0-8 cm.)
A1 RSL 7021	0 to 3 inches, grayish brown (10YR 5/2) gravelly sandy loam, dark reddish brown (5YR 3/3) moist; weak coarse platy and weak coarse granular structure; soft, very friable, nonsticky and nonplastic; many very fine, fine, common medium and few coarse roots; common very fine tubular and interstitial pores; 15 percent gravel; medium acid (pH 6.0); clear smooth boundary.
	(8-28 cm.)
A3 RSL	3 to 11 inches, pinkish gray (7.5YR 6/2) gravelly sandy loam, reddish brown (5YR 4/4) moist; massive; slightly hard, friable, nonsticky and nonplastic; many very fine,

25 percent gravel and 5 percent cobbles; strongly acid (pH 5.5); clear wavy

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Dystric Cryochrept, fine-loamy, mixed

SOIL Umpa gravelly loam, deep variant SOIL Nos. S68Calif-31-10 LOCATION Placer County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. R7026 - 7031

UMPA GRAVELLY LOAM, DEEP VARIANT 1/
S68Calif-31-10

Location: Tahoe Basin, Placer County, California; 2,300 feet north and 650 feet east of E¼ corner sec. 35, T. 16 N., R. 16 E., 30 feet east of logging road and 2¼ miles NNW of Tahoe City. Date of Sampling: September 19, 1968.
Description by: J. H. Rogers. Collectors: J. H. Rogers and P. G. Nazar.
Classification: Dystric Cryochrept, fine-loamy, mixed.
Vegetation: Red fir, whitebark and western white pine, prostrate manzanita. Climate: Precipitation is 35 to 40 inches. Mean annual temperature is 40°F., January 23°F., July 57°F. Parent Material: Latite. Topography: Slightly convex upland with 18 percent slope and north aspect. Elevation: 7120 feet. Drainage: Well drained; runoff is medium; permeability is moderate. Soil Moisture: Soil dry to 15 inches, slightly [illegible]

moist below.
Remarks: This is not the series type location. Profile taken in Jorge-Tahoma complex. Surface stones cover about 6 percent of area. Field determination of pH by Hellige-Truog triplex indicator. Question as to whether the B23t formed by weathering in place. Coarse fragments estimated as percent by volume. *Depth to a lithic contact is more than 40 inches. Clay content is higher and content of coarse fragments is lower than defined for the Umpa series.

HORIZON	DESCRIPTION
01	(1-0 cm.) 1/2 to 0 inch, fresh and decomposed conifer needles. Not sampled.
A11 RSL 7026	(0-8 cm.) 0 to 3 inches, brown and dark brown (10YR 4/3, 3/3) gravelly loam, very dark brown (10YR 2/2) moist; weak fine granular structure; soft, very friable, nonsticky, nonplastic; many very fine and common fine roots; many very fine tubular and interstitial pores; 20 percent gravel; slightly acid (pH 6.2); abrupt smooth boundary.
A12 RSL 7027	(8-33 cm.) 3 to 13 inches, brown (7.5YR 5/4) gravelly loam, reddish brown (5YR 4/4) moist; weak fine and medium subangular blocky structure; slightly hard, friable, slightly sticky, slightly plastic; common very fine and fine and many medium roots; common very fine, fine and medium pores; 20 percent gravel; medium acid (pH 5.8); clear smooth boundary.
B1t RSL 7028	(33-51 cm.) 13 to 20 inches, light brown (7.5YR 6/4) gravelly loam, brown (7.5YR 4/4) moist; moderate fine and medium subangular blocky structure; slightly hard, friable, slightly sticky and slightly plastic; common very fine, fine, medium and coarse roots; common very fine, fine and medium pores; few thin clay films in pores; 20 percent gravel and 5 percent cobbles; strongly acid (pH 5.5); clear smooth boundary.
B21t RSL 7029	(51-79 cm.) 20 to 31 inches, light brown (7.5YR 6/4) gravelly clay loam, brown (7.5YR 4/4) moist; moderate fine and medium subangular blocky structure; slightly hard, friable, slightly sticky and slightly plastic; common very fine, fine and medium roots; common very fine, fine and medium pores; common thin clay films line pores and on faces of peds; 20 percent gravel, 10 percent cobbles; strongly acid (pH 5.5); clear smooth boundary.
B22t RSL 7030	(79-130 cm.) 31 to 52 inches, light brown (7.5YR 6/4) gravelly clay loam, brown (7.5YR 4/4) moist; moderate fine subangular blocky structure; hard, firm, slightly sticky and slightly plastic; few very fine, fine and medium roots; few very fine and fine tubular pores; many thin clay films line pores and on faces of peds; 25 percent gravel, 10 percent cobbles, 2 percent stones; very strongly acid (pH 4.8); clear wavy boundary.
B23t RSL 7031	(130-165 cm.) 52 to 65 inches, light brown (7.5YR 6/4) clay loam, reddish brown (5YR 4/4) moist; moderate fine and medium subangular blocky structure; very hard, firm, slightly sticky, and plastic; no roots; few very fine and fine tubular pores; many thin clay films line pores and on faces of peds; no gravel; very strongly acid (pH 4.8); clear wavy boundary.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Durochrept, coarse-loamy, mixed, thermic
SOIL Arlington loam (taxad junct) SOIL Nos. S66Calif-37-3 LOCATION San Diego County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66243 - 66248

Depth (in.)	Horizon	1B1h Total Sand (2–0.05)	Total Silt (0.05–0.002)	Total Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)			1/ Coarse fragments 1A2a > 2 Pct.	2 – 19 (Pct of < 76mm)	19 – 76 (Pct of < 76mm)	
		Size class and particle diameter (mm) 3A1																	
		Pct. of < 2 mm																	
0- 1	A11	50.9	38.2	10.9	3.4	15.8	15.4	11.8	4.5	17.8	20.4	26.7	46.4	0	11	10	8	2	2/
1- 5	A12	54.5	34.7	10.8	2.7	14.9	17.9	13.5	5.5	15.6	19.1	26.2	49.0	0	11	11	8	3	2/
5-10	A2	51.5	37.6	10.9	2.2	13.3	16.9	13.6	5.5	16.9	20.7	27.9	46.0	0	11	15	9	6	2/
11-21	B2tsim	65.5	9.5	25.0	11.3	22.2	15.7	12.5	3.8	5.8	3.7	15.3	61.7	0	25	9	0	1	3/
21-24	IIB31sim	82.7	9.0	8.3	16.6	27.8	19.5	14.0	4.8	5.2	3.8	16.2	77.9	0	8	27	4	11	4/
24-58+	IIIB32sim	63.4	10.5	26.1	5.0	19.0	23.5	11.9	4.0	5.5	5.0	14.2	59.4	0	26	80	10	59	4/

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct	Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content Pct	4B1c 1/3 bar Pct	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1		
0- 1	5.11														5.2		
1- 5	0.96					1.70	1.73		0.93		12.9	4.1	0.005	0.004	6.0		
5-10	0.31					1.81	1.84		0.89		10.5	3.2	0.004	0.004	5.4		
11-21	0.22					1.52	1.64		0.94		23.3	13.0	0.027	0.026	4.5		
21-24	0.13														4.3		
24-58+	0.13														4.0		

Depth (in.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1a NH_4OAc	5A3a Sum	6G1d KCl Ext. Al^{+++}	Base saturation: 5C3 Sum	5C1 NH_4OAc
	meq/100 g									Pct	Pct.
0- 1	4.7	1.9	0.5	0.7	7.8	11.0	18.4	18.8	0.2	41	42
1-5	4.4	2.1	0.4	0.4	7.3	3.9	8.3	11.2	-	65	88
5-10	2.2	2.0	0.4	0.4	5.0	3.9	6.7	8.9	0.4	56	75
11-21	4.0	6.2	2.3	0.4	12.9	13.7	23.5	26.6	7.4	48	55
21-24	2.9	4.1	2.7	0.2	9.9	10.7	20.7	20.6	5.1	48	48
24-58+	3.8	6.8	4.8	0.3	15.7	8.7	19.0	24.4	3.6	64	83

Depth (in.)	Clay mineralogy (<0.002 mm) 7A: Mont.	Verm.	Mica	Kaolin	
	X-ray				
0- 1					
1- 5	t	-	x	xx	

1/ From characterization sample (5Kg)(except as noted).
2/ 2-19 mm. from characterization sample, 19-76 mm. from field weight (>50 Kg).
3/ 2-19 mm. from characterization sample, 19-76 mm. from field weight; >2 mm. includes volume estimate of >76 mm. fraction.

ARLINGTON LOAM (taxadjunct) 1/
S66Calif-37-3

Location: San Diego County, California; Miramar Transect No. 1, station 44.1; 950 feet west and 750 feet north of the southeast corner of section 24, T. 15 S., R. 3 W. Date of sampling: June 10, 1966.
Description by: George Borst, David Estrada, Gerald Kester, Reuben Nelson, and Dennis Nettleton. Samples collected by: Klaus Flach, Benny Brasher, and Dennis Nettleton.
Classification: Typic Durochrept, coarse-loamy, mixed, thermic.
Vegetation: Flattop buckwheat. Climate: Weather record for San Diego, near the sampling site, is mean winter air temperature 55° F., mean summer air temperature 68° F., mean annual precipitation 10 inches. Parent Material: Stream deposit over Pleistocene marine terrace. Topography: Flat marine terrace of less than 1 percent slope. Elevation: 425 to 450 feet. Drainage: Moderately well drained; runoff is medium; permeability is very slow. Soil Moisture: Dry.

Remarks: Very cobbly below 24 inches.

HORIZON	DESCRIPTION
A11 RSL No. 66243	0 to 1 inch, grayish brown (10YR 5/2) loam, very dark grayish brown (10YR 3/2) when moist; weak coarse platy structure; soft, friable, nonsticky, slightly plastic; abundant very fine roots; very strongly acid (pH 5.0); abrupt smooth boundary.
A12 RSL No. 66244	1 to 5 inches, strong brown (7.5YR 5/6) loam, dark yellowish red (5YR 4/6) when moist; massive; hard, friable, nonsticky, plastic; few very fine to medium roots; few very fine tubular pores; strongly acid (pH 5.5); abrupt smooth boundary.
A2 RSL No. 66245	5 to 10 inches, light brown (7.5YR 6/4) loam, brown (7.5YR 5/4) when moist; massive; hard, friable, nonsticky, plastic; few very fine to medium roots; many very fine tubular pores; very strongly acid (pH 5.0); abrupt smooth boundary.
B1	10 to 11 inches, pink (7.5YR 7/4) loam, light brown (7.5YR 6/4) when moist; massive; hard, friable, slightly sticky, plastic; few very fine to medium roots; many very fine tubular pores; few thin colloid stains on mineral grains; very strongly acid (pH 4.6); abrupt wavy boundary.*
B2tsim RSL No. 66246	11 to 21 inches, grayish brown (2.5Y 5/2) duripan, dark grayish brown (2.5Y 4/2) when moist with dark red (2.5YR 3/6) coatings of iron oxide and clay on cleavage planes; strong medium angular blocky structure; indurated; very few, very fine to fine roots; very strongly acid (pH 4.5); clear smooth boundary.
IIB31sim RSL No. 66247	21 to 24 inches, pale brown (10YR 6/3) gravelly duripan, brown (10YR 5/3) when moist; massive; indurated; very few fine roots; very strongly acid (pH 4.5); clear smooth boundary.
IIIB32sim RSL No. 66248	24 to 58 inches +, very cobbly duripan; massive; indurated, but somewhat weaker than above.

* Strongly degraded at contact with underlying duripan.

1/ The Arlington series is in a coarse-loamy, mixed, thermic family of Haplic Durixeralfs. This pedon has an indurated duripan. It also has clay skins and a clay increase sufficient for an argillic horizon in the duripan horizon. However, the classification system does not provide for horizons having properties of both argillic horizons and duripans.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Pentz sandy loam (taxadjunct) LOCATION Amador County, California

SOIL FAMILY: Typic Durochrept, coarse-loamy, mixed, thermic.

SOIL NOS. S60Calif-3-20 LAB. NOS. 6085 - 6089

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 2/	
0- 5	A11	1.9	13.7	13.4	27.9	13.2	21.2	8.7	40.8	10.0	0	fsl-sl
5-10	A12	1.5	14.7	13.1	28.4	13.0	20.6	8.7	40.0	10.2	0	fsl-sl
10-16	C1	1.9	13.9	12.4	27.6	13.3	21.2	9.7	40.3	10.5	1	fsl-sl

PENTZ SANDY LOAM (taxadjunct) 1/
S60Calif-3-20

Location: On Strohm's Ranch, 0.1 mile S of the center of the NW1/4 of sec. 30, T. 5 N., R. 10 E., 1/4 mile SW of Easterling's house, 3/4 mile SW of Buena Vista Peaks, Photo DDE-26-114, Amador County, California.
Date of Sampling: May 2, 1960 Description by: Jack McClelland.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Pentz fine sandy loam (taxadjunct) **LOCATION** Amador County, California

SOIL FAMILY: Typic Durochrept, coarse-loamy, mixed, thermic.

SOIL NOS. 60Calif-3-21 **LAB. NOS.** 6090 - 6095

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 5	A11	1.4	5.3	9.7	31.3	13.6	28.5	10.2	44.1	16.0	3	fsl
5-10	A12	1.2	4.7	9.6	30.8	14.5	28.4	10.8	45.5	14.7	4	fsl-sl
10-19	C1	1.7	5.2	9.5	29.7	12.9	29.7	11.3	43.7	15.5	1	fsl
19-22	IIC2m	Not	determined									
22-28	IIIC3	3.2	8.5	12.2	29.4	12.0	24.5	10.2	41.3	11.6	1	fsl
28-46+	IIIC4	2.9	10.1	13.3	38.5	11.6	18.7	4.9	43.1	8.4	0	ls

pH 8C1a H_2O 1:1	8C1c KC1 (1 N) 1:1	1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC x 10³ MILLIMHOS PER CM 25°C.	$CaCO_3$ equivalent %	6C1b Free iron as % Fe_2O_3	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.2	5.0		0.93	0.096	9.7				0.6			7.3
5.9	4.6		0.49	0.041	12.0				0.6			6.8
5.7	4.3		0.23	0.027					0.6			7.7
		Not	determined									
5.5	3.9		0.13						0.5			10.0
5.3	3.5		0.02						0.4			11.3

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Na 6P2a	5E1a K 6Q2a	BASE SAT. % 5C3	Clay mineralogy (<0.002 mm.) Mont.	Verm.	Mixed layer Mica-Verm. X-ray 7A2	Mica	% Kaolin DTA 7A3	% Gibb. DTA 7A3
		milliequivalents per 100g. soil										
11.9	6.2	1.2	3.4	0.1	1.0	71						
10.1	4.7	1.2	2.7	0.1	1.0	72	xx	-	-	x	30	-
9.4	3.8	1.4	2.3	0.2	1.1	74	x	-	-	xx	35	-
		Not	determined									
10.9	2.9	2.0	2.2	0.3	2.1	77						
13.2	3.8	3.0	2.2	0.5	2.8	82						

1/ Percent of total soil

- = looked for but not found
tr = detected, very small amount
x = detected, small amounts
xx = moderate amounts
xxx = large amounts
xxxx = predominant amounts

PENTZ FINE SANDY LOAM (taxadjunct) 1/
S60Calif-3-21

Location: Arroyo Seco Land Grant (section lines projected) 0.2 mile N of the center of the SE1/4 of sec. 9, T. 5 N., R. 9 E., 100 feet SE of Highway 88, 0.6 mile SW of Jackson Creek bridge on Highway 88, on Orr Ranch, Amador County.
Date of Sampling: May 2, 1960 Description by: Jack McClelland.
Collectors: Grant Kennedy, Reuben Nelson, Robert Sketchley, Clifford Henry, John Rogers, Paul Nazar, Jack McClelland
Vegetation: Annual grasses and forbs including soft chess, filaree, lupine, and clovers
Climate: Mean annual rainfall - 19.6 inches; average temperatures: January-46°F.; July - 88° F.; mean - 61° F. (Ione Station)
Parent Material: Rhyolitic tuff of the Valley Springs formation containing numerous hardpan lenses in the upper portion at least.
Relief: Gently sloping foothills on a SW facing plane slope of 5 percent.
Elevation: 220 feet
Drainage: Well drained; moderate runoff; moderately permeable above the parent rock or hardpans. Considerable seepage occurs during the winter months in these soils above the parent rock.
Remarks: This soil is a Lithosol. Bureau of Public Roads and California Division of Highway samples were obtained of the A11 (0-5 inches) and IIIC4 (28-46 inches) horizons. The rhyolitic tuff in the roadcut nearby when dry and baked is extremely hard throughout. Soil was moist when described.
Classification: Typic Durochrept, coarse-loamy, mixed, thermic.

HORIZON		DESCRIPTION
A11 RSL No. 6090	0- 5"	Light brownish gray (10YR 6/5/2) sandy loam, dark grayish brown (10YR 4/2) when moist; massive; slightly hard, friable, nonsticky and nonplastic; abundant very fine roots; many very fine interstitial and tubular pores; slightly acid (pH 6.2); clear smooth boundary.
A12 RSL No. 6091	5-10"	Similar to above but medium acid (pH 6.0); clear smooth boundary.
C1 RSL No. 6092	10-19"	Light brownish gray (10YR 6/2) sandy loam, brown (10YR 5/3) when moist; massive; slightly hard, friable, nonsticky and nonplastic; abundant very fine roots; many very fine interstitial and tubular pores, many fine tubular pores; discontinuous hardpan layer near the upper boundary which varies from 1 to 3 inches thick; strongly acid (pH 5.5); abrupt smooth boundary.
IIC2m RSL No. 6093	19-22"	Very pale brown (10YR 7/3) indurated hardpan with silica coating on upper surface; brown (10YR 5/3) when moist; extremely hard; few very fine roots along cleavage planes; occasional fine crack; abrupt smooth boundary.
IIIC3 RSL No. 6094	22-28"	Light gray (10YR 7/2) loamy sand, grayish brown (10YR 5/2) when moist; massive; slightly hard, friable, nonsticky and nonplastic between indurated discontinuous hardpan lenses (0 to 3); few very fine roots; many very fine interstitial, few very fine tubular pores; strongly acid (pH 5.5); abrupt smooth boundary.
IIIC4 RSL No. 6095	28-46"+	Light gray (5Y 7/2) loamy sand between a succession of hardpan layers 2 - 4" thick, the upper surfaces of which are coated with silica and the lower portions relatively uncemented, olive (5Y 5/3) when moist with yellowish brown (10YR 5/4 - 5/6) staining in lower portions of hardpan layers; indurated in upper hardpan lenses, slightly hard and firm in lower portions of lenses, nonplastic and nonsticky; no roots; many very fine interstitial pores in lower portion of hardpan; very strongly acid (pH 5.0).

1/ The Pentz series is in a loamy, mixed, nonacid, thermic, shallow family of Typic Xerorthents. This pedon is a taxadjunct because it has a duripan.

CONSECUTIVE PEDON NUMBER C 69 13 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES TRIGO HARDPAN VARIANT SOIL NO. S69 CALI. 31- 3 SAMPLE NO. 9 63-9 67 PLACER COUNTY AREA

SOIL FAMILY TYPIC DUROCHREPT COARSE-LOAMY,MIXED,THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SANDS VCS 2-1	CS 1-.05	MS .5-.25	FS .25-.10	VFS .1-.05	CSI .05-.02	FSI .02-.002	CLAYS CO. 2-.2U	FINE LT .2U	CARBONATE 3A1A	INT. II .2-.02	COARSE FRAGMENTS PCT. OF WHOLE SOIL I 75-20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75		
9 63	0- 23	AP	38.8	50.3	10.9	0.4	3.1	5.1	17.9	12.3	20.9	29.4				44.0	--A --	--	--
9 64	23- 36	B	39.2	48.8	12.0	0.3	3.1	5.2	16.7	13.9	17.8	31.0				41.9	--A --	--	--
9 65	36- 51	2C1SIM	22.7	59.8	17.5	0.8	3.1	4.2	8.8	5.8	10.5	49.3				21.2	--G --	--	--
9 66	51- 79	2C2SI	57.4	36.5	6.1	0.2	3.3	5.9	25.1	22.9	14.3	22.2		--		54.7	--A --	--	--
9 67	79-105	2C3	84.5	13.8	1.7	0.1	6.1	13.6	47.1	17.6	8.1	5.7				52.0	--A --	--	--

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A -PERCENT--	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B -PERCENT-	19-2 6E1B	EXT. IRON FE 6C2B PCT.	ATTERBERG LIMIT LL 4F2 PCT.	PL 4F1 NO.40	BULK DENSITY EST 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4AH	CM 3B2	WATER CONTENT 1/10 BAR 4B1C -----PERCENT---	1/3 BAR 4B1C	15 BAR 4B2	EXTENSIBLITY LEF 4D1 -PERCENT--	LE 4D1	PH H2O 1/1 8C1A	PH H2O 1/10 8C1A	PH CACL2 8C1E
									GRAMS PER CC											
9 63	0- 23	0.63	0.061						1.62	1.68	1.00		18.5	4.9	1.2	1.2	5.5		4.8
9 64	23- 36	0.32	0.038						1.60	1.68	1.00		17.5	5.5	1.6	1.6	5.7		5.1
9 65	36- 51	0.24	0.042						1.30	1.36	1.00		39.9	14.2	1.5	1.5	6.7		6.4
9 66	51- 79								1.54	1.59	1.00		20.8	7.1	1.1	1.1	6.8		6.5
9 67	79-105								1.48	1.51	1.00		10.6	5.4	0.7	0.7	6.6		6.2

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT.	C E C NA-OAC 5A2A	NH4-OAC 5A6A	EC 8A1A	WATER EXTRACT FROM SATURATED PASTE CA 6N1B	MG 6O1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	B 6T1A	PH SAT. PASTE 8C1B
		MEQ/100G					PCT.	MEQ/100G		MMHO	MEQ/1000 ML								PPM	
9 63	0- 23	5.4	2.2	0.1	0.1	7.8	76		10.3											
9 64	23- 36	6.5	2.8	0.2	0.1	9.6	87		11.0											
9 65	36- 51	16.6	7.7	1.2	0.2	25.7	100+		22.5											
9 66	51- 79	9.2	5.3	0.5	0.2	15.2	100+		15.0											

SAMPLE NO.	DEPTH CM.	WATER AT SAT. 8A PCT.	EXCHANGEABLE NA 5B4B MEQ.	NA 5D2 PCT.	SAR 5E	GYPSUM AMT. REQ. 6F1B PERCENT	6P3A	CLAY MINERALOGY LESS THAN 0.002 MM X RAY I 7A2E	II 7A2E	III 7A2E	DTA IV 7A2E	PCT. 7A3	LT. .0002 MM X RAY I 7A2E	II 7A2E	III 7A2E	MINERAL CODE	RELATIVE AMOUNT
9 63	0- 23		0.1	1				MI 4	KK 3	MT 1						MI MICA	1 TRACE
9 64	23- 36		0.2	2				MI 4	KK 3	MT 1						KK KAOLINIT	2 SMALL
9 65	36- 51		1.2	5												MT MONTMORT	3 MODERATE
9 66	51- 79		0.5	3													4 ABUNDANT
9 67	79-105		0.4	4				MT 4	MI 3	KK 3							5 DOMINANT
																	6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
G=GROUND WHOLE. HORIZON INDURATED OR GRAVEL TOO SOFT TO BE SEPARATED. ALL DATA FOR WHOLE SOIL.

TRIGO LOAM, HARDPAN VARIANT* 1/

S69Calif-31-3

Location: Placer County, California; 38 feet north and 700 feet east of SW corner sec. 30, T. 10 N., R. 6 E. John Fiddyment ranch approximately 3.7 miles northwest of Roseville, California. Date of Sampling: May 6, 1969.

Description by: G. M. Kennedy, T. M. Ryan. Collectors: K. W. Flach, W. D. Nettleton, G. M. Kennedy, L. A. Brittan, T. M. Ryan.

Classification: Typic Durochrept, coarse-loamy, mixed, thermic.

Vegetation: Filaree, crimson clover, rose clover, burr clover, soft chess. Climate: Average annual precipitation is 17 to 32 inches; mean annual air temperature is 61°F. Parent Material: Surficial sediments (fine grained sandstone and siltstone) deposited over Laguna Formation. Topography: Undulating terrace on a southeasterly slope of 5 percent. Elevation: 106 feet. Drainage: Well to moderately well drained; rapid and ponded runoff; moderate permeability. Soil Moisture: Slightly moist below surface.

Remarks: Roots matted horizontally at upper boundary of IIC1sim horizon. Top of duripan has 2 to 4 mm thick silica layer with dark reddish brown (5YR 3/3 moist) clay films overlying silica. Approximately 1.5 meters to right of described profile there is a clay layer with strong fine prismatic structure below loam material which abruptly overlies IIC1sim horizon. Silt layer below IIC3 horizon at 42 inches. Occasional smooth well rounded gravel in IIC1sim horizon. Temperature at 20 inches is 64°F.

* The Trigo series is not currently classified as having a duripan.

HORIZON	DESCRIPTION
Ap RSL 6963	(0-23 cm.) 0 to 9 inches, light yellowish brown (10YR 6/4) loam, brown and dark yellowish brown (7.5YR 4/4, 10YR 4/4) moist; common fine faint very pale brown (10YR 7/3) mottles; weak medium subangular blocky structure; slightly hard, friable, slightly sticky, plastic; common very fine and fine vertical roots; many very fine tubular and interstitial pores; medium acid (pH 6.0); clear smooth boundary.
B RSL 6964	(23-36 cm.) 9 to 14 inches, pale brown (10YR 6/3) loam, brown (7.5YR 4/4) moist; common fine distinct very pale brown (10YR 7/4) mottles; massive; slightly hard, friable, slightly sticky, plastic; few very fine vertical roots; many very fine and common fine tubular and interstitial pores; slightly acid (pH 6.5); abrupt smooth boundary.
IIC1sim RSL 6965	(36-51 cm.) 14 to 20 inches, very pale brown (10YR 8/3) silt loam, light olive brown (2.5Y 5/4) moist; very pale brown (10YR 7/4) stains; massive; extremely hard, very firm, brittle, slightly sticky, slightly plastic; very few very fine and fine tubular pores; very few thick clay films in root channels;moderately alkaline (pH 8.0); abrupt smooth boundary.
IIC2si RSL 6966	(51-79 cm.) 20 to 31 inches, light gray (2.5Y 7/2) sandy loam, yellowish brown (10YR 5/4) moist; massive; slightly hard, firm, nonsticky, nonplastic; few very fine and fine tubular pores; very few moderately thick clay films line pores and bridge mineral grains; mildly alkaline (pH 7.7); clear smooth boundary.
IIC3 RSL 6967	(79-105 cm.) 31 to 42 inches, light gray (2.5Y 7/2) loamy sand, dark brown (10YR 4/3) moist; massive; hard, very firm, nonsticky, nonplastic; abundant mica; mildly alkaline (pH 7.5); clear wavy boundary.

1/ The Trigo series is in a loamy, mixed, nonacid, thermic, shallow family of Typic Xoerorthents. This pedon has a duripan.

CONSECUTIVE PEDON NUMBER C 69 19 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES TRIGO HARDPAN VARIANT SOIL NO. S69 CALI. 31-11 SAMPLE NO. 9 91-9 95 PLACER COUNTY AREA

SOIL FAMILY TYPIC DUROCHREPT COARSE-LOAMY,MIXED,THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SANDS VCS 2-1	CS 1-.5	MS .5-.25	FS .25-.10	VFS .1-.05	SILTS CSI .05-.02	FSI .02-.002	CLAYS CO. 2-.2U	FINE LT .2U	CARBONATE 3A1A	INT. II .2-.02	1A2A COARSE FRAGMENTS PCT. OF WHOLE SOIL I 75-20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75		
9 91	0- 15	AP	49.3	38.4	12.3	2.9	6.5	6.4	22.0	11.5	18.9	19.5				42.9	--A --	--	--
9 92	15- 36	B	44.2	40.1	15.7	3.1	5.1	4.8	18.5	12.7	18.2	21.9				41.9	--A --	--	--
9 93	36- 392C	1	22.9	60.8	16.3	1.7	3.6	2.8	9.4	5.4	19.1	41.7				30.4	--A --	--	--
9 94	39- 512C	2SIM	33.1	52.4	14.5	2.9	5.8	5.3	11.0	8.1	15.4	37.0				29.3	--G --	--	--
9 95	0- 0	AP															--A --	--	--

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	19-2 6E1B	EXT. IRON FE 6C2B	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4AH	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTENSBLTY LEF 4D1	LE 4D1	PH H2O 1/1 8C1A	H2O 1/10 8C1A	CACL2 8C1E
		-PERCENT--		-PERCENT-		PCT.	PCT.NO.40		GRAMS PER CC				----PERCENT----			-PERCENT--				
9 91	0- 15	0.98	0.079							1.65	1.71	1.00		15.8	6.0	1.2	1.2	6.0		5.3
9 92	15- 36	0.36	0.039						1.60			1.00			6.3			6.1		5.4
9 93	36- 39	0.29							1.70			1.00			10.6			5.8		5.3
9 94	39- 51									1.83	1.92	1.00		18.5	11.1	1.6	1.6	6.6		6.0
9 95	0- 0	0.50	0.052						1.65			1.00			5.8					

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2O	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT. PCT.	C E C NA- OAC 5A2A	NH4- OAC 5A6A	EC 8A1A	WATER EXTRACT FROM SATURATED PASTE 8A1 CA 6N1B	MG 6O1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	B 6T1A	PH SAT. PASTE 8C1B
		---MEQ/100G---						-MEQ/100G		MMHO	---MEQ/1000 ML---								PPM	
9 91	0- 15	7.2	4.5	0.1	0.2	12.0	98		12.3											
9 92	15- 36	8.4	5.0	0.2	0.2	13.8	100+		13.6											
9 93	36- 39	14.0	5.9	0.3	0.2	20.4	91		22.4											
9 94	39- 51	13.3	6.8	0.4	0.2	20.7	97		21.3											
9 95	0- 0																			

SAMPLE NO.	DEPTH CM.	WATER AT SAT. 8A PCT.	EXCHANGEABLE NA 5B4B MEQ.	K 5D2 PCT.	SAR 5E	GYPSUM AMT. 6F1B PERCENT	REQ. 6P3A	CLAY MINERALOGY LESS THAN 0.002 MM X RAY I 7A2E	II 7A2E	III 7A2E	DTA IV 7A2E	PCT. 7A3	LT. .0002 MM X RAY I 7A2E	II 7A2E	III 7A2E	MINERAL CODE	RELATIVE AMOUNT
9 91	0- 15		0.1	1				MT 3	MI 3	KK 3						MT MONTMORT	1 TRACE
9 92	15- 36		0.2	1												MI MICA	2 SMALL
9 93	36- 39		0.3	1				MT 3	MI 3	KK 3						KK KAOLINIT	3 MODERATE
9 94	39- 51		0.4	2													4 ABUNDANT
9 95	0- 0																5 DOMINANT
																	6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
G=GROUND WHOLE. HORIZON INDURATED OR GRAVEL TOO SOFT TO BE SEPARATED. ALL DATA FOR WHOLE SOIL.

TRIGO LOAM, HARDPAN VARIANT* 1/
S69Calif-31-11

Location: Placer County, California. Approximately 3.5 miles southwest of Lincoln. 300 feet west and 70 feet north of SE corner of sec. 33, T.12N., R.6E. Date of Sampling: May 7, 1969.

Description by: L. A. Brittan, G. M. Kennedy. Collectors: K. W. Flach, W. D. Nettleton, G. M. Kennedy, T. M. Ryan, L. A. Brittan.

Classification: Typic Durochrept, coarse-loamy, mixed, thermic.

Vegetation: Rose clover, soft chess, ripgut brome, filaree. Climate: Average annual precipitation 20 inches. Mean annual temperature 61°F. Parent Material: Surficial sediments deposited over Laguna Formation (fine grained sandstone and siltstone). Topography: Undulating terrace on a southeast slope of 4 percent. Elevation: 120 feet. Drainage: Well to moderately well drained; medium runoff; moderate permeability. Soil Moisture: Essentially dry; below wilting point when sampled.

Remarks: Roots mat horizontally along upper boundary of IIC1 horizon. At approximately 21 inches there is clay banding in the IIC2sim horizon. There is an occasional gravel in the profile. The pores in the IIC1 horizon may be root channels. Top of IIC1 horizon has common thick dark reddish brown (5YR 3/3) moist clay films; also colloid stains inside root channels and pores. *The Trigo series is not currently classified as having a duripan.

HORIZON	DESCRIPTION
	(0-15 cm.)
Ap RSL 6991	0 to 6 inches, light brownish gray (2.5Y 6/2) and pale brown (10YR 6/3) loam, dark brown (10YR 4/3) moist; few fine distinct yellowish brown (10YR 5/6) mottles; massive; hard, friable, slightly sticky, slightly plastic; common very fine vertical roots; common very fine pores; medium acid (pH 6.0); abrupt smooth boundary.
	(15-36 cm.)
B RSL 6992	6 to 14 inches, pale brown (10YR 6/3) heavy loam, dark brown (10YR 3/3) moist; massive; hard, friable, slightly sticky, slightly plastic; common very fine vertical roots; common very fine and fine and few medium pores; slightly acid (pH 6.5); abrupt wavy boundary.
	(36-39 cm.)
IIC1 RSL 6993	14 to 15½ inches, light gray (2.5Y 7/2) light loam, light gray (2.5Y 7.2) moist; common fine distinct dark yellowish brown (10YR 4/4) and yellowish brown (10YR 5/8) mottles; massive; very hard, extremely firm, slightly sticky, slightly plastic; common very fine tubular pores; few moderately thick clay films line pores; slightly acid (pH 6.3); abrupt wavy boundary.
	(39-51 cm.)
IIC2sim RSL 6994	15½ to 20 inches, light gray (10YR 7/2) sandy loam, dark brown (10YR 4/3) moist; common fine distinct yellowish brown (10YR 5/6) mottles; massive; extremely hard, extremely firm, brittle, slightly sticky, slightly plastic; top of horizon has 1 to 2 mm thick silica layer; mildly alkaline (pH 7.6).

CONSECUTIVE PEDON NUMBER C 69 20 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES TRIGO HARDPAN VARIANT SOIL NO. S69 CALI. 31-12 SAMPLE NO. 9 96-9 97 PLACER COUNTY AREA

SOIL FAMILY TYPIC DUROCHREPT COARSE-LOAMY,MIXED,THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	CSI .05- .02	FSI .02- .002	CO. 2- .2U	FINE LT .2U	CARBONATE 3A1A.02	INT. II .2- OF 75- 20-5 5-2	COARSE FRAGMENTS- PCT. I	II	III
			SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1													1A2A			
			PERCENT OF LESS THAN 2MM													WHOLE 20 SOIL	PCT. OF LT 75		
9 96	0- 18	AP	40.5	46.1	13.4	1.9	4.1	6.9	16.2	11.4	17.1	29.0			37.4	--A	--	--	--
9 97	18- 30	B	41.2	45.2	13.6	0.9	3.5	6.8	16.4	13.6	15.4	29.8			39.0	--A	--	--	--

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A -PERCENT--	NITROGEN 6B1A -PERCENT--	CARBONATE AS CACO3 LT 2 6E1B -PERCENT-	19-2 6E1B -PERCENT-	EXT. IRON FE 6C2B PCT.	ATTERBERG LIMIT LL 4F2 PCT.	PL 4F1 PCT.	BULK DENSITY EST 1/3 4A1F NO.40 GRAMS PER CC	1/3 BAR 4A1F	DRY 4AH	CM 3B2	WATER CONTENT 1/10 BAR 4B1C -----PERCENT-----	1/3 BAR 4B1C	15 BAR 4B2	EXTENSIBILITY LEF 4D1 -PERCENT--	LE 4D1	PH H2O 1/1 8C1A	PH H2O 1/10 8C1A	PH CACL2 8C1E
9 96	0- 18	0.81	0.066						1.62			1.00			6.0			5.7		5.2
9 97	18- 30	0.39							1.60			1.00			5.8			5.7		5.1

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6Q2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT. PCT.	C E C NA- OAC 5A2A -MEQ/100G	NH4- OAC 5A6A	WATER EXTRACT FROM SATURATED PASTE EC 8A1A -MEQ/100G	CA 6N1B MMHO	MG 6O1B	NA 6P1A -MEQ/1000 ML	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	B 6T1A PPM	8A1 PH SAT. PASTE 8C1B
9 96	0- 18	7.5	3.4	0.1	0.2	11.2	85		13.2											
9 97	18- 30	7.5	3.0	0.1	0.1	10.7	87		12.3											

SAMPLE NO.	DEPTH CM.	WATER AT SAT. 8A PCT.	EXCHANGEABLE NA. 5B4B MEQ.	NA. 5D2 PCT.	SAR 5E	GYPSUM- ANT. REQ. 6F1B PERCENT	6P3A	CLAY MINERALOGY --LESS THAN 0.002 MM--- X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 PCT.	-L.T. .0002 MM- X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
9 96	0- 18		0.1	1				MI 4	KK 3	MT 2						MI MICA	1 TRACE
9 97	18- 30		0.1	1				MI 4	KK 3	MT 2						KK KAOLINIT	2 SMALL
																MT MONTMORT	3 MODERATE
																	4 ABUNDANT
																	5 DOMINANT
																	6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.

TRIGO LOAM, HARDPAN VARIANT* 1/
S69Calif-31-12

Location: Placer County, California. Approximately 4.7 miles south of Lincoln on Highway 65 and 150 feet east along embankment of PG&E substation, 750 feet west and 500 feet south of NE corner of sec. 9, T. 11 N., R.6E. Date of Sampling: May 7, 1969. Description by: T. M. Ryan, L. A. Brittan. Collectors: K. W. Flach, W. D. Nettleton, G. M. Kennedy, L. A. Brittan, T. M. Ryan.

[redacted]

Vegetation: Filaree, rose clover, soft chess, lupine, and purple vetch. Climate: Average annual precipitation 17 to 32 inches. Mean annual air temperature 61°F. Parent Material: Surficial sediments deposited over Laguna Formation. Topography: Undulating terrace on a south aspect of 2 to 9 percent slope. Elevation: 125 feet. Drainage: Well to moderately well drained, rapid and ponded runoff, moderate permeability. Soil Moisture: Almost dry in control section.

Remarks: Roots matted horizontally at top boundary of IIC1sim horizon. IIC1sim breaks out in plates and is interbedded with thick dark red (2.5YR 3/6) clay films. *The Trigo series is not currently classified as having a duripan.

HORIZON	DESCRIPTION
Ap RSL 6996	(0-18 cm.) 0 to 7 inches, light yellowish brown (10YR 6/4) and pale brown (10YR 6/3) loam, dark yellowish brown (10YR 3/4) moist; weak very fine and fine subangular blocky structure; slightly hard, friable, slightly sticky, slightly plastic; few very fine vertical roots; many very fine tubular and interstitial pores; neutral (pH 6.7); clear wavy boundary.
B RSL 6997	(18-30 cm.) 7 to 12 inches, brown (10YR 5/3) loam, dark brown (7.5YR 4/4) moist; massive; slightly hard, friable, slightly sticky, slightly plastic; common very fine vertical roots; many very fine tubular and interstitial pores; few thin clay films bridge mineral grains; slightly acid (pH 6.3); abrupt smooth boundary.
IIC1sim	(30-66 cm.) 12 to 26 inches, light gray (2.5Y 7/2) silt loam, light yelloiwsh brown (2.5Y 5/4) moist; massive; extremely hard, very firm, brittle, slightly sticky, slightly plastic; continuous thick clay films in bedding planes; neutral (pH 6.6); abrupt wavy boundary.
IIC2si	(66-155 cm. 26 to 62 inches, stratified sand and silt layers ranging in thickness from a few mm to several cm thick.

1/ The Trigo series is in a loamy, mixed, nonacid, thermic, shallow family of Typic Xerorthents. This pedon has a duripan.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Crouch stony fine sandy loam (taxadjunct) **LOCATION** San Diego County, California

SOIL FAMILY: Typic Xerochrept, coarse-loamy, mixed, mesic

SOIL NOS. S60Calif-37-6 **LAB. NOS.** 60425 - 60430

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 3	A11	8.2	14.3	10.6	26.8	14.5	19.5	6.1	42.6	8.2	17	sl
3- 9	A12	7.0	13.5	10.7	28.2	15.4	19.0	6.2	43.7	8.0	10	sl-fsl-ls
9-18	B1	7.6	14.7	11.7	29.6	14.4	15.5	6.5	41.3	6.8	3	ls-sl-fsl
18-30	B2	11.6	18.9	11.7	24.9	11.5	14.8	6.6	34.4	6.9	2	lcos
30-43	B3		not		det	ermi	ned					
43-49+	C1		not		det	ermi	ned					

pH 8C1a H_2O 1:1	8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	Atterberg limits Plastic limit	Liquid limit	Plastic index	Free Fe_2O_3 6C1b %	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
						Pct. of 0.4 mm.						
7.0	6.0		2.38	0.150	15.9				1.9			5.8
6.9	5.9		0.98	0.064	15.3				1.9			4.4
6.8	5.3		0.29	0.019		N.P.	-	N.P.	1.8			4.1
6.6	4.9		0.26	0.013					2.0			4.5
6.5	4.7		0.28	0.012					-			
not	det	ermi	ned									

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 6N2b Ca	6O2c Mg	6H2a H	6P2a Na	5B1a 6Q2a K	BASE SAT. % 5C3	Sum of Cations me./100 g. soil 5A3a	g. K				MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →							milliequivalents per liter →				
15.5	11.0	1.4	3.7	0.2	0.7	78	17.0					
10.5	7.2	1.2	2.6	0.1	0.3	77	11.4					
8.3	4.8	1.2	1.7	0.1	0.2	79	8.0					
9.5	5.6	1.5	1.9	0.1	0.2	80	9.3					
	not		det	ermi	ned							

1/ Percent of the whole soil

CROUCH STONY FINE SANDY LOAM (taxadjunct) 1/
S60Calif-37-6

Description: George Borst
Date of Sampling: November 15, 1960
Collectors: Reuben Nelson, Fred C. Eggers, Cliff Henry, James Lockard, and George Borst
Location: Approximately 200 feet west of California State Highway 78 [illegible]

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Hugo sandy loam variant **LOCATION** San Mateo County, California

SOIL FAMILY: Typic Xerochrept, coarse-loamy, mixed, mesic.

SOIL NOS. S57-Calif-41-1 **LAB. NOS.** 57447-450

PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1

HUGO SANDY LOAM variant 1/
S57Calif-41-1

Collectors: T. H. Arkley, R. Ulrich, and R. J. Wagner.
Date of sampling: May, 1957. **Description by:** Rudolph Ulrich.
Location: 1.8 miles north of Sky Londe along California Highway 5 (Skyline Blvd.)

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Xerochrept, coarse-loamy, mixed, mesic

SOIL Stoner loam SOIL Nos. S65Calif-27-19 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65552 - 65558

Depth (In.)	Horizon	1B1b Total Sand (2–0.05)	Total Silt (0.05–0.002)	Total Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	* Coarse fragments 1A2a > 2	2 - 19	19 - 76	
		Pct. of < 2 mm														Pct.	Pct. of < 76mm		
0-5	A11	48.9	35.1	16.0	8.5	10.6	6.7	13.1	10.0	13.8	21.3	31.3	38.9	0	16	5	3	2	**
5-13	A12	52.1	32.0	15.9	7.5	11.6	6.8	14.1	12.1	12.1	19.9	31.9	40.0	0	16	9	7	2	**
13-24	B1t	50.9	33.4	15.7	7.9	11.6	6.5	13.2	11.7	13.4	20.0	32.5	39.2	0	16	11	9	2	**
24-37	B2t	59.9	28.3	11.8	10.7	13.9	7.7	15.9	11.7	12.5	15.8	32.9	48.2	0	12	18	16	2	**
37-47	C1	57.2	33.9	8.9	6.3	12.0	7.9	18.2	12.8	15.0	18.9	38.1	44.4	0	9	9	7	2	**
47-54	C2	66.4	27.3	6.3	12.6	16.4	9.2	17.8	10.4	13.4	13.9	33.5	56.0	0	6	15	13	2	**
54-60+	C3	63.5	31.6	4.9	8.0	16.0	9.3	19.1	11.1	13.3	18.3	34.8	52.4	0	5	10	8	2	**

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLE in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1		
0-5	6.73	0.290	23	1.3	-	1.04	1.07		0.98		24.4	16.1	0.007	0.007	6.1		
5-13	1.19	0.070	17	1.4	-	1.26	1.30		0.96		22.7	9.5	0.011	0.011	5.1		
13-24	0.50			1.3	-	1.39	1.43		0.94		20.8	9.7	0.012	0.011	5.5		
24-37	0.21			0.8	-	1.47	1.53		0.89		21.3	8.4	0.013	0.011	5.7		
37-47	0.40			1.4	-	1.37	1.44		0.95		23.3	9.2	0.015	0.014	5.8		
47-54	0.34			1.5	-	1.54	1.60		0.91		17.2	7.5	0.012	0.011	5.9		
54-60+	0.25			1.2	-	1.45	1.50		0.94		19.7	5.5	0.010	0.009	6.0		

Depth (In.)	Extractable bases 5B1a 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A1a NH_4OAc	5A2a NaOAc		6G1d KCl Ext. Al^{+++}		Atterberg limits Plastic limit	Liquid limit	Plastic index			Base saturation 5C1 NH_4OAc	
	meq/100 g											Pct. of <0.4 mm					Pct.	Pct.
0-5	25.3	7.3	0.2	1.2	34.0	14.9		41.2									82	
5-13	5.9	3.3	0.2	0.3	9.7	14.1	18.5					28	34	6			52	
13-24	9.9	4.0	0.2	0.2	14.3	8.4	19.4										74	
24-37	8.5	4.5	0.4	0.2	13.6	5.1	16.2										84	
37-47	10.8	6.6	0.4	0.1	17.9	5.6	21.6										83	
47-54	10.9	5.9	0.4	0.1	17.3	5.1	20.8										83	
54-60+	9.0	4.1	0.6	0.1	13.8	3.5	15.9						-	N.P.			87	

STONER LOAM
S65Calif-27-19

Location: Monterey County, California, 650 feet NW of Bouchers Gap, 8.5 miles up Palo Colorado Canyon from Hwy #1, about 100 feet up slope from culvert on small drainage way. Date of Sampling: May 24, 1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich; site selection by T. D. Cook.
Classification: Typic Xerochrept, coarse-loamy, mixed, mesic. (Yellowish Brown Lateritic).
Vegetation: Dense canopy of hardwood trees, mostly tanbark oak. Climate: Precipitation is 40 inches. Mean annual temperature 56°F., mean January 48°F., mean July 60°F. Parent Material: Deeply weathered granite. Topography: Slightly convex steep SW slope of 45 percent, not subject to accretions from above or serious natural erosion losses. Elevation: 2050 feet. Drainage: Well drained, moderately to moderately slow permeability, medium runoff. Soil Moisture: Moisture near field capacity at time of sampling.
Remarks: Field pH determinations by Truog Kit. Asterisks indicate horizons sampled for California Division of Highways.

HORIZON	DESCRIPTION
O1 Not sampled	2½ to 2 inches, surface cover of leaves and twigs. (pH 7.0).
O2 Not sampled	2 to 0 inches, porous. Loose duff, very dark brown (10YR 2/2m). (pH 6.0).
A11 RSL 65552	0 to 5 inches, dark grayish brown (10YR 4/2) loam, dark brown (7.5YR 3/2) moist; moderate medium subangular blocky structure; slightly hard, friable, sticky and slightly plastic; plentiful fine and medium roots; many very fine interstitial and few fine tubular pores; strongly acid (pH 5.5); clear smooth boundary.
*A12 RSL 65553	5 to 13 inches, brown (7.5YR 5/4) loam, darker brown (7.5YR 4/4) moist; moderate coarse subangular blocky structure; slightly hard, friable, slightly sticky and slightly plastic; plentiful fine medium and coarse roots; many very fine interstitial and few fine tubular pores; strongly acid (pH 5.5); gradual smooth boundary.
B1t RSL 65554	13 to 24 inches, light yellowish brown (10YR 6/4) light clay loam, strong brown (7.5YR 5/6) with faint yellowish red mottles or veins (5YR 4/6) moist; weak medium and coarse subangular blocky structure; hard, friable, sticky and plastic; plentiful fine medium and coarse roots; many very fine interstitial and few fine tubular pores; medium acid (pH 6.0); gradual wavy boundary.
B2t RSL 65555	24 to 37 inches, light yellowish brown (10YR 6/4) sandy clay loam, yellowish brown (10YR 5/4) moist, with some strong brown (7.5YR 5/6) moist mixed colors; weak coarse subangular blocky structure; hard, friable, sticky and plastic; plentiful fine medium and coarse roots; many very fine interstitial pores; appears to have few thin clay films in bridges, clay not generally oriented; medium acid (pH 5.8); clear irregular boundary.
C1 RSL 65556	37 to 47 inches, light yellowish brown (10YR 6/4) fine sandy loam, yellowish brown (10YR 5/6) moist; massive; material shows definite rock structure but rock is quite rotten; hard, friable, slightly sticky and slightly plastic; few fine medium and coarse roots; many very fine interstitial pores; medium acid (pH 5.6); clear smooth boundary.
C2 RSL 65557	47 to 54 inches, light yellowish brown (10YR 6/4) fine sandy loam, yellowish red (5YR 5/6) moist; massive; clearly shows rock structure but rock is strongly weathered; hard, friable, slightly sticky and slightly plastic; few fine roots; common very fine interstitial pores; many thin clay films in bridges; slightly acid (pH 6.3); most of these profile differences due to differences in the parent material, not genetic; clear smooth boundary.
*C3 RSL 65558	54 to 60 inches, mixed light yellowish brown (10YR 6/4) and yellowish red (5YR 5/6) moist; mostly brown (10YR 5/3) dry, weathered granite showing rock structure; hard, friable; no roots; many very fine interstitial pores; slightly acid (pH 6.3).

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Xerochrept, coarse-loamy, mixed, mesic

SOIL Stoner gravelly sandy loam SOIL Nos. S66Calif-47-10 LOCATION Siskiyou County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66401 - 66408

Depth (In.)	Horizon	1B1b Total Sand (2–0.05)	Total Silt (0.05–0.002)	Total Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)		3A1a Non-carbonate clay	1/ Coarse fragments 1A2a > 2 Pct.	2–19 (Pct. of < 76mm)	19–76 (Pct. of < 76mm)
		Pct. of < 2 mm																
0-1	A11	65.8	24.8	9.4	28.9	17.0	6.4	8.5	5.0	8.9	15.9	17.9	60.8		9	39	38	1
1-3	A12	59.9	27.6	12.5	23.5	15.4	6.1	9.1	5.8	8.2	19.4	18.7	54.1		13	35	34	1
3-12	A13	56.7	30.3	13.0	15.8	15.9	7.1	10.5	7.4	13.4	16.9	26.2	49.3		13	26	26	-
12-21	B21t	58.0	28.3	13.7	13.5	15.6	7.8	12.4	8.7	11.1	17.2	26.4	49.3		14	27	27	-
21-27	B22t	59.1	28.2	12.7	15.1	15.8	7.6	12.2	8.4	11.3	16.9	26.1	50.7		13	27	27	-
27-36	B23t	59.9	27.9	12.2	16.6	15.6	7.5	11.9	8.3	10.5	17.4	25.5	51.6		12	29	28	1
36-60+	B24t	50.1	28.4	21.5	11.9	12.4	6.8	11.1	7.9	9.7	18.7	23.8	42.2		22	80	64	16
0-6	A1 2/															24	23	1

Depth (In.)	6A1a Organic carbon Pct	6B1a Nitrogen Pct	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 C'	Water content: Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1a H_2O 1:1		
0-1	2.30	0.147	16										6.5			6.4		
1-3	1.52	0.105	14										5.7			6.4		
3-12	0.86	0.088	10				1.54	1.57		0.83		16.8	6.5	0.007	0.005	6.0		
12-21	0.53						1.63	1.65		0.82		13.8	6.4	0.005	0.004	5.9		
21-27	0.28						1.71	1.73		0.81		13.3	6.3	0.005	0.004	5.9		
27-36	0.20						1.80	1.84		0.78		13.1	6.3	0.007	0.006	5.8		
36-60+	0.26												8.4			5.9		
0-6	1.52	0.118	13										6.4			6.5		

Depth (In.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1a NH_4OAc	5A3a Sum		6G1d KCl Ext. Al^{+++}						Base saturation: 5C1 NH_4OAc Pct	5C3 Sum Pct
	meq/100 g																
0-1	8.3	0.9	0.1	0.5	9.8	2.8	10.5	12.6								93	78
1-3	7.1	0.8	0.1	0.4	8.4	3.0	9.6	11.4								88	74
3-12	7.1	0.9	0.1	0.4	8.5	3.0	9.8	11.5								87	74
12-21	4.6	0.8	0.1	0.4	5.9	3.2	7.7	9.1								77	65
21-27	3.8	0.9	0.1	0.4	5.2	2.8	7.0	8.0								74	65
27-36	3.6	1.3	0.1	0.3	5.3	2.6	6.8	7.9								78	67
36-60+	4.9	1.9	0.1	0.4	7.3	2.6	9.0	9.9								81	74
0-6	8.3	0.8	0.1	0.6	9.8	3.4	11.0	13.2								89	74

Depth (In.)	Clay mineralogy (<0.002 mm) 7A: Mont.	Verm.	Mica	Kaolin	
	X-ray				

1/ From characterization sample (5 kg).

2/ Composite of 20 subsamples taken in a 15x12' grid adjacent to the master pedon.

STONER GRAVELLY SANDY LOAM
S66Calif-47-10

Location: Siskiyou County, California. A pit located on the Madeline Brown Ranch in McConnaughy Gulch in Scott Valley, $\frac{1}{4}$ mile north of the McConnaughy Gulch Road up Dockery Gulch. 1600 feet west and 2500 feet north of the SW corner of Sec. 5, T41N, R8W. Date of Sampling: May 26, 1966.
Description by: J. J. Newlun. Collectors: J. Newlun, D. Lucas, and L. Mann.
Classification: Typic Xerochrept, coarse-loamy, mixed, mesic.
Vegetation: In the order of dominance: Buckbrush (Ceanothus cuneatus); Juniper, white oak, annual bromes and other grasses and forbs, an occasional ponderosa pine; some oak had been cleared from the site. The oak canopy is approximately 25 percent at the present time. Climate: The average annual precipitation is approximately 16 inches with much of it in the form of snow. The average January temperature is 32°F., the average July temperature 71°F., and a mean annual temperature of 51°F. Parent Material: The soil has formed in gravelly alluvium from mixed sources, dominantly metamorphosed sedimentary and igneous rocks and some ultra-mafic rocks. The alluvium was deposited in fan, bottom and terrace positions along drainageways from the adjacent mountains. The soil is commonly somewhat stratified at depths greater than 36 inches. It contains 20 to 60 percent cobbles, stones and gravels but most of the material $>$ 2 mm is fine to medium sized gravels. Topography: The site is located about midslope on a single convex slope of a terrace which has a gradient of 5 percent and a south aspect. Elevation: 3160 feet. Drainage: Well drained, moderate permeability, medium runoff. Soil Moisture: Air dry throughout profile.
Remarks: Reactions determined by Truog soil reaction test. Profile S66Calif-47-9 is the

tation zone in which these soils occur in this soil survey area. The soil has an ochric epipedon and a cambic horizon. Most of the Stoner soils are cultivated but this profile and the paired profile have not been disturbed as far as could be ascertained.

HORIZON	DESCRIPTION
A11 RSL 66401	0 to 1 inch, brown (10YR 5/3) gravelly sandy loam, dark brown (10YR 3/3) moist and rubbed with dark yellowish brown (10YR 3/4) moist ped faces; weak fine granular structure; soft, very friable, slightly sticky, slightly plastic; many fine and very fine roots; strongly acid (pH 5.5); 30 percent fine and medium gravels (included in sample); abrupt smooth boundary.
A12 RSL 66402	1 to 3 inches, brown (10YR 5/3) gravelly sandy loam, dark yellowish brown (10YR 3/4) moist; massive; slightly hard, friable, slightly sticky, slightly plastic; many fine and very fine roots; common fine and few medium tubular pores; medium acid (pH 5.8); 25 percent fine and medium gravels (included in sample); abrupt smooth boundary.
A13 RSL	3 to 12 inches, brown (10YR 5/3) gravelly sandy loam, dark yellowish brown (10YR 3/4) moist; weak medium subangular blocky structure; slightly hard, friable

OIL SURVEY LABORATORY Riverside, California

OIL TYPE Vista sandy loam **LOCATION** Riverside County, California

SOIL FAMILY: Typic Xerochrept, coarse-loamy, mixed, thermic

SOIL NOS. S55Calif-33-6 **LAB. NOS.** 55554-559

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-1	A11	11.4	15.4	10.6	21.4	13.9	22.0	5.3	42.0	6.3	11	cosl
1-9	A12	10.9	17.4	11.1	21.1	12.6	20.9	6.0	40.0	5.9	9	cosl
9-15	A	11.1	16.3	10.8	20.2	12.3	21.7	7.6	38.3	7.5	12	cosl
15-20	A3	11.1	16.9	10.4	19.5	12.1	19.6	10.4	36.9	6.0	13	cosl
20-32	B2	11.3	15.0	10.4	19.7	11.1	21.3	11.2	34.6	8.9	20	cosl
32-43	Cir	not analysed										

pH 8C1b SATURATED	8C1a	8C1a	ORGANIC MATTER 6A1a ORGANIC	6B1a NITRO-		EST% SALT (BUREAU	8A1a Elec. Condu.c. EC x 10³	6E1b $CaCO_3$ equiv.	6F1a GYPSUM me./100g.	MOISTURE TENSIONS 4B1a 1/10	4B1a 1/3	4B2 15

VISTA SANDY LOAM
S55Calif-33-6

Collectors: Fred Eggers, Arnold Knecht, Floyd G. Stephens, W. G. Harper, R. E. Nelson, F. F. Harradine. Sampling date: December 5, 1955.

Description by: W. G. Harper

Location: 700' S of NE corner of Sec. 34, T5S, R1W. On the McSweeney Ranch, about 4 miles S of Hemet, Riverside County, California.

Elevation: 1650 feet

Precipitation: 14 inches

Classification: Typic Xerochrept, coarse-loamy, mixed, thermic.

California Group: VII

Vegetation: Wild buckwheat, California sage, wild oats, cheat grass and some giant wild rye.

Parent Material: Granodiorite.

Physiography: Rolling upland with many outcrops of granite 3 to 25 feet across and 1 to 6 feet above surface of ground.

Drainage: Well drained.

HORIZON		DESCRIPTION
A_{11}	0- 1"	Grayish brown (10YR 5/2 dry, very dark grayish brown 3/2 moist) sandy loam; very weakly defined medium to fine crumb; loose, very friable; moderately rapid permeability; many plant roots; pH 7.2; noncalcareous; abrupt smooth boundary.
A_{12}	1- 9"	Brown (10YR 5/3 dry, dark brown 3/3 moist) sandy loam; very weakly defined medium to fine crumb; slightly hard, friable; moderately rapid permeability; moderate number of plant roots; pH 6.5; abrupt smooth boundary.
A	9-15"	Similar to horizon above in color, texture and structure; slightly hard, friable; moderately rapid permeability; few plant roots; noncalcareous; pH 6.0; abrupt smooth boundary.
A_3	15-20"	Brown (10YR 5/3 dry, dark brown 3/3 moist) sandy loam; moderately defined medium to fine subangular blocky; slightly hard, friable, nonsticky; moderately permeable; few roots; pH 6.0; abrupt smooth boundary.
B2	20-32"	Yellowish brown (10YR 4.5/4 dry, dark yellowish brown 3.5/4 moist) gritty loam; massive; slightly hard, friable, slightly sticky; moderately permeable; very few roots; pH 6.8; clear, irregular boundary.
CDr	32-43"	Light yellowish brown and yellowish brown (10YR 6/4 to 5/4 dry, dominantly brown 4/3 moist) decomposing granodiorite; massive; very hard, firm; slowly permeable; noncalcareous; pH 6.8

Remarks: The entire profile is highly micaceous.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Vista sandy loam **LOCATION** Riverside County, California

SOIL FAMILY: Typic Xerochrept, coarse-loamy, mixed, thermic

SOIL NOS. S55Calif-33-7 **LAB. NOS.** 55560-564

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-1	A11	14.6	18.4	10.5	20.4	10.7	20.5	4.9	36.0	7.0	12	cosl
1-9	A12	15.7	19.9	11.0	20.1	9.9	17.5	5.9	33.2	5.9	10	lcos
9-15	A	18.5	21.0	10.2	18.8	8.1	16.2	7.2	28.7	6.4	16	cosl
15-24	B2	16.6	21.1	9.5	19.8	9.4	14.5	9.1	30.0	5.3	26	cosl
24-30	CDr			not analysed								

8C1b pH SATURATED PASTE 1:1	8C1a 1:5	8C1d 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B1a 1/10 ATMOS. %	4B1a 1/3 ATMOS. %	4B2 15 ATMOS. %
6.4		6.4	1.09	0.074	14.7		0.37			22.0	13.3	3.5
6.2		6.3	0.65	0.050	13.0		0.34			19.4	11.7	3.4
6.7		6.2	0.20				0.15			16.1	9.9	3.8
6.9		6.8	0.14				0.22			16.2	9.4	4.0

5A2a CATION EXCHANGE CAPACITY ←(Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	5B Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg		6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
10.3	6.1	0.9	2.3	0.2	0.4	74	0.7	0.5	2.7			31.5
9.3	5.3	1.0	2.3	0.1	0.3	74	0.5	0.3	2.4			27.8
7.5	4.3	1.2	1.0	0.2	0.3	86	0.3	0.2	0.8			24.7
8.5	4.9	1.5	0.8	0.1	0.2	89	0.5	0.1	1.2			24.9

1/- Percent of the whole soil

VISTA SANDY LOAM
S55Calif-33-7

Collectors: Fred Eggers, Arnold Knecht, F. G. Stephens, W. G. Harper, R. E. Nelson, F. F. Harradine. Sampling date: December 5, 1955

Description by: W. G. Harper

Location: On E side of Hwy. 79 in NW 1/4, Sec. 18, T7S, R1E, about 12 miles south of Hemet, Riverside County, California.

Elevation: 2550 feet

Precipitation: 16 inches

Classification: Typic Xerochrept, coarse-loamy, mixed, thermic.

California Group: VII

Vegetation: Chamise 6-10' high, other shrubs, much cheat grass and alfileria.

Parent Material: Granodiorite

Physiography: Rolling upland with numerous rock outcrops from 3 to 20 feet in diameter and up to 14 feet high. Sample taken near top of ridge which is about 200 feet wide.

Drainage: Well drained

HORIZON		DESCRIPTION
A11	0- 1"	Brown (10YR 5/3 dry, very dark brown 2.5/2 moist) sandy loam; very weakly defined medium to fine crumb; slightly hard, very friable; moderately permeable; many plant roots; high in organic matter; pH 6.0; abrupt smooth boundary.
A12	1- 9"	Brown (10YR 5/3 dry, very dark grayish brown 3/2 moist) fine sandy loam or sandy loam; moderately defined medium to fine crumb; hard, very friable; moderately permeable; many roots; high in organic matter; pH 6.2; clear smooth boundary.
A	9-15"	Grayish brown (10YR 5/2 dry, very dark grayish brown 3/2 moist) gritty sandy loam; massive; slightly hard, very friable; moderately permeable; moderate number of plant roots; pH 6.5; clear smooth boundary.
B2	15-24"	Brown (10YR 5/3 dry, dark grayish brown 3.5/2 moist) gritty sandy loam; weakly defined, medium, subangular blocky; very hard, friable; moderately permeable; moderate number of plant roots; pH 6.5; clear smooth boundary.
CDr	24-30"+	Yellow, white and black feldspars, quartz and biotite of decomposing granodiorite; massive; extremely hard, firm to friable;

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Xerochrept, coarse-loamy, mixed, thermic

SOIL: Vista coarse sandy loam — SOIL Nos. S64Calif-37-1 — LOCATION: San Diego County, California

SOIL SURVEY LABORATORY: Riverside, California — LAB. Nos. 641 - 647

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	* Clay <.002	** Clay <.002	1/ Coarse fragments 1A2a >2	2 - 19	19 - 76
		Size class and particle diameter (mm) 3A1 — Pct of < 2 mm														Pct	Pct. of < 76 mm	
0- 3	A11	72.7	17.1	10.2	19.5	21.2	9.0	14.8	8.2	8.4	8.7	24.2	64.5	11.5	9.3	19	19	0
3- 9	A12	70.9	18.5	10.6	17.8	18.0	9.0	17.2	8.9	9.0	9.5	26.8	62.0	10.9	8.5	22	22	0
9-19	A13	70.3	18.2	11.5	18.2	18.4	8.9	16.0	8.8	8.4	9.8	25.8	61.5	12.2	9.3	23	23	0
19-28	B21	68.7	19.4	11.9	16.2	17.0	8.6	16.3	10.6	9.5	9.9	28.2	58.1	12.5	9.5	24	24	0
28-35	B22	70.6	18.5	10.9	19.5	18.0	8.5	16.1	8.5	8.7	9.8	26.0	62.1	12.4	8.9	28	28	0
35-44	C1	82.3	12.5	5.2	35.3	23.2	6.9	11.8	5.1	6.0	6.5	17.0	77.2	4.1	2.4	41	41	0
44-61+	C2	82.9	12.5	4.6	32.5	23.9	7.8	12.9	5.8	6.0	6.5	18.1	77.1	4.5	2.7	40	40	0

Depth (In.)	* 6A1a Organic carbon Pct.	*6B1a Nitrogen Pct	C/N	* 6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 C'	Water content Pct.	4B1c 1/3 bar Pct.	*4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0- 3	0.96	0.080	12	1.0			1.55	1.89		0.88		8.7	5.4	0.066	0.059	6.7	6.9	
3- 9	0.56	0.055	10	0.9			1.51	1.86		0.86		9.8	4.0	0.072	0.061	6.5	6.6	
9-19	0.34			1.0			1.46	1.51		0.86		12.0	3.1	0.010	0.010	6.6	6.8	
19-28	0.28			0.9			1.43	1.52		0.85		10.1	3.7	0.021	0.017	6.6	6.7	
28-35	0.10			1.1			1.46	1.52		0.82		10.9	6.4	0.014	0.011	6.7	6.9	
35-44	0.03			0.7			1.92	1.97		0.67		6.7	4.6	0.008	0.006	7.2	7.1	
44-61+	0.03			0.8									4.2			7.3	7.4	

Depth (In.)	Extractable bases 6N2a * Ca	6O2a* Mg	6P2a* Na	5B1a 6Q2a* K	Sum of bases	6H2a* Ext. Acidity	Cation Exch. Capacity 5A2a* NaOAc	5A3a* Sum	Water extract from saturated paste 8A1 6N1a Ca	6O1a Mg	6P1a ** Na	6Q1a ** K	*	6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0- 3	10.5	3.4	1.4	0.2		1.7	16.9	17.2										
3- 9	10.5	2.8	0.4	0.1		2.3	18.6	16.1										
9-19	13.0	4.0	0.4	0.1		2.2	19.8	19.7										
19-28	11.1	4.0	0.4	0.1		1.8	19.1	17.4										
28-35	13.5	5.1	0.4	0.1		2.1	22.8	21.2										
35-44	9.1	3.7	0.5	tr.		1.2	14.9	14.5			0.7	tr.						0.08
44-61+	9.0	3.7	0.7	0.2		0.8	16.2	14.4										

Depth (In.)	* 8A Water at Saturation Pct.	5D2 Exchangeable Na Pct.	5B1b Exch. Na meq / 100g	Base Sat. Cations Pct.	Base Sat. NaOAc CEC Pct.	Clay mineralogy (<0.002 mm) 7A: Relative abundance based on height of diagnostic X-ray peak 7A2 Mont.	Verm.	Hydrobiotite	Illite	Kaolin	% Kaolin DTA 7A3	% Kaolin in ground whole soil (DTA)
						X-ray						
0- 3		8		90	92	-	-	t	xxxx	xx	7	-
3- 9		2		86	74	-	xxx	x	-	xx		
9-19		2		89	89	-	xxx	x	-	xx		
19-28		2		89	81	-	xxx	x	-	xx		
28-35		2		90	84	-	xx	x	t	xxx	12	2
35-44	44.4	3		92	90	-	xxxx	t	-	xx	6	-

1/ From characterization sample; determined by shaking overnight in a sodium hexametaphosphate solution.

* Analysis of ground whole soil; results expressed on <2-mm basis.

** Analysis of ground whole soil; results expressed on whole-soil basis.

\- = looked for but not found

t = trace

VISTA SANDY LOAM
S64Calif-37-1

Location: San Diego County, California. A soil profile pit located 2-1/4 miles south and 3/4 mile east of Fallbrook in the NE1/4 of the SW1/4 of section 31, T. 9 S., R. 3 W.
Date of Sampling: February 17, 1964.
Description By: David C. Estrada and Gerald Kester. Collectors: Klaus Flach, Gerald Kester, George Borst, Clifford Henry, Roy Bowman and Gerald Anderson.
Classification: Typic Xerochrept, coarse-loamy, mixed, thermic.
Vegetation: Annual weeds and grasses. Climate: The annual precipitation is 15 inches with a mean annual temperature of 60° F., a mean January temperature of 54° F., and a mean July temperature of 70° F. Parent Material: Tonalite grus. Topography: Southwest-facing, smooth 8 percent slopes in hilltop and upper backslope positions on rolling upland. Elevation: 590 feet above sea level. Drainage: Well drained; medium runoff; moderately rapid permeability. Soil Moisture: This profile was moist to about 15 inches.
Remarks: There is much rodent activity by ground squirrels. The irregular boundary between the B22 and C1 horizons may be the result of this activity. In a nearby pit burrow holes occur in the C1 horizon. Isolated bodies of C horizon occur within, or surrounded by, material of the B horizon.

HORIZON	DESCRIPTION
A11 RSL No. 641	0 to 3 inches, dark grayish brown (10YR 4/2) coarse sandy loam, dark brown (10YR 3/3) moist; moderate, fine to medium crumb structure; soft, very friable, nonsticky and nonplastic; plentiful very fine and fine roots; many very fine and fine random pores; neutral (pH 6.7); abrupt smooth boundary.
A12 RSL No. 642	3 to 9 inches, dark brown (10YR 4/3) coarse sandy loam, dark brown (10YR 3/3) moist; weak fine to medium granular structure; slightly hard, very friable, nonsticky and nonplastic; plentiful very fine and fine roots; common very fine and fine tubular and interstitial pores; neutral (pH 6.7); diffuse to clear irregular boundary. Krotovinas filled with this material, 3 to 4 inches wide, extend into the underlying horizons, in places extending into the upper part of the C horizon.
A13 RSL No. 643	9 to 19 inches, dark brown (10YR 4/3) coarse sandy loam, dark brown (10YR 3/3) moist; weak fine to medium granular structure; slightly hard, very friable, nonsticky and nonplastic; few very fine and fine roots; common very fine and fine tubular and interstitial pores; slightly acid (pH 6.5); clear wavy boundary.
B21 * RSL No. 644	19 to 28 inches, dark brown (10YR 4/3) coarse sandy loam, dark brown (10YR 3/3) moist; massive; hard, friable, nonsticky and nonplastic; few very fine and fine roots; common very fine and fine tubular and interstitial pores; slightly acid (pH 6.3); clear smooth boundary.
B22 RSL No. 645	28 to 35 inches, yellowish brown (10YR 5/4) coarse sandy loam, dark yellowish brown (10YR 3/4) moist; massive; hard, friable, nonsticky and nonplastic; very few very fine and fine roots; common very fine and fine interstitial, few fine tubular pores; slightly acid (pH 6.3); abrupt irregular to broken boundary.
C1 RSL No. 646	35 to 44 inches, yellowish brown (10YR 5/4) and very pale brown (10YR 7/4) tonalite grus composed mostly of quartz, feldspar, hornblende, and accessory dark-colored minerals; clear irregular boundary.
C2 * RSL No. 647	44 to 61 inches +, brown (10YR 5/3) and very pale brown (10YR 7/3) tonalite grus composed mostly of quartz, feldspar, hornblende, and other dark-colored minerals, mostly unweathered.

*Soil horizons sampled for BPR

U S DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Xerochrept, coarse-loamy, mixed, thermic.

SOIL Vista sandy loam SOIL Nos. S64Calif-37-4 LOCATION San Diego County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 648 - 6413

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (<0.002)	Sand Very coarse (2-1)	Sand Coarse (1-0.5)	Sand Medium (0.5-0.25)	Sand Fine (0.25-0.1)	Sand Very fine (0.1-0.05)	Silt 0.05-0.02	Silt Int. III (0.02-0.002)	Int II (0.2-0.02)	(2-0.1)	* Clay <.002	** Clay <.002	1/ Coarse fragments 1A2a >2	2 - 19	19 - 76
		Size class and particle diameter (mm) 3A1															Pct of <76 mm	
		Pct. of <2 mm														Pct		
0- 8	Ap	64.0	22.3	13.7	13.2	15.3	9.1	17.0	9.4	9.6	12.7	28.2	54.6	14.3	12.2	15	15	0
8-15	B12	64.0	22.7	13.3	13.3	15.5	8.7	16.8	9.7	10.6	12.1	29.4	54.3	14.6	12.9	12	12	0
15-24	B21	62.8	23.8	13.4	12.8	13.2	8.3	17.9	10.6	11.1	12.7	31.7	52.2	15.1	12.5	18	18	0
24-36	B22	64.8	22.0	13.2	13.1	15.6	8.8	17.7	9.6	9.8	12.2	28.4	55.2	13.9	11.7	16	16	0
36-47	C1	77.3	16.3	6.4	24.0	20.1	7.5	17.1	8.6	6.9	9.4	24.7	68.7	5.1	3.3	35	35	0
47-56+	C2	70.8	22.3	6.9	21.6	17.6	7.3	15.4	8.9	8.3	14.0	25.6	61.9	6.4	3.5	45	45	0

Depth (In.)	* 6A1a Organic carbon Pct.	6B1a * Nitrogen Pct.	C/N	* 6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	Bulk density 4A1h Oven dry g/cc	g/cc	3B2 C'	Water content Pct	Water content 4B1c 1/3 bar Pct.	Water content 4B2* 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1b Saturated Paste	pH 8C1a H_2O 1:1	pH 8C1a H_2O 1:10
0- 8	0.53	0.064	8	0.9									6.1			6.4	6.7	
8-15	0.41	0.038	11	0.9									6.5			6.8	7.0	
15-24	0.29			0.9									6.8			6.9	7.2	
24-36	0.23			1.0									6.5			6.9	7.2	
36-47	0.05			0.8									4.8			7.1	7.2	
47-56+	0.07			1.1									4.4			7.0	7.2	

Depth (In.)	Extractable bases 6N2a * Ca	6O2a * Mg	6P2a * Na	5B1a 6Q2a * K	Sum * of bases	6H2a * Ext. Acidity	Cation Exch. Capacity 5A2a * NaOAc	5A3a * Sum	Water extract from saturated paste 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0- 8	11.9	3.0	0.8	0.2	15.9	2.0	17.9	17.9										
8-15	12.2	3.0	0.8	0.1	16.1	1.7	18.0	17.8										
15-24	13.3	3.5	1.0	0.1	17.9	1.6	20.3	19.5										
24-36	13.0	4.0	1.0	0.1	18.1	1.5	20.8	19.6										
36-47	10.3	4.2	0.9	0.2	15.6	0.9	18.0	16.5										
47-56+	12.2	4.4	1.5	0.2	18.3	1.1	24.6	19.4										

Depth (In.)	8A Water at Saturation Pct.	5D2 Exchangeable Na Pct.	5B1b Exch. Na meq / 100g	Base Sat. Cations Pct.	Base Sat. NaOAc CEC Pct.	Clay mineralogy (<0.002 mm) 7A Mont.	Verm.	Mica	Kaolin.
						X-ray			
0- 8		5		89	89				
8-15		4		90	89				
15-24		5		92	89				
24-36		5		92	87				
36-47		5		94	86				
47-56+		6		94	74				

1/ From characterization sample; determined by shaking overnight in a sodium hexametaphosphate solution.

* Analysis of ground whole soil; results expressed on <2-mm basis.

** Analysis of ground whole soil; results expressed on whole-soil basis.

VISTA SANDY LOAM
S64Calif-37-4

Location: San Diego County, California. A soil profile pit located 1.5 miles south and 1.5 miles east of Fallbrook in the SE1/4 of the NE1/4 of section 30, T. 9 S., R. 3 W.
Date of Sampling: February 19, 1964.
Description By: David C. Estrada and Gerald Kester. Collectors: Klaus Flach, Gerald Kester.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Xerochrept, coarse-loamy, mixed, thermic

SOIL Vista aspect study SOIL Nos. S65Calif-27-7 S65Calif-27-29 S65Calif-27-30 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65599 - 65601

		1Blb	Size class and particle diameter (mm) 3A1

VISTA SANDY LOAM
S65Calif-27-7

Location: Monterey County, California, about 10 miles NE of King City on Montgomery Ranch, 300 feet due west from Pinalito VABM, about 75 feet down from ridge top. (Site #2 South Slope). Date of Sampling: May 18, 1965.

Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, and T. D. Cook.

Classification: Typic Xerochrept, coarse-loamy, mixed, thermic. (minimal) Noncalcic Brown.

Vegetation: Chamise, wild oats, red brome, many small weeds. Climate: Precipitation is 14 inches; mean annual temperature 60°F., January mean 58°F., July mean is 67°F. Parent Material: Diorite. Topography: Steep slopes, locally 47 percent, slightly convex, southern exposure, about 125 feet below ridge crest. Elevation: 1500 feet. Drainage: Somewhat excessively drained, moderately rapid permeability, medium runoff. Soil Moisture: Dry.

[redacted]

posite sample included individual samples (borings) from a plot with 4 rows, 12 feet apart. Each row was sampled 6 times at 6-foot intervals. Surface has weak fine pavement of fine gravels. The N pair at this location is S65Calif-27-6-1. Other units of this study are samples #65Calif-27-29-1 on South slope and S65-Calif-27-30-1 on North exposure. Twenty-four samples were taken for the composite.

HORIZON	DESCRIPTION
A1 RSL 65599	0 to 12 inches, pale brown (10YR 6/3) fine gravelly coarse sandy loam, dark brown (10YR 4/3) moist; massive; slightly hard, very friable, nonsticky and nonplastic; plentiful very fine roots; many very fine interstitial and common very fine tubular pores; medium acid (pH 6.0) 0 to 3

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Xerochrept, coarse-loamy, mixed, S65Calif-27-7

SOIL Vista aspect study thermic SOIL Nos. S65Calif-27-29 S65Calif-27-30 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65599 - 65601

Lab. No.	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)			* Coarse fragments 1A2a > 2 Pct.	2-19	19-76
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm															Pct. of < 76mm	
65599	A1***															28	26	2 **
65600	A1***															23	23	0 **
65601	A1***															22	22	0 **

Lab. No.	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc		Water content: Pct	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1a H_2O 1:1		
65599	1.35	0.075	18															
65600	1.00	0.076	13															
65601	2.48	0.120	21															

Depth (In.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1a NH_4OAc	5A3a Sum		6G1d KCl Ext. Al^{+++}						Base saturation	
	meq/100 g															Pct.	Pct.

Depth (In.)	Clay mineralogy (<0.002 mm) 7A: Mont.	Verm.	Mica	Kaolin	
	X-ray				

* From characterization sample (5Kg) (except as noted).
** 2-19 mm. from characterization sample, 19-76 mm. from field volume estimate.
*** Composite A horizon sample.

VISTA SANDY LOAM
S65Calif-27-29

Location: Monterey County, California. 350 feet west of center SW¼ SW¼, Sec. 2, T17S, R6E, 4.5 miles up Bryant Canyon Road from Metz Road out of Soledad. (Site #1, south slope). Date of Sampling: May 27, 1965.

Description by: T. D. Cook and W. D. Nettleton. Collectors: W. D. Nettleton and T. D. Cook.

Classification: Typic Xerochrept, coarse-loamy, mixed, thermic. (minimal) Noncalcic Brown.

Vegetation: Red brome, wild oats, few chamise plants, some forbs. Climate:

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Xerochrept, coarse-loamy, mixed, thermic S65Calif-27-7

SOIL Vista aspect study SOIL Nos. S65Calif-27-29 S65Calif-27-30 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65599 - 65601

Lab. No.	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)			* Coarse fragments 1A2a > 2 Pct.	2-19	19-76	
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm															Pct. of < 76mm		
65599	A1***															28	26	2	**
65600	A1***															23	23	0	**
65601	A1***															22	22	0	**

Lab. No.	6A1a Organic carbon Pct.	6B1a Nitrogen Pct	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	* g/cc		Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1		
65599	1.35	0.075	18															
65600	1.00	0.076	13															
65601	2.48	0.120	21															

Depth (in.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1a NH_4OAc	5A3a Sum		6G1d KCl Ext. Al^{+++}						Base saturation Pct.	Pct.
	meq/100 g																

Depth | Clay mineralogy (<0.002 mm) 7A

* From characterization sample (5Kg) (except as noted).
** 2-19 mm. from characterization sample, 19-76 mm. from field

VISTA SANDY LOAM
S65Calif-27-30

Location: Monterey County, California. 600 feet SW from center of SW¼ Sec. 2, T17S, R6E, 4.5 miles up Bryant Canyon Road from Metz Road out of Soledad. (Site #2 North Slope). Date of Sampling: May 27, 1965.

Description by: T. D. Cook and W. D. Nettleton. Collectors: W. D. Nettleton and T. D. Cook.

Classification: Typic Xerochrept, coarse-loamy, mixed, thermic. (minimal) Noncalcic Brown.

Vegetation: Thick stand of chamise, a little coffee berry and understory of soft chess, ripgut and red brome. Climate: Precipitation is 10 inches. Mean annual temperature is 65°F., mean January 55°F., and mean July 70°F. Parent Material: Slightly weathered granodiorite. Topography: Smooth even hill slope of 25 percent on north-facing slope, about 300 feet below ridge crest. Elevation: 1550 feet. Drainage: Well drained, moderately rapid permeability, medium runoff. Soil Moisture: 0 to 15 inches dry.

Remarks: For this study the Vista north slopes are assumed to be mollic, though the epipedon at this site may be too thin. The related north Vista sample is S65Calif-27-6, the south slope about 400 feet away is S65Calif-27-29 and the more distant south slope is S65Calif-27-7. Composite sample for a few determinations obtained by auger from A1 to a depth of about 8 inches. Composite sample included individual samples (borings) from a plot with 4 rows, 12 feet apart. Each row was sampled 6 times at 6-foot intervals.

HORIZON	DESCRIPTION
A1 RSL 65601	0 to 5 inches, dark grayish brown (10YR 4/2) fine gravelly sandy loam, very dark grayish brown (10YR 3/2) moist; moderate fine granular structure; slightly hard, very friable, nonsticky and nonplastic; abundant very fine roots nearly form a mat; many very fine interstitial pores; medium acid (pH 5.8); 25 percent fine gravel; clear wavy boundary.
B1	5 to 15 inches, brown (7.5YR 5/3) gravelly light loam, reddish brown (5YR 4/4) moist; moderate medium subangular blocky structure; slightly hard, very friable, slightly sticky and slightly plastic; many very fine interstitial and common very fine tubular pores; common thin clay films in bridges and pores; slightly acid (pH 6.3).

CONSECUTIVE PEDON NUMBER C 69 28 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES CIENEBA TAXADJUNCT SOIL NO. S69 CALI. 31-23 SAMPLE NO. 9 137-9 141 PLACER COUNTY AREA

SOIL FAMILY TYPIC XEROCHREPT COARSE-LOAMY,MIXED,THERMIC,SHW RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1-.05	MS .5-.25	FS .25-.10	VFS .1-.05	CSI .05-.02	FSI .02-.002	SILTS	CLAYS CO. 2-.2U	FINE LT .2U	CARBONATE 3A1A	INT. II .2-.02	1A2A COARSE FRAGMENTS PCT. OF WHOLE SOIL I 75-20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM															PCT. OF LT 75		
9 137	0- 13	A11	74.9	15.9	9.2	24.2	19.3	10.1	14.7	6.6	1.4	14.5					15.1	19W	--	--	19
9 138	13- 33	A12	68.7	22.1	9.2	18.4	18.4	10.2	14.4	7.3	8.5	13.6					22.9	12W	--	--	12
9 139	33- 51	2B2	65.5	22.8	11.7	21.3	16.1	8.6	13.2	6.3	8.1	14.7					21.0	18W	--	--	18
9 140	51- 94	3C 1	66.4	16.7	16.9	20.9	17.9	8.7	13.3	5.6	6.2	10.5					18.5	24W	--	--	24
9 141	94-130	3C 2	81.3	9.4	9.3	32.6	23.9	9.6	11.7	3.5	3.5	5.9				--	12.2	32W	--	--	32

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	19-2 6E1B	EXT. IRON FE 6C2B	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4AH	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTENSBLTY LEF 4D1	LE 4D1	PH H2O 1/1 8C1A	H2O 1/10 8C1A	CACL2 8C1E
		PERCENT		PERCENT		PCT.	PCT.	NO.40	GRAMS PER CC				PERCENT			PERCENT				
9 137	0- 13	1.35	0.098							1.60	1.62	0.88		10.3	3.9	0.4	0.4	5.6		4.9
9 138	13- 33	0.40	0.041						1.65			0.92			4.1			5.5		4.9
9 139	33- 51	0.14								1.74	1.76	0.87		9.2	4.4	0.4	0.3	5.5		4.9
9 140	51- 94								1.82			0.82			7.9			5.7		5.2
9 141	94-130									1.82	1.88	0.76		11.2	4.7	1.1	0.8	5.9		5.4

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT.	C E C NA-OAC 5A2A	NH4-OAC 5A6A	EC 8A1A	CA	MG 6N1B	NA 6O1B	K 6P1A	CO3 6Q1A	HCO3 6I1A	CL 6J1A	SO4 6K1A	B 6L1B	PH SAT. PASTE 6T1A 8C1B
		MEQ/100G					PCT.	MEQ/100G		MMHO	WATER EXTRACT FROM SATURATED PASTE MEQ/1000 ML									PPM
9 137	0- 13	4.2	0.7	0.1	0.2	5.2	68		7.6											
9 138	13- 33	3.8	0.4	0.1	0.2	4.5	83		5.4											
9 139	33- 51	4.7	0.8	0.1	0.1	5.7	100+		4.1											
9 140	51- 94	4.5	1.0	0.1	0.1	5.7	83		6.9											
9 141	94-130	6.1	1.1	0.1	0.1	7.4	100+		6.7											

SAMPLE NO.	DEPTH CM.	WATER EXCHANGEABLE AL. SAT. 8A PCT.	NA 5B4B MEQ.	NA 5D2 PCT.	SAR 5E	GYPSUM AMT. 6F1B PERCENT	REQ. 6P3A	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 PCT.	LT. .0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
9 137	0- 13		0.1	1				MI 3	KK 3	VR 1						MI MICA	1 TRACE
9 138	13- 33		0.1	2												KK KAOLINIT	2 SMALL
9 139	33- 51		0.1	2				MI 3	KK 3							VR VERMICLT	3 MODERATE
9 140	51- 94		0.1	1													4 ABUNDANT
9 141	94-130		0.1	1													5 DOMINANT
																	6 INDETER.

SAMPLE PREPARATION CODES
W=DATA DETERMINED ON WHOLE SOIL AND RECALCULATED TO 2 MM BASIS. GRAVEL FROM SLAKED SUBSAMPLE.

CIENEBA COARSE SANDY LOAM (TAXADJUNCT) 1/
S69Calif-31-23

Location: Placer County, California. From Rocklin Road, approximately 2,200 feet south on Sierra College Blvd. and 30 feet west to pit; 1,200 feet east and 400 feet north from SW corner of sec. 21, T. 11 N., R. 7 E. Date of Sampling: May 9, 1969.
Description by: T. M. Ryan, L. A. Brittan. Collectors: K. W. Flach, W. D. Nettleton, G. M. Kennedy, L. A. Brittan, T. M. Ryan.
Classification: Typic Xerochrept, coarse-loamy, mixed, thermic, shallow.

Granodiorite. Topography: Rolling upland, 13 percent slope on west aspect. Elevation: 500 feet. Drainage: Somewhat excessively drained, slow to medium runoff, moderately rapid permeability. Soil Moisture: Slightly moist.
Remarks: C1 horizon digs with difficulty with spade. Stone line or thin discontinuous horizontally tilted quartzite dike in upper part of third horizon. Less than 10 percent rounded gravel.

HORIZON DESCRIPTION

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Millsholm silt loam variant **LOCATION** Alameda County, California

SOIL FAMILY: Typic Xerochrept, fine-silty, mixed, thermic.

SOIL NOS. S59Calif-01-5 **LAB. NOS.** 59230 thru 59233

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0-7	A_{11}	0.1	0.5	0.6	2.9	7.2	65.9	22.8	33.4	41.7	0	sil
7-14	A_{12}	0.0	0.3	0.4	1.8	7.5	63.1	26.9	31.3	40.7	1	sil-sicl
14-22	B_2	0.0	0.3	0.4	1.9	7.2	63.3	26.9	31.3	40.6	6	sil-sicl
22-60+	D_r			not analysed								

pH 8C1b Saturated Paste 1:1	8C1a 1:10	Organic Matter 6A1a Organic Carbon Pct.	6B1a Nitrogen Pct.	C/N	6E1b $CaCO_3$ equivalent Pct.	8A1a Elec. conduc. mmhos/cm. at 25°C.						4B2 Soil water 15 atmos. Pct.
6.0	6.6	1.43	0.131	10.9		0.5						10.1
5.7	6.6	0.64	0.077	8.3		0.2						11.4
5.7	6.6	0.53	0.066	8.0		0.1						11.7

5A2a CATION EXCHANGE CAPACITY (+ Na)	EXTRACTABLE CATIONS 5B 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a Ca	6O1a Mg	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
28.3	13.4	8.3	6.7	0.2	0.4	77	1.0	0.1				44.0
29.1	14.4	10.1	6.8	0.3	0.3	79	0.8	<0.1				38.0
31.2	13.5	10.5	6.5	0.2	0.3	79	0.7	<0.1				38.2

MILLSHOLM SILT LOAM variant 1/
S59Calif-01-5

Collectors: Dick Huff, Larry Welch, Lothair Grant.
Description by: Larry E. Welch.
Date of Sampling: May 5, 1959.

Location: Alameda County, California, in the SE 1/4 of the SE 1/4 of the NE 1/4 of Sec. 28, T3S, R1W, near the junction of Palomaris and Stony-

Relief: Strongly sloping uplands on a convex SW-facing slope of 35 percent.

Drainage: Well drained, moderate runoff, moderate permeability.

Parent Material: Shale.

Climate: Rainfall about 20 inches.

Vegetation: Annual grasses.

Classification: Typic Xerochrept, fine-silty, mixed, thermic.

DESCRIPTION

HORIZON		
A_{11}	0- 7"	Olive (5Y 5/3) silt loam, olive brown (2.5Y 4/3) when moist; massive; very hard when dry, friable when moist, slightly sticky and slightly plastic when wet; abundant very fine roots;

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE San Emigdio loam (taxadjunct) **LOCATION** Riverside County, California

SOIL FAMILY: Calcixerollic Xerochrept, coarse-loamy, mixed, th rmic.

SOIL NOS. S55Calif-33-10 **LAB. NOS.** 55573-577

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	3A1 2A2 > 2 1/	TEXTURAL CLASS
0-8	A11	1.7	4.8	5.9	17.9	15.7	38.2	15.8	44.6	21.1	3	1
8-23	A12	2.6	5.9	6.1	17.8	14.3	38.0	15.3	42.0	22.0	4	1
23-41	C1	2.6	5.7	5.8	17.1	14.1	40.2	14.5	45.6	23.9	3	1
41-54	C2	7.2	13.0	10.4	22.0	13.0	25.5	8.9	40.5	13.6	19	s1
54-60	C3	2.4	4.5	5.4	19.3	17.6	40.1	10.7	51.1	19.7	4	1

pH 8C1b SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
6.6		6.7	0.44	0.049	9.0		0.39					7.5
7.2		7.6	0.27	0.030	9.0		0.43					7.4
7.6		8.4	0.18				0.57	1				8.0
7.9		8.6	0.10				0.42	1				5.1
7.9		8.5	0.08				0.30	2				6.4

5A2a CATION EXCHANGE CAPACITY ←(Na)	EXTRACTABLE CATIONS 5B 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a Ca+Mg 6O1a		6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
18.4	13.4	2.9	0.7	0.1	0.6	96	0.5	0.2	3.2			29.6
20.5	16.7	3.0	0.8	0.1	0.3	96	0.5	0.1	3.6			30.5
21.6	30.4	2.9	-	0.3	0.2	100+	0.8	0.1	5.1			32.5
13.8	19.8	3.7	-	0.2	0.1	100+	0.9	0.1	3.5			24.3
18.2	25.5	2.5	-	0.3	0.2	100+	1.4	Tr.	1.8			31.1

1/ Percent of the whole soil

SAN EMIGDIO LOAM (taxadjunct) 1/
S55Calif-33-10

Location: 300' W and 1000' S of NE corner of Sec. 12, T3S, R3W, near Moreno, Riverside County, California.
Sampled by: Fred Eggers, Arnold Knecht, F. G. Stephens, W. G. Harper, R. E. Nelson and F. F. Harradine. **Description by:** W. G. Harper
Elevation: 1625 feet. **Sampling date:** December 6, 1955.
Precipitation: 14 inches

Classification: Calcixerollic Xerochrept, coarse-loamy, mixed, thermic.

California Group: I

Vegetation: This dry-farmed area had a crop of wheat in 1955.

Parent Material: Alluvium from San Timoteo Badlands

Physiography: Gently sloping.

Drainage: Well drained.

HORIZON		DESCRIPTION
A_{11}	0- 8"	Brown (10YR 5/3 dry; very dark grayish brown 3.5/2.5 moist) loam; weakly defined medium to fine granular; hard, friable; moderately permeable; many roots; noncalcareous; pH 7.5; clear smooth boundary.
A_{12}	8-23"	Brown (10YR 5/3 dry; dark brown 4/2 moist) loam; weakly defined medium to fine angular blocky or subangular blocky; extremely hard, friable; moderately permeable; many roots; noncalcareous; pH 8.0; clear smooth boundary.
C 1	23-41"	Light yellowish brown (10YR 6/4 dry; dark brown 4/3 moist) loam; weakly defined medium to fine angular blocky; hard, friable; moderately permeable; many roots; effervesces slightly; contains slight amount of mycelium-like lime; pH 8.0; clear smooth boundary.
C 2	41-54"	Pale brown (10YR 6/3 dry; dark brown 4/3 moist) fine sandy loam; massive; slightly hard to hard, very friable; moderately permeable; effervesces slightly; pH 8.0; about 50 percent rounded pebbles 2 to 6" in diameter, some coated with white lime; clear smooth boundary.

CONSECUTIVE PEDON NUMBER C 68 92 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES INVILLE TAXADJUNCT SOIL NO. S68 CALI. 9-17 SAMPLE NO. 8 748-8 753 TAHOE BASIN AREA

SOIL FAMILY DYSTRIC XEROCHREPT COARSE-LOAMY,MIXED,FRIGID RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SANDS VCS 2-1	CS 1-.05	MS .5-.25	FS .25-.10	VFS .1-.05	SILTS CSI .05-.02	FSI .02-.002	CLAYS CO. 2-.2U	FINE LT .2U	CARBONATE 3A1A	1A2A INT. COARSE FRAGMENTS II .2-.02 PCT. OF WHOLE SOIL	I 75-20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM													PCT. OF LT 75			
8 748	0- 13	A11	74.2	17.2	8.6	12.9	21.1	12.8	18.2	9.2	7.9	9.3				25.5	26A	9	17Z	N
8 749	13- 46	A12	73.2	16.4	10.4	14.4	20.6	12.5	17.8	7.9	7.4	9.0				23.7	26A	10	16Z	N
8 750	46- 69	B21T	70.5	16.7	12.8	11.3	18.7	12.9	19.4	8.2	7.1	9.6				24.6	23A	11	12Z	N
8 751	69-120	B22T	70.9	15.0	14.1	10.7	20.5	13.4	18.7	7.6	5.4	9.6				22.1	19A	5	14Z	N
8 752	120-175	C1 SIM	81.9	8.7	9.4	14.3	22.7	18.1	20.6	6.2	3.8	4.9				18.6	34A	8	26Z	N
8 753	175-200	C 2	84.0	11.7	4.3	8.9	35.0	15.2	18.3	6.6	4.3	7.4				20.0	7A	--	7Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		PERCENT		PERCENT		PERCENT				PCT.NO.40		GRAMS PER CC				PERCENT			PERCENT	
8 748	0- 13	1.50	0.053			2.6				NP	NP	1.25			0.86			6.0		
8 749	13- 46	0.49	0.026			2.6						1.30			0.85			5.8		
8 750	46- 69	0.25				2.5				NP	NP	1.35			0.87			6.1		
8 751	69-120	0.20				2.4						1.35			0.89			7.0		
8 752	120-175					1.2						1.40			0.79			6.0		
8 753	175-200					0.6				NP	NP	1.40			0.96			4.4		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 584A 6N2E	MG 584A 6O2D	NA 584A 6P2A	K 584A 6Q2A	SUM	EXT. ACIDITY 6H2A	KCL EXT. AL 6G1E	CEC SUM 5A3A	CEC NH4-OAC 5A6A	BASE SATURATION SUM+ ACIDITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CACL2 8C1E
		MEQ/100G									PERCENT					
8 748	0- 13	2.8	0.3	TR	0.4	3.5	8.0		11.5	9.5	31		37		5.7	5.2
8 749	13- 46	1.8	0.3	0.1	0.4	2.6	5.7		8.3	9.2	31		28		5.6	5.0
8 750	46- 69	1.7	0.4	0.1	0.4	2.6	5.7	0.4	8.3	5.8	31	87	45		5.5	5.0
8 751	69-120	1.7	0.4	0.1	0.3	2.5	4.5	0.2	7.0	5.5	36	93	45		5.6	5.1
8 752	120-175	1.3	0.2	0.1	0.2	1.8	3.3	0.1	5.1	4.6	35	95	39		5.5	5.0
8 753	175-200	2.1	0.8	0.1	0.1	3.1	3.0	0.2	6.1	5.7	51	94	54		5.4	4.9

SAMPLE NO	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY I 7A2E	II 7A2E	III 7A2E	IV 7A2E	DTA 7A3	7A3	L.T. 0.0002 MM X RAY I 7A2E	II 7A2E	III 7A2E	MINERAL CODE	RELATIVE AMOUNT
		PERCENT										
8 748	0- 13	KK 3	VR 2	MI 2							KK KAOLINIT	1 TRACE
8 749	13- 46										VR VERMICLT	2 SMALL
8 750	46- 69										MI MICA	3 MODERATE
8 751	69-120											4 ABUNDANT
8 752	120-175											5 DOMINANT
8 753	175-200	KK 2										6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

INVILLE GRAVELLY COARSE SANDY LOAM (TAXADJUNCT) 1/

S68Calif-9-17

Location: Tahoe Basin, El Dorado County, California. 0.2 mile northeast of south ¼ corner, sec. 11, T. 12 N., R. 18 E.; three miles south of Bijou, ¼ mile inside Forest Service boundary, 50 feet south of High Meadow Road, South Lake Tahoe. Date of Sampling: October 26, 1967.

Description by: R. C. Huff. Collectors: G. M. Kennedy, G. L. Anderson, J. H. Rogers, D. Palmquist.

Classification: Dystric Xerochrept, coarse-loamy, mixed, frigid.

Vegetation: Not cultivated; Jeffrey pine, white fir, greenleaf manzanita, squaw carpet. Climate: Precipitation is 30 inches. Mean annual temperature is 43°F., January 27°F., July 61°F. Parent Material: Mixed alluvium. Topography: Gently sloping terrace, 2 to 5 percent north slope. Elevation: 6500 feet. Drainage: Well drained; runoff is medium; permeability is slow to 48 inches and very slow below. Soil Moisture: Below wilting point.

Remarks: Field determinations for pH by Hellige-Truog triplex indicator. Portion of the C1m horizon with 7.5YR colors not cemented. Horizons A11, A12 (0-18") and B21t, B22t (18 -48") were sampled for State Division of Highways.

HORIZON	DESCRIPTION
	(5-0 cm.)
01 & 02	2 to 0 inches, not sampled. Pine needles and partly decomposed needles.
	(0-13 cm.)
A11 RSL 68748	0 to 5 inches, brown (7.5YR 5/4) gravelly coarse sandy loam, dark brown (7.5YR 3/2) moist; weak fine crumb structure; soft, very friable, nonsticky and nonplastic; many very fine, fine, medium and coarse roots; many very fine interstitial pores; 20 percent gravel; slightly acid (pH 6.5); clear smooth boundary.
	(13-46 cm.)
A12 RSL 68749	5 to 18 inches, strong brown (7.5YR 5/6) gravelly coarse sandy loam, yellowish red (5YR 4/6) moist; weak fine crumb structure; soft, very friable, slightly sticky and slightly plastic; many very fine, fine, medium and coarse roots; many very fine interstitial pores; 20 percent gravel, less than 2 percent cobbles; medium acid (pH 6.0); diffuse wavy boundary.
	(46-69 cm.)
B21t RSL 68750	18 to 27 inches, reddish yellow (7.5YR 6/6) cobbly coarse sandy loam, yellowish red (5YR 4/6) moist; massive; very hard, firm, slightly sticky and slightly plastic; many very fine, few fine and medium roots; common very fine interstitial, few very fine and fine tubular pores; many thin clay films in pores and bridges; 20 percent cobbles, 10 percent gravel; medium acid (pH 6.0); clear wavy boundary.
	(69-120 cm.)
B22t RSL 68751	27 to 48 inches, reddish yellow (7.5YR 6/6) very gravelly coarse sandy loam, strong brown (7.5YR 5/6) moist; massive; very hard, firm, slightly sticky and slightly plastic; common very fine, few fine and medium roots; common very fine interstitial, few very fine and fine tubular pores; common thin clay films in pores and bridges; 50 percent gravel and less than 1 percent cobbles; medium acid (pH 5.7); abrupt wavy boundary.
	(120-175 cm.)
C1sim RSL 68752	48 to 68 inches, light yellowish brown and very pale brown (10YR 6/4, 7/4) very gravelly loamy sand, yellowish brown and brownish yellow (10YR 5/4, 6/6) moist; reddish yellow (7.5YR 6/8) blotches, strong brown (7.5YR 5/8) moist; massive; very hard, very firm, nonsticky and nonplastic; roots are matted on 2 lenses about a foot apart; very few very fine interstitial pores; 50 percent gravel and 3 percent cobbles; reddish yellow (7.5YR) blotches are not cemented; medium acid (pH 5.7); abrupt wavy boundary.
	(175-200 cm.)
C2 RSL 68753	68 to 78 inches, very pale brown (10YR 7/3) loamy sand, light yellowish brown (10YR 6/4) moist; yellow (10YR 7/6) blotches; massive; soft, very friable; common fine and medium roots; common very fine interstitial pores; strongly acid (pH 5.5).

1/ The Inville series is in a loamy-skeletal, mixed, frigid family of Ultic Haploxeralfs. This pedon lacks an argillic horizon and has less than 35 percent coarse fragments by volume.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Hugo sandy loam variant LOCATION San Mateo County, California

SOIL FAMILY: Dystric Xerochrept, coarse-loamy, mixed, mesic.

SOIL NOS. S57-Calif-41-2 LAB. NOS.

HUGO SANDY LOAM variant 1/
S57Calif-41-2

Location: San Mateo County, California. 4.9 miles north of Sky Londa along California Highway No. 5 (Skyline Blvd.), thence 2.35 miles southwest along Swett Road to junction with Double K Ranch Road, then 1/4 mile southerly along Double K Ranch Road to sample site in road cut (uphill) on west side of road, SW1/4 NW1/4 sec. 17, T. 6 S., R. 4 W. Date of Sampling: May, 1957.
Description by: Rudolph Ulrich. Collectors: T. H. Arkley, R. Ulrich, and R. J. Wagner.
Classification: Dystric Xerochrept, coarse-loamy, mixed, mesic.
Vegetation: Second growth Redwood and Douglas fir overstory, with an understory of tanbark oak and madrone trees, scattered poison oak, and little grass in ground cover due to opening produced by the roadway. Parent Material: Weathered Vaqueros sandstone. Topography: Mountainous with mostly steep and very steep slopes generally in excess of 45 percent and narrow crests and divides. F slope of 45 percent, convex toward east (estimated). Elevation: About 1,800 feet. Drainage: Runoff rapid to very rapid; permeability rapid; somewhat excessive general drainage.
Remarks: Bulk samples only. Condition of soil throughout extremely wet following two heavy rains each in excess of 2 inches.

HORIZON		DESCRIPTION
A_{00} & A_0 (No sample taken)	2-0"	Fresh and rotting leaves, needles, and twigs. Very dark colored wet and soggy and spongy. Abrupt smooth boundary.
A_1 (S57-Cal-41-2-1)	0-8"	Pale brown (10YR 6/3) dry, dark brown, (10YR 4/3) moist sandy loam with few whitish mycelial strands; weak fine granular; slightly hard, very friable, nonsticky, nonplastic; common reddish brown subrounded to subangular shot (2-10 mm.) between A_0 and A_1, few (2-5 mm.) in horizon; pH 6.0; common worm casts upper few inches; common fine and medium roots, few large ones; many fine interstitial pores, common fine continuous simple tubular pores; gradual smooth boundary.
B_1 (S57-Cal-41-2-2)	8-26"	Very pale brown (10YR 7/4) strong brown (7.5YR 5/5) moist, gravelly (sorted out and discarded about 25 percent by volume as gravel and small stones) sandy loam; very weak coarse subangular blocky; slightly hard, very friable, nonsticky, nonplastic; clay skins, if any, are very thin and very patchy; pH 5.6; common fine and medium roots, few larger ones; many fine interstitial pores, common fine and few coarse simple tubular pores; diffuse boundary.
B_2 (S57-Cal-41-2-3)	26-38"	Very pale brown (10YR 7/4) strong brown (7.5YR 5/5) moist, very stony (sorted out and discarded about 65 percent by volume as angular stones and coarse gravel) heavy sandy loam with few thin patchy stronger brown (7.5YR 5/8) moist, clay skins and thin pore linings; very weak medium subangular blocky; slightly hard to hard, friable, nonsticky, very slightly plastic; pH 5.0; common medium and fine roots, few large ones; many fine interstitial pores, few fine and coarse simple tubular pores; abrupt broken boundary.
D_r	38"+	Yellow (10YR 7/5) brownish yellow (10YR 6/6) moist, soft sandstone (Vaqueros Formation) with thin patchy reddish

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Temescal very fine sandy loam **LOCATION** Riverside County, California

SOIL FAMILY: Lithic Xerochrept, loamy, mixed, thermic

SOIL NOS. S55Calif-33-9 **LAB. NOS.** 55569-572

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-2	A11	5.5	8.8	8.3	30.8	19.5	20.9	6.2	55.0	6.5	5	fsl
2-9	A12	3.9	8.6	8.5	31.0	19.9	20.5	7.6	54.8	6.8	10	fsl
9-19	C1	3.8	9.8	9.2	27.5	17.1	17.2	15.4	42.4	9.9	16	fsl
19+	Dr			not analysed								

8C1b pH SATURATED PASTE 1:1	8C1a pH 1:5	8C1a pH 1:10	6A1a ORGANIC MATTER ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	4B1a MOISTURE TENSIONS 1/10 ATMOS. %	4B1a 1/3 ATMOS. %	4B2 15 ATMOS. %
6.2		6.2	2.14	0.134	16.0		0.37					5.5
6.7		6.7	0.71	0.057	12.5		0.35					4.5
6.6		6.6	0.29	-	-		0.25					7.8

5A2a CATION EXCHANGE CAPACITY (Na)	5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg		6K1a Cl	8A MOISTURE AT SATURATION %
	EXTRACTABLE CATIONS 5B milliequivalents per 100g. soil						milliequivalents per liter					
15.8	9.0	1.4	4.6	0.2	0.4	71	0.7	0.4	2.8			39.1
11.5	6.6	1.2	2.0	0.1	0.3	80	0.5	0.3	2.5			29.9
12.4	4.4	1.5	2.1	0.1	0.5	75	0.6	0.2	1.5			32.1

1/ - Percent of the whole soil

TEMESCAL VERY FINE SANDY LOAM
S55Calif-33-9

Collectors: Fred Eggers, Arnold Knecht, F. G. Stephens, W. G. Harper, R. E. Nelson and F. F. Harradine.

Sampling date: December 7, 1955. Description by: W. G. Harper

Location: Above road cut on west side of Highway 79, 1 mile S of Sage near N line of Sec. 18, T7S, R1E, about 12 miles south or Hemet, Riverside County, California.

Elevation: 2400 feet

Precipitation: 16 inches

Classification: Lithic Xerochrept, loamy, mixed, thermic.

Vegetation: Chemise, cheatgrass, alfileria, wild buckwheat, manzanita; chamise is taller and more dense than at Site 5.

Parent Material: Mica schist. Undivided pre-Cretaceous metamorphics.

Physiography: Rolling upland

Drainage: Well drained

HORIZON		DESCRIPTION
A_{11}	0- 2"	Brown (10YR 5/3 dry, dark brown 3/3 moist) very fine sandy loam; very weakly defined very fine crumb structure; soft, very friable; moderately permeable; many roots and much organic matter; some moss on surface; micaceous; pH 6.0; abrupt smooth boundary.
A_{12}	2- 9"	Dark brown (10YR 4/3 dry, dark brown 3/3 moist) very fine sandy loam; very weakly defined fine crumb structure; slightly hard, very friable; moderately permeable; moderate amounts of roots and organic matter; very strongly micaceous with a few angular pebbles; pH 6.5; abrupt smooth boundary.
C_1	9-19"	Dark brown (10YR 4/3 dry, dark brown 3/3 moist) very fine sandy loam or loam (may have a few clay films); moderately defined medium to fine subangular blocky; very hard, friable; roots common to few, many root channels and some worm holes; high in mica; pH 6.5; abrupt irregular boundary. This horizon contains many angular mica schist fragments. This differs but little from a C horizon even though it has some structure and may have a few clay films.
Dr	19"+	Decomposing to hard mica schist bedrock. The mica schist may be somewhat harder than at Site 5.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Toomes loam **LOCATION** Placer County, California

SOIL FAMILY: Lithic Xerochrept, loamy, mixed, thermic

SOIL NOS. S61Calif-31-1 **LAB. NOS.** 6173 - 6175

TOOMES LOAM
S61Calif-31-1

Collectors: Reuben Nelson, Grant Kennedy, John Rogers, Lynn Brittan, David Estrada, Paul Nazar, Lothair Grant Description by: Grant Kennedy.
Date of Sampling: April 26, 1961
Location: Placer County, California, 5.5 miles north of Roseville, Whitney Ranch, SW1/4SE1/4SE1/4, sec. 34, T. 12 N., R. 6 E.
Elevation: 170 feet
Relief: Hummocky uplands with a 3 percent slope to the northeast.
Drainage: Well drained; rapid runoff; moderate permeability
Parent Material: Andesitic breccia
Climate: Mean annual rainfall 24 inches, T 61.3°, J 45.6°, J 74.7°
Vegetation: Soft chess, wild oats, filaree, bur clover

[redacted]

Remarks: The hummocks are about 20 feet apart and 9 feet in diameter and 18 to

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Toomes loam **LOCATION** Placer County, California

SOIL FAMILY: Lithic Xerochrept, loamy, mixed, thermic

SOIL NOS. S61Calif-31-2 **LAB. NOS.** 6176 - 6178

DEPTH INCHES	HORIZON	1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	3A1 2A2 > 2 1/	TEXTURAL CLASS
		PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)										
0- 1	A_{11}	3.2	9.6	6.6	17.3	10.1	41.4	11.8	36.0	25.9	13	1
1-11	A_{12}	4.7	8.6	5.6	16.4	11.0	42.6	11.1	37.3	27.1	13	1
11-17+	R											

pH 8C1b Saturated Paste 1:1	pH 8C1c KCl 1:1	Organic Matter 6A1a Organic Carbon Pct.	Organic Matter 6B1a Nitrogen Pct.	C/N	$CaCO_3$ equivalent %	Elec. Conduc. Mmhos/cm. %	6C1b Free Fe_2O_3 %	4A3a Bulk Den. gm/cc			Soil Water 15 ATMOS. %
5.8	5.1	4.61	0.338	13.6			1.8				12.4
6.3	5.0	0.61	0.060	10.2			1.8	1.75			6.7
											-

5A2a CATION EXCHANGE CAPACITY (Na)	6N2b Ca	EXTRACTABLE CATIONS 6O2c Mg	6H2a H	6P2a Na	5B1a 6Q2a K	BASE SAT. % 5C3	5A3a Sum of Cations soil		Calc. Sum of Cations clay		Calc. Sum of Cations clay + fine silt
		milliequivalents per 100g. soil					milliequivalents per 100 g.				
20.1	10.2	2.5	7.6	0.2	1.0	65	21.5		182		40
13.5	5.9	2.7	3.1	0.2	0.3	77	13.2		118		25

1/ Percent of the whole soil

TOOMES LOAM
S61Calif-31-2

Collectors: Reuben Nelson, Grant Kennedy, John Rogers, Lynn Brittan, David Estrada, Paul Nazar, Lothair Grant **Description by: Grant Kennedy.**
Date of Sampling: April 26, 1961
Location: Placer County, California, 5.5 miles north of Roseville, Whitney Ranch, 60 feet south of entrance road, 1/8 mile east of 99E, NW1/4NE1/4SE1/4 sec. 33, T. 12 N., R. 6 E.
Elevation: 120 feet
Relief: Hummocky uplands with a 2 percent slope to the northwest.
Drainage: Well drained; rapid runoff; moderate permeability
Parent Material: Andesitic breccia
Climate: Mean annual rainfall 23 inches, T. 61.3°, J. 45.6°, J 74.4°

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Auburn silt loam **LOCATION** Amador County, California

SOIL FAMILY: Ruptic-Lithic Xerochrept, loamy, mixed, thermic

SURVEY NOS. S59Calif-3-7 **LAB. NOS.** 59306 - 59309

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-1½	A_{11}	2.5	4.4	3.4	6.7	5.9	63.4	13.7	31.5	41.9	14	sil
1½-9	A_{12}	2.3	3.7	2.9	6.7	6.2	64.7	13.5	31.5	43.5	13	sil
9-14	B_2	2.6	4.3	2.7	6.1	6.2	64.1	14.0	32.1	42.2	13	sil
14-24+	D_r	Partially weathered amphibolite										

pH 8C1a H_2O 1:1	pH 8C1c KCl 1:1	pH 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	"Free" Fe_2O_3 6C1b %	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	GYPSUM me./100g. SOIL	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.4	5.3		3.79	0.247	15.3	2.5						9.8
6.4	4.8		1.18	0.098	12.0	2.7						7.0
6.5	4.5		0.42	0.043	9.8	3.1						7.7

5A2a CATION EXCHANGE CAPACITY ←(Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6Q2c	6H2a H	Na 6P2a	K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE Na	K				MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
23.2	11.7	5.3	6.4	0.1	0.5	73						
19.6	9.3	5.0	5.1	<0.1	0.1	74						
20.7	8.8	7.5	4.6	<0.1	<0.1	78						

1/ Percent of total soil

AUBURN SILT LOAM
S59Calif-3-7

Collectors: Grant Kennedy, Reuben Nelson, Robert Sketchley, John Rogers
Date of sampling: June 16, 1959 Description by: Grant Kennedy.
Location: Amador County, California - 1/4 mile west and 100 feet north of the SE corner of Section 6, T6N, R9E.
Elevation: 620 feet
Relief: Moderately steep foothills on a 10% slope to the east.
Drainage: Well drained; moderate runoff; moderately permeable above the parent rock.
Parent Material: Amphibolite schist.
Climate: Mean annual rainfall about 24 inches.
Vegetation: Annual grasses, oaks, scattered Digger pine.
Remarks: This soil is a very weakly developed Brunizem near the Brunizem-Noncalcic Brown zonal boundary. Scattered rock outcrops occur in this area.
Classification: Ruptic-Lithic Xerochrept, loamy, mixed, thermic.

HORIZON		DESCRIPTION
A_{11}	0-1½"	Strong brown (7.5YR 5/6) silt loam. reddish brown (5YR 4/4) [illegible] moist, slightly sticky and nonplastic when wet; abundant very fine roots; many very fine and fine pores; slightly acid, pH 6.4; clear, smooth lower boundary.
A_{12}	1½-9"	Color texture, structure and consistence as above, except yellowish red (5YR 5/6) when dry and slightly plastic when wet; abundant very fine and fine, plentiful medium roots; many very fine and fine, plentiful medium pores; few thin, discontinuous clay films in pores; slightly acid, pH 6.4; gradual, smooth lower boundary.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Auburn silt loam **LOCATION** Amador County, Calif.

SOIL FAMILY: Ruptic-Lithic Xerochrept, loamy, mixed, thermic

SOIL NOS. S59Calif-3-8 **LAB. NOS.** 59310 - 59313

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-1	A_{11}	2.9	4.5	3.3	5.5	4.5	65.9	13.4	35.1	38.5	4	sil
1-7	A_{12}	1.4	3.6	2.9	5.7	5.8	66.7	13.9	37.9	38.0	19	sil
7-14	B_2	2.0	4.6	3.2	5.7	5.5	65.5	13.5	33.3	41.1	8	sil
14-26+	D_r	Decomposing amphibolite schist										

pH		ORGANIC MATTER		"Free" Fe_2O_3	ELECTRICAL CONDUC-			MOISTURE TENSIONS		
8C1a	8C1c	6A1a	6B1a							4B2

AUBURN SILT LOAM
S59Calif-3-8

Collectors: Grant Kennedy, Reuben Nelson, Robert Sketchley and John Rogers.
Date of Sampling: June 16, 1959. Description by: Grant Kennedy.
Location: Amador County, California on the Kuhl property - 300 feet NE of the SW corner of Section 7, T7N, R9E.
Elevation: 760 feet.
Relief: Very gently sloping foothills, on a 3 percent slope to the northwest.
Drainage: Well drained; moderate runoff; moderately permeable above the D...

Vegetation: Annual grasses, forbs, and oaks including filaree, soft chess, ripgut and fiddle-neck used for range.
Remarks: The soil is a (minimal) Brunizem. The profile is nearly free of stones and the bedrock is weathered irregularly.
Classification: Ruptic-Lithic Xerochrept, loamy, mixed, thermic.

CONSECUTIVE PEDON NUMBER C 68 82 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES CELIO. SOIL NO. S68 CALI. 9-14 SAMPLE NO. 8 678-8 684 TAHOE BASIN AREA

SOIL FAMILY AQUENTIC CRYUMBREPT, SANDY-SKELETAL, MIXED RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SANDS VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBONATE 3A1A	INT. II .2-.02	1A2A COARSE FRAGMENTS PCT. OF WHOLE SOIL	I 75- 20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75			
8 678	0- 20	A11	85.7	10.5	3.8	30.8	27.9	10.5	11.7	4.8	4.7	5.8				14.9	46A	14	32Z	N
8 679	20- 41	A12	81.6	14.5	3.9	23.3	26.6	12.1	13.1	6.5	6.0	8.5				18.7	46A	17	29Z	N
8 680	41- 58	C 1	83.4	13.1	3.5	21.2	28.8	13.3	13.9	6.2	5.4	7.7				17.9	56A	21	35Z	N
8 681	58-115	2C 2	94.3	3.4	2.3	19.0	40.8	20.4	11.8	2.3	1.4	2.0				8.2	51A	9	42Z	N
8 682	115-140	2C 3	SIM 97.2	1.3	1.5	36.0	40.9	13.7	6.3	0.3	0.3	1.0				2.4	63A	20	43Z	N
8 683	140-170	2C 4	98.5	0.7	0.8	67.0	27.0	3.2	1.1	0.2	0.1	0.6				0.7	46A	7	39Z	N
8 684	170-205	2C 5	98.5	0.8	0.7	16.1	39.8	21.6	18.9	2.1	0.6	0.2				10.7	27A	6	21Z	N

SAMPLE NO.	DEPTH CM.	ORG. MATTER CARBON 6A1A -PERCENT-	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B -PERCENT-	2-20 6E1B	SESQUIOXIDES DI-CIT EXT FE 6C2B	AL 6G7A	PYROP EXT FE 6C5A	AL 6G5A	ATTERBERG LIMIT LL 4F2 PCT.	PL 4F1 NO.40	BULK DENSITY EST. 1/3 BAR 4A1F GRAMS	DRY 4A1H PER CC	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
						DEPTH PERCENT									PERCENT			PERCENT	
8 678	0- 20	3.53	0.145			2.7				NP	NP	1.30		0.70			6.0		
8 679	20- 41	2.12	0.107			2.8						1.35		0.69			6.4		
8 680	41- 58	1.19	0.081			2.7				NP	NP	1.40		0.60			5.5		
8 681	58-115	0.47				1.6						1.50		0.63			2.3		
8 682	115-140	0.08				1.2						1.80		0.46			1.8		
8 683	140-170					0.4						1.70		0.65			0.7		
8 684	170-205					0.3						1.70		0.81			0.7		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6Q2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACIDITY 6H2A	KCL EXT. AL 6G1E	CEC SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACIDITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CACL2 8C1E
		MEQ/100G									PERCENT					
8 678	0- 20	1.7	0.1	0.1	0.2	2.1	14.8	0.7	16.9	16.4	12	75	13		5.4	4.9
8 679	20- 41	0.5	0.1	TR	0.2	0.8	12.7	0.1	13.5	9.7	6	89	9		5.5	5.2
8 680	41- 58	0.5	0.1	TR	0.1	0.7	10.7	0.1	11.4	7.5	7	88	10		5.6	5.2
8 681	58-115	0.1	TR	TR	0.1	0.3	3.8	0.1	4.1	2.7	7	74	11		5.5	5.4
8 682	115-140	0.1	TR	TR	TR	0.2		0.1	0.2	2.1	100+	70	11		5.6	5.4
8 683	140-170	0.1	TR	TR	TR	0.2		0.1	0.2	1.8	100+	70	13		5.5	5.3
8 684	170-205	0.1	TR	0.1	0.1	0.3		0.1	0.3	1.3	100+	77	26		5.5	5.2

CLAY MINERALOGY

SAMPLE NO	DEPTH CM.	LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 678	0- 20	VR 1	MI 1								VR VERMICLT	1 TRACE
8 679	20- 41										MI MICA	2 SMALL
8 680	41- 58	MI 1	KK 1	VM 1								3 MODERATE
8 681	58-115										KK KAOLINIT	4 ABUNDANT
8 682	115-140										VM VM/MICA	5 DOMINANT
8 683	140-170											6 INDETER.
8 684	170-205											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

CELIO GRAVELLY LOAMY COARSE SAND
S68Calif-9-14

Location: Tahoe Basin, El Dorado County, California. 0.1 mile north of southeast corner sec. 30, T. 12 N., R. 18 E. 0.25 mile west of Meyers checking station, west side of Highway 89 extension. Date of Sampling: October 25, 1967.
Description by: R. C. Huff Collectors: C. M. Kennedy, J. H. Rogers, G. L. Anderson

D. P[redacted]a Imquist.

CONSECUTIVE PEDON NUMBER C 68 83 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES TALLAC SOIL NO. S68 CALI. 9-16 SAMPLE NO. 8 685-8 690 TAHOE BASIN AREA

SOIL FAMILY ENTIC CRYUMBREPT LOAMY-SKELETAL,MIXED RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1-.5	MS .5-.25	FS .25-.10	VFS .1-.05	SILTS CSI .05-.02	FSI .02-.002	CLAYS CO. 2-.2U	FINE LT .2U	CARBO NATE 3A1A	1A2A INT. 11 .2-.02	COARSE FRAGMENTS PCT. OF WHOLE SOIL 1A2A	I 75-20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM																	
8 685	3- 0	01+02															--A	--	--	--
8 686	0- 38	A11	68.5	26.4	5.1	11.2	16.9	10.5	16.7	13.2	13.1	13.3				35.9	14A	11	3Z	N
8 687	38- 53	A12	67.3	28.5	4.2	9.1	17.0	10.8	16.9	13.5	14.7	13.8				37.8	43A	21	22Z	N
8 688	53- 79	C 1	63.6	32.2	4.2	10.8	16.9	9.8	14.4	11.7	15.1	17.1				34.8	46A	20	26Z	N
8 689	79-105	C 2	76.5	19.7	3.8	17.4	23.2	12.7	14.8	8.4	10.5	9.2				26.5	48A	11	37Z	N
8 690	105-165	C 3SIM	73.6	23.3	3.1	7.5	18.7	13.5	20.4	13.5	13.1	10.2				37.0	35A	12	23Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR-BON 6A1A	NITRO-GEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP FE 6C5A	EXT AL 6G5A	ATTERBERG L I M I T LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 4A1F	1/3 BAR 4A1F	DRY 4A1H 3B2	CM	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		-PERCENT-		-PERCENT-		PERCENT				PCT.NO.40		GRAMS PER CC				PERCENT			-PERCENT-	
8 685	3- 0											0.40								
8 686	0- 38	3.92	0.195			1.2				NP	NP	1.35		0.93				8.2		
8 687	38- 53	2.97	0.162			1.1						1.40		0.71				7.4		
8 688	53- 79	1.61	0.110			1.1				NP	NP	1.50		0.67				6.5		
8 689	79-105	0.79				1.0						1.50		0.66				4.3		
8 690	105-165	0.29				1.1						1.70		0.75				4.2		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACI-DITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 685	3- 0															
8 686	0- 38	7.1	0.4	0.1	0.5	8.1	15.9		24.0	19.0	34		43		5.9	5.4
8 687	38- 53	4.5	0.4	0.1	0.5	5.5	13.2		18.7	15.5	29		35		6.0	5.4
8 688	53- 79	2.4	0.2	TR	0.5	3.1	12.6		15.7	12.4	20		25		6.0	5.4
8 689	79-105	1.3	0.1	TR	0.4	1.8	7.2		9.0	7.6	20		24		5.9	5.4
8 690	105-165	1.4	0.1	0.1	0.6	2.2	5.3		7.5	7.8	29		28		5.7	5.3

CLAY MINERALOGY

SAMPLE NO	DEPTH CM.	LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE		RELATIVE AMOUNT
8 685	3- 0												1 TRACE
8 686	0- 38	VR 2	KK 2								VR	VERMICLT	2 SMALL
8 687	38- 53										KK	KAOLINIT	3 MODERATE
8 688	53- 79	VR 2	KK 2	MI 1							MI	MICA	4 ABUNDANT
8 689	79-105												5 DOMINANT
8 690	105-165												6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

TALLAC GRAVELLY COARSE SANDY LOAM
S68Calif-9-16

Location: Tahoe Basin, El Dorado County, California; 0.2 mile north and 0.1 mile west

Lot 18 of the Forest Service Spring Creek Tract. Date of Sampling: October 26, 1967.
Description by: R. C. Huff. Collectors: G. L. Anderson, G. M. Kennedy, J. H. Rogers, D. Palmquist.
Classification: Entic Cryumbrept, loamy-skeletal, mixed.
Vegetation: Not cultivated. Jeffrey pine, white fir, some stipa and annual weeds.
Climate: Precipitation is 35 to 40 inches. Mean annual temperature about 41°F., January about 30°F., July about 63°F. Parent Material: Glacial deposit, predominantly metabasic. Topography: Undulating moraine; 2 to 5 percent east slope. Elevation: 6480 feet. Drainage: Well drained; runoff is medium; permeability is moderate to 42 inches, slow below. Soil Moisture: Below wilting point.
Remarks: Field determinations of pH by Hellige-Truog triplex indicator. Horizons A11, A12 (0-21"); C1, C2 (21-42"); and C3si (42-65") were sampled for State Division of Highways.

HORIZON	DESCRIPTION
01&02 RSL 68685	(3-0 cm.) 1 to 0 inches, fresh and decomposed conifer needles. Not sampled.
A11 RSL 68686	(0-38 cm.) 0 to 15 inches, dark grayish brown (10YR 4/2) gravelly coarse sandy loam, very dark brown (10YR 2/2) moist; weak fine crumb structure; soft, very friable, nonsticky, nonplastic; many very fine, fine, medium and coarse roots; many very fine interstitial pores; 14 percent gravel; medium acid (pH 6.0); diffuse wavy boundary.
A12 RSL 68687	(38-53 cm.) 15 to 21 inches, dark brown (10YR 4/3) gravelly coarse sandy loam, very dark grayish brown (10YR 3/2) moist; massive; soft, very friable, nonsticky, nonplastic; many very fine, fine, medium and coarse roots; many very fine inter-

CONSECUTIVE PEDON NUMBER C 68 84 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES TALLAC SHALLOW VARIANT SOIL NO. 568 CALI. 9-15 SAMPLE NO. 8 691-8 696 TAHOE BASIN AREA

SOIL FAMILY DURIC HAPLUMBREPT COARSE-LOAMY,MIXED,FRIGID RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1-	MS .5-	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A	INT. II .02	1A2A COARSE FRAGMENTS PCT. OF WHOLE SOIL	I 75- 20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM															PCT. OF LT 75		
8 691	8- 4	01															--A	--	--	--
8 692	4- 0	02															--A	--	--	--
8 693	0- 13	A1	64.2	31.7	4.1	16.0	12.2	8.3	15.9	11.8	15.7	16.0				36.2	43A	16	27Z	N
8 694	13- 63	AC	64.3	31.1	4.6	11.6	13.9	9.1	16.6	13.1	15.9	15.2				38.4	35A	12	23Z	N
8 695	63- 97	C1	68.5	29.6	1.9	11.4	15.7	10.4	18.9	12.1	15.9	13.7				38.2	37A	12	25Z	N
8 696	97-150	C2	SIM62.9	34.9	2.2	10.8	13.0	8.6	17.1	13.4	19.3	15.6				42.5	42A	12	30Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR-BON 6A1A	NITRO GEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES FE 6C2B	AL 6G7A	FE 6C5A	AL 6G5A	ATTERBERG LL 4F2	PL 4F1	BULK DENSITY 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		PERCENT		PERCENT		PERCENT				PCT.NO.40		GRAMS PER CC				PERCENT			PERCENT	
8 691	8- 4														1.00					
8 692	4- 0														1.00					
8 693	0- 13	4.85	0.204			0.7						1.35			0.73			9.2		
8 694	13- 63	2.17	0.113			0.7						1.40			0.78			5.8		
8 695	63- 97	0.52	0.035			1.0						1.50			0.75			2.7		
8 696	97-150	0.28				0.5						1.70	2.30	2.31	0.69		5.9	2.3	0.1	

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACI-DITY	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 691	8- 4															
8 692	4- 0															
8 693	0- 13	11.3	1.4	0.1	0.6	13.4	11.5		24.9	21.0	54		64		6.0	5.5
8 694	13- 63	4.4	0.5	0.1	0.4	5.4	10.4	0.4	15.8	12.8	34	93	42		5.4	4.9
8 695	63- 97	1.3	0.3	0.1	0.2	1.9	4.5		6.4	5.3	30		36		5.8	5.1
8 696	97-150	4.1	0.8	0.1	0.1	5.1			5.1	6.0	100+		85		5.9	5.2

SAMPLE NO	DEPTH CM.	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM MINERAL X RAY 7A2E I	7A2E II	7A2E III	CODE	RELATIVE AMOUNT
8 691	8- 4											1 TRACE
8 692	4- 0											2 SMALL
8 693	0- 13											3 MODERATE
8 694	13- 63											4 ABUNDANT
8 695	63- 97											5 DOMINANT
8 696	97-150											6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

TALLAC GRAVELLY COARSE SANDY LOAM, SHALLOW VARIANT 1/

S68Calif-9-15

Location: Tahoe Basin, El Dorado County, California. Near center of sec. 10, T. 12 N., R. 18 E., 100 feet west of Fallen Leaf Lake road. Date of Sampling: October 26, 1967.
Description by: G. M. Kennedy. Collectors: J. H. Rogers, R. C. Huff, G. L. Anderson, D. Palmquist.
Classification: Duric Haplumbrept, coarse-loamy, mixed, frigid. This soil has the duripan at slightly shallower depths than Tallac official.
Vegetation: Not cultivated. White fir, Jeffrey pine, incense cedar. Climate: Precipitation is 35 to 40 inches. Mean annual temperature is 41°F., January 26°F., July 60°F. Parent Material: Mixed, predominantly metamorphic alluvium. Topography: Moderately sloping lateral moraine, 8 percent north slope. Elevation: 6500 feet. Drainage: Well drained; runoff is medium; permeability is moderately rapid to duripan which has slow permeability. Soil Moisture: Dry.
Remarks: Field determinations of pH by Hellige-Truog triplex indicator. C2m horizon is compacted till? and is very hard in place, but will disintegrate under pressure. The AC horizon may have had some mixing due to tree falls. Horizons AC (5-25") and C2m (38-60") were sampled for State Division of Highways.

HORIZON	DESCRIPTION
	(8-4 cm.)
01 RSL 68691	3¼ to 1½ inches, not sampled.
	(4-0 cm.)
02 RSL 68692	1½ to 0 inches, not sampled.
	(0-13 cm.)
A1 RSL 68693	0 to 5 inches, dark grayish brown (10YR 4/2) gravelly coarse sandy loam, very dark grayish brown and very dark brown (10YR 3/2, 2/2) moist; strong very fine granular structure; soft, very friable, nonsticky and nonplastic; many very fine, fine, medium and coarse roots; many very fine tubular and interstitial pores; about 30 percent gravel; medium acid (pH 6.0); clear smooth boundary.
	(13-63 cm.)
AC RSL 68694	5 to 25 inches, brown (10YR 4/3) gravelly coarse sandy loam, dark brown (10YR 3/3) moist; moderate fine and medium granular structure; soft, very friable, nonsticky and nonplastic; many very fine, fine, medium and coarse roots; many very fine interstitial, common very fine, fine and medium tubular pores; about 25 percent gravel; medium acid (pH 5.8); clear wavy boundary.
	(63-97 cm.)
C1 RSL 68695	25 to 38 inches, light yellowish brown (2.5Y 6/4) cobbly coarse sandy loam, olive brown (2.5Y 4/4) moist; common large prominent strong brown (7.5YR 5/6) mottles, yellowish red and strong brown (5YR 5/6, 7.5YR 5/6) moist; massive; slightly hard, friable, nonsticky and nonplastic; many very fine, fine, medium and coarse roots; common very fine tubular pores; about 26 percent pebbles and cobblestones; medium acid (pH 6.0); clear wavy boundary.
	(97-150 cm.)
C2sim RSL 68696	38 to 60 inches, light gray (2.5Y 7/2) gravelly coarse sandy loam, grayish brown (2.5Y 5/2) moist; common large prominent brown (7.5YR 4/4) moist mottles; massive; extremely hard, nonsticky and nonplastic; few very fine, fine and medium roots; about 29 percent pebbles and cobblestones; slightly acid (pH 6.2).

1/ Tallac is in a loamy-skeletal, mixed, family of Entic Cryumbrepts. This pedon has a duripan, less than 35 percent coarse fragments by volume, and summer temperatures that are greater than 8° C. (47° F.) at 50 cm. (20 inches).

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Fiddletown gravelly loam (taxadjunct) **LOCATION** Amador County, California

SOIL FAMILY: Pachic Xerumbrept, loamy-skeletal, mixed, mesic.

SOIL NOS. S60Calif-3-27 **LAB. NOS.** 60120 - 60123

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 8	A1	9.0*	8.8*	3.7	7.8	11.3	49.9	9.5	39.0	27.4	44	1-sil
8-22	AC	14.4*	11.8*	4.0	7.3	11.3	43.2	8.0	36.3	23.1	43	1
22-46	C1	11.4*	12.6*	4.6	6.5	14.5	41.5	8.9	38.0	21.6	41	1
46+	R	Rock										

pH		ORGANIC MATTER			ELECTRICAL CONDUCTIVITY		6C1b	MOISTURE TENSIONS
	KCl	6A1a	6B1a	EST% SALT		CaCO3	Free Iron	4B2

FIDDLETOWN GRAVELLY LOAM
S60Calif-3-27

Location: 0.12 mile SE of the center of the NE1/4 of sec. 1, T. 7 N., R. 11 E., [redacted]

200 feet south of the east dam on Bateman Ranch, 3/8 mile east of the ranch house; 200 feet up the slope from the north fork of Deadman Creek.

Date of Sampling: May 6, 1960 Amador County, California

Collectors: Grant Kennedy, Reuben Nelson, Robert Sketchley, Clifford Henry, Paul Nazar, Jack McClelland. Description by: Grant Kennedy.

Vegetation: Bracken fern and annual grasses - dominantly soft chess (area was cleared but has not been cultivated - original vegetation included ponderosa pine, black oak, incense cedar)

Climate: Mean annual precipitation - 33"; average temperatures (Tiger Creek Station): January - 40° F.; July - 70° F.; mean annual - 55° F.

Parent Material: Dark-colored schists of the Calaveras formation, nearly vertical folding

Relief: Steep uplands, about 100 feet above the base of a north-facing slope of 20 percent.

Elevation: 2,160 feet

Drainage: Well drained; moderately slow runoff; moderate permeability

Remarks: This soil is a dark-colored, Ando-like Regosol. The soil was moist when described.

Classification: Pachic Xerumbrept, loamy-skeletal, mixed, mesic.

DESCRIPTION

A1 RSL No. 60120	0- 8"	Dark grayish-brown (10YR 4/2) gravelly loam containing about 25% gravel and 5% cobbles and stones, very dark brown (7.5YR 2/2) when moist; moderate very fine granular structure; soft, friable, slightly sticky and slightly plastic; abundant very fine roots; many very fine interstitial and tubular, few fine tubular pores; medium acid (pH 6.0); clear smooth boundary.
AC RSL No. 60121	8-22"	Brown (10YR 5/3) gravelly loam containing about 25% gravel and 40% cobbles and stones, dark brown (7.5YR 3/3) when moist; otherwise similar to the above except few fine and medium roots; slightly acid (pH 6.2); diffuse smooth boundary.
C1 RSL No. 60122	22-46"	Grayish-brown (10YR 5/2) gravelly loam about 80% of which consists of gravel, cobbles, stones, and rock outcrop, dark brown (10YR 3/3) when moist; moderate fine subangular blocky structure; soft, very friable; slightly sticky and slightly plastic; plentiful very fine, few fine to coarse roots; many very fine interstitial and tubular, few fine tubular pores; no clay films observed even on surfaces of rock fragments; slightly acid (pH 6.5); abrupt very irregular boundary.
R	46"+	Gray (N 5/) dark-colored, fine-grained, hard schist, very

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Fiddletown gravelly loam **LOCATION** Amador County, California

SOIL FAMILY: Pachic Xerumbrept, loamy-skeletal, mixed, mesic

SOIL NOS. S60Calif-3-28 **LAB. NOS.** 60124 - 60127

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 8	A1	9.1*	9.3*	3.8	7.4	11.7	47.9	10.8	40.1	24.3	39	1
8-18	AC	9.8*	8.9*	3.8	8.3	12.4	45.9	10.9	40.5	23.4	30	1
18-45	C1	8.8*	9.5*	4.2	8.6	12.9	44.8	11.2	41.0	22.4	27	1
45+	R	not analysed										

pH H_2O 8C1a 1:1	pH KCl (1 N) 8C1c 1:1	ORGANIC MATTER 1:10	6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	6C1b Free Iron as % Fe_2O_3	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.3	5.3		3.40	0.215	15.8				3.8			10.6
6.2	5.2		1.89	0.133	14.2				4.1			9.6
6.3	5.2		1.13	0.084	13.5				4.0			8.9

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Na 6P2a	K 6Q2a	BASE SAT. % 5C3	Clay mineralogy (<0.002 mm.) Mont.	Verm. X-ray	Mixed layer Mica-Verm. 7A2	Mica	% Kaolin DTA 7A3	% Gibb. DTA 7A3
	milliequivalents per 100g. soil											
23.1	11.7	1.6	13.0	0.2	0.6	52						
18.5	8.1	0.8	10.4	0.1	0.5	48						
13.8	5.7	0.8	8.5	0.1	0.4	45	-	xx	x	-	15	2
							-	x	x	x	35	5

* Many concretions in these fractions.

1/ Percent of 1 inch fraction

- = looked for but not found
tr = detected, very small amounts
x = detected, small amounts
xx = moderate amounts
xxx = large amounts
xxxx = predominant amounts

FIDDLETOWN GRAVELLY LOAM
S60Calif-3-28

Location: 0.15 mile N of the S1/4 corner of sec. 35, T. 8 N., R. 11 E., 120 feet south of the Fiddletown-Silver Lake Road, 0.2 mile east of the Fiddle Hill Ranch; 1.5 miles east of Fiddletown, Amador County, California.
Date of Sampling: May 6, 1960
Collectors: Grant Kennedy, Reuben Nelson, Robert Sketchley, Clifford Henry, Paul Nazar, Jack McClelland Description by: Grant Kennedy.
Vegetation: Black oak, incense cedar, poison oak
Climate: Mean annual precipitation - 32"; average temperatures (Tiger Creek Station): January - 40° F.; July - 70° F.; mean annual - 55° F.
Parent Material: Dark-colored schists of the Calaveras formation with nearly vertical folding
Relief: Steep uplands, in the mid-portion of a north-facing slope of
Elevation: 1,980 feet
Drainage: Well drained; moderately slow runoff; moderate permeability
Remarks: This soil is a dark-colored, Ando-like Regosol. It was moist when described. Classification: Pachic Xerumbrept, loamy-skeletal, mixed, mesic.

HORIZON		DESCRIPTION
O1 & O2	0- 1"	Loose litter of leaves, twigs, etc. partly decomposed in the lower portion.
A1 RSL No. 60124	0- 8"	Dark grayish-brown (10YR 4/2) gravelly (25%) loam, very dark brown (7.5YR 2/2) when moist; moderate very fine granular structure; soft, friable, slightly sticky and slightly plastic; abundant very fine, few fine to medium roots; many very fine interstitial and tubular, few fine and medium tubular pores; slightly acid (pH 6.2); gradual smooth boundary.
AC RSL No. 60125	8-18"	Dark-brown (7.5YR 4/3) gravelly loam (25%) with about 15% cobbles and stones, very dark brown (7.5YR 2/2) when moist; moderate fine granular structure; otherwise similar to the A1; gradual irregular boundary.
C1 RSL No. 60126	18-45"	Similar to above except about 80% of the volume is gravel, cobbles, stones, and rock outcrop; abrupt irregular boundary.
R RSL No. 60127	45"+	Gray (N 5/) somewhat fractured schist, very dark gray (N 3/) when moist; neutral (pH 7.0 - ground rock).

NOTE: The depth to rock varies from 25 to 50 inches, but dominantly ranges around 36 inches.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Pachic Xerumbrept, clayey-skeletal, montmorillonitic, thermic

SOIL Santa Lucia shaly clay loam (taxadjunct) SOIL Nos. S62Calif-42-9 LOCATION Santa Barbara County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 62122 - 62125

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002) **	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)			Coarse fragments 2A2 > 2 Pct.	2-19 (Pct. of <19 mm)	19-76 (Pct. of <19 mm)
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm																
0-8	A11	32.3	30.1	37.6	7.8	9.3	4.0	6.7	4.5	7.7	22.4	15.5	27.8				23	
8-22	A12	27.2	29.1	43.7	3.6	6.4	3.8	7.7	5.7	8.4	20.7	18.2	21.5				18	
22-36	A13	25.8	25.2	49.0	4.6	7.0	3.2	6.2	4.8	7.0	18.2	15.2	21.0				50	
36-46+	R		Rock															

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.	6C2a Ext. Iron as Fe Pct.	Bulk density: 4A1f 1/3 bar clods* g/cc	4A1h Oven-dry clods* g/cc	g/cc		Water content: Pct.	4B1c 1/3 bar clods* Pct.	4B2 15 bar Pct.	4D1 Extensibility COLE in./in.		pH: 8C1a H_2O 1:1	8C1c 1 N KCl 1:1	(1:1)
0-8	10.30	0.693	15			1.3	0.89	0.95				47.9	40.7	0.020		5.5	4.7	
8-22	4.35	0.347	13			1.6	1.02	1.06				54.4	38.0	0.014		4.4	3.4	
22-36	3.32	0.262	13			1.8	1.16	1.18				43.0	40.7	0.007		4.3	3.2	
36-46+	Rock						1.20	1.22				39.2		0.007				

Depth (In.)	Extractable bases 5B1a: 6N2b Ca	6O2c Mg	6P2a Na	6Q2a K		6H2a Ext. Acidity	Cation Exch. Capacity 5A3a Sum									Base saturation 5C3 Sum Pct.	Pct.
	meq/100 g																
0-8	19.7	11.1	0.5	3.7		26.2	61.2									57	
8-22	5.1	0.2	0.7	2.6		38.7	47.3									18	
22-36	3.1	7.8	0.8	1.8		41.5	51.9									20	
36-46+	Rock																

Depth (In.)

* Clods of A12, A13, and R horizons were fragments of shale.

** Shale fragments of sand and silt size break down in particle size distribution analysis. The soil has the chemical properties of a material of higher clay content and the mechanical properties of one of lower clay content.

SANTA LUCIA SHALY CLAY LOAM (taxadjunct) 1/
S62Calif-42-9

Location: Santa Barbara County, California. SE¼ Sec. 4, T9N, R35W. Very near watershed divide in Casmalia Hills, reached by trail road east southeast out of Corralitos Ranch headquarters; 5.4 airline miles south and slightly east of Guadalupe. Date of Sampling: April 19, 1962.
Description by: L. C. Leifer. Collectors: R. E. Nelson, Lothair Grant, G. E. Shipman, C. J. Hilliker, L. C. Leifer.
Classification: Pachic Xerumbrept, clayey-skeletal, montmorillonitic, thermic.
Vegetation: Grass-oak used for range; mostly ripgut brome, a little bur clover and a little Spanish clover, 50 percent coast live-oak canopy, sample site in small opening in trees. Climate: Precipitation 16 inches. Mean annual temperature 57°F., January 49°F., July 63°F. Parent Material: Diatomaceous shale, Monterey. Topography: Generally steep hills, locally smooth slightly convex 25 percent east slope. Elevation: 1400 feet. Drainage: Well drained. Runoff is medium. Permeability is moderate. Soil Moisture: All is nearly dry and moderately moist below.
Remarks: Field determinations of pH by Hellige-Truog triplex indicator. Paired site sample numbers are Santa Lucia S62Calif-42-10-1 through 4.

HORIZON	DESCRIPTION
A11 RSL 62122	0 to 8 inches, very dark gray (10YR 3/1d, 3/1m) shaly clay loam; moderate medium subangular blocky structure, upper 1" strong fine and medium granular on 3/4 surface, where granular lacking due to trampling; hard, friable, sticky and plastic; abundant very fine roots and plentiful medium roots; many very fine and few fine interstitial pores; medium acid (pH 6.0); gradual smooth boundary.
A12 RSL 62123	8 to 22 inches, very dark grayish brown (10YR 3/2) very shaly clay loam, very dark brown (10YR 2/2) when moist; 65 percent by volume of shale fragments larger than 3/4" diameter; strong fine granular structure; slightly hard, friable, sticky and plastic; plentiful very fine and medium, and few fine and coarse roots; many very fine and few fine interstitial pores; very strongly acid (pH 5.0); gradual wavy boundary.
A13 RSL 62124	22 to 36 inches, very dark grayish brown (10YR 3/2) very shaly clay loam, very dark brown (10YR 2/2) when moist; 75 percent by volume of rock fragments larger than 3/4" diameter; strong fine granular structure; slightly hard, friable, sticky and plastic; few very fine, fine, and coarse and plentiful medium roots; many very fine and few fine interstitial pores; very strongly acid (pH 4.5); abrupt irregular boundary.
R RSL 62125	36 to 46 inches + very pale brown (10YR 8/3d) shale, pale brown (10YR 6/3) when moist; shale is fractured and joints contain up to 5 percent A1 soils in joints to depth of 46 inches, less than 1 percent soil below 46 inches.

NOTE: Rock fragments larger than 3/4" discarded from samples going to Riverside Laboratory.

1/ The Santa Lucia series is in a clayey-skeletal, mixed, thermic family of Pachic Ultic Haploxerolls. This pedon has an umbric epipedon with a base saturation percentage of 18 and a montmorillonitic control section.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Pomponio fine sandy loam variant **LOCATION** San Mateo County, California

SOIL FAMILY: Xeric Argialboll, fine, montmorillonitic, mesic

SOIL NOS. S55Ca1.41-3 **LAB. NOS.** 55110-55116

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0- 7	A_{11}	0.4	3.6	13.7	31.6	7.0	25.7	18.0	32.1	12.7	0	fsl
7-13	A_{12}	0.3	2.9	10.9	28.0	8.6	30.2	19.1	37.0	13.6	0	l
13-18	A_{21}	0.2	2.7	9.2	25.8	9.6	36.8	15.7	42.7	15.4	0	l
18-26	A_{22}	0.3	2.8	9.2	26.3	10.5	34.5	16.4	40.6	16.3	0	l
26-37	B_{21}	0.2	1.2	3.0	11.5	7.7	30.0	46.4	29.8	14.3	0	c
37-52	B_{22}	0.1	1.1	2.8	10.9	7.8	32.0	45.3	30.5	15.4	0	c
52-62	C	0.2	1.5	3.6	11.6	7.2	36.9	39.0	31.9	18.5	0	cl

pH 8C1b SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	Free Fe_2O_3 6C1b %	GYPSUM me./100g. SOIL	4A3a Bulk Density g/cc.		4B2 Soil water 15 atmos. Pct.
5.7		6.5	3.12	.238	13.1		0.4	1.0		1.26		8.6
5.7		6.6	2.13	.174	12.2		0.4	1.1				6.7
6.0		6.9	0.67	.071	9.4		0.3	1.1		1.56		9.0
6.4		7.2	0.44				0.3	1.1				7.1
6.0		7.1	0.39				1.1	1.6		1.43		22.6
5.6		6.5	0.30				2.8	1.3		1.55		22.0
5.7		6.8	0.20				2.8	1.6		1.60		18.6

5A2a CATION EXCHANGE CAPACITY (+ Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	5B Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg	6M1a HCO_3 1/	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
24.5	9.0	5.1	11.8	0.7	0.3	56	2.5	0.1	2.0	1.4	1.3	36.7
25.3	7.8	5.3	9.1	0.7	0.2	61	2.5	0.1	1.8	1.4	0.4	31.1
23.6	4.4	4.0	6.5	0.5	0.1	58	2.0	<0.1	0.7	0.3	1.3	22.2
14.8	4.2	4.3	3.7	0.7	0.1	72	2.5	<0.1	0.5	0.4	0.7	24.3
36.3	10.5	15.3	4.4	2.4	0.5	91	7.4	<0.1	2.2	0.3	8.8	93.1
38.6	9.9	15.2	4.5	3.2	0.5	86	15.3	0.1	7.4	0.2	24.1	82.9
31.8	11.4	13.2	3.5	3.2	0.4	89	19.7	0.1	9.8	0.2	31.3	64.5

1/ no $CO_3^{=}$ found

POMPONIO FINE SANDY LOAM variant
S55Calif-41-3

Location: San Mateo County, California. On Deeney Ranch, about 3/4 mile east of California highway No. 1. 75 feet west of old picket fence on east edge of pasture, 1/3 mile east of abandoned farm buildings. Date of Sampling: February 23, 1955.

Description by: W. M. Johnson. Collectors: Reuben Nelson, Ralph Nelson, Richard Wagner, Melvin Mello, and W. M. Johnson.

Classification: Xeric Argialboll, fine, montmorillonitic, mesic.

Vegetation: Annual fescue, rye grass, a little bur clover and scattered blackberry. Land Use: Pasture that apparently was once cultivated. Climate: Mean annual rainfall estimated at about 23 inches. June, July, and August are very dry. Fogginess is high in many parts of the year, particularly during summer and fall. Average January temperature is about 48° F., average July temperature about 60° F. On-shore winds carrying salt spray are common. Parent Material: Appears to be clayey sediments of the coastal plain. May be weathered Purissima formation with some local alluvium in the upper layers. Relief: In hilly and steep upland, a slightly concave slope in "cove" position, about 150 feet downslope from crest of hill. Slope about 17 percent, gently concave, facing west. Elevation: About 435 feet. Drainage: Runoff rapid. Permeability moderate in A horizon, very slow in B horizon.

Remarks: A horizon wet, B and upper C horizons moist, lower C slightly moist at time of sampling. Erosion at this site appears to be negligible.

HORIZON		DESCRIPTION
A_{11}	0-7"	Very dark gray (10YR 3/0.8, dry; 2/0.8, moist) fine sandy loam. Weak coarse and medium granular. Slightly hard, friable, slightly plastic, nonsticky. Roots very numerous. Peds slightly porous. pH 5.7. Clear, wavy lower boundary.
A_{12}	7 - 13"	Light loam or heavy fine sandy loam of the same color as the A_{11}. Weak very coarse, coarse, and medium granular. Hard, friable, slightly plastic, slightly sticky. Roots very numerous. Peds slightly porous. pH 5.5. Clear, wavy lower boundary.
A_{21}	13 - 18"	Finely mottled dark gray and gray (10YR 4/1 and 6/1, dry; 2/1 and 4/1, moist) loam. Weak medium and fine subangular blocky. Hard, friable, plastic, slightly sticky. Roots numerous. Moderately porous. pH 5.8. Clear, wavy lower boundary.
A_{22}	18 - 26"	Gray (10YR 5/1, dry; 3.5/1.5, moist) loam with common very fine mottles of light gray and few fine mottles of yellowish brown (dry). Weak medium subangular blocky. Very hard, friable, plastic, slightly sticky. Roots numerous. Moderately porous. pH 6.1. Abrupt, wavy lower boundary.
B_{21}	26 - 37"	Very finely mottled light brownish gray and yellowish brown (2.5Y 6/2 and 10YR 5/6, dry; 2.5Y 5/2 and 10YR 4/4, moist) clay. Moderate coarse prismatic, breaking to moderate medium blocky. Extremely hard, very firm, very plastic, very sticky. Moderate continuous clay skins. Few roots, mainly along cracks. Peds nonporous. pH 6.8. Clear, wavy lower boundary.
B_{22}	37 - 52"	Finely, indistinctly mottled light brownish gray and light yellowish brown (2.5Y 6/2 and 1Y 6/4, dry; 2.5Y 5/0 and 1Y 5/4, moist) clay. Moderate medium

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Pomponio fine sandy loam variant **LOCATION** San Mateo County, California

SOIL FAMILY: Xeric Argialboll, fine, montmorillonitic, mesic

SOIL NOS. S55Ca1.41-4-1 thru 8 **LAB. NOS.** 55117-55124

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0- 7	A_{1p}	0.1	0.8	2.2	34.0	18.4	24.7	19.8	58.9	12.2	0	fsl
7-13	A_{12}	0.0	0.7	2.6	33.8	18.2	25.8	18.9	58.5	13.4	0	fsl
13-17	A_{21}	0.0	0.8	2.8	34.7	18.8	25.2	17.7	61.0	11.4	0	fsl
17-21	A_{22}	0.1	1.1	3.1	36.0	20.3	25.8	13.6	63.5	11.9	0	fsl
21-30	B_{21}	0.0	0.5	1.8	25.6	14.7	20.2	37.2	45.0	10.9	0	cl
30-41	B_{22}	0.0	0.5	1.5	26.4	13.0	22.3	36.3	46.5	10.6	0	cl
41-50	B_{23}	0.0	0.4	1.6	30.3	13.1	21.8	32.8	50.8	8.7	0	scl
50-60	C_1	0.0	0.4	1.8	32.3	13.5	24.0	28.0	53.0	10.6	0	scl

8C1b pH SATURATED PASTE 1:1	8C1a pH 1:5	8C1a pH 1:10	6A1a ORGANIC MATTER ORGANIC CARBON %	6B1a ORGANIC MATTER NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C	Free Fe_2O_3 6C1b %	GYPSUM me./100g. SOIL	4A3a Bulk Density g/cc		4B2 Soil wat r 15 atmos. Pct.

POMPONIO FINE SANDY LOAM variant 1/
S55Calif-41-4

Location: About 3/4 mile south of Tunitas Creek, 300 yards east of Highway No. 1 on south side of Star Hill Road, San Mateo County, California. See Sheet 21, Soil Survey Report of San Mateo Area, California, 1961. Date of Sampling: February 24, 1955.

Description by: W. M. Johnson. Collectors: Reuben E. Nelson, Ralph E. Nelson, Richard Wagner, Melvin Mello, and W. M. Johnson.

Classification: Xeric Argialboll, fine, montmorillonitic, mesic.

Vegetation: Formerly cultivated, now in grass pasture. Annual fescue, ryegrass, annual weeds, iris, geranium, and wild buckwheat. Climate: Mean annual rainfall estimated at about 23 inches. June, July and August are very dry. Fogginess is high in many parts of the year, particularly during summer and fall. Average January temperature is about 48° F., average July temperature about 60° F. On-shore winds carrying salt spray are common. Parent Material: Clayey coastal plains sediments. May have some local alluvium in the upper part of the profile. Relief: Hilly and steep upland, in eroded marine terrace breaks. Site about 150 feet down slope from the crest of a hill. Slope about 14 percent, slightly concave, toward the west. Elevation: About 475 feet. Drainage: Runoff is rapid. Permeability moderate in A horizon, very slow in B horizon.

Remarks: Soil appears to have suffered little or no erosion of A horizon. At time samples were taken, soil was wet in the A horizon, moist below.

DESCRIPTION

A_{1p}	0 - 7"	Very dark gray (10YR 3/1, dry; 2/1, moist) fine sandy loam. Weak coarse and medium granular. Slightly hard, friable, slightly plastic, nonsticky. Roots very numerous. Slightly porous. Many worm casts. pH 5.2. Abrupt smooth lower boundary.
A_{12}	7 - 13"	Fine sandy loam of the same color as the A_{1p}. Weak coarse and medium granular. Hard, friable, slightly plastic, nonsticky. Roots very numerous. Peds are moderately porous. Very many worm casts. pH 5.5. Clear, slightly wavy lower boundary.
A_{21}	13 - 17"	Very dark gray (10YR 3.5/1, dry; 2/1, moist) fine sandy loam with common medium and fine mottles of light gray (dry). Weak coarse and medium granular. Hard, friable, slightly plastic, nonsticky. Roots are numerous. Peds are moderately porous. pH 5.5. Clear, wavy lower boundary.
A_{22}	17 - 21"	Dark gray (10YR 4/1, dry; 2.5/1, moist) fine sandy loam with common medium and fine mottles of light gray (dry). Weak medium and fine subangular blocky. Hard, friable, slightly plastic, nonsticky. Roots common. Moderately porous. pH 5.7. Abrupt, wavy lower boundary.
B_{21}	21 - 30"	Dark grayish brown (10YR 4/1.5, dry; 3/2, moist) clay loam. Moderate coarse prismatic, breaking to moderate medium blocky. Extremely hard, firm, very plastic, sticky. Moderate continuous clay skins. Has a few conspicuous white spots that are probably gypsum. Very few roots. Peds are nonporous. pH 6.2. Clear, wavy lower boundary.
B_{22}	30 - 41"	Light olive brown (2.5Y 5/3, dry; 4/3, moist) clay loam with common fine black mottles and a few conspicuous, medium, white spots (probably gypsum). Moderate medium blocky. Extremely hard, firm, very plastic, sticky. Moderate continuous clay skins. Very few roots. Peds are nonporous. pH 6.5. Clear, wavy lower boundary.
B_{23}	41 - 50"	Finely mottled light brownish gray and yellowish brown (2.5Y 6/2 and 10YR 5/6, dry; 2.5Y 5/2 and 10YR 4/6, moist) sandy clay loam with a few conspicuous medium spots of white. Weak coarse blocky. Extremely hard, firm, very plastic, sticky. Moderate continuous clay skins. Very few roots. Peds are nonporous. Contains [illegible]
C_1	50 - 60"+	Finely mottled light brownish gray and yellowish brown (2.5Y 6/2 and 10YR 5/4, dry; 2.5Y 5/2 and 10YR 4/4, moist) sandy clay loam. Very weak coarse blocky, almost massive. Very hard, firm, very plastic, sticky. Thin [illegible]

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Watsonville loam (taxadjunct) **LOCATION** San Mateo County, California

SOIL FAMILY: Abruptic Argiaquoll, fine, montmorillonitic, mesic

SOIL NOS. S55Ca1. 41-11 **LAB. NOS.** 55380-55386

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 2/	TEXTURAL CLASS
0- 9	A_{11}	2.4	3.4	3.8	12.3	15.4	47.1	15.6	49.3	21.3	10	l
9-14	A_{12}	2.7	3.2	4.0	11.6	15.6	46.7	16.2	49.5	20.4	10	l
14-24	B_{21}	1.4	3.0	3.1	9.3	11.4	30.0	41.8	32.6	14.9	11	c
24-33	B_{22}	1.5	3.2	3.4	11.2	12.4	31.5	36.8	37.5	13.8	4	cl
33-44	B_3	2.5	6.1	5.8	12.3	9.8	25.5	38.0	29.6	13.1	10	cl
44-54	C_{11}	3.6	17.0	16.4	14.7	5.2	17.3	25.8	18.1	10.9	23	scl
54-60	C_{12}	0.1	0.3	0.5	1.4	1.0	39.6	57.1	11.0	30.5	0	c

8C1b pH SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	4A3a Bulk Density g./cc.	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
5.3		5.9	3.82	.259	14.7	1.22	0.7					10.3
5.5		6.2	1.46	.109	13.4		0.7					6.9
5.0		6.3	0.80	.075	10.7	1.56	0.9					17.7
4.8		5.9	0.56				2.3					15.0
4.6		5.4	0.38				4.7					16.0
4.4		5.3	0.26			1.58	7.8					10.9
4.3		5.2	0.23				5.8					25.8

5A2a CATION EXCHANGE CAPACITY ←(Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg	6J1a HCO_3 1/	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
30.6	6.1	4.5	10.2	1.0	0.4	54	4.8	0.3	4.2	1.8	3.5	48.6
16.4	4.1	4.2	6.6	0.9	0.3	59	5.0	0.2	2.5	1.7	2.5	30.3
35.3	7.3	11.5	7.7	2.5	0.6	74	6.2	0.1	1.3	0.7	4.8	60.0
29.6	7.6	11.4	7.1	3.1	0.4	76	16.8	0.1	5.4	0.4	21.0	53.8
37.6	6.7	13.1	7.3	3.9	0.4	77	33.8	0.3	12.7	0.1	47.8	63.3
23.7	5.3	8.3	6.4	3.0	0.3	73	52.0	0.2	21.2	0.2	24.8	39.9
51.1	11.1	15.0	9.7	25.3	0.5	84 2/	42.0	0.1	11.5	0.1	55.9	89.7

1/ No CO_3 found

2/ Percent of the whole soil

WATSONVILLE LOAM (taxadjunct) 1/
S55Calif-41-11

Location: San Mateo County, California in the San Mateo County SCD. 1.1 miles south and 0.8 miles west of Half Moon Bay on bluff overlooking the Pacific Ocean. Date of Sampling: May 19, 1955.
Description by: Ralph E. Nelson. Collectors: Reuben E. Nelson, Ralph E. Nelson, Rudolph Ulrich, R. J. Wagner.
Classification: Abruptic Argiaquoll, fine, montmorillonitic, mesic.
Vegetation: Annual grasses; soft chess, wild oat; California oat grass (Danthonia Californica), lupine, wild strawberry, coyote brush

nearly level. Elevation: 70 feet. Drainage: Surface drainage very slow; internal drainage very slow, restricted due to claypan.
Remarks: Few worms in the A11 horizon.

OIL SURVEY LABORATORY Riverside, California

SOIL TYPE Watsonville loam (taxadjunct) **LOCATION** San Mateo County, California

SOIL FAMILY: Abruptic Argiaquoll, fine, montmorillonitic, mesic

SOIL NOS. S55Cal. 41-12 **LAB. NOS.** 55387-55393

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 2/	TEXTURAL CLASS
0- 9	A_{11}	0.9	2.2	3.1	14.9	11.7	49.9	17.3	62.3	10.2	2	1
9-12	A_{12}	1.0	2.1	3.1	14.2	16.7	43.4	19.5	52.3	18.1	4	1
12-21	B_{21}	0.7	1.7	2.4	8.8	12.2	30.4	43.8	36.8	11.8	2	c
21-36	B_{22}	0.7	1.7	2.4	8.9	12.8	31.5	42.0	36.5	13.9	3	c
36-46	B_{23}	0.4	1.5	1.9	7.5	13.2	36.7	38.8	39.3	15.9	1	cl
46-54	B_3	0.0	0.4	0.5	4.3	19.6	41.5	33.7	47.9	16.8	0	cl
54-64	C	1.8	5.5	5.4	16.9	17.4	30.9	22.1	45.9	13.7	8	1

8C1b pH SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	4A3a Bulk Density g./cc.	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
5.4		6.1	2.53	.185	13.7	1.45	1.2					8.6
5.5		6.3	1.87	.146	12.8		0.7					8.6
5.2		6.4	0.95	.097	9.8	1.58	0.8					17.9
5.0		6.0	0.65				2.4					17.6
4.7		5.9	0.48				4.1					16.6
4.7		5.7	0.28				4.3					16.0
4.7		5.8	0.13			1.62	4.1					11.0

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg	6J1a HCO_3 1/-	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
20.5	5.2	4.8	9.1	1.0	0.5	56	7.7	0.3	4.1	2.1	6.4	37.7
18.9	6.2	5.1	8.2	1.2	0.2	61	5.9	0.1	1.8	2.3	2.5	36.2
35.9	7.7	13.9	8.6	2.2	0.4	74	5.8	0.1	1.4	0.7	4.2	58.2
35.2	6.1	10.4	8.0	3.5	0.4	72	8.1	0.1	4.7	0.4	20.0	62.6
35.7	6.5	12.6	8.0	4.5	0.4	75	29.8	0.1	6.6	0.3	33.2	58.0
33.8	4.4	9.3	6.9	4.2	0.2	72	34.0	0.1	5.7	0.4	36.6	61.5
24.6	6.5	12.7	4.2	6.9	0.4	86	32.3	0.2	4.6	0.2	34.9	46.3

1/ No CO_3 found

2/ Percent of the whole soil

3/ ESP 28

WATSONVILLE LOAM (taxadjunct) 1/
S55Calif-41-12

Location: San Mateo County, California in the San Mateo County SCD, 100 feet north of the end of Rose Avenue at ocean. Just off edge of gully.
Date of Sampling: May 19, 1955.
Description by: Ralph E. Nelson. Collectors: Reuben E. Nelson, Ralph E. Nelson, Rudolph Ulrich, and R. J. Wagner.
Classification: Abruptic Argiaquoll, fine, montmorillonitic, mesic.
Vegetation: Annual grasses; soft chess, wild oat, and California oat grass (Danthonia Californica), lupine, wild strawberry, coyote brush, Indian soap plant. Climate: 22 inches rainfall. Parent Material: Marine sediments. Relief: First terrace above ocean, smooth and nearly level. Elevation: 70 feet. Drainage: Surface drainage very slow; internal drainage very slow, restricted due to claypan. Remarks: Few worms in the A11 horizon.

HORIZON		DESCRIPTION
A_{11}	0-9"	Gray (10YR 5/1, very dark brown 2/2 moist) loam. Weak, fine subangular blocky. Very hard, friable, very slightly plastic and nonsticky. There are fine rock particles of siliceous material scattered throughout this horizon and throughout the entire profile. There are a few fine iron and manganese shot concretions. Many roots and few worm casts. The reaction is 6.0. The lower boundary is clear and smooth.
A_{12}	9-12"	Gray (10YR 5/1, very dark brown 2/2 mottled with yellowish brown 10YR 5/6 and 5/8 moist) loam. Weak, fine subangular blocky. Hard, friable, very slightly plastic and nonsticky. There are many fine manganese and iron shot concretions and there is a whitish siliceous coating just above the horizon below. The reaction is 6.0. The lower boundary is abrupt and smooth.
B_{21}	12-21"	Very dark grayish brown (10YR 3/2, dark brown and very dark brown 10YR 3/3-2/2 moist) clay. Strong medium to coarse angular blocky. Extremely hard, very firm, very plastic and sticky. Thick continuous clay skins which are nearly black in color (10YR 2/1 moist). Some fine iron and manganese shot. Few roots down the cracks between the structural aggregates. The reaction is 5.8. The lower boundary is clear and smooth.
B_{22}	21-36"	Prominently mottled gray brown (2.5Y 5/2) and brownish yellow (10YR 6/6, 6/8, very dark grayish brown 2.5Y 3/2 and yellowish brown 10YR 5/6-5/8 moist) clay. Moderate medium angular blocky. Extremely hard, very firm, very plastic and sticky. The clay skins are nearly black (10YR 2/1) and very dark grayish brown (2.5Y 3/2) and are thinner and less continuous than in the horizon above. There are a few roots going down cracks between structural aggregates. The reaction is 5.5. The lower boundary is clear and smooth.
B_{23}	36-46"	This horizon is prominently mottled dark grayish brown and grayish brown (2.5Y 4/2, 5/2; very dark grayish brown and dark

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Logan silt loam **LOCATION** Lassen County, California

SOIL FAMILY: Typic Calciaquoll, fine-silty, mixed, mesic.

SOIL NOS. S55-Cal. 18-5 **LAB. NOS.** 55516-523

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-6	A1p	0.6	2.0	2.0	9.7	10.7	46.5	28.5	30.4	34.0	0	cl
6-9	A12	0.4	1.6	1.6	8.9	11.0	47.4	29.1	31.1	33.9	0	cl
9-16	A13	0.3	1.2	1.6	9.3	10.6	46.5	30.5	30.4	33.7	0	cl
16-22	Cca	0.2	1.1	1.4	8.3	9.5	44.5	35.0	28.6	31.6	0	cl
22-35	C1	0.6	1.9	2.0	8.6	10.2	48.7	28.0	29.8	35.4	0	cl
35-42	C2	2.1	3.0	2.2	7.8	9.0	52.6	23.3	29.7	37.4	2	sil
42-53	C3	1.8	2.3	1.3	6.1	9.2	57.3	22.0	28.7	42.3	3	sil
53-65+	C4	1.0	2.3	2.1	10.9	14.5	53.7	15.5	40.4	35.8	7	sil

8C1b pH SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8C1a Elec. Condc. EC×10³ MILLIMHOS PER CM @25°C.	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
7.8		8.4	2.88	.258	11.2		1.0	2	-			20.1
7.9		8.4	2.42	.222	10.9		0.8	2	-			20.9
7.9		8.5	1.43	.164	8.7		0.9	5	-			22.3
7.9		8.6	0.76				0.7	30	-			21.4
7.9		8.5	0.47				1.5	26	-			19.4
7.8		8.5	0.32				0.7	33	-			18.0
7.7		8.4	0.30				0.6	30	-			26.1
7.7		8.5	0.24				0.6	36	-			20.3

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	5D1 EXCH. Na %	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg		6K1a	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						milliequivalents per liter →					
34.7				1.2	4.0	3	2.3	1.0	6.8			62.2
35.0				2.1	3.6	6	2.9	0.7	5.9			64.6
34.9				1.8	3.3	5	4.4	0.6	5.7			59.1
27.0				1.1	1.9	4	3.4	0.4	3.9			62.7
27.1				1.1	1.6	4	5.4	0.4	9.4			55.3
27.1				1.3	1.1	5	2.6	0.2	3.9			52.4
35.3				1.9	1.1	5	2.6	0.2	3.7			55.4
35.2				0.9	1.0	3	2.5	0.2	3.9			58.5

1/ Percent of the whole soil

LOGAN SILT LOAM
S55Calif-18-5

Location: Lassen County, California. About 6 miles southeast of Susanville. 150 feet west and 20 feet south of the east quarter corner of Sec. 18, T. 29 N., R. 13 E., Honey Lake Valley SCD. Date of Sampling: August, 1955.
Description by: W. M. Johnson. Collectors: S. B. Johnson, Ralph E. Nelson, Stuart

Classification: Typic Calciaquoll, fine-silty, mixed, mesic.
Vegetation: Judging by fence rows, original vegetation probably includes salt grass, greasewood, rabbitbrush, giant wild rye, and horsetail rush. Land Use: Cultivated, irrigated alfalfa field. Climate: At Susanville the January average temperature is 30.7° F., the July average is 69.5° F., and the frost free period averages 142 days. Summers are hot and dry, winters are cold and snowy. Mean annual rainfall at Susanville is 17.8 inches, most of which falls during the period October to May. Rainfall at this site is estimated to be about 14 inches per year.
Parent Material: Calcareous silty and clayey alluvium, including reworked Lake Lahontan sediments and alluvium from the mountains lying to the south.
Relief: Smooth, nearly level basin floor of old Lake Lahontan. Slope less than 1/2 percent. Elevation: About 4,100 feet. Drainage: Runoff slow or very slow (water must pond sometimes). Permeability moderate down to the water table. Water table sometimes rises to the surface, dropping to below 6 feet at other times. Imperfectly drained.
Remarks: Soil is moist below the upper 1 or 2 inches down to about 36 inches; below 36 inches it is wet at this time.

HORIZON		DESCRIPTION
A_{1p}	0 - 6"	Gray (10YR 5.5/1, dry; 3/1, moist) silt loam. Moderate coarse and medium granular. Very hard, friable, plastic, sticky. Roots are profuse and well distributed. Many worm casts and burrows. Moderately calcareous.- pH 8.0. Abrupt, smooth lower boundary.
A_{12}	6 - 9"	Gray (10YR 5/1, dry; 3/1, moist) silt loam. Weak coarse prismatic, breaking to weak coarse and medium granular. Very hard, friable, plastic, sticky. Roots profuse and well distributed. Many worm casts and burrows. Moderately calcareous. pH 8.2. Clear, smooth lower boundary.
A_{13}	9 - 16"	Gray (10YR 5/1.5, dry; 2.5/2, moist) silty clay loam with common white specks and common medium and coarse mottles of grayer material mixed from the horizon above by worm activity. Weak, coarse prismatic structure, the peds being riddled with worm burrows and containing many medium sized worm casts. No clay skins observed. Roots very numerous. Moderately calcareous; lime is mostly in specks and threads. pH 8.4. Clear, wavy lower boundary.
C_{ca}	16 - 22"	Light gray (1Y 7/1, dry; 4/2, moist) silty clay loam. Massive. Friable, plastic, very sticky. Roots are numerous. One krotovina observed. Common worm casts and worm burrows. Very strongly calcareous, the lime being

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haplaquoll, fine-loamy, mixed, noncalcareous, mesic

SOIL Ramelli silty clay loam SOIL Nos. S61Calif-32-13 LOCATION Sierra Valley, Plumas County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 61686 - 61694

Depth (in.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	3A1 Int. II (0.2-0.02)	(2-0.2)			Coarse fragments 2A2 > 2 Pct.	2 - 19	19 -76
		Pct. of < 2 mm															Pct. of < 76 mm	
0-2	A11	14.4	36.7	48.9	1.1*	2.2*	1.8*	6.3	3.0	9.1	27.6	16.1	7.4			0		
2 -7	A12	22.4	34.4	43.2	0.9	2.4	2.5	9.8	6.8	8.8	25.6	22.0	9.2			0		
7-14	B21	21.7	31.2	47.1	0.9	2.3	2.3	8.8	7.4	7.7	23.5	21.1	8.3			0		
14-21	B22	34.0	31.8	34.2	0.9	3.1	4.0	14.8	11.2	12.4	19.4	33.7	12.7			0		
21-27	IIC1	42.5	32.9	24.6	0.8	3.4	5.8	17.9	14.6	11.4	21.5	37.5	16.4			0		
27-37	IIIC2g	60.7	26.2	13.1	1.5	3.3	4.3	31.5	20.1	11.2	15.0	54.8	17.1			1		
37-42	IIIC3g	67.1	22.5	10.4	0.7	3.1	3.2	37.6	22.5	11.4	11.1	62.8	15.7			0		
42-47	IVC4g	82.0	8.1	9.9	1.7	9.9	23.3	40.0	7.1	3.4	4.7	29.2	56.2			3		
47-64+	VC5g	12.8	36.6	50.6	0.2	1.3	1.6	4.7	5.0	9.4	27.2	17.5	4.7			0		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		6E1b Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	4B2 15 bar Pct.			pH 8C1b Sat. Paste	8C1a 1:10
0-2	9.63	0.662	14		-								29.4			6.4	6.9
2 -7	3.35	0.293	11		-								23.1			6.3	7.2
7-14	1.07	0.108	10		-								23.2			6.8	7.8
14-21	0.42	0.043	10		-								19.1			7.1	8.0
21-27	0.12				tr.								15.9			7.4	8.3
27-37	0.10				-								13.2			7.4	8.1
37-42	0.08				-								12.1			7.0	7.6
42-47	0.07				-								8.4			7.0	7.4
47-64+	0.06				-								24.8			6.9	7.9

Depth (in.)	Extractable bases 5B1a: 6N2c Ca	602c Mg	6P2a Na	6Q2a K	6H2a Ext. Acidity	Cat. Exch. Capa. 5A2a NaOAc		5B1b Exch. Na	Water extract from saturated paste 8A1: 6N1a Ca	601a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-2	29.3	10.7	0.7	1.0	8.2	51.8												
2 -7	23.4	9.5	0.6	0.4	4.4	40.6												
7-14	25.4	11.9	0.7	0.3	3.0	40.4		0.6	2.7	2.0	1.9	-		-	4.7	0.2		0.56
14-21	21.0	10.4	0.7	0.3	1.9	33.6												
21-27	20.8	10.3	0.8	0.2	1.3	31.2												
27-37	15.9	8.2	0.7	0.1	1.3	26.5												
37-42	13.9	7.4	0.6	0.1	1.8	24.4												
42-47	9.7	5.1	0.4	0.1	1.2	17.6												
47-64+	23.4	14.2	0.7	0.4	2.3	39.7												

Depth (in.)	8A Water at Saturation Pct.	5C3 Base Sat. (Sum) Pct.	Clay Mineralogy (<0.002 mm.) X-ray: Montmorillonite	Vermiculite	Mica	Kaolinite	Kaolinite % DTA
0-2		84	xxx	-	-	t	3
2 -7		88					
7-14	69.8	93					
14-21		94	xxx	x	-	x	8
21-27		96					
27-37		95					
37-42		92					
42-47		92					
47-64+		94					

* Sands are mostly organic matter

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

RAMELLI SILTY CLAY LOAM

S61Calif-32-13

Location: Sierra Valley, Plumas County, California. SE corner of the SW1/4 of the NW1/4 of section 35, T. 22 N., R. 15 E., 150 feet south of Baldock loam sample site, Sample No. S61Calif-32-12. Approximately 2 3/4 miles north and 1 3/4 miles west of Loyalton. Date of Sampling: September 14, 1961.
Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, James McLaughlin, Eugene Begg.
Vegetation: Mostly sedges and bluegrass. Climate: Estimated annual precipitation 13 inches. Nearest weather station (Portola): Mean annual temperature 45.1° F.; January average 28.7° F.; July average 62.6° F.; mean annual precipitation 20.62 inches. Parent Material: Alluvium from mixed sources.
Relief: Bottomland areas; continuous depressed areas between mounds with a smooth slope of less than 1 percent.
Elevation: 4888 feet. Drainage: Imperfectly drained; runoff very slow; permeability moderately slow.
Soil Moisture: Soil is moist at 2½ inches with moisture increasing with depth.
Remarks: This soil is a Humic Gley soil with a structural B horizon. This site was sampled in the driest part of an abnormally dry year. In wetter years it is expected the water table might stand at about

42 inches. The VC5g horizon has manganese shot and manganese stains. The VC5g appears to be old fine-textured lake sediments. The A12 (2½ to 7 inches), the B21 (7 to 14 inches), and the IIIC3g (37 to 42 inches) were sampled for the Bureau of Public Roads and the California State Division of Highways.
Classification: Typic Haplaquoll, fine-loamy, mixed, noncalcareous, mesic.

HORIZON	DESCRIPTION
A11 RSL No. 61686	0 to 2½ inches, gray (10YR 5/1) silty clay loam, black (10YR 2/1) when moist; moderate very fine platy structure; soft, friable, sticky and slightly plastic; abundant very fine and few fine to medium roots; many very fine tubular and interstitial, few fine to medium tubular pores; slightly acid (pH 6.5); clear, smooth boundary.
A12 RSL No. 61687	2½ to 7 inches, very dark gray (10YR 3/1) silty clay loam, black (N 2/) when moist; moderate fine to medium subangular blocky structure; slightly hard, friable, sticky and plastic; abundant very fine to medium roots; many very fine tubular and interstitial, common fine to medium tubular pores; few thin clay films in pores; neutral (pH 7.0); clear, smooth boundary.
B21 RSL No. 61688	7 to 14 inches, dark gray (N 4/) silty clay, black (N 2/) when moist; strong medium prismatic to strong medium angular blocky structure; very hard, firm, very sticky and plastic; abundant very fine common fine to medium mostly exped roots; common very fine and few fine to medium tubular pores; continuous thin clay films in pores and on ped faces; moderately alkaline (pH 7.9); gradual, smooth boundary.
B22 RSL No. 61689	14 to 21 inches, gray (10YR 5/1) clay loam (near clay), very dark gray (10YR 3/1) when moist; strong fine to medium prismatic breaking to strong fine to medium angular blocky structure; very hard, firm, sticky and plastic; plentiful very fine to medium mainly exped roots; common very fine and few fine to medium tubular pores; many thin clay films in pores and on ped faces; moderately alkaline (pH 8.0) (pH 8.1 with thymol blue); clear, wavy boundary.
IIC1 RSL No. 61690	21 to 27 inches, light gray (10YR 7/1) clay loam, dark grayish brown (10YR 4/2) when moist; weak fine to medium prismatic breaking to strong fine to medium angular blocky structure; very hard, very firm, slightly sticky and plastic; few fine to medium roots; common very fine and few fine to medium tubular pores; many moderately thick clay films in pores and on ped faces; moderately alkaline (pH 8.2) (pH 8.0 with thymol blue); gradual, smooth boundary.
IIIC2g RSL No. 61691	27 to 37 inches, light gray (10YR 7/1) sandy clay loam with common medium distinct white (N 8/) and very pale brown (10YR 7/4) mottles, light olive brown (2.5Y 5/4) with common medium distinct mottles, dark yellowish brown (10YR 4/4) when moist; massive breaking to weak medium subangular blocky structure; hard, very firm, sticky and slightly plastic; very few fine roots; few very fine tubular and interstitial pores; common thin clay films in pores; moderately alkaline (pH 8.0)

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haplaquoll, clayey over loamy, montmorillonitic, noncalcareous, mesic

SOIL Ramelli silty clay loam (taxadjunct) SOIL Nos. S61Calif-32-14 LOCATION Sierra Valley, Plumas County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 61695 - 61705

Depth (in.)	Horizon	1B1b Size class and particle diameter (mm) 3A1														Coarse fragments 2A2		
		Total			Sand					Silt		Int. II (0.2–0.02)	(2–0.2)			> 2	2 – 19	19 – 76
		Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	0.05–0.02	Int. III (0.02–0.002)					Pct.	Pct. of < 76 mm	
		Pct. of < 2 mm																
0–2	A11	10.1	42.2	47.7	0.7*	1.7*	1.6*	3.4	2.7	6.5	35.7	11.1	5.5			2		
2–7	A12	15.2	37.3	47.5	0.9	2.5	3.4	5.6	2.8	6.7	30.6	12.5	9.4			0		
7–14	A13	14.0	36.2	49.8	0.5	2.4	2.9	4.8	3.4	7.5	28.7	13.4	8.1			0		
14–18	B21	27.0	26.7	46.3	1.2	5.8	6.0	10.1	3.0	7.6	19.1	15.9	[illegible]			[illegible]		

RAMELLI SILTY CLAY LOAM (taxadjunct) 1/

S61Calif-32-14

Location: Sierra Valley, Plumas County, California. 100 feet north and 600 feet east of the center of section 28, T. 22 N., R. 15 E. 600 feet east of Hariot Lane. Approximately two miles north and three miles west of Loyalton. Date of Sampling: September 14, 1961.
Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, James McLaughlin, Eugene Begg.
Vegetation: Meadow vegetation with a mixture of sedges and big blue grass. Climate: Estimated annual precipitation 13 inches. Nearest weather station (Portola): Mean annual temperature 45.1° F.; January average 28.7° F., July average 62.6° F.; mean annual precipitation 20.62 inches. Parent Material: Alluvium from mixed sources. Relief: This is a continuously depressed bottomland area with a smooth slope of less than 1 percent. Elevation: 4890 feet. Drainage: The soil is imperfectly drained; runoff is very slow; permeability is moderately slow. Soil Moisture: Moist below 7 inches.
Remarks: This is a Humic Gley soil with a structural B horizon. The VC6g consists of fine textured lake sediments. At 90 inches blue (5G 4/1) clay occurs. The A13 (7-14 inches), the B21 (14-18 inches) and the IIC2g (34-45 inches) were sampled for the Bureau of Public Roads and the California State Division of Highways. **Classification: Typic Haplaquoll, clayey over loamy, montmorillonitic, noncalcareous, mesic.**

HORIZON	DESCRIPTION
A11 RSL No. 61695	0 to 2 inches, dark gray (10YR 4/1) silty clay loam, black (10YR 2/1) when moist; moderate very fine platy structure; soft, friable, sticky and slightly plastic; abundant very fine, few fine to medium roots; many very fine tubular and few very fine interstitial pores; medium acid (pH 6.0); clear, smooth boundary.
A12 RSL No.	2 to 7 inches, dark grayish brown (10YR 4/2) silty clay loam, black (10YR 2/1) when moist; weak coarse platy breaking to moderate fine granular structure; soft, friable, sticky and slightly [illegible] medium tubular and few very fine interstitial pores; few thin clay films in pores; medium acid (pH 6.0); gradual, smooth boundary.
A13 RSL No. 61697	7 to 14 inches, gray (N 5/) silty clay loam, black (N 2/) when moist, moderate medium subangular blocky structure; slightly hard, friable, sticky and plastic; plentiful very fine to medium roots; many very fine and few fine to medium tubular pores and few very fine interstitial pores; few thin clay films in pores; slightly acid (pH 6.5); clear, smooth boundary.
B21 RSL No. 61698	14 to 18 inches, dark gray (N 4/) silty clay, black (N 2/) when moist; strong medium prismatic breaking to strong coarse angular blocky structure; very hard, firm, very sticky and plastic; plentiful very fine and few fine to medium mainly exped roots; few very fine and fine tubular pores; continuous thin clay films in pores and on ped faces; neutral (pH 7.0); clear, slightly wavy boundary.
IIB2 RSL No. 61699	18 to 26 inches, gray (10YR 5/1) sandy clay, dark gray (10YR 4/1) when moist; strong medium prismatic breaking to strong coarse angular blocky structure; extrememly hard, very firm, sticky and plastic; few very fine to medium mainly exped roots; pores and clay films the same as the horizon above; mildly alkaline (pH 7.5); strongly effervescent, lime occurs in soft masses; clear, slightly wavy boundary.
IIIC1 RSL No. 61700	26 to 28 inches, light brownish gray (10YR 6/2) sandy clay loam, grayish brown (10YR 5/2) when moist; weak medium prismatic structure; extremely hard, very firm, sticky and plastic; few fine to medium roots; plentiful very fine tubular and interstitial and few fine tubular pores; common thin clay films in pores and on ped faces; moderately alkaline (pH 8.0)(pH 8.0 with thymol blue);

SCS-420
10-64

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Vertic Haplaquoll, very-fine, montmorillonitic, noncalcareous, thermic

SOIL Sacramento clay SOIL Nos. S64Calif-57-10 LOCATION Yolo County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64495 - 64502

Depth (In.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Noncarbonate clay	Coarse fragments 1A2a: > 2	2 - 19	19 - 76
		Pct. of < 2 mm														Pct.	Pct. of < 76 mm	
0-7	Ap	0.7	42.4	56.9	0.1	0.1	0.1	0.2	0.2	3.2	39.2	3.5	0.5			0		
7-16	A12	0.8	41.5	57.7	tr.	0.1	0.1	0.3	0.3	3.5	38.0	4.0	0.5			0		
16-31	A13	0.7	38.4	60.9	tr.	0.1	0.2	0.2	0.2	4.0	34.4	4.3	0.5			0		
31-38	A14	1.5	35.2	63.3	tr.	0.1	0.3	0.6	0.5	2.1	33.1	2.9	1.0			0		
38-53	A15	3.5	33.6	62.9	tr.	0.2	0.7	1.7	0.9	1.7	31.9	3.5	2.6			0		
53-60	C1	4.3	37.0	58.7	0.7	1.1	0.8	1.1	0.6	3.3	33.7	4.5	3.7			0		
60-69	C2	2.6	44.9	52.5	0.1	0.3	0.4	1.0	0.8	4.2	40.7	5.6	1.8			0		
69-77+	C3	1.4	59.7	38.9	0.1	0.2	0.2	0.4	0.5	12.6	47.1	13.3	0.9			0		

Depth (In.)	6A1a Organic carbon	6B1a Nitrogen	C/N	6C2a Ext. Iron as Fe	6E1b Carbonate as $CaCO_3$	Bulk density: 4A1f 1/3 bar	4A1h Oven dry		3B2 Cm	Water content:	4B1c 1/3 bar	4B2 15 bar	4D1 Extensibility COLEf	4D1 Extensibility COLE	pH: 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
	Pct.	Pct.		Pct.	Pct.	g/cc	g/cc	g/cc		Pct.	Pct.	Pct.	in./in.	in./in.			
0-7	1.77	0.161	11	1.6	-	1.43	1.76		1.00		29.3	25.2	0.072	0.072		6.0	6.6
7-16	1.38	0.130	11	1.9	-	1.33	1.64		1.00		33.3	27.1	0.070	0.070		6.7	7.4
16-31	0.79	0.110	7	1.4	tr.	1.29	1.63		1.00		32.9	25.8	0.081	0.081		7.5	8.2
31-38	0.84	0.107	8	2.1	tr.	1.30	1.65		1.00		33.1	29.3	0.082	0.082		7.9	8.5
38-53	0.82	0.069	12	2.4	tr.	1.26	1.56		1.00		36.5	30.9	0.075	0.075		7.9	8.4
53-60	0.59			1.9	tr.	1.25	1.59		1.00		35.6	27.9	0.085	0.085		7.8	8.5
60-69	0.59			2.0	tr.	1.30	1.65		1.00		34.7	26.8	0.084	0.084		7.7	8.4
69-77+	0.36			1.3	-	1.48	1.73		1.00		26.5	20.1	0.052	0.052		7.9	8.5

Depth (In.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	5B1a Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K	6I1a CO_3	6J1a HCO_3	8A1 6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter								mmho/cm
0-7	20.7	15.8	0.4	0.8	37.7	7.6	42.6		2.1	2.5	1.3	0.2	-	0.1	1.5		0.62
7-16	22.6	17.4	0.5	0.5	41.0	5.0	48.8		0.6	0.6	0.7	0.1	-	0.9	0.3		0.16
16-31	27.0	21.1	1.1	0.6	49.8		53.6		1.2	1.2	1.7	0.1	-	1.8	0.4		0.41
31-38	30.0	23.3	1.4	0.4	55.1		54.8		1.0	0.8	2.2	0.2	-	1.8	0.2		0.40
38-53	30.2	22.2	1.4	0.4	54.2		54.8		1.0	1.0	2.2	0.2	-	1.4	0.7		0.41
53-60	31.5	22.2	1.5	0.4	55.6		58.3		1.4	1.2	2.6	tr.	-	1.3	1.6		0.52
60-69	29.9	21.9	1.3	0.4	53.5		52.4		0.9	0.9	2.0	0.1	-	1.2	0.5		0.34
69-77+	21.8	16.3	1.1	0.4	39.6		41.4		0.8	0.6	2.0	0.1	-	1.2	0.6		0.33

Depth (In.)	8A Water at Saturation	Exch. bases: 5B3b Ca	5B3b Mg	5B1b Na	5D2 Exchangeable Na	5E SAR	Gypsum	Atterberg limits: Plastic limit	Liquid limit	Plastic index
	Pct.	meq./100 g.			Pct.		Pct.	Pct. of <0.4 mm		
0-7	64.5			0.3	1	1		25	62	37
7-16	61.3			0.4	1	1				
16-31	67.3			1.0	2	2		33	66	33
31-38	70.3			1.3	3	2				
38-53	72.4			1.2	3	2				
53-60	72.9			1.3	3	2				
60-69	69.2			1.2	2	2				
69-77+	54.6			1.0	3	2		30	43	13

SACRAMENTO CLAY
S64Calif-57-10

Location: Yolo County, California. A pit located about 100 feet east of the intersection of Hwy 45 and Road 112, T.12N, R.2E., (sections not surveyed), located about 2.25 miles northwest of Knights Landing. Photo ABB-41T-131. Date of Sampling: October 1, 1964.
Description by: G. J. Staidl Samples Collected by: R. E. Nelson, B. R. Brasher, G. M. Kennedy, W. F. Andrews, L. A. Bates, G. J. Staidl, C. B. Goudey.
Classification: Vertic Haplaquoll, very-fine, montmorillonitic, noncalcareous, thermic.
Vegetation: Cultivated, disked honey dew melon crop. Climate: Average annual precipitation is 16.36 inches; mean annual temperature 61°F. Parent Material: Mixed fine alluvium.
Topography: Basin; plane; less than one percent slope. Elevation: 26 feet above sea level.
Drainage: Runoff is slow; permeability is slow; drainage class, altered drainage with the soil having been formed under poorly drained conditions and altered to somewhat poorly drained.
Soil Moisture: The profile was moist below 7 inches.
Remarks: (1) Tucker soil reaction test used in field pH determination. (2) Paired Sample No. 64

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Vertic Haplaquoll, very-fine, montmorillonitic, noncalcareous, thermic

SOIL: Sacramento clay SOIL Nos. 864Calif-57-11 LOCATION Yolo County, California

SOIL SURVEY LABORATORY: Riverside, California LAB. Nos. 64503 - 64510

Depth (in.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	Coarse fragments 1A2a: > 2	2 - 19	19 - 76
		Pct. of < 2 mm														Pct.	Pct. of < 76 mm	
0-8	Ap	0.9	30.7	68.4	0.1	0.1	0.1	0.3	0.3	2.1	28.6	2.6	0.6			0		
8-22	A12	1.3	30.4	68.3	tr.	0.1	0.2	0.5	0.5	2.0	28.4	2.8	0.8			0		
22-30	A13	3.0	33.2	63.8	tr.	0.1	0.3	1.3	1.3	3.2	30.0	5.3	1.7			0		
30-41	A14	2.0	39.2	58.8	tr.	0.1	0.2	1.0	0.7	1.9	37.3	3.2	1.3			0		
41-48	A15	1.3	45.8	52.9	0.1	0.3	0.2	0.4	0.3	3.0	42.8	3.5	1.0			0		
48-62	C1	2.9	48.4	48.7	0.7	0.6	0.3	0.5	0.8	2.9	45.5	4.0	2.1			0		
62-74	C2	3.9	50.6	45.5	0.9	0.9	0.4	0.8	0.9	4.2	46.4	5.5	3.0			0		
74-84+	C3	3.6	52.3	44.1	0.5	0.7	0.5	0.9	1.0	4.3	48.0	5.7	2.6			0		

Depth (in.)	6A1a Organic carbon	6B1a Nitrogen	C/N	6C2a Ext. Iron as Fe	6E1b Carbonate as $CaCO_3$	Atterberg limits: Plastic limit	Liquid limit	Plastic index	Water content:	4B1c 1/3 bar	4B2 15 bar	4D1 Extensibility COLE	4D1 Extensibility COLE	pH: 8C1b Saturated paste	8C1a H_2O 1:1	8C1a H_2O 1:10
	Pct.	Pct.		Pct.	Pct.	Pct. of <0.4 mm			Pct.	Pct.	Pct.	in./in.	in./in.			
0-8	2.12	0.204	10		-	34	69	35			28.5				6.7	7.1
8-22	1.67	0.176	9		-						29.8				7.0	7.3
22-30	1.35	0.208	6		tr.						30.9				7.8	8.2
30-41	0.79	0.059	13		tr.	41	72	31			29.3				7.7	8.2
41-48	0.61	0.035	17		tr.						29.6				7.9	8.5
48-62	0.48				2						19.3				8.0	8.6
62-74	0.45				2						18.6				8.1	8.7
74-84+	0.43				2	30	55	25			18.2				8.1	8.7

Depth (in.)	Extractable bases: 6N2a Ca	6O2a Mg	6P2a Na	5B1a 6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste: 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	8A1 6K1a Cl	6L1a SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-8	26.4	19.6	1.2	0.8	48.0		49.0		2.5	2.8	2.8	0.1		-	0.5	4.1		0.82
8-22	26.9	20.1	1.4	0.6	49.0		43.9		2.4	2.7	3.3	0.1		-	0.8	3.3		0.84
22-30	31.9	24.4	4.2	0.4	60.9		59.1		11.2	14.0	14.9	tr.		-	1.1	13.4	26.8	3.56
30-41	28.1	23.5	3.7	0.5	55.8		48.4		9.0	10.6	13.2	tr.		-	1.1	12.2	21.0	2.97
41-48	26.6	23.3	2.8	0.4	53.1		44.5		4.5	6.3	7.8	tr.		-	1.4	5.8	12.0	1.83
48-62	23.8	19.4	2.1	0.4	45.7		36.0		2.8	4.2	6.0	0.1		-	1.5	3.5	8.5	1.28
62-74	25.5	18.0	1.4	0.4	45.3		40.1		2.7	3.2	4.4	tr.		-	1.5	2.9	6.3	1.05
74-84+	25.9	19.3	1.2	0.4	46.8		39.3		2.2	2.6	3.0	0.1		-	1.6	1.8	4.3	0.78

Depth (in.)	8A Water at Saturation	Exch. bases: 5B3b Ca	5B3b Mg	5B1b Na	5D2 Exchangeable Na	5E SAR	6F1a Gypsum	Clay mineralogy (<0.002 mm.): Mont.	Verm.	Mica	Kaolin.	
	Pct.	meq./100 g.			Pct.		Pct.	X-ray				
0-8	68.5			1.0	2	2						
8-22	75.5			1.1	2	2						
22-30	80.5			3.0	5	4	-					
30-41	82.3			2.6	5	4	-					
41-48	71.8			2.2	4	3						
48-62	62.8			1.7	5	3						
62-74	60.7			1.2	3	3						
74-84+	58.5			1.0	2	2						

SACRAMENTO CLAY
S64Calif-57-11

Location: Yolo County, California. A pit located 3/4 mile northeast of State Hwy. 45 and County Road 109A, 50 feet southeast of Road 109A, T.12N., R.2E. (section lines not surveyed), 4 miles northwest of Knights Landing. Photo ABB-41T-134. Date of Sampling: October 1, 1964.
Description by: C. B. Goudey. Collectors: R. E. Nelson, B. R. Brasher, G. M. Kennedy, W. F. Andrews, L. A. Bates, G. J. Staidl, C. B. Goudey.
Classification: Vertic Haplaquoll, very-fine, montmorillonitic, noncalcareous, thermic.
Vegetation: Fallow, chiseled and planed. Climate: 16 inches average annual precipitation and mean annual temperature of 61°F. Parent Material: Mixed fine alluvium. Topography: Basin; plane; less than one percent, field was leveled (very slight cuts and fills). Elevation: 24 feet. Drainage: Runoff-slow; permeability - slow; altered drainage class (soil formed under poor drainage, presently imperfectly drained). Soil Moisture: Profile was moist throughout below 8 inches.
Remarks: (1) Truog field pH determinations; (2) Paired profile 64 Cal. 57-10; (3) Diagnostic features--Thick mollic epipedon; common slickensides; more than 30 percent clay; profile mottled throughout (surface mottling due to rice culture); (4) Soil temperature - Ap-30°C, C3-19°C.; (5) B.P.R. samples taken from the Ap, A14 and C3 horizons; (6) Grayish brown lenses banded about one inch wide; occur in the A14 horizon.

HORIZON	DESCRIPTION
Ap RSL 64503	0 to 8 inches, gray (5Y 5/1) silty clay with common fine distinct yellowish brown (10YR 5/8) mottles, very dark gray (5Y 3/1) with common fine distinct dark yellowish brown (10YR 4/4) mottles when moist; plowed; hard, friable, sticky and plastic; abundant micro and very fine roots; common micro and very fine tubular pores; slightly acid (pH 6.5); clear smooth boundary.
A12 RSL 64504	8 to 22 inches, mottled gray (5Y 5/1) light olive gray (5Y 6/2) and yellowish brown (10YR 5/8) silty clay, very dark gray (5Y 3/1) olive gray (5Y 5/2) and dark yellowish brown (10YR 4/4) when moist; weak coarse angular blocky structure; hard, very firm, sticky and plastic; abundant micro and very fine vertical roots; common micro to very fine tubular pores; few slickensides; moderately alkaline (pH 8.0); clear wavy boundary.

CONSECUTIVE PEDON NUMBER C 68112 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES SECCA SHALLOW VARIANT SOIL NO. S68 CALI. 29-35 SAMPLE NO. 8 839-8 843 NEVADA COUNTY AREA

SOIL FAMILY TYPIC ARGIXERDLL LOAMY-SKELETAL,MIXED,MESIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1-.5	MS .5-.25	FS .25-.10	VFS .1-.05	CSI .05-.02	FSI .02-.002	SILTS CO. 2-.2U	CLAYS FINE LT .2U	CARBONATE 3A1A.02	1A2A INT. II .2-.02	COARSE FRAGMENTS PCT. OF WHOLE SOIL 75-20	I 20-5	II 5-2	III PCT. OF LT 75
			PERCENT OF LESS THAN 2MM																	
8 839	0- 5	A1	44.0	42.7	13.3	6.9	9.7	4.8	10.7	11.9	23.9	18.8				42.3	19A	--	3	16
8 840	5- 15	A3	43.4	40.1	16.5	8.3	9.5	5.0	10.1	10.5	21.3	18.8				37.8	75*	47	13	5
8 841	15- 48	B T	52.7	23.7	23.6	16.3	15.1	5.7	8.4	7.2	10.9	12.8				22.4	69*	48	4	13
8 842	48- 79	C	46.9	36.7	16.4	8.1	12.6	6.0	10.6	9.6	14.2	22.5				29.8	28A	5	13	10
8 843	0- 15	A1															29A	--	5	24

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT EXT FE 6C2B	AL 6G7A	PYROP EXT FE 6C5A	AL 6G5A	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4A1H	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		PERCENT		PERCENT		PERCENT				PCT.NO.40		GRAMS PER CC				PERCENT			PERCENT	
8 839	0- 5	1.93	0.100			4.3	0.2						1.38	1.39	0.89		14.4	8.9	0.2	0.2
8 840	5- 15	1.59	0.084			4.7	0.1					1.40			0.39			12.1		
8 841	15- 48	0.53	0.040			4.1	0.2			41	23		1.15	1.84	0.51		47.4	13.0	17.0	7.3
8 842	48- 79	0.19				1.8	0.2						1.46	1.72	0.82		29.3	14.8	5.6	4.5
8 843	0- 15	1.33	0.070									1.39			0.82			7.4		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6Q2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	NH4-OAC 5A6A	BASE SUM+ ACI-DITY	SATURATION SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 839	0- 5	7.7	6.9	TR	0.4	15.0	5.9		20.9	18.6	72		81		6.0	5.6
8 840	5- 15	9.1	9.0	TR	0.1	18.2	7.7		25.9	22.4	70		81		6.0	5.5
8 841	15- 48	13.1	12.7	0.1	0.1	26.0			26.0	27.5	100+		95		6.4	6.0
8 842	48- 79	22.7	20.1	TR	TR	42.9			42.9	44.3	100+		97		6.7	6.2
8 843	0- 15														6.2	5.7

CLAY MINERALOGY

SAMPLE NO	DEPTH CM.	LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 839	0- 5											1 TRACE
8 840	5- 15										MT MONTMORT	2 SMALL
8 841	15- 48	MT 4	VR 3	KK 1							VR VERMICLT	3 MODERATE
8 842	48- 79										KK KAOLINIT	4 ABUNDANT
8 843	0- 15											5 DOMINANT
												6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
*=THE 75-250MM FRACTION IS INCLUDED.

SECCA GRAVELLY LOAM, SHALLOW VARIANT 1/
S68Calif-29-35

Location: Nevada County, California; 1,000 feet north of the S 1/4 corner of sec. 3, T. 16 N., R. 8 E., 600 feet north of California Highway 49, 2½ miles WNW of Nevada City, California. Date of Sampling: September 9, 1968.
Description by: T. M. Ryan and L. A. Brittan. Collectors: L. A. Brittan, G. Staidl, C. B. Goudey, T. M. Ryan, G. M. Kennedy.
Classification: Typic Argixeroll, loamy-skeletal, mixed, mesic.
Vegetation: Scrub live oak, yerba santa, Digger pine, vinegar weed, rabbit brush (?) orchardgrass, perennial rye, toyon. Cultivated. Climate: Average annual precipitation is 46 inches. Mean annual temperature is 56°F. Parent Material: Ultrabasic rock. Topography: Moderately sloping, plain slope, 20 percent, east aspect, mid-slope. Elevation: 2,425 feet. Drainage: Well drained; runoff is medium; permeability is moderate over very slow in the subsoil. Soil Moisture: Profile dry to 2 inches, very slightly moist below.
Remarks: pH taken by Hellige method. This area has been cleared and planted to orchardgrass and perennial ryegrass. This is the second planting of these grasses. In neither case has the stand established itself. This is very possibly due to the erosion of the A1 horizon and the fact that this soil is on ultrabasic rock. There is a gravel mulch or a pavement on the surface. In the A3 horizon there are 25 percent cobbles by volume.

HORIZON	DESCRIPTION
A1 RSL 68839	(0-5 cm.) 0 to 2 inches, reddish brown (5YR 4/4) gravelly loam, dark reddish brown (5YR 3/3) moist; weak coarse and very coarse platy structure; soft, very friable, slightly sticky, slightly plastic; few very fine and fine random roots; common very fine interstitial pores; 25 percent by volume gravel; medium acid (pH 6.0); abrupt wavy boundary.
A3 RSL 68840	(5-15 cm.) 2 to 6 inches, reddish brown (5YR 4/4) cobbly clay loam, dark reddish brown (5YR 3/3) moist; moderate fine and medium subangular blocky structure; slightly hard, friable, sticky, plastic; common very fine and fine random roots; common very fine interstitial and tubular pores; common thin clay films on faces of peds, lining pores, and as bridges; 20 percent by volume gravel; slightly acid (pH 6.5); clear wavy boundary.
Bt RSL 68841	(15-48 cm.) 6 to 19 inches, reddish brown (5YR 4/4) cobbly clay, dark reddish brown (5YR 3/3) moist; moderate medium and coarse subangular blocky structure; hard, firm, sticky, plastic; common very fine and fine random and common medium horizontal roots; few very fine interstitial and few fine tubular pores; many moderately thick clay films line pores and as bridges; slightly acid (pH 6.5); clear wavy boundary.
C RSL 68842	(48-79 cm.) 19 to 31 inches, light yellowish brown (2.5Y 6/4) clay loam, dark yellowish brown (10YR 4/4) moist; massive; hard, firm, slightly sticky, plastic; few very fine and fine random and common medium horizontal roots; common very fine and fine interstitial and few very fine tubular pores; common moderately thick brown and dark brown (7.5YR 4/4 moist) clay films line pores and as bridges; neutral (pH 6.7); clear wavy boundary.
R	(79-97 cm.) 31 to 38 inches, slightly weathered ultrabasic rock.

1/ The Secca series is in a fine, mixed, mesic family of Mollic Haploxeralfs. This pedon has a mollic epipedon and is in a loamy-skeletal family.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE: Liv rmore very gravelly loam (taxadjunct) **LOCATION** Alameda County, California

SOIL FAMILY: Typic Argixeroll, loamy-skeletal, mixed, thermic.

SOIL NOS. S59Calif-01-4 **LAB. NOS.** 59224 thru 59229

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)							3A1		2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 **	
0-5	A_{11p}	12.3	12.2	6.2	8.8	8.7	38.9	12.9	32.6	20.1	61	1
5-13	A_{12p}	13.8	13.0	5.9	7.6	7.8	37.3	14.6	29.6	19.8	59	1
13-23	A_{13}	11.1	14.6	8.8	10.1	7.3	30.2	17.9	26.0	17.0	51	1 - cosl
23-30	B_2	21.6	22.6	11.9	10.3	4.0	15.7	13.9	15.1	9.3	61	cosl
30-38	C_1	21.3	33.1	14.2	11.6	2.6	8.7	8.5	11.1	5.1	77	lcos
38-60	C_2	44.2	31.7	4.8	3.4	1.6	7.4	6.9	6.7	4.0	74	lcos
pH		Organic Matter			6E1b	8A1a						4B2

LIVERMORE VERY GRAVELLY LOAM (taxadjunct) 1/
S59Calif-01-4

Location: Alameda County, California. 36 feet east of Tesla Road at the junction with Marina Avenue, about 1.5 miles southeast of the center of Livermore.
Description by: Larry E. Welch. Collectors: R. C. Huff, L. E. Welch, and L. H. Grant. Date of Sampling: May 5, 1959.
Classification: Typic Argixeroll, loamy-skeletal, mixed, thermic.
Vegetation: Cultivated vineyard. Climate: Mean annual rainfall about 15 inches. Parent Material: Alluvium from sedimentary rocks. Relief: Nearly level, slightly convex slope of less than 1 percent, on a low terrace. Elevation: 550 feet. Drainage: Well drained, slow runoff, moderate to moderately rapid permeability. Soil Moisture: Moist throughout.
Remarks: This is a minimal Noncalcic Brown soil. Numerous types of soil amendments and fertilizers have been added to this field which probably explains the high pH of the surface horizons. Some cobbles and stones throughout.

DESCRIPTION

HORIZON		
A_{11p}	0-5"	Grayish brown (10YR 5/2) loam, very dark grayish brown (10YR 3/2) when moist; very weak very fine granular structure; slightly hard when dry, friable when moist, slightly plastic and slightly sticky when wet; abundant fine and very fine roots; many fine and very fine pores; moderately alkaline, pH 8.0; gradual smooth lower boundary.
A_{12p}	5-13"	Dark grayish brown (10YR 4/2) loam, very dark grayish brown (10YR 3/2) when moist; massive; slightly hard when dry, friable when moist; slightly plastic and slightly sticky when wet; abundant fine and very fine roots; many fine and very fine pores; moderately alkaline, pH 8.0; gradual, smooth lower boundary.
A_{13}	13-23"	Dark grayish brown (10YR 4/2) heavy loam, very dark grayish brown (10YR 3/2) when moist; weak fine granular structure; slightly hard when dry, friable when moist; slightly sticky and slightly plastic when wet; plentiful fine and very fine roots; many fine and very fine pores; neutral, pH 7.0; gradual smooth lower boundary.
B_2	23-30"	Dark grayish brown (10YR 4/2) gravelly sandy clay loam, very dark grayish brown (10YR 3/2) when moist; hard when dry, friable when moist, sticky and plastic when wet; few fine or very fine roots; many fine and very fine pores; colloid mainly in bridges and films covering sand and gravel; slightly acid, pH 6.5; gradual, wavy lower boundary.
C_1	30-38"	Dark grayish brown (10YR 4/2) gravelly sandy loam, dark brown (10YR 3/3) when moist; massive; slightly hard when dry, friable when moist; few roots; many fine and very fine pores; colloid mainly in bridges and on sand grains; slightly acid, pH 6.5; gradual smooth lower boundary.
C_2	38-60"	Dark grayish brown (10YR 4/2) gravelly sandy loam, very dark grayish brown (10YR 3/2) when moist; massive; similar to above but slightly less colloid present.

1/ The Livermore series is in a loamy-skeletal, mixed, thermic family of Typic Haploxerolls. This pedon has a thicker mollic epipedon than is permitted in the series.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Argixeroll, coarse-loamy, mixed, thermic
SOIL Chualar sandy loam (taxadjunct) SOIL Nos. S65Calif-27-17 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65434 - 65442

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand Very coarse (2-1)	Sand Coarse (1-0.5)	Sand Medium (0.5-0.25)	Sand Fine (0.25-0.1)	Sand Very fine (0.1-0.05)	Silt 0.05-0.02	Silt Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Noncarbonate clay	* Coarse fragments 1A2a > 2 Pct.	2 - 19 Pct. of < 76 mm	19 -76 Pct. of < 76 mm	
		Size class and particle diameter (mm) 3A1																	
		Pct. of < 2 mm																	
0-8	Ap1	68.6	22.5	8.9	16.2	19.9	10.5	14.3	7.7	12.1	10.4	27.2	60.9	0	9	20	17	3	
8-19	Ap2	63.2	25.6	11.2	13.3	18.7	9.3	14.0	7.9	13.6	12.0	29.3	55.3	0	11	18	18	0	**
19-29	A13	64.2	25.3	10.5	15.6	19.4	8.6	12.6	8.0	13.2	12.1	27.5	56.2	0	11	19	19	0	**
29-35	A3	63.5	25.9	10.6	15.1	19.0	8.8	13.1	7.5	13.9	12.0	28.1	56.0	0	11	17	17	0	**
35-43	B1t	67.8	21.9	10.3	17.4	18.0	8.8	15.3	8.3	11.6	10.3	28.1	59.5	0	10	23	23	0	**
43-58	B21t	63.0	20.0	17.0	18.0	16.3	8.0	13.6	7.1	9.7	10.3	23.7	55.9	0	17	24	24	0	**
58-79	B22t	57.4	25.6	17.0	11.3	14.9	8.3	14.2	8.7	12.9	12.7	29.1	48.7	0	17	24	24	0	**
79-105	C1	67.4	17.9	14.7	16.7	17.5	9.0	15.3	8.9	9.6	8.3	26.7	58.5	0	15	30	30	0	**
105-114	C2	53.2	29.0	17.8	12.8	13.4	7.4	11.8	7.8	13.0	16.0	27.0	45.4	0	18	27	27	0	**

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	Bulk density 4A1f 1/3 bar g/cc	Bulk density 4A1h Oven dry g/cc	g/cc	Water content Pct.	Water content 4B1c 1/3 bar Pct.	Water content 4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1b Saturated Paste	pH 8C1a H_2O 1:1	pH 8C1a H_2O 1:10
0-8	0.96	0.099	10	0.5	-						4.8				5.3	
8-19	0.48	0.048	10	0.7	-						4.9				6.9	
19-29	0.40			0.6	-						4.7				7.1	
29-35	0.22			0.6	-						4.7				7.0	
35-43	0.12			0.6	-						4.9				7.6	
43-58	0.09			0.8	-						8.0				8.5	
58-79	0.06			0.8	-						7.1				8.6	
79-105	0.06			0.6	-						7.6				7.9	
105-114	0.06			1.0	-						8.6				7.8	

Depth (in.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	5B1a 6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 8A1 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-8	5.8	2.0	0.3	1.2	9.3			7.2										
8-19	5.6	1.7	0.5	0.2	8.0			7.8										
19-29	5.6	1.6	0.6	0.1	7.9			7.9										
29-35	4.2	1.8	0.5	0.1	6.6			6.9										
35-43	3.6	2.6	0.6	0.1	6.9			7.0										
43-58	7.2	4.6	1.4	0.2	13.4			12.7										
58-79	7.1	4.6	1.9	0.2	13.8			12.5	0.7	0.6	8.6	0.1		-	4.4	3.2		1.0
79-105	7.3	4.7	1.9	0.2	14.1			13.9										
105-114	7.4	4.8	2.0	0.3	14.6			14.6										

Depth (in.)	8A Water at Saturation Pct.	5D2 Exchangeable Na Pct.	5B1b Exch. Na meq./100g.	Gypsum Pct.	5C1 Base Sat. NH_4OAc Pct.	Clay mineralogy (<0.002 mm.) 7A Mont.	Verm.	Mica	Kaolin.
						X-ray			
0-8					100+			xxx	x
8-19		6			100+				
19-29		7			100				
29-35		7			96			xxx	x
35-43		9			98				
43-58		11			100+	xx		xxx	x
58-79	25.1	14	1.7		100+				
79-105		14			100+				
105-114		14			100	t		xxx	t

* From characterization sample (5Kg) (except as noted).
** 2-19 mm. from characterization sample, 19-76 mm. from field volume estimate.

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

CHUALAR SANDY LOAM (taxadjunct) 1/
S65Calif-27-17

Location: Monterey County, California, 0.35 mile northwest from Gloria Road, 0.57 mile along Gloria Road from Iverson Road, 20 feet southwest into field from dirt road. Date of Sampling: May 22, 1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich; site selection by T. D. Cook.
Classification: Typic Argixeroll, coarse-loamy, mixed, thermic. Noncalcic Brown (medial with brunizemic properties).
Vegetation: Irrigated cropland, beans. Climate: Precipitation 12 inches; mean annual temperature 58°F., mean January 50°F., mean July 65°F. Parent Material: Alluvium from phaneritic, intermediate igneous rocks. Topography: Slightly convex, smooth alluvial fan with a gently sloping surface of 1½ percent. Elevation: 450 feet. Drainage: Well drained, moderately slow subsoil permeability, medium runoff. Soil Moisture: Moist throughout. About field capacity.
Remarks: Field determinations by Truog Kit. Paired samples are S65Calif-27-13-1 through 7. Reaction of Ap2 horizon unnaturally high, may be due to fertilizers or amendments, all of horizon somewhat more alkaline than typical.

HORIZON	DESCRIPTION
Ap1 RSL 65434	0 to 8 inches, dark grayish brown (10YR 4/2) heavy coarse sandy loam, very dark grayish brown (10YR 3/2) moist; moderate medium angular blocky structure; somewhat compacted by tillage and somewhat stronger brown (7.5YR 4/3) moist when broken and rubbed; hard, very friable, slightly sticky and slightly plastic; few very fine roots; many very fine interstitial and few very fine tubular pores; neutral (pH 7.0); clear smooth boundary.
Ap2 RSL 65435	8 to 19 inches, dark grayish brown (10YR 4/2) heavy coarse sandy loam, very dark grayish brown (10YR 3/2) moist; moderate coarse angular blocky structure, somewhat compacted by tillage; very hard, very friable, slightly sticky and slightly plastic; few very fine roots; many very fine interstitial and few very fine tubular pores; moderately alkaline (pH 8.0), noncalcareous; gradual smooth boundary.
A13	19 to 29 inches, dark grayish brown (10YR 4/2) heavy coarse sandy loam, very
65436	slightly hard, very friable, slightly sticky and slightly plastic; few very fine roots; many very fine interstitial, few very fine and few fine tubular pores; mildly alkaline (pH 7.5); clear wavy boundary.
A3 RSL 65437	29 to 35 inches, brown (10YR 5/3) heavy sandy loam, dark brown (7.5YR 3/2) moist; a little lighter in color (7.5YR 4/3) moist when rubbed; weak coarse subangular blocky structure; hard, very friable, slightly sticky and slightly plastic; few very fine roots; many very fine interstitial pores; few thin clay films in bridges; mildly alkaline (pH 7.5); gradual smooth boundary.
B1t RSL 65438	35 to 43 inches, brown (10YR 5/3) heavy coarse sandy loam, brown (7.5YR 4/3) moist; massive; slightly hard, very friable, slightly sticky and slightly plastic; no roots; common very fine interstitial and few very fine tubular pores; few thin clay films in bridges; moderately alkaline (pH 8.0), noncalcareous; horizon feels somewhat brittle; clear smooth boundary.
B21t	43 to 58 inches, brown (7.5YR 5/4) sandy clay loam, darker brown (7.5YR 4/3)

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Carlsbad gravelly sandy loam variant **LOCATION** San Diego County, Calif.

SOIL FAMILY: Typic Argixeroll, fine-loamy, mixed, thermic.

SOIL NOS. S56-Cal.-37-1 **LAB. NOS.** 56206-213

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0- 6	A1	1.4	21.8	25.5	17.7	3.8	21.8	8.0	24.2	9.1	23	sl
6-12	A12	0.4	17.6	26.7	22.1	4.4	21.3	7.5	28.2	7.6	32	sl
12-18	A21	0.6	17.2	27.2	25.0	4.4	19.2	6.4	28.6	6.8	54	sl
18-24	A22	1.0	15.9	25.9	26.4	4.9	20.3	5.6	30.3	7.8	46	sl
24-30	B1	4.0	17.4	23.6	21.2	3.7	16.8	13.3	20.2	10.2	21	sl
30-36	B21	0.6	16.5	24.4	20.6	2.9	13.8	21.2	17.9	8.2	3	scl
36-45	B22	0.2	16.3	23.4	19.7	2.9	11.8	25.7	16.6	7.2	0	scl
45-55	B3	0.3	15.0	22.2	17.5	2.9	8.6	33.5	13.8	5.6	0	scl
84-96	B'2m	10.1	38.0	23.9	18.4	4.3	2.0	3.3		2.4	0	cos

8C1b pH SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	6C1b Free Fe_2O_3 Pct.	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C *	8A1 6P1a No	8A MOISTURE AT SATURATION %	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
6.7		7.6	2.24	0.144	15.6	0.4	1.3	3.2	27.6	17.4	11.8	4.3
6.9		7.6	0.83	0.051	16.3	0.4	0.5	2.3	20.6	14.4	8.6	3.2
6.7		7.1	0.40	0.031	12.9		0.5	2.5	17.2	11.6	6.8	2.6
6.9		7.1	0.24	0.022		0.3	0.5	3.5	14.8	10.8	6.7	2.4
6.6		7.1	0.11			0.8	0.8	5.9	17.5	13.2	9.7	4.2
6.7		7.2	0.11			1.4	0.9	7.0	22.9			7.0
6.8		7.3	0.11			1.0	1.1	8.9	26.2			8.2
6.9		7.7	0.09			1.5	1.5	11.9	34.0			12.5
						1.5						

5A2a CATION EXCHANGE	EXTRACTABLE CATIONS 5B 5B1a	5B1a	6H2a	Exch. 5B1b	Exch. 5B1b	BASE SAT. %	Clay mineralogy (<0.002 mm.) 1/ Mont.	Verm.	Mixed layer	Mica	Kaolin	Other

CARLSBAD GRAVELLY SANDY LOAM variant 1/
S56Calif-37-1

Collectors: L. D. Whittig and R. E. Nelson on May 31, 1956.
Location: Approximately two miles west of Linda Vista.San Diego County.

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65427 - 65433

Depth (cm)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: (0.05-0.02)	Int. III (0.02-0.002)	Int. II (0.2-0.02)	2-0.1			Coarse fragments 1A2a: > 2 Pct. of whole soil	2-19	19-76
		Size class and particle diameter (mm) 3A1 — Pct. of <2 mm														Pct. of <76mm		
0-18	Ap	62.1	27.7	10.2	15.4	16.7	8.7	13.1	8.2	16.2	11.5	31.4	53.9			11	11	0
18-53	A12	62.0	27.0	11.0	13.0	16.9	9.0	13.7	9.4	15.8	11.2	32.6	52.6			14	14	0
53-76	B1t	57.3	28.5	14.2	12.2	15.2	7.8	13.1	9.0	16.3	12.2	32.7	48.3			10	10	0
76-110	B21t	49.2	28.4	22.4	9.1	11.0	5.3	12.8	11.0	14.7	13.7	33.9	38.2			16	16	0
110-140	B22t	57.9	24.3	17.8	17.5	14.8	6.2	10.8	8.6	12.6	11.7	27.6	49.3			21	21	0
140-150	B3	79.8	11.5	8.7	25.7	24.0	10.8	13.7	5.6	6.3	5.2	18.7	74.2			32	29	3
150-205	IIC	91.1	4.4	4.5	20.6	31.1	17.8	17.8	3.8	2.6	1.8	14.1	87.3			35	32	3

Depth (cm)	6A1a Organic carbon 8/ Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. iron as Fe Pct.	Carbonate as $CaCO_3$: 6E1b <2 mm Pct.	3A1a <0.002 mm Pct.	Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	Est. 1/3 bar g/cc	3B2 Cm	Water content:	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEf cm/cm	4D1 Extensibility COLE cm/cm	pH: 8C1b Saturated paste	8C1a H_2O 1:1	8C1c IN KCl 1:1
0-18	0.89	0.080	11	0.6	-				1.60	0.93			4.6				7.6	
18-53	0.70	0.063	11	0.7	-		1.61	1.64		0.91		10.7	5.0	0.006	0.005		8.1	
53-76	0.32			0.9	-		1.79	1.82		0.93		12.0	5.8	0.006	0.005		8.0	
76-110	0.21			1.4	-		1.81	1.89		0.88		15.1	10.0	0.015	0.013		7.3	
110-140	0.16			1.1	-		1.85	1.91		0.84		13.3	8.0	0.011	0.009		7.2	
140-150	0.07			0.6	-		1.65	1.68		0.77		8.4	4.2	0.006	0.004		6.9	
150-205	0.06			0.4	-				1.65	0.75			2.6				7.0	

Depth (cm)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	5B1a 6Q2a K	Sum of bases	6H2a Ext. acidity	Cation exch. capacity: 5A1a NH_4OAc	5A3a Sum of cations		6G1a KCl ext. Al +++	Ratios to clay 8D1: CEC NH_4OAc	Ext. iron	15 bar water	Atterberg limits: Plastic limit	Liquid limit	Plastic index	Base saturation: 5C3 Sum of cations	5C1 NH_4 OAc
	meq/100g													Pct. of <0.4 mm			Pct	Pct
0-18	7.3	2.0	0.3	0.6	10.2		8.0				0.78	0.06	0.45					100+
18-53	7.7	1.8	0.4	0.4	10.3		8.1				0.74	0.06	0.45	15	18	3		100+

CHUALAR SANDY LOAM
S65Calif-27-13

Location: Monterey County, California, 0.45 mile SW on Chualar Road from Old Stage Road to pumphouse on north side of road, 830 feet northwest on dirt road, 20 feet northeast into field. Date of Sampling: May 20, 1965.

Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich; site selection by T. D. Cook.

Classification: Typic Argixeroll, fine-loamy, mixed, thermic family. (Noncalcic Brown (medial with brunizemic properties).

Vegetation: Irrigated cropland, beans. Climate: Precipitation 12 inches; mean annual temperature 58°F., mean January 50°F., and mean July 65°F. Parent Material: Alluvium from phaneritic, intermediate igneous rocks. Topography: Very gently sloping smooth fan of 1½ percent slope, facing west. Elevation: 180 feet. Drainage: Well drained, moderately slow subsoil permeability, medium runoff. Soil Moisture:

Remarks: Field pH determinations by Truog Kit. Paired samples are S65Calif-27-17-1 through 9. Asterisks indicate samples taken for California Division of Highways.

Alkaline reaction of the A and possibly the upper B due to fertilizers or amendments.

HORIZON	DESCRIPTION
Ap RSL No. 65427	0-7 inches (0-18 cm.), dark grayish brown (10YR 4/2) coarse sandy loam, very dark grayish brown (10YR 3/2) moist; weak medium granular and weak medium subangular blocky structure; hard (dry), friable (moist), slightly sticky and slightly plastic (wet); plentiful very fine roots; many very fine interstitial pores; 7 percent gravel by volume; clear smooth boundary.
A12 * RSL No. 65428	7-21 inches (18-53 cm.), dark grayish brown (10YR 4/2) coarse sandy loam, very dark grayish brown (10YR 3/2) moist; moderate medium and coarse granular structure; hard (dry), friable (moist), slightly sticky and slightly plastic (wet); few very fine roots; many very fine interstitial pores; 9 percent gravel by volume; clear wavy boundary.
B1t * RSL No. 65429	21-30 inches (53-76 cm.), yellowish brown (10YR 5/4) heavy coarse sandy loam, dark brown (10YR 3/3) moist; massive; hard (dry), friable (moist), sticky and plastic (wet); few very fine roots; many very fine interstitial, common very fine and few fine tubular pores; common thin clay skins in bridges and pores; considerable rodent activity; 7 percent gravel by volume; clear smooth boundary.
B21t RSL No. 65430	30-44 inches (76-110 cm.), brown (7.5YR 5/4) heavy loam, dark brown (7.5YR 3/2) moist; brown to dark brown (7.5YR 4/4) crushed; massive, distinct vertical joints suggest some structure with continuous thin clay skins on major joints; very hard (dry), firm (moist), very sticky and plastic (wet); no roots; common very fine interstitial pores; continuous thin clay skins in bridges and pores; 12 percent gravel by volume; diffuse smooth boundary.
B22t * RSL No. 65431	44-55 inches (110-140 cm.), brown (7.5YR 5/4) heavy finely gravelly coarse sandy loam, dark brown (7.5YR 3/2) moist, brown to dark brown (7.5YR 4/4) rubbed; massive; very hard (dry), firm (moist), very sticky and plastic (wet); common very fine interstitial and few very fine tubular pores; continuous thin

CONSECUTIVE PEDON NUMBER C 69 54 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES CHUALAR TAXADJUNCT SOIL NO. S69 CALI. 36-21 SAMPLE NO. 9 287-9 290 SAN BERNARDINO AREA

SOIL FAMILY TYPIC ARGIXEROLL FINE-LOAMY,MIXED,THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SANDS VCS 2-1	CS 1- .5	MS .5- .25	FS .25- .05	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A	1A2A INT. II .2- .02	COARSE FRAGMENTS PCT. OF WHOLE 20 I 75-	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														SOIL PCT. OF LT 75		
9 287	0- 18	AP	32.1	45.5	22.4	0.8	1.9	2.4	7.3	19.7	27.0	18.5	--	--	51.4	--A	--	--	--
9 288	18- 30	B1	26.7	45.4	27.9	0.2	1.0	1.6	5.4	18.5	24.6	20.8			46.6	--A	--	--	--
9 289	28- 42	B23T	25.6	46.4	28.0	0.3	0.9	1.2	5.3	17.9	24.9	21.5			46.3	--A	--	--	--
9 290	105-155	2B3 T	15.8	37.1	47.1	1.3	1.4	1.1	3.5	8.5	15.9	21.2			26.7	--A	--	--	--

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	19-2 6E1B	EXT. IRON FE 6C2B	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST 1/3 BAR 4A1F	DRY 4A1F	CM 4AH 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTENSIBLITY LEF 4D1	LE 4D1	PH H2O 1/1 8C1A	H2O 1/10 8C1A	CACL2 8C1E
		-PERCENT--		-PERCENT-		PCT.	PCT.NO.40		GRAMS PER CC			-----PERCENT---			-PERCENT--				
9 287	0- 18	1.88	0.193			0.8				1.34	1.57	1.00		24.9	9.9	5.4	5.4	6.7	6.6
9 288	18- 30	0.79	0.084			0.8			1.40			1.00			11.3			7.0	6.5
9 289	28- 42	0.36		TR		0.9				1.47	1.80	1.00		25.6	13.4	7.0	7.0	7.9	7.0
9 290	105-155	0.22		5		0.7			1.50			1.00			21.0			8.0	9.3	7.6

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6D2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT.	C E C NA- OAC 5A2A	NH4- OAC 5A6A	EC 8A1A	WATER EXTRACT FROM SATURATED PASTE CA 6N1B	MG 6O1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	B 6T1A	PH SAT. PASTE 8C1B
		--------MEQ/100G--------					PCT.	-MEQ/100G		MMHO	----------MEQ/1000 ML----------								PPM	
9 287	0- 18	15.2	4.8	0.5	2.0	22.5	100+		21.3											
9 288	18- 30	20.4	7.6	0.6	0.6	29.2	100+		28.0											
9 289	28- 42	21.5	8.6	3.9	0.5	34.5	100+		30.9	1.01										
9 290	105-155	37.2	11.6	10.1	0.5	59.4	100+		43.3	1.85	1.8	1.1	19.2	0.1		3.9	4.2	10.4		7.8

SAMPLE NO.	DEPTH CM.	WATER AT SAT. 8A PCT.	EXCHANGBLE NA 5B4B MEQ.	NA 5D2 PCT.	SAR 5E	GYPSUM AMT. 6F1B PERCENT	REQ. 6P3A	CLAY MINERALOGY LESS THAN 0.002 MM X RAY I 7A2E	II 7A2E	III 7A2E	DTA IV 7A2E	PCT. 7A3	L.T. .0002 MM X RAY I 7A2E	II 7A2E	III 7A2E	MINERAL CODE	RELATIVE AMOUNT
9 287	0- 18		0.5	2				MI 5	MM 2							MI MICA	1 TRACE
9 288	18- 30		0.6	2												MM MT/MICA	2 SMALL
9 289	28- 42		3.9	13				MT 4	MI 2								3 MODERATE
9 290	105-155	84.1	8.5	20	16											MT MONTMORT	4 ABUNDANT
																	5 DOMINANT
																	6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.

CHUALAR LOAM (TAXADJUNCT) 1/
S69Calif-36-21

Location: San Bernardino County, California, 75 feet east and 600 feet south of Pine and Central Avenue intersection, NW¼ SW¼ of sec. 31, T.2S., R.7W. Date of Sampling: June 9, 1969.
Description by: A. A. Knecht, W. Z. Brock. Collectors: A. A. Knecht, W. Z. Brock.
Classification: Typic Argixeroll, fine-loamy, mixed, thermic.
Vegetation: Small grain stubble, some bindweeds, mustard and thistle. Climate: Average annual precipitation about 16 inches; mean temperature about 61°F. Parent Material: Fine textured alluvium. Topography: Alluvial fan, nearly level, about 1 percent slope. Elevation: 550 feet. Drainage: Moderately well drained; slow runoff, slow permeability. Area above flood plain and water table is below five feet. Soil Moisture: Profile was dry throughout when sampled.
Remarks: Field reaction was determined by Truog method. Four horizons (0-7"; 7-12"; 28-42" and 42-60") were analyzed in the Riverside Soil Survey Laboratory.
*Some soil properties slightly outside the range for the series as currently defined.

HORIZON	DESCRIPTION
	(0-18 cm.)
Ap RSL 69287	0 to 7 inches, dark grayish brown (2.5Y 4/2) loam, very dark grayish brown (2.5Y 3/2) moist; weak very coarse angular blocky structure; very hard, firm, sticky, plastic; many very fine roots; many very fine continuous and vertical inped simple pores; slightly acid (pH 6.5); clear wavy boundary.
	(18-30 cm.)
B1 RSL 69288	7 to 12 inches, brown (10YR 4/3) clay loam, very dark grayish brown (10YR 3/2) moist; weak coarse angular blocky structure; very hard, firm, very sticky, very plastic; common very fine roots; many very fine continuous vertical inped simple pores; neutral (pH 7.0); clear smooth boundary.
	(30-46 cm.)
B21t	12 to 18 inches, brown (10YR 4/3) clay loam, very dark grayish brown (10YR 3/2) moist; weak coarse angular blocky structure; very hard, very firm, very sticky, very plastic; common very fine roots; common very fine and few fine continuous vertical inped simple pores; common thin clay films on faces of peds; mildly alkaline (pH 7.5); clear smooth boundary.
	(46-71 cm.)
B22t	18 to 28 inches, brown (10YR 4/3) clay loam, dark grayish brown (10YR 4/2) moist; weak coarse subangular blocky structure; extremely hard, very firm, very sticky,very plastic; few very fine roots; common very fine and few fine continous vertical inped simple pores; many thin clay films on faces of peds and common moderately thick clay films line tubular pores; mildly alkaline (pH 7.5); gradual smooth boundary.
	(71-105 cm.)
B23t RSL 69289	28 to 42 inches, brown (10YR 4/3) clay loam, very dark grayish brown (10YR 3/2) moist; moderate very coarse prismatic structure; extremely hard, very firm, very sticky, very plastic; few very fine roots; common very fine and few fine continuous vertical inped simple pores; many thin clay films on faces of peds and common moderately thick clay films line tubular pores; moderately alkaline (pH 8.0); gradual smooth boundary.
	(105-155 cm.)
IIB3t RSL 69290	42 to 62 inches, olive brown (2.5Y 4/4) clay, dark yellowish brown (10YR 3/4) moist; strong very coarse prismatic parting to moderate very coarse angular blocky structure; extremely hard, very firm, very sticky, very plastic; few very fine roots; few very fine inped simple pores; continuous thick clay films occur on faces of peds and line tubular pores; strongly effervescent; medium generally rounded segregated lime occurs in soft masses; moderately alkaline (pH 8.0); clear wavy boundary.
	(155-170 cm.)
IIC	62 to 67 inches, light gray (2.5Y 7/2) clay loam, light olive brown (2.5Y 5/4) moist; massive; extremely hard, very firm, sticky, very plastic; few very fine roots; few very fine discontinuous inped open pores; strongly effervescent; moderately alkaline (pH 8.0).

1/ The Chualar series is in a fine-loamy, mixed, thermic family of Typic Argixerolls. This pedon has more than 35 percent clay in the lower part of the argillic horizon.

OIL SURVEY LABORATORY Riverside, California

SOIL TYPE Lobitos loam (taxadjunct) **LOCATION** San Mateo County, California

SOIL FAMILY: Typic Argixeroll, fine-silty, mixed, mesic.

SOIL NOS. S55Cal. 41-13 **LAB. NOS.** 55394-55398

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 9	A_{11}	0.0	0.2	0.2	1.7	29.0	46.7	22.2	57.6	19.6	0	l
9-18	A_{12}	0.0	0.1	0.2	1.7	31.0	41.9	25.1	58.3	16.1	3	l
18-29	B_2	0.0	0.1	0.2	1.2	25.5	41.9	31.1	52.6	15.8	38	cl
29-34	C_2	0.0	0.1	0.2	1.5	26.0	46.3	25.9	55.8	17.8	75	l
34+	D_r			not analysed								
		pH		ORGANIC MATTER			ELECTRI-			MOISTURE TENSIONS		

LOBITOS LOAM (taxadjunct) 1/
S55Calif-41-13

Location: 2,300 feet northeast of junction of Irish Creek Road and Tunitas Road on west side of Irish Creek Road, San Mateo County, California. See Sheet 15, Soil Survey Report, San Mateo Area, California, May, 1961. Date of sampling: May 19, 1955.
Description by: Ralph E. Nelson. Collectors: Reuben E. Nelson, Ralph E. Nelson, Rudolph Ulrich, and R. J. Wagner.
Classification: Typic Argixeroll, fine-silty, mixed, mesic.
Vegetation: Annual grasses and a few scattered perennial grasses as understory to poison oak, baccharis, California sage, and ceanothus. Climate: 25 to 40 inches rainfall.
Parent material: Silty sandstone or shale of the Purisima formation. Topography: Steep hills to mountainous; slope at sampling location about 60 percent; plain toward southeast.
Elevation: 300 feet. Drainage: Runoff very rapid; internal drainage moderate.
Remarks: Worms and worm casts down to 9 inches.

HORIZON	DESCRIPTION
Aoo and Ao	1 or ½-0 inches (no sample taken), loose, dark colored fresh and rotting twigs, leaves, and grass blades; abrupt smooth lower boundary.
A11 RSL No. 55394	0-9 inches, dark grayish brown (10YR 4/2, very dark brown 2/2 moist) loam or silt loam. Weak, medium to fine subangular blocky, hard friable, slightly plastic and nonsticky. There are some worm casts and numerous roots. The reaction is 6.5. The lower boundary is clear and wavy.
A12	9-18 inches, dark grayish brown (10YR 4/2, indistinctly mottled very dark brown
RSL No. 55395	and very dark grayish brown 10YR 2/2-3/2 moist) loam or silt loam. Weak medium to fine subangular blocky, hard, friable, slightly plastic and nonsticky. There are many roots and some rock fragments. The reaction is 6.1. The lower boundary is clear and wavy.
B2 RSL No. 55396	18-29 inches, light olive brown (2.5Y 5/4, olive brown 4/4 moist) silty clay loam; moderate fine subangular blocky, hard firm, plastic and slightly sticky. There are thin patchy olive colored clay skins. Rotten rock fragments give the soil a rather angular gravelly appearance. Numerous roots. The reaction is 6.5. The lower boundary is clear and wavy.
C2 RSL No. 55397	29-34 inches, light olive brown (2.5Y 5/4, olive brown 4/4 moist) loam. Massive. Numerous pieces of rotten rock throughout the horizon. Hard, friable, very slightly plastic and nonsticky. Reaction is 6.3. Few roots. Lower boundary clear and wavy.
Dr RSL No. 55398 (shattered)	34 inches +, light olive brown, coarsely angular, slightly weathered fragments of silty Purisima sandstone or shale and some loosened soil material. Many feet thick.

1/ The Lobitos series is in a fine-loamy, mixed, mesic family of Typic Argixerolls. This pedon has more than 50 percent silt + very fine sand.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aquic Argixeroll, coarse-loamy, mixed, mesic

SOIL Beckwourth coarse sandy loam SOIL Nos. S61Calif-32-15 LOCATION Sierra Valley, Plumas County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 61706 - 61711

Depth (In.)	Horizon	1B1b Total: Sand (2–0.05)	Total: Silt (0.05–0.002)	Total: Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	3A1 Int. II (0.2–0.02)	3A1 (2–0.2)			Coarse fragments 2A2: ≥ 2 Pct.	2 – 19 (Pct. of < 76 mm)	19 – 76 (Pct. of < 76 mm)
		Pct. of < 2 mm																
0–2½	A11	81.2	11.4	7.4	13.6	27.9	18.1	17.5	4.1	5.0	6.4	16.6	69.6			6		
2½–15	A12	84.5	9.0	6.5	13.5	25.6	20.0	20.4	5.0	4.2	4.8	18.2	70.5			7		
15–23	B1	82.3	10.8	6.9	6.0	20.9	20.6	27.5	7.3	5.5	5.3	25.9	61.9			8		
23–34	B2t	77.0	11.6	11.4	6.2	21.9	20.2	22.5	6.2	5.5	6.1	22.1	60.4			7		
34–48	C1	87.4	5.2	7.4	13.9	26.1	19.6	23.7	4.1	2.7	2.5	16.6	73.5			14		
48–58+	C2	94.1	2.2	3.7	10.5	32.2	28.6	21.5	1.3	1.1	1.1	9.2	86.0			9		

BECKWOURTH COARSE SANDY LOAM
Calif-32-15

Location: Sierra Valley, Plumas County, California. 0.6 mile north and 0.2 mile west of the southeast corner of section 21, T. 22 N., R. 15 E. 0.1 mile north of road. About 4.7 miles north and 3.2 miles west of Loyalton. Date of Sampling: September 15, 1961.
Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, James McLaughlin, Eugene Begg.
Vegetation: The vegetation is silver sage. The silver sage has been sprayed with almost 100 percent kill.
Climate: Estimated annual precipitation 15 inches. Nearest weather station (Portola): Mean annual temperature 45.1° F.; January average 28.7° F.; July average 62.6° F.; mean annual precipitation 20.62 inches. Parent Material: Alluvium dominantly from granitic sources. Relief: Very slightly hummocky flood plain with a slope of less than 1 percent. Elevation: 4880 feet. Drainage: Moderately well drained; very slow runoff; the permeability is rapid in the upper part of the profile and moderate in the lower part. Soil Moisture: Slightly moist below 2½ inches; moist below 15 inches.
Remarks: This is a Chestnut soil with a weakly developed Bt horizon. A water table occurred at 60 inches. This is a perched water table held up by fine textured lake sediments. At this site auger borings in the pit located these sediments at 66 inches. Krotovinas are common in the B1 extending to the top of the B2t. This profile was quite moist considering the date of the sampling. It was slightly moist below 2½ inches and moist below 15 inches. The vegetation in this field was silver sage which had been sprayed. There was very little living vegetation, consequently the moist condition. The A12 (2½ to 15 inches), the B2t (23 to 34 inches) and the C1 (34 to 48 inches) were sampled for the Bureau of Public Roads and the California State Division of Highways.
Classification: Aquic Argixeroll, coarse-loamy, mixed, mesic. 1/

HORIZON	DESCRIPTION
A11	0 to 2½ inches, very dark gray (10YR 3/1) coarse sandy loam, black (10YR 2/1) when moist;

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aridic Argixeroll, ashy, mesic, shallow

SOIL Casuse sandy loam (taxadjunct) SOIL Nos. S66Calif-25-2 LOCATION Modoc County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66319 - 66322

Depth (in.)	Horizon	1B1b Size class and particle diameter (mm) Total: Sand (2–0.05)	Total: Silt (0.05–0.002)	Total: Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)		3A1a Non-carbonate clay	1/ Coarse fragments 1A2a > 2 Pct.	2 - 19	19 - 76
		Pct. of < 2 mm															Pct of < 76mm	
0-1½	A1	60.2	24.1	15.7	7.3	14.2	10.5	18.4	9.8	10.5	13.6	29.5	50.4		16	9	8	1
1½-6	B21t	50.9	23.0	26.1	4.2	11.8	9.3	16.3	9.3	9.8	13.2	27.6	41.6		26	7	6	1
6-11	B22t	43.7	21.2	35.1	4.6	8.9	7.5	14.6	8.1	9.0	12.2	25.0	35.6		35	26	20	6
11+	R																	

CASUSE SANDY LOAM (taxadjunct) 1/
S66Calif-25-2

Location: Modoc County, California. SE¼ SW¼ of Sec. 23, T42N, R12E; 1130 feet due east of the west side road; 190 feet west of the east boundary fence line and 630 feet north of the south boundary fence line. 1-3/4 miles SW of Alturas. Date of Sampling: April 7, 1966.

Description by: G. J. Badura. Collectors: K. W. Flach, S. B. Johnson, G. J. Badura.

Classification: **Aridic** Argixeroll, ashy, mesic, **shallow.**

Vegetation: Native vegetation, used for range; few Junipers, big sagebrush (Artemisia tridentata), quackgrass (Agropyron repens), squirreltail (Sitanion hystrix), cheatgrass (Bromus tectorum) and numerous annual forbs. Climate: Precipitation 12 to 14 inches, much of which is snow. Mean annual temperature is 47°F., January 29°F., July 66°F. Parent Material: Pumice tuff, nonindurated (Rhyolitic tuff). Topography: Rolling terrace-like, 4 percent west slope. Elevation: 4428 feet. Drainage: Well drained, A horizons have moderately rapid permeability and B horizons are moderately permeable, runoff is medium. Soil Moisture: The entire profile was moist at the time of sampling.

Remarks: Field determinations of pH by bromthymol blue indicator.

HORIZON	DESCRIPTION
A1 RSL 66319	0 to 1½ inches, brown (10YR 5/3) sandy loam, dark brown (10YR 3/3) moist; massive; slightly hard, very friable, nonsticky, nonplastic; many very fine to medium roots; many very fine and fine vesicular pores; slightly acid (pH 6.4); clear smooth boundary.
B21t RSL 66320	1½ to 6 inches, brown (10YR 4/3) light sandy clay loam, dark brown (10YR 3/3) moist; weak medium subangular blocky structure that breaks to weak very fine platy; slightly hard, friable, slightly sticky, slightly plastic; many very fine to medium roots; many very fine and fine interstitial and tubular pores; common thin clay bridges; neutral (pH 6.7); gradual to diffuse smooth boundary.
B22t RSL 66321	6 to 11 inches, brown (10YR 4/3) sandy clay loam, dark brown (10YR 3/3) moist; weak medium subangular blocky structure; hard, friable, sticky, plastic; many very fine to medium roots; many very fine and fine interstitial and tubular pores; few moderately thick clay films in pores and many thin clay bridges; neutral (pH 7.0); abrupt to clear boundary.
R RSL 66322	11 inches plus, weakly fractured soft tuff with pumice.

NOTE: Pumice in the tuff ranges in size mainly from 1/8 inch to 1 inch.

1/ **The Casuse series is in a loamy, mixed, mesic, shallow family of Aridic Argixerolls. This pedon is in the same subgroup but has 28 percent volcanic glass and more weathered glass than other sand grains.**

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aridic Argixeroll, loamy, mixed, mesic, shallow
SOIL Casuse sandy loam SOIL Nos. S66Calif-25-3 LOCATION Modoc County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66323 - 66326

Depth (in.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)		3A1a Non-carbonate clay	1/ Coarse fragments 1A2a > 2	2-19	19-76
		Size class and particle diameter (mm) 3A1																
		Pct. of < 2 mm														Pct.	Pct. of < 76mm	
0-1½	A1	65.1	25.3	9.6	10.4	15.6	10.7	17.3	11.1	12.5	12.8	32.1	54.0		10	25	23	2
1½-6	B21t	47.8	26.4	25.8	7.2	9.8	7.4	13.8	9.6	10.8	15.6	27.5	38.2		26	18	17	1
6-14	B22t	49.6	22.3	28.1	6.3	10.4	7.9	14.7	10.3	8.9	13.4	27.2	39.3		28	14	14	-
14+	R																	

Depth (in.)	6A1a Organic carbon Pct	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct		Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 C'	Water content: Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1a H_2O 1:1	
0-1½	0.73	0.067	11				1.30	1.32		0.86		17.6	7.6	0.006	0.005	6.3	
1½-6	0.62	0.069	9				1.19	1.27		0.91		28.0	15.4	0.022	0.020	6.4	
6-14	0.47	0.056	8				1.15	1.26		0.93		28.3	16.5	0.031	0.029	6.4	
14+																-	

Depth (in.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1a NH_4OAc	5A3a Sum		6G1d KCl Ext. Al^{+++}						Base saturation: 5C1 NH_4OAc	5C3 Sum
	meq/100 g															Pct.	Pct
0-1½	6.3	3.1	0.1	1.1	10.6	2.6	13.6	13.2								78	80
1½-6	14.0	6.7	0.4	1.9	23.0	3.8	26.7	26.8								86	86
6-14	14.0	6.8	0.4	1.7	22.9	3.8	28.9	26.7								79	86
14+																	

Depth (in.)	Clay mineralogy (<0.002 mm) 7A: Mont.	Verm.	Mica	Kaolin	
	X-ray				

1/ From characterization sample (5 kg).

CASUSE SANDY LOAM
S66Calif-25-3

Location: Modoc County, California. NE¼ SE¼ of Sec. 30, T42N, R11E; 1110 feet due north of the railroad intersection and about 40 feet east of the road; 11 miles west by southwest of Alturas and 7 miles due east of Canby. Date of Sampling: April 7, 1966.

Description by: G. J. Badura. Collectors: K. W. Flach, S. B. Johnson, G.J. Badura.

Classification: **Aridic Argixeroll, loamy, mixed, mesic, shallow.**

Vegetation: Native vegetation, used for range; scattered Junipers, big sagebrush (Artemisia tridentata), cheatgrass (Bromus tectorum), and annual forbs.

Climate: Precipitation 12 to 14 inches, much of which is snow. Mean annual temperature 47°F., January 29°F., July 66°F. Parent Material: Pumice tuff, nonindurated (Rhyolitic tuff). Topography: Rolling terrace-like, 6 percent west slope. Elevation: 4380 feet. Drainage: Well drained; A horizons have moderately rapid permeability and B horizons are moderately permeable; runoff is medium. Soil Moisture: The top 1½ inches were dry but moist below at the time of sampling.

Remarks: Field determinations of pH by bromthymol blue indicator.

HORIZON	DESCRIPTION
A1 RSL 66323	0 to 1½ inches, grayish brown (10YR 5/2) sandy loam, dark brown (10YR 3/3) moist; massive to weak medium platy structure; slightly hard, very friable, nonsticky and nonplastic; common very fine to medium roots; many vesicular and interstitial pores; neutral (pH 6.6); clear smooth boundary.
B21t RSL 66324	1½ to 6 inches, dark grayish brown (10YR 4/4) light sandy clay loam, dark brown (10YR 3/3) moist; weak medium to coarse subangular blocky structure that breaks to weak medium platy; slightly hard, friable, slightly sticky, slightly plastic; many very fine to medium roots; many very fine tubular and interstitial pores; many thin clay bridges; neutral (pH 6.6); clear smooth boundary.
B22t RSL 66325	6 to 14 inches, brown (10YR 4/3) with some pale yellow pumice, sandy clay loam, brown to dark brown (10YR 4/3 to 3/3) moist; moderate medium subangular blocky structure; hard, friable, sticky, plastic; many very fine to medium roots; many very fine tubular pores; common moderately thick clay films on ped faces and many moderately thick clay bridges; neutral (pH 6.7); clear smooth boundary.
R RSL 66326	14 inches plus, weakly fractured tuff and pumice.

NOTE: Pumice in the tuff ranges in size but is mainly between ¼ and 1 inch.

CONSECUTIVE PEDON NUMBER C 69 58 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES CHUALAR VARIANT SOIL NO. S69 CALI. 36-27 SAMPLE NO. 9 304-9 308 SAN BERNARDINO AREA

SOIL FAMILY CALCIC ARGIXEROLL FINE-LOAMY,MIXED,THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	CSI .05- .02	FSI .02- .002	CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A	INT. II .2- .02	PCT. OF WHOLE SOIL	COARSE FRAGMENTS I 75- 20	II 20-5	III 5-2
			SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1												1A2A					
			PERCENT OF LESS THAN 2MM															PCT. OF LT 75		
9 304	0- 28	AP	29.5	45.8	24.7	1.1	1.0	1.2	7.2	19.0	27.8	18.0				51.8	--A	--	--	--
9 305	28- 48	B21T	27.0	42.4	30.6	0.5	0.7	1.2	6.9	17.7	26.4	16.0				49.1	--A	--	--	--
9 306	48- 97	B22T	24.3	42.9	32.8	0.4	0.7	0.9	5.8	16.5	24.7	18.2				45.4	--A	--	--	--
9 307	97-150	B23T	44.3	30.4	25.3	1.4	2.7	4.0	16.0	20.2	18.5	11.9				49.2	--A	--	--	--
9 308	150-185	B3	27.2	41.4	31.4	0.5	1.0	1.1	9.6	15.0	20.8	20.6				43.1	--A	--	--	--

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	19-2 6E1B	EXT. IRON FE 6C2B	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4AH	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTENSBLTY LEF 4D1	LE 4D1	PH H2O 1/1 8C1A	H2O 1/10 8C1A	CACL2 8C1E
		-PERCENT--		-PERCENT-		PCT.	PCT.NO.40		GRAMS PER CC				----PERCENT----			-PERCENT--				
9 304	0- 28	2.02	0.156							1.43	1.68	1.00		26.6	11.6	5.5	5.5	6.7		6.3
9 305	28- 48	0.52	0.051						1.44			1.00			13.5			7.6		6.9
9 306	48- 97	0.35								1.44	1.84	1.00		27.3	15.0	8.5	8.5	7.8		7.0
9 307	97-150	0.07			1				1.46			1.00			12.5			7.7	8.7	7.3
9 308	150-185	0.06							1.48			1.00			18.7			7.6	8.4	7.3

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6Q2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT.	C E C NA- OAC 5A2A	NH4- OAC 5A6A	EC 8A1A	WATER EXTRACT FROM SATURATED PASTE CA 6N1B	MG 6O1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	8A1 B 6T1A	PH SAT. PASTE 8C1B
		--------MEQ/100G--------					PCT.	-MEQ/100G		MMHO	----------MEQ/1000 ML----------								PPM	
9 304	0- 28	17.7	6.8	0.3	0.7	25.5	100+		25.3											
9 305	28- 48	21.6	9.7	1.4	0.4	33.1	100+		31.0											
9 306	48- 97	22.3	11.6	2.2	0.4	36.5	100+		34.1	0.83										
9 307	97-150	27.6	9.7	2.2	0.4	39.9	100+		26.4	2.42	13.3	8.9	12.9	0.2		2.2	1.0	29.5		7.5
9 308	150-185	28.9	11.5	3.6	0.5	44.5	100+		34.2	2.96	15.9	9.4	16.0	0.1		2.7	6.4	29.5		7.2

SAMPLE NO.	DEPTH CM.	WATER AT SAT. 8A PCT.	EXCHANGBLE NA 5B4B MEQ.	NA 5D2 PCT.	SAR 5E	GYPSUM AMT. 6F1B PERCENT	REQ. 6P3A	CLAY MINERALOGY LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 PCT.	L.T. .0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
9 304	0- 28		0.3	1				MI 5	MT 2	MM 2						MI MICA	1 TRACE
9 305	28- 48		1.4	5												MT MONTMORT	2 SMALL
9 306	48- 97		2.2	6				MI 5	MT 2	MM 2						MM MT/MICA	3 MODERATE
9 307	97-150	56.4	1.5	6	4												4 ABUNDANT
9 308	150-185	64.9	2.6	7	4												5 DOMINANT
																	6 INDETER.

A=STANDARD PREPARATION.

CHUALAR LOAM, CALCAREOUS VARIANT 1/

S69Calif-36-27

Location: San Bernardino County, California; about 90 feet north of Carbon Canyon Road, about 1/4 mile west of Pipeline Road, SW¼ SE¼ NE¼ of sec. 21, T. 2 S., R. 8 W. Date of Sampling: June 11, 1969.
Description by: A. A. Knecht, W. Z. Brock. Collectors: A. A. Knecht, W. Z. Brock.
Classification: Calcic Argixeroll, fine-loamy, mixed, thermic.
Vegetation: Annual grasses and bindweeds. Climate: Average annual precipitation about 16 inches; mean temperature about 61°F. Parent Material: Mixed alluvium. Topography: Alluvial fan, gently sloping, about four percent slopes. Elevation: 750 feet. Drainage: Well drained; medium runoff, moderately slow permeability. Soil Moisture: This profile was dry throughout when sampled.
Remarks: Field reaction was determined by the Truog method. All horizons were analyzed by the Riverside Soil Survey Laboratory. *The Chualar series does not have secondary lime or sodium increasing with depth in the lower part of the solum. There are other minor differences.

HORIZON	DESCRIPTION
	(0-28 cm.)
Ap RSL 69304	0 to 11 inches, dark grayish brown (10YR 4/2) loam, very dark grayish brown (10YR 3/2) moist; moderate very coarse angular blocky structure; very hard, very firm, sticky, plastic; many very fine roots; many very fine vertical discontinuous tubular pores; slightly acid (pH 6.5); clear smooth boundary.
	(28-48 cm.)
B21t RSL 69305	11 to 19 inches, grayish brown (10YR 5/2) clay loam, very dark grayish brown (10YR 3/2) moist; strong very coarse prismatic structure; very hard, very firm, sticky, plastic; many very fine roots; many very fine interstitial pores; strongly alkaline (pH 8.5); clear smooth boundary.
	(48-97 cm.)
B22t RSL 69306	19 to 38 inches, brown (10YR 5/3) clay loam, dark grayish brown (10YR 4/2) moist; strong very coarse prismatic structure; extremely hard, extremely firm, very sticky, very plastic; few very fine roots; common very fine random interstitial pores; few thin clay films on faces of peds and lining pores; strongly alkaline (pH 8.5); gradual wavy boundary.
	(97-150 cm.)
B23t RSL 69307	38 to 59 inches, yellowish brown (10YR 5/4) clay loam, brown (10YR 4/3) moist; moderate very coarse prismatic structure; very hard, firm, very sticky, very plastic; common very fine pores; common moderately thick clay films on faces of peds and lining pores; strongly effervescent; fine irregularly shaped lime occurs in filaments and in soft masses; strongly alkaline (pH 8.5); gradual smooth boundary.
	(150-185 cm.)
B3 RSL 69308	59 to 73 inches, brownish yellow (10YR 6/6) clay loam, dark yellowish brown (10YR 4/4) moist; moderate medium subangular blocky structure; very hard, firm, very sticky, very plastic; few very fine tubular pores; few moderately thick clay films line pores; strongly effervescent; fine disseminated lime occurs in soft masses; strongly alkaline (pH 8.5).

1/ The Chualar series is in a fine-loamy, mixed, thermic family of Typic Argixerolls. This pedon has secondary lime and increasing amounts of sodium in the lower solum.

SOIL SURVEY LABORATORY Riv rside, California

SOIL TYPE Vallecitos gravelly loam (taxadjunct) **LOCATION** Alameda County, California

SOIL FAMILY: Lithic Argixeroll, clayey, montmorillonitic, thermic

SOIL NOS. S59Calif-01-8 **LAB. NOS.** 59238 thru 59242

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	
0-1½	A_{11}	11.1	13.4	7.1	11.0	7.2	37.1	13.1	29.9	20.5	30	1
1½-6	A_{12}	10.1	12.5	6.7	11.0	7.6	41.4	10.7	29.7	25.6	29	1
6-12	B_2	7.8	8.6	3.9	6.8	5.5	30.3	37.1	21.9	18.0	8	cl
12-16	B_3	6.4	7.9	4.3	7.9	6.3	29.7	37.5	21.4	19.4	8	cl

VALLECITOS GRAVELLY LOAM (taxadjunct) 1/
S59Calif-01-8

Collectors, R. C. Huff, L. E. Welch, Lothair Grant.
Description by: L. E. Welch.
Date of Sampling: May 7, 1959.

Location: Alameda County, California, in the SW1/4 of the SW1/4 of the NW 1/4 of Sec. 9, T5S, R4E, about .05 mile north and .1 mile east of the W1/4 corner of Sec. 9 in the SE corner of the county.

Elevation: About 2,480 feet.

Relief: Sloping uplands on 25 percent convex, southeast facing slope.

Drainage: Well drained, moderate runoff, moderately slow permeability.

Parent Material: Metasedimentary rocks.

Climate: About 25 inches of rainfall.

Vegetation: Grass with scattered oaks and some rabbitbrush and chamise.

Remarks: The soil is a medial Noncalcic Brown soil.
Classification: Lithic Argixeroll, clayey, montmorillonitic, thermic.

DESCRIPTION

HORIZON		
A_{11}	0-1 1/2"	Brown (10YR 5/3) light loam, dark brown (10YR 3/3) when moist; weak fine granular structure; soft when dry, friable when moist; abundant very fine roots; many very fine pores; medium acid, pH 6.0; clear, smooth lower boundary.
A_{12}	1 1/2- 6"	Brown (10YR 5/3) clay loam, dark brown (10YR 3/3) when moist; massive; hard when dry, friable when moist, plastic and slightly sticky when wet; abundant very fine roots; many very fine pores; slightly acid, pH 6.5; abrupt, wavy lower boundary.
B_2	6-12"	Reddish brown (5YR 4/3) clay, dark reddish brown (5YR 3/3) when moist; nearly massive; very hard when dry, firm when moist, very plastic and slightly sticky when wet; plentiful very fine roots; many very fine pores; thin continuous clay films in pores and in bridges; neutral, pH 7.0; clear, smooth lower boundary.
B_3	12-16"	Brown (10YR 5/3) clay, dark brown (10YR 4/3) when moist; massive; very hard when dry, firm when moist, very plastic and slightly sticky when wet; few very fine roots; common very fine pores; thin and continuous clay films in pores; colloid mainly in bridges; neutral, pH 7.0; abrupt, irregular lower boundary.
D_r	16"+	Bluish gray (5B 5/1) dry, metamorphosed fine-grained sandstone with calcite seams; clay films along cleavage planes.

1/ The Vallecitos series is in a clayey, montmorillonitic, thermic family of Ruptic-Entic-Lithic Mollic Haploxeralfs. This pedon has a mollic epipedon and a continuous argillic horizon.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Vallecitos gravelly loam (taxadjunct) **LOCATION** Alameda County, California

SOIL FAMILY: Lithic Argixeroll, clayey, montmorillonitic, thermic

SOIL NOS. S 59Calif-01-9 **LAB. NOS.** 59243 thru 59247

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	
0-1	A_{11}	9.8	10.8	6.0	8.8	6.0	46.0	12.6	37.1	19.8	17	l
1-7	A_{12}	7.8	11.8	6.5	10.2	7.9	41.0	14.8	31.7	23.1	15	l
7-14	B_{21}	5.4	7.2	3.7	6.1	5.2	30.3	42.1	21.0	18.1	14	c
14-18	B_{22}	4.8	8.6	3.9	6.5	5.4	27.7	43.1	19.5	17.4	38	c
18+	D_r		Not sampled									

VALLECITOS GRAVELLY LOAM (taxadjunct) 1/

S59Calif-01-9

Collectors: R. C. Huff, L. E. Welch, Lothair Grant.
Description by: L. E. Welch.
Date of Sampling: May 5, 1959.

Location: Alameda County, California, in the NE1/4 of the SE1/4 of the SW 1/4 of Sec. 4, T5S, R4E, in the southeast corner of the county.

Elevation: 2,350 feet.

Relief: Sloping uplands on a convex slope of 20 percent facing south.

Drainage: Well drained, moderate runoff, moderately slow permeability.

Parent Material: Metasedimentary rocks.

Climate: Rainfall about 25 inches.

Vegetation: Grass with scattered oaks and some rabbitbrush and chamise.

Remarks: This is a medial Noncalcic Brown soil.

Classification: Lithic Argixeroll, clayey, montmorillonitic, thermic.

DESCRIPTION

HORIZON		
A_{11}	0- 1"	Brown (10YR 5/3) loam, dark brown (10YR 3/3) when moist; weak fine granular structure; slightly hard when dry, friable when moist, slightly sticky when wet; abundant very fine roots; many very fine pores; slightly acid, pH 6.5; clear, smooth lower boundary.
A_{12}	1- 7"	Brown (10YR 5/3) loam, dark brown (10YR 3/3) when moist; massive; hard when dry, friable when moist, slightly sticky when wet; abundant very fine roots; many very fine pores; slightly acid, pH 6.5; clear, smooth lower boundary.
B_{21}	7-14"	Brown (7.5YR 5/4) clay loam, reddish brown (5YR 4/3) when moist; massive; very hard when dry, firm when moist, slightly plastic and slightly sticky when wet; plentiful very fine roots; many very fine pores; thin continuous clay films in pores; neutral, pH 6.8; clear, smooth lower boundary.
B_{22}	14-18"	Brown (10YR 5/3) clay, dark brown (7.5YR 4/3) when moist, massive; very hard when dry, firm when moist, plastic and slightly sticky when wet; few very fine roots; many very fine pores; moderately thick continuous clay films in pores and bridges; neutral, pH 7.0; abrupt, irregular lower boundary.
D_r	18"+	Dark greenish gray schist.

1/ The Vallecitos series is in a clayey, montmorillonitic, thermic family of Ruptic-Entic Lithic Mollic Haploxeralfs. This pedon has a mollic epipedon and a continuous argillic horizon.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Denison clay loam (taxadjunct) **LOCATION** San Mate County, Calif rnia

SOIL FAMILY: Pachic Argixeroll, fine-loamy, mixed, isomesic.

SOIL NOS. S55Cal.41-2 **LAB. NOS.** 55103-55109

		1B1b	PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)	3A1	

DENISON CLAY LOAM (taxadjunct) 1/
S55Calif-41-2

Location: 0.41 miles north of Miramar Hotel, Miramar Village, San Mateo County, California. About 125 feet east of edge of sea cliff.
Date of Sampling: February 23, 1955.
Description by: W. M. Johnson. Collectors: Reuben Nelson, Ralph Nelson, Richard Wagner, Melvin Mello, and W. M. Johnson.
Classification: Pachic Argixeroll, fine-loamy, mixed, isomesic.
Land Use: Cultivated field now lying fallow. Commonly used for Brussels sprouts and artichokes, with some irrigation. Climate: Mean annual rainfall is about 25 inches, most of which comes during the months from November through March. June, July, and August are very dry. Fogginess is high in many parts of the year, particularly during summer and fall. Average January temperature is about 48° F., average July temperature about 60° F. On-shore winds carrying salt spray are common. Parent Material: Alluvium derived mainly from soils of the Coast Range Mountains developed in quartz diorite. High in quartz sand, and contains considerable mica. Relief: Smooth, very gently sloping alluvial fan overlying marine terrace. Slope about ½ percent, plane, toward the west. Elevation: About 18 feet. Drainage: Runoff, slow. Permeability, moderate in A1, slow below. Moderately well drained.
Remarks: Soil moist throughout when sampled.

HORIZON		DESCRIPTION
A_{1p}	0 - 4"	Black (10YR 2.5/0.8, dry; 1.5/0.8, moist) heavy loam with common fine white specks due to quartz grains. Cloddy due to tillage; breaks to weak fine granular. Very hard, friable, plastic, sticky. Very many roots. Numerous worm casts. pH 6.8. Abrupt, smooth lower boundary.
A_{12}	4 - 8"	Heavy loam of the same color as the A_{1p}. Weak coarse prismatic, breaking to weak medium granular. Extremely hard, plastic, sticky. Very many roots. pH 6.5. Clear, wavy lower boundary.
B_{21}	8 - 16"	Clay loam of the same color as the A horizon. Moderate very coarse prismatic, breaking to weak coarse and medium blocky. Very firm, very plastic, very sticky. Moderate continuous clay skins. Slickensides are common. Many roots. pH 6.3. Clear, wavy lower boundary.
B_{22}	16 - 29"	Clay loam of the same color as the A horizon. Moderate very coarse prismatic, breaking to weak medium and coarse blocky. Moderate continuous clay skins. Very firm, very plastic, very sticky. Moderate number of roots. Many slickensides. pH 6.3. Clear, irregular lower boundary.
B_3	29 - 45"	Finely mottled yellowish brown and pale brown (10YR 5/6 and 6/3, dry; 4/6 and 5/3, moist) clay loam with common medium mottles of dark gray (dry) on ped surfaces, and common fine white specks. Moderate very coarse prismatic, breaking to weak medium and coarse blocky. Moderate clay skins on vertical faces. Very firm, very plastic, very sticky. Many slickensides. Few roots. pH 7.0. Clear, irregular lower boundary.
C_1	45 - 55"	Finely mottled light gray and yellowish brown (2.5Y 7/2 and 10YR 5/6, dry; 2.5Y 5/3 and 10YR 4/6, moist) light clay loam with common dark gray skins on ped surfaces, and fine white specks. Weak very coarse prismatic. Friable, plastic, sticky. Moderate patchy clay skins on vertical surfaces. Very few roots. Slickensides are common. pH 7.0. Gradual, wavy lower boundary.
C_2	55 - 63"+	Finely mottled light gray and yellowish brown (2.5Y 7/2 and 10YR 5/4, dry; 2.5Y 5/2 and 10YR 4/4, moist) loam with common dark gray films on ped surfaces and fine white specks. Massive, with coarse pattern of vertical carcks. Friable, plastic, sticky. Moderate patchy clay skins on vertical surfaces and in linings of tubular pores. pH 7.4.

1/ The Denison series is in a fine, montmorillonitic, isomesic family of Pachic Argixerolls. This pedon has slightly less than the 35 percent clay needed for the series.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Pachic Argixeroll, fine-loamy, mixed, thermic

SOIL Lockwood shaley loam SOIL Nos. S65Calif-27-1 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65482 - 65488

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	1/ Coarse fragments > 2 Pct.	2 - 19	19-76
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm															Pct. of < 76mm	
0-3	Ap1	33.0	45.8	21.2	4.7	6.0	3.6	6.5	12.2	27.3	18.5	43.2	20.8	0	21			tr.
3-16	Ap2	37.0	43.6	19.4	6.2	6.9	3.9	6.5	13.5	26.4	17.2	43.6	23.5	0	19			tr.
16-26	A13	34.8	44.6	20.6	4.2	6.0	3.8	6.1	14.7	28.2	16.4	46.4	20.1	0	21			1
26-40	A14	35.8	44.6	19.6	4.8	6.0	3.7	6.3	15.0	29.0	15.6	47.7	20.8	0	20			12
40-57	B21t	55.2	26.7	18.1	17.0	13.2	5.8	8.3	10.9	14.5	12.2	29.6	44.3	0	18			24
57-82	B22t	56.8	18.7	24.5	15.9	14.8	6.6	9.5	10.0	9.1	9.6	23.9	46.8	0	25			26
82+	IIC	30.0	41.5	28.5	3.4	4.0	2.4	6.5	13.7	23.0	18.5	40.7	16.3	0	29			0

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1		
0-3	2.21	0.211	10		-								19.5			4.9		
3-16	1.48	0.078	19		-		1.17	1.20		1.00		30.0	19.8	0.008	0.008	5.5		
16-26	0.79				-		1.11	1.14		1.00		31.5	19.0	0.009	0.009	6.8		
26-40	0.62				-		1.11	1.14		1.00		31.2	18.7	0.010	0.010	7.2		
40-57	0.21				-		1.18	1.23		1.00		34.4	28.6	0.013	0.013	7.0		
57-82	0.21				-		1.14	1.17		1.00		38.0	32.8	0.009	0.009	6.9		
82+	0.19				-		1.08	1.14		1.00		41.2	26.2	0.018	0.018	6.8		

Depth (in.)	Extractable bases 5B1a 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A1a NH_4OAc	5A3a Sum		6G1d KCl Ext. Al^{+++}		Atterberg limits Plastic limit	Liquid limit	Plastic index	5D2 Exch. Na		Base saturation 5C1 NH_4OAc	
	meq/100 g											Pct. of <0.4 mm			Pct.		Pct	Pct.
0-3	13.2	4.0	0.5	1.4	19.1		20.7										92	
3-16	12.3	4.8	0.7	1.4	19.2		18.8										100+	
16-26	12.5	4.6	0.8	1.0	18.9		17.8					26	32	6			100+	
26-40	11.8	4.8	1.1	0.5	18.2		17.8								6		100+	
40-57	10.7	5.7	1.9	0.3	18.6		18.4					28	40	12	10		100+	
57-82	11.2	7.5	2.8	0.4	21.9		23.3								12		94	
82+	12.0	7.5	1.9	0.3	21.7		22.0								9		99	

Depth (in.)	Clay mineralogy (<0.002 mm) 7A Mont.	Verm.	Mica	Kaolin	
	X-ray				

1/ Apparent coarse fragments dispersed readily. Hence the <19 mm. material was ground whole and included in the characterization sample. All chemical and physical data are on <19 mm. basis. The 19-76 mm. fraction was determined from field weights.

LOCKWOOD SHALY LOAM
S65Calif-27-1

Location: Monterey County, California, 100 feet SW, 50 feet NW, from corner of Teague and Central Avenue, about 7 miles NW of King City. Date of Sampling: May 17, 1965. Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich. Site selection by T. D. Cook.
Classification: Pachic Argixeroll, fine-loamy, mixed, thermic. (minimal) Brunizem.
Vegetation: Mature walnut orchard, cultivated. Climate: Precipitation 12 inches, mean annual temperature 59°F., January mean 50°F., July mean 65°F. Parent Material: Alluvium. Almost wholly from diatomaceous (Monterey) shale. Topography: Gently sloping smooth fan, slightly over 1 percent slope. Elevation: 400 feet. Drainage: Well drained, moderately slow permeability, medium runoff. Soil Moisture: Moist (field capacity) throughout below 3 inches.
Remarks: Field determination of pH by Hellige-Truog triplex indicator. Paired site sample number is S65Calif-27-2-1 through 7. Soils are marginally shaly and probably perform on moisture relations as non-shaly soils to a depth of 26 inches. Asterisks indicate horizons sampled for California Division of Highways.

HORIZON	DESCRIPTION
Ap1 RSL 65482	0 to 3 inches, gray (10YR 5/1) shaly loam, very dark grayish brown (10YR 3/2) moist; moderate fine and medium subangular blocky structure; slightly hard, very friable, slightly sticky and slightly plastic; few very fine roots; many very fine interstitial pores; very strongly acid (pH 5.0); A horizons contain about 10 to 15 percent by volume of fine gravel-size shale fragments, low surface pH due to amendments or pesticides, low pH temporary; abrupt smooth boundary.
Ap2 RSL 65483	3 to 16 inches, gray (10YR 5/1) shaly loam, very dark brown (10YR 2/2) moist; weak very coarse angular blocky crushing to moderate medium granular structure; slightly hard, friable, slightly sticky and slightly plastic; few very fine and plentiful fine roots; common very fine interstitial and very few fine tubular pores; slightly acid (pH 6.5); horizon strongly compacted due to tillage; gradual smooth boundary.
*A13 RSL 65484	16 to 26 inches, gray (10YR 5/1) shaly loam, very dark brown (10YR 2/2) moist; strong medium granular structure; slightly hard toward soft, very friable, slightly sticky and slightly plastic; few very fine roots; many very fine interstitial, common fine and medium tubular pores; neutral (pH 7.0); gradual smooth boundary.
A14 RSL 65485	26 to 40 inches, gray (10YR 5/1) heavy shaly loam, very dark grayish brown (10YR 3/2) moist; moderate medium granular structure; soft, very friable, slightly sticky and slightly plastic; few fine and medium roots; many very fine interstitial and common fine tubular pores due to worms; neutral (pH 7.0);

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Pachic Argixeroll, fine-loamy, mixed, thermic

SOIL Lockwood shaly loam SOIL Nos. S65Calif-27-2 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65489 - 65497

Depth (in.)	Horizon	1B1b Size class and particle diameter (mm) 3A1														1/ Coarse fragments 1A2a		
		Total			Sand					Silt								
		Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	0.05–0.02	Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)			> 2 Pct.	2 - 19	19 - 76
		Pct. of < 2 mm														Pct.	Pct. of < 76mm	
0-4	Ap1	39.8	41.1	19.1	6.7	8.6	4.7	8.3	11.5	25.4	15.7	41.4	28.3	0	19			tr.
4-13	Ap2	33.7	43.7	22.6	5.6	7.6	4.0	6.2	10.3	25.8	17.9	39.6	23.4	0	23			tr.
13-34	A13	35.4	43.4	21.2	6.6	8.5	4.0	6.3	10.0	25.2	18.2	38.8	25.4	0	21			4
34-47	A3	46.3	33.7	20.0	11.0	11.5	5.2	8.1	10.5	17.0	16.7	31.9	35.8	0	20			20
47-57	B2t	48.6	29.4	22.0	9.6	11.7	5.9	10.0	11.4	13.0	16.4	29.9	37.2	0	22			20
57-71	B3t	24.2	45.0	30.8	4.6	4.1	1.8	4.2	9.5	17.8	27.2	30.0	14.7	0	31			2
71-85	IIC	32.4	42.4	25.2	4.6	4.4	2.7	6.7	14.0	18.9	23.5	37.2	18.4	0	25			7
57-71*	bands	8.8	50.8	40.4	0.2	0.4	0.3	1.8	6.1	15.5	35.3	22.9	2.7	0	40			
57-71**	Inter-bands	25.2	50.1	24.7	0.4	0.8	1.1	7.6	15.3	22.2	27.9	42.7	9.9	0	25			

LOCKWOOD SHALY LOAM
S65Calif-27-2

Location: Monterey County, California, .6 mile northwest, .45 mile southwest from Central Avenue and Hobson Avenue, NW of King City, 8th row NW, 5½ rows NE in walnut grove from SW corner of grove. Date of Sampling: May 17, 1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton and B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich. Site selection by T. D. Cook.
Classification: **Pachic Argixeroll, fine-loamy, mixed, thermic. (minimal) Brunizem.**
Vegetation: Mature irrigated walnut grove in fallow condition. Climate: Precipitation is 12 inches, mean annual temperature 59°F., January mean 50°F., July mean 65°F. Parent Material: Alluvium. Almost wholly from diatomaceous (Monterey) shale. Topography: Gently sloping smooth fan, slightly over 1 percent slope. Elevation: 525 feet. Drainage: Well drained; **moderately slow permeability; medium runoff.** Soil Moisture: Moist throughout (field capacity).
Remarks: Field pH by Truog Kit. Matched profile S65Calif-27-1-1 through 7. This site slightly more shaly than first profile and not quite 1 value darker. Low pH of Ap1 due to fertilizer or amendments, probably temporary; pH was 6.5 to 7.0 a few weeks before sampling.

HORIZON	DESCRIPTION
Ap1 RSL 65489	0 to 4 inches, dark gray (10YR 4/1) shaly heavy loam, very dark brown (10YR 2/2) moist; weak medium granular structure; slightly hard, friable, slightly sticky and slightly plastic; abundant very fine roots; many very fine interstitial pores; very strongly acid (pH 5.0); about 25 percent by volume fine gravel shale fragments; clear smooth boundary.
Ap2 RSL 65490	4 to 13 inches, dark gray (10YR 4/1) shaly heavy loam, very dark brown (10YR 2/2) moist; moderate medium granular structure; hard, friable, slightly sticky and slightly plastic; many very fine interstitial and few fine and few medium tubular pores; slightly acid (pH 6.5); clear smooth boundary.
A13 RSL 65491	13 to 34 inches, dark gray (10YR 4/1) shaly heavy loam, very dark brown (10YR 2/2) moist; moderate medium granular structure; hard, friable, slightly sticky and slightly plastic; few very fine and few medium roots; many very fine interstitial pores, common fine and few medium tubular pores; slightly acid (pH 6.5); gradual smooth boundary.
A3 RSL 65492	34 to 47 inches, grayish brown (10YR 5/2) gravelly light clay loam, dark grayish brown (10YR 4/2) moist; weak medium subangular blocky structure; slightly hard, slightly sticky and slightly plastic; very few very fine roots; many very fine interstitial pores, common very fine tubular pores; few thin clay films in tubular pores; moderately alkaline (pH 8.0), noncalcareous; A3, B2t, and B3t horizons have large irregular pockets, due to gophers, filled with A soil material; clear wavy boundary.
B2t RSL 65493	47 to 57 inches, very pale brown (10YR 7/3) gravelly light clay loam, dark brown (10YR 4/3) moist; massive; slightly hard, friable, sticky and plastic; no roots; common very fine interstitial and few very fine tubular pores; continuous thin and moderately thick clay films in pores in bands; moderately alkaline (pH 8.0), noncalcareous; contains 3 clay bands 1/2 inch thick 5 inches apart, (7.5YR 3/2m); clear smooth boundary.
B3t RSL 65494	57 to 71 inches, very pale brown (10YR 7/3) gravelly light clay loam, dark yellowish brown (10YR 4/4) moist; massive; slightly hard, friable, sticky and plastic; common very fine interstitial and common very fine tubular pores; clay films in bands only; moderately alkaline (pH 8.0), noncalcareous; two 1/2 inch bands; clear smooth boundary.
IIC RSL 65495	71 to 85 inches, very pale brown (10YR 7/3) heavy silt loam, brown (10YR 5/3) moist; massive; slightly hard, firm, sticky, slightly plastic; few very fine interstitial and many very fine and common fine tubular pores; few thin clay films on joints and very few thin films in some tubular pores; moderately alkaline (pH 8.0), noncalcareous; 3 weak 1/2 inch clay bands.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Denison clay loam **LOCATION** San Mateo, California

SOIL FAMILY: Pachic Argixeroll, fine, montmorillonitic, isomesic.

SOIL NOS. S55Ca1.41-1 **LAB. NOS.** 5594-55102

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 2/	TEXTURAL CLASS
0- 5	A_{1p}	4.9	6.3	3.8	8.8	7.9	34.6	33.7	30.9	17.2	1	cl
5-10	A_{12}	4.8	5.7	3.9	9.3	8.2	35.1	33.0	33.2	16.0	2	cl
10-23	B_{21}	2.3	4.5	3.0	6.7	6.0	32.3	45.2	23.1	19.4	1	c
23-34	B_{22}	3.5	5.3	3.5	7.1	6.3	30.8	43.5	24.6	16.8	1	c
34-45	B_3	5.0	6.3	3.6	6.4	6.7	32.3	39.7	26.1	16.8	3	cl
45-52	C_1	2.9	4.3	2.6	5.1	6.8	39.1	39.2	27.9	21.2	1	cl
52-61	C_2	2.1	3.3	2.0	4.1	5.8	45.8	36.9	27.2	27.1	0	sicl
61-70	D_{g1}	13.3	13.3	6.0	10.7	7.7	32.5	16.5	31.7	15.0	5	l
70-77	D_{g2}	5.0	6.6	4.3	8.6	9.2	39.5	26.8	37.8	16.2	2	l

8C1b SATURATED PASTE 1:1	pH 8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM 25°C.	Free Fe_2O_3 Pct. 6C1b	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
5.8		6.6	3.67	.266	13.8		1.0	0.9				16.4
6.3		7.2	3.42	.260	13.2		0.7	0.8				16.6
6.2		7.3	3.15	.189	16.7		0.7	0.6				21.3
6.5		7.6	2.19	.121	18.1		0.7	0.5				19.6
7.0		8.2	0.48	.030	16.0		0.6	0.6				17.2
7.3		8.5	0.34	.018	18.9		0.5	1.0				17.9
7.5		8.6	0.16	.017			0.5	1.3				18.4
7.3		8.5	0.12	.030			0.7	1.2				11.1
7.2		8.6	0.17	.027			0.6	1.7				13.2

5A2a CATION EXCHANGE CAPACITY (+Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg	6P1a 6I1a CO_3+HCO_3 1/	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
45.8	25.2	12.8	10.1	1.3	0.4	80	3.9	0.1	8.6	1.4	1.9	53.0
44.1	29.5	14.5	7.6	1.5	0.4	86	3.3	0.1	3.6	2.4	1.9	53.5
53.8	30.0	20.3	8.4	1.9	0.3	86	3.7	0.1	3.2	3.7	2.3	67.2
45.4	19.8	19.8	2.8	1.9	0.3	94	3.8	<0.1	2.4	1.4	2.3	63.3
37.3	18.5	16.6	4.9	1.7	0.3	88	3.6	<0.1	1.6	0.6	2.1	59.5
37.7	19.6	17.3	2.8	2.1	0.3	93	3.6	<0.1	1.3	0.7	1.9	61.6
41.2	21.3	17.2	3.5	2.3	0.3	92	3.5	<0.1	1.1	0.7	2.1	60.8
33.2	16.5	11.2	2.4	1.9	0.2	93	4.4	<0.1	1.5	0.6	3.2	40.0
27.8	13.4	9.9	2.7	1.4	0.2	90	4.2	<0.1	1.1	0.4	3.3	42.7

1/ only HCO_3^- found

2/ Percent of whole soil

DENISON CLAY LOAM
S55Calif-41-1

Location: 1/4 mile east of Princeton, 40 feet east of the sea cliff, 1/8 mile SW of junction of Highway 1 and Piose Rd. on east side of Piose Rd., San Mateo County, California. See Sheet 6, Soil Survey Report of San Mateo Area, California, May 1961. Date of Sampling: February 23, 1955.
Description by: W. M. Johnson. Collectors: Reuben Nelson, Ralph Nelson, Richard Wagner, Melvin Mello, and W. M. Johnson.
Classification: Pachic Argixeroll, fine, montmorillonitic, isomesic.
Use: Cultivated; lying fallow at time of sampling. Fields are usually irrigated for some crops. Common crops are Brussels sprouts and artichokes. Climate: Mean annual rainfall is about 25 inches, most of which comes during the months from November through March. June, July, and August are very dry. Fogginess is high in many parts of the year, particularly during summer and fall. Average January temperature is about 48°F, average July temperature, about 60° F. On-shore winds carrying salt spray are common. Parent Material: Alluvium derived from soils and rocks of adjacent hills, mainly from quartz diorite. Alluvium is clay loam or sandy clay loam in texture, rather high in sharp quartz sand. Relief: Smooth, very gently sloping alluvial fan overlying marine terrace. Slope about ½ percent,

Other Features: Entire profile was moist when samples were taken. History of fertilization not known.
Reaction: pH's were measured with Truog Triplex indicator.

HORIZON	DESCRIPTION
A1p RSL No. 5594	0 to 5 inches, black (10YR 2/1, dry; 1.5/1, moist) clay loam with common fine white specks (these are quartz grains). Cloddy, due to tillage. Clods break to weak fine granules. Very hard, friable, plastic, sticky. Worms and worm casts are numerous. Roots very numerous. pH 6.0. Abrupt, smooth lower boundary.
A12 RSL No. 5595	5 to 10 inches, black (10YR 2/1, dry; 1.5/1, moist) clay loam. Weak very coarse and coarse prismatic, breaking to weak medium granular. Friable, plastic, sticky. Many earthworms and worm casts. Roots are numerous. pH 6.5. Clear, wavy lower boundary.
B21 RSL No. 5596	10 to 23 inches, black (10YR 2/1, dry; 1.5/1, moist) clay. Strong very coarse prismatic, breaking to weak coarse blocky. Extremely hard, firm, very plastic, very sticky. Moderate continuous clay skins. Contains a very few hard shot concretions 1/8 inch in diameter. Roots moderately numerous. pH 6.3. Clear, wavy lower boundary.
B22 RSL No. 5597	23 to 34 inches, black (10YR 2/1, dry; 1.5/1, moist) clay. Moderate very coarse prismatic, breaking to weak coarse and medium blocky. Moderate continuous clay skins. Contains an occasional hard black shot concretion about 1/8 in diameter. Extremely hard, very firm, very plastic, very sticky. Few roots. Many slickensides. pH 6.3. Clear, irregular lower

OIL URVEY LABORATORY Riverside, California

SOIL TYPE Lobitos silt loam (taxadjunct) LOCATION San Mateo County, California

SOIL FAMILY: Pachic Ultic Argixeroll, fine, montmorillonitic, mesic.

SOIL NOS. S55Cal. 41-14 LAB. NOS. 55399-55403

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 8	A_{11}	0.2	2.0	5.8	11.5	5.8	38.7	36.0	27.2	21.9	4	cl
8-15	A_{12}	0.1	1.9	5.3	10.7	4.9	39.2	37.9	25.7	22.7	16	cl
15-28	B_2	0.7	2.1	2.9	5.7	3.7	42.9	42.0	19.2	30.2	52	sic
28-36	C_2	0.7	2.5	2.0	3.9	3.8	52.4	34.7	12.8	45.7	57	sicl
36+	D_r		not	analyzed								

pH 8C1b SATURATED PASTE 1:1	pH 8C1a 1:5	pH 8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	ORGANIC MATTER 6B1a NITROGEN %	ORGANIC MATTER C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	GYPSUM me./100g. SOIL	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
5.8		6.4	3.84	.310	12.4							17.6
5.7		6.4	2.42	.231	10.5							19.9
5.4		6.4	1.01	.112	9.0							26.3
5.3		6.3	0.69	.099	7.0							28.2

LOBITOS SILT LOAM (taxadjunct) 1/
S55Calif-41-14

Location: San Mateo County, California, San Mateo County SCD, 1/4 mile southeast of Lobitos. Date of Sampling: May 20, 1955.
Description by: Ralph E. Nelson. Collectors: Reuben E. Nelson, Ralph E. Nelson, Rudolph Ulrich, R. J. Wagner.
Classification: Pachic Ultic Argixeroll, fine, montmorillonitic, mesic.
Vegetation: Annual grasses, poison oak, wild blackberries, ferns and miscellaneous forbs. Climate: 25 to 40 inches rainfall. Parent Material: Silty sandstone or shale of the Purisima formation. Relief: Rolling hills to mountainous. Sample site located on 30 percent slope, slightly convex, toward northeast. Elevation: 320 feet. Drainage: Runoff rapid; internal drainage moderate.
Remarks: Many worms down to 15 inches.

HORIZON		DESCRIPTION
S55 Calif-41-14-1		
A_{11}	0- 8"	Very dark grayish brown to very dark gray (10YR 3/2-3/1, very dark brown to almost black 10YR 2/2-2/1 moist) silt [redacted]
		[redacted] friable, slightly plastic and very slightly sticky. There [redacted] There are abundant worm casts and roots. The reaction is 6.4. The lower boundary is clear and smooth.
S55 Calif-41-14-2		

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Supan cobbly gravelly loam (taxadjunct) **LOCATION** Amador County, California

SOIL FAMILY: Ultic Argixeroll, loamy-skeletal, mixed, mesic.

SOIL NOS. S 59Calif-3-1 **LAB. NOS.** 59265 - 59270

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-3	A_{11}	1.2	4.2	4.2	10.5	8.0	48.6	23.3	31.0	32.0	16	l
3-11	A_{12}	0.8	3.5	3.8	10.6	8.1	49.1	24.1	42.6	21.4	10	l-sil
11-18	B_1	0.8	3.3	3.4	9.0	7.5	43.9	32.1	28.1	29.0	17	cl
18-37	B_{21}	0.8	4.1	4.2	9.1	7.2	38.5	36.1	26.1	25.2	20	cl
37-49	B_{22}	1.0	7.1	5.7	11.6	7.9	38.2	28.5	27.2	25.8	9	cl
49-68	R	not analysed										

pH 8C1a H_2O 1:1	8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	6C1b "Free" Fe_2O_3 %	ELECTRICAL CONDUCTIVITY EC x 10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	GYPSUM mm./100g. SOIL	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.6	5.7		4.26	0.269	15.8	4.4						15.6
6.0	4.9		1.66	0.139	11.9	4.9						13.3
5.9	4.8		0.63	0.060	10.5	5.4						15.5
5.8	4.9		0.43	0.048	9.0	5.4						17.2
6.0	4.8		0.22	0.034		5.1						17.4

5A2a CATION EXCHANGE CAPACITY ← Na →	EXTRACTABLE CATIONS 6N2b Ca	6O2c Mg	6H2a H	6P2a Na	5B1a 6Q2a K	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE Na	K				MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
30.1	17.7	3.4	7.8	<0.1	2.2	75						
21.4	11.3	2.4	5.7	<0.1	2.1	74						
24.3	8.7	2.7	12.1	0.2	0.1	49						
21.3	8.1	3.7	9.7	<0.1	0.8	57						
21.7	7.4	5.3	8.2	<0.1	0.8	62						

1/ Percent of the whole soil

SUPAN COBBLY GRAVELLY LOAM (taxadjunct) 1/
S59 Calif-3-1

Collectors: Grant Kennedy, Reuben Nelson, Bob Sketchley, John Rogers.
Date of Sampling: June 15, 1959. Description by: Grant Kennedy.
Location: Amador County, California, near the County Airport, 200 feet E. of Highway 49, and 50 feet south of Access Road, in the SW1/4 of the NW1/4 of Section 17, T6N, R11E.
Elevation: 1,660 feet.
Relief: Nearly level ridgetop with steep sloping sides, slope 1 percent to SW.
Drainage: Well drained; slow runoff; moderately slowly permeable subsoils
Parent Material: Andesitic conglomerate of the Mehrten formation.
Biological Activity: Earthworms and burrowing rodents are active.
Climate: Mean annual rainfall 28 inches.
Vegetation: Annual grasses, forbs, a few oaks including filaree, clovers, wild radish, live oak.
Remarks: Medial developed Brunizem. The soil is relatively inextensive and was not sampled for the Bureau of Public Roads. Below about 11 inches about 50 percent of the mass consists of rounded cobbles and stones.
Classification: Ultic Argixeroll, loamy-skeletal, mixed, mesic.

HORIZON		DESCRIPTION
A_{11}	0-3"	Reddish brown (5YR 4/3) cobbly gravelly loam, dark reddish brown (5YR 2/2) when moist; moderate, medium and fine granular structure; soft when dry, friable when moist, nonsticky and slightly plastic when wet; abundant very fine and fine roots; many very fine and fine pores; few fine shot; neutral, pH 6.6; clear, smooth lower boundary.
A_{12}	3-11"	Reddish brown (5YR 4/3) heavy loam, dark reddish brown (5YR 2/3) when moist; massive; slightly hard when dry, friable when moist, slightly sticky and slightly plastic when wet; abundant very fine and fine roots; many very fine and fine pores; few fine shot; medium acid, pH 6.0; clear, wavy lower boundary.
B_1	11-18"	Reddish brown (5YR 4/4) gravelly clay loam, dark reddish brown (2.5YR 3/4) when moist; massive; hard when dry, friable when moist, slightly sticky and slightly plastic when wet; plentiful very fine and fine roots; many very fine, fine and few medium pores; few fine shot; common thin discontinuous clay films in pores; medium acid, pH 5.9; clear, wavy lower boundary.
B_{21}	18-37"	Reddish brown (5YR 4/4) gravelly clay loam, dark reddish brown (2.5YR 3/4) when moist; massive; hard when dry, firm to friable when moist, slightly sticky and plastic when wet; plentiful very fine and fine roots; common very fine and fine pores; thin continuous clay films in pores; medium acid, pH 5.8; gradual, wavy lower boundary.
B_{22}	37-49"	Reddish brown (5YR 4/4) slightly gritty clay loam, dark reddish brown (2.5YR 3/4) when moist; massive; hard when dry, friable to firm when moist, slightly sticky and plastic when wet; few very fine and fine roots; common very fine and fine pores; thin continuous clay films in pores; medium acid, pH 6.0; clear, wavy lower boundary.
R	49-68"	Pale brown (10YR 6/3) loamy sand (crushed), yellowish brown (10YR 5/4) when moist; massive; extremely hard when dry, brittle when moist; consolidated sandstone containing rounded andesitic cobbles; few fine pores; new clay films along fracture planes; medium acid, pH 6.0.

1/ The Supan series is a member of the fine-loamy, mixed, mesic family of Pachic Argixerolls. This pedon has slightly less than 75 percent base saturation in parts of the upper 30 inches, the mollic epipedon is less than 20 inches thick, and the argillic horizon has somewhat more than 35 percent rock fragments

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Supan cobbly loam (taxadjunct) **LOCATION** Amador County, California

SOIL FAMILY: Ultic Argixeroll, loamy-skeletal, mixed, thermic

SOIL NOS. S59Calif-3-2 **LAB. NOS.** 59271 - 59277

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	3A1 0.02-0.002	2A2 >2 1/	TEXTURAL CLASS
0-8	A_1	0.7	2.3	2.8	9.0	7.9	49.6	27.7	30.9	32.3	11	SYI cl-l-
8-16	B_{11}	0.8	2.0	2.6	8.6	7.9	47.1	31.0	29.5	31.4	9	cl
16-22	B_{12}	0.6	2.1	2.8	8.7	8.0	45.1	32.7	29.2	29.7	17	cl
22-38	B_2	0.6	2.9	3.4	9.4	7.7	39.2	36.8	26.9	26.0	14	cl
38-58	B_{31}	1.7	11.7	9.2	15.3	8.8	28.8	24.5	28.2	18.1	13	l-scl
58-70	B_{32}	1.1	9.5	8.1	13.0	8.0	35.2	25.1	27.1	23.6	30	l
70-75	R	3.2	19.1	12.8	13.5	7.3	27.5	16.6	24.4	17.9	58	sl

pH 8C1a	8C1c KCl	ORGANIC MATTER 6A1a	6B1a	6C1b "Free"	ELECTRICAL CONDUCTIVITY	$CaCO_3$	GYPSUM	MOISTURE TENSIONS		4B2

SUPAN COBBLY LOAM (taxadjunct) 1/
S59Calif-3-2

Collectors: Grant Kennedy, Reuben Nelson, Bob Sketchley, John Rogers.
Date of Sampling: June 15, 1959. Description by: Grant Kennedy.
Location: Amador County, California, on the Cazazzas property, along Highway 88, 100 feet north of the highway in the NE1/4 of the NW1/4 of Section 26, T6N, R10E.
Classification: Ultic Argixeroll, loamy-skeletal, mixed, thermic.
Elevation: 1,200 feet.

Drainage: Well drained, slow runoff; moderately slow permeability.
Parent Material: Andesitic conglomerate of the Mehrten formation.
Biological Activity: Burrowing rodents and earthworms have been active in these soils, particularly in the upper 2 feet where many krotovinas are present.
Climate: Mean annual rainfall about 25 inches.
Vegetation: Annual grasses, oak, and forbs including filaree, bur clover, soft chess and ripgut.
Remarks: This is a Reddish Brunizem near the Noncalcic Brown soil zone. This soil may have been cultivated at one time. It appears that some of the cobbles above the B_2 may have been removed. The profile contains about 20 to 30 percent cobbles.

HORIZON		DESCRIPTION
A_1	0-8"	Reddish brown (5YR 4/3) loam, dark reddish brown (5YR 3/2)
		massive below; soft when dry, friable when moist, nonsticky and nonplastic when wet; abundant very fine roots in top 2 inches, plentiful below; many very fine pores; slightly acid, pH 6.3; clear, smooth lower boundary.
B_{11}	8-16"	Reddish brown (5YR 4/3) clay loam, dark reddish brown (5YR 3/2) when moist; massive between krotovinas, strong fine granular within them; soft when dry, friable when moist, nonsticky and

OIL SURVEY LABORATORY Riverside, California

OIL TYPE Kitchen Creek sandy loam **LOCATION** San Diego County, California

SOIL FAMILY: Ultic Argixeroll, coarse-loamy, mixed, mesic.

SOIL NOS. S60Calif-37-1 **LAB. NOS.** 60391 - 60397

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 3	A11	16.2	16.0	8.5	18.9	13.4	20.1	6.9	37.2	8.0	23	cosl
3-12	A12	15.7	18.1	9.2	18.0	12.1	19.4	7.5	33.5	8.8	22	cosl
12-19	B21t	15.3	18.6	9.5	19.1	11.1	17.1	9.3	31.9	7.4	24	cosl
19-34	B22t	9.7	16.3	8.0	20.4	13.8	20.6	11.2	38.8	8.0	20	cosl
34-48	B3	12.6	21.8	10.5	21.0	11.1	15.2	7.8	32.7	6.1	21	cosl-lcos
48-64	C1	25.6	35.8	11.4	12.1	2.8	4.7	7.6	11.0	2.1	14	lcos-cos
64-72+	C2	14.8	42.5	13.9	12.8	3.1	4.5	8.4	12.0	1.8	21	lcos

pH 8C1a H_2O 1:1	pH 8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC x10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	Free Fe_2O_3 6C1b %	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.0	5.4		2.76	0.187	14.8				0.6			6.6
6.1	5.1		1.04	0.071	14.6				0.8			4.3
6.2	4.8		0.31	0.028					0.7			4.2
6.3	4.5		0.20	0.019					0.7			4.6
6.6	4.7		0.11						0.5			3.3
6.9	5.1		0.05						0.5			2.6
7.0	5.4		0.06						0.4			2.3

5A2a CATION EXCHANGE CAPACITY (+(Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Na 6P2a	K 6Q2a	BASE SAT. % 5C3	Sum of Cations 5A3a me./100g. soil	K	MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →							milliequivalents per liter →	
16.3	11.6	1.5	5.6	0.2	0.4	71	19.3		
11.2	6.5	0.6	3.9	0.2	0.2	66	11.4		
7.3	3.8	1.0	2.0	0.2	0.1	72	7.1		
8.1	5.1	1.6	1.8	0.2	0.1	80	8.8		
5.6	3.3	1.2	1.2	0.2	<0.1	80	5.9		
4.6	2.6	1.0	0.7	0.2	<0.1	84	4.5		
4.1	1.7	0.8	0.6	0.2	<0.1	82	3.3		

1/ Percent of whole soil

KITCHEN CREEK SANDY LOAM
S60Calif-37-1

Description: George Borst and Fred C. Eggers

Date of Sampling: November 8, 1960
Collectors: Reuben Nelson, Lothair Grant, George Borst, and Fred C. Eggers
Location: Approximately 0.15 mile east of southwest corner, sec. 18S, T. 11 S., R. 5 E., S.B.B.M., 75 feet north of Montezuma Valley Road, **San Diego County, Calif.**
Elevation: Approximately 4,100 feet Relief: Gently sloping, 4 percent.
Vegetation: Native chaparral Use: Range
Drainage: Well drained
Parent Material: Granitic alluvium
Climate: This soil occurs under about 17" annual rainfall. The mean annual temperature is about 57°, the mean January temperature about 44°, and the mean July temperature about 73°.
Remarks: The A horizon at this location is relatively thick and erosion is relatively slight, as compared with most soils in this series in San Diego County.
Reaction: pH is determined in the field with a Truog (Hellige) soil reaction kit.
Classification: **Ultic Argixeroll, coarse-loamy, mixed, mesic.**

HORIZON		DESCRIPTION
A11	0- 3"	Dark grayish brown (10YR 4/2) sandy loam, very dark grayish brown (10YR 3/2) when moist; moderate fine crumb structure; soft, very friable, nonsticky and nonplastic; abundant very fine random roots; neutral (pH 7.0); clear smooth boundary.
A12	3-12"	Dark grayish brown (10YR 4/2) sandy loam, very dark grayish brown (10YR 3/2) when moist; weak fine to medium crumb structure; slightly hard, friable, nonsticky and nonplastic; abundant fine random roots; few fine random tubular pores; medium acid (pH 6.0); clear wavy boundary.
B21t	12-19"	Brown (10YR 5/3) coarse sandy loam, dark brown (7.5YR 4/4) when moist; massive; slightly hard, friable, nonsticky and nonplastic; plentiful medium random roots; many fine random tubular pores; colloid in bridges between mineral grains; slightly acid (pH 6.2); gradual smooth boundary.
B22t	19-34"	Brown (10YR 5/3) coarse sandy loam, dark brown (7.5YR 4/4) when moist; massive; very hard, friable, nonsticky and nonplastic; plentiful medium random roots; many fine random tubular pores; colloid in bridges between mineral grains; slightly acid (pH 6.5); gradual smooth boundary.
B3	34-48"	Brown (10YR 5/3) stratified fine gravelly sandy loam, dark brown (10YR 4/3) when moist; massive; hard, friable, nonsticky and nonplastic; plentiful medium random roots; many fine random tubular pores; colloid in bridges between mineral grains; neutral (pH 6.7);

SOIL SURVEY LABORATORY Riverside, California

OIL TYPE: Kitchen Creek sandy loam **LOCATION** San Diego Co., California

SOIL FAMILY: Ultic Argixeroll, coarse-loamy, mixed, mesic.

SOIL NOS. S60Calif-37-2 **LAB. NOS.** 60398 - 60405

DEPTH INCHES	HORIZON	1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
		PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)										
0- 2	A11	10.7	20.8	8.9	20.7	11.7	20.8	6.4	35.4	9.4	15	cosl
2- 7	A12	11.8	18.6	7.6	16.7	10.1	21.4	13.8	32.4	9.1	13	cosl
7-13	B21t	12.7	18.0	6.8	13.6	8.4	22.2	18.3	27.2	11.4	14	cosl
13-32	B22t	12.0	17.7	7.3	16.1	9.7	24.6	12.6	30.4	13.5	16	cosl
32-43	B23t	12.4	18.4	7.1	16.6	10.0	23.4	12.1	30.4	13.1	20	cosl
43-55	B31	12.0	20.5	9.0	18.7	9.8	20.6	9.4	30.0	11.4	24	cosl
55-63	B32	12.8	24.0	9.1	18.7	9.7	18.9	6.8	29.5	9.8	27	cosl
63-73+	C1	18.8	23.6	8.6	16.9	9.4	18.2	4.5	28.1	9.2	25	lcos

8C1a pH H_2O 1:1	8C1c pH KCl 1:1		6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM 25°C.	$CaCO_3$ equivalent %	Free Fe_2O_3 6C1b %	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.0	5.2		5.47	0.200	27.4				0.7			8.0
6.7	5.6		1.00	0.064	15.6				0.9			5.7
6.5	5.0		0.56	0.044	12.7				1.1			6.7
6.7	4.4		0.17	0.015	[illegible]				0.9			5.9
6.7	4.3		0.10						0.7			5.5
6.7	4.4		0.10						0.7			5.4
6.8	4.4		0.06						0.6			4.2
6.9	4.5		0.04						0.5			3.6

5A2a CATION EXCHANGE CAPACITY (Na)	6N2b Ca	6O2c Mg	6H2a H	6P2a Na	6Q2a K	BASE SAT. % 5C3	5A3a Sum of Cations me./100g. soil	K				MOISTURE AT SATURATION %
	EXTRACTABLE CATIONS 5B1a											
	milliequivalents per 100g. soil →							milliequivalents per liter →				
19.0	10.5	1.7	8.3	0.2	0.5	61	21.2					
12.2	6.7	1.6	3.5	0.1	0.5	72	12.4					
13.0	7.2	1.6	3.7	0.2	0.3	72	13.0					
11.1	6.8	1.9	2.4	0.2	0.1	79	11.4					
9.9	5.9	2.2	1.8	0.2	0.1	82	10.2					
8.1	4.9	2.3	1.3	0.2	0.1	85	8.8					
7.4	4.1	2.2	0.9	0.2	0.1	88	7.5					
6.8	3.2	2.2	0.6	0.4	0.1	91	6.5					

1/ Percent of whole soil

KITCHEN CREEK SANDY LOAM
S60Calif-37-2

Description: George Borst and Fred C. Eggers
Date of Sampling: November 8, 1960
Collectors: Reuben Nelson, Lothair Grant, George Borst, and Fred C. Eggers
Location: Approximately 400 feet south-southwest of the center of sec. 13, T. 11 S., R. 4 E., S.B.B.M., about 0.8 mile north of Montezuma Valley Road, San Diego County, California.
Elevation: Approximately 4,100 feet
Relief: Strongly sloping, 10 percent Use: Range Vegetation: Native chaparral
Parent Material: Granitic alluvium Drainage: Well drained
Climate: This soil occurs under about 17" annual rainfall. The mean annual temperature is about 57°; the mean January temperature about 44°, and the mean July temperature about 73°.
Remarks: The A horizon at this location is somewhat thinner and erosion relatively more active than at site S60Calif37-2; these are considered to be representative of the soil as it occurs in San Diego County.

HORIZON		DESCRIPTION
A11	0- 2"	Brown (10YR 5/3) sandy loam, dark brown (10YR 3/3) when moist;

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Kitchen Creek loamy coarse sand **LOCATION** San Diego County, California

SOIL FAMILY: Ultic Argixeroll, coarse-loamy, mixed, mesic

SOIL NOS. S60Calif-37-4 **LAB. NOS.** 60411 - 60417

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 2	A11	18.1	27.7	11.8	16.1	6.4	14.1	5.8	22.3	6.3	33	lcos
2- 9	A12	17.1	23.3	12.1	20.5	8.2	12.7	6.1	26.2	5.9	18	lcos
9-13	A13	12.0	23.0	12.9	21.7	7.8	13.3	9.3	26.0	6.6	16	cosl-lcos
13-24	C1	11.3	24.8	12.0	20.7	7.5	12.1	11.6	24.6	5.9	32	cosl
24-34	C2	13.4	29.5	10.1	18.1	6.9	12.7	9.3	23.1	6.2	26	cosl-lcos
34-41	C3	18.0	35.0	10.6	17.6	5.2	7.9	5.7	19.1	3.1	28	lcos-cos
41-49	C4	12.8	34.8	11.3	20.2	7.0	9.4	4.5	24.2	3.4	33	" "

8C1a pH H_2O 1:1	8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	Free Fe_2O_3 6C1b %	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.8	6.0		2.95	0.156	18.9				0.5			5.4
7.2	6.2		1.40	0.090	15.6				0.7			4.0
7.2	5.9		0.70	0.051	13.7				0.6			4.4
6.9	5.2		0.26	0.025					0.8			6.0
6.9	4.1		0.28	0.018					0.8			6.1
6.7	4.2		0.18						0.7			4.2
6.5	3.8		0.11						0.6			7.4

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 6N2b Ca	6O2c Mg	6H2a H	6P2a Na	6Q2a K	BASE SAT. % 5C3	5A3a Sum of Cations me./100 g. soil					MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						milliequivalents per liter →					
14.5	11.1	1.2	2.9	0.1	0.5	72	15.8					
12.4	9.5	1.3	1.9	0.1	0.2	85	13.0					
12.0	7.6	1.1	1.9	0.1	0.3	83	11.0					
12.9	9.0	1.7	3.0	0.2	0.2	79	14.1					
18.0	13.2	2.6	2.3	0.2	0.1	87	18.4					
12.8	9.1	2.0	1.8	0.2	0.1	86	13.2					
11.4	8.0	2.1	1.5	0.2	<0.1	87	11.8					

1/ Percent of the whole soil

KITCHEN CREEK LOAMY COARSE SAND
S60Calif-37-4

Description: George Borst
Date of Sampling: November 10, 1960
Collectors: Reuben Nelson, Lothair Grant, Fred C. Eggers, John Wachtell, and George Borst
Location: Approximately 1,000 feet southeast of center of sec. 19, T. 11 S., R. 5 E., 800 feet east of private road along center of sec. 19, and 3,400 feet south of Montezuma Valley Road, **San Diego County, California.**
Elevation: 4,000 feet **Relief:** Strongly sloping, 12 percent
Vegetation: Native chaparral **Use:** Range
Drainage: Somewhat excessively drained
Parent Material: Deeply weathered granitic rock
Climate: This soil occurs under about 17" annual rainfall. The mean annual temperature is about 57°, the mean January temperature is about 44°, and the mean July temperature about 73°.
Remarks: A layer of angular quartz fragments, 2 to 6 mm. in size, occurs on

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Kitchen Creek loamy coarse sand **LOCATION** San Diego County, California

SOIL FAMILY: Ultic Argixeroll, coarse-loamy, mixed, mesic.

SOIL NOS. S61Calif-37-4 **LAB. NOS.** 6124 - 6131

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 4	A11	11.4	20.2	11.0	25.1	10.9	14.4	7.0	33.6	6.0	16	lcos
4-12	A12	12.3	19.8	10.2	24.8	10.9	14.7	7.3	33.6	6.5	12	lcos-cosl
12-21	A13	10.3	19.2	9.9	25.8	10.6	15.8	8.4	33.2	8.4	13	cosl
21-28	B1	10.0	19.4	8.8	24.5	10.1	18.1	9.1	32.2	10.7	11	cosl
28-42	B2	9.3	16.7	9.1	23.1	9.5	19.3	13.0	31.0	11.1	7	cosl
42-54	B3	11.5	21.1	10.1	22.0	8.5	16.6	10.2	28.2	9.8	9	cosl
54-67	C1	9.7	25.3	12.2	27.3	9.1	13.4	3.0	31.3	7.1	15	lcos
67-73+	C2	17.5	23.6	12.2	24.1	7.6	12.2	2.8	25.4	7.2	12	lcos-s

pH 8C1a H_2O 1:1	8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	Free Fe_2O_3 6C1b %	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.4	5.5		2.29	0.152	15.1				0.5			4.8
6.3	5.4		1.34	0.091	14.7				0.5			4.0
6.3	5.2		0.55	0.042	13.1				0.6			3.9
6.8	5.1		0.27	0.024					0.5			4.1
6.6	4.6		0.13						0.6			5.8
6.5	4.2		0.11						0.5			5.5
6.5	4.0		0.05						0.5			3.2
6.2	3.8		0.05						0.5			3.0

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 6N2b Ca	6O2c Mg	6H2a H	6P2a Na	5B1a 6Q2a K	BASE SAT. % 5C3	Sum of Cations me./100g soil 5A3a	K				MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →							milliequivalents per liter →				
14.9	9.5	0.6	4.7	0.2	0.3	69	15.3					
11.2	7.7	0.6	2.8	0.1	0.3	76	11.5					
8.9	5.4	0.4	1.9	0.2	0.4	77	8.3					
7.7	4.3	0.5	1.5	0.1	0.2	77	6.6					
8.7	5.6	1.0	2.0	0.2	0.1	77	8.9					
10.3	6.2	1.4	1.6	0.3	0.1	83	9.6					
7.0	3.9	1.1	1.0	0.3	<0.1	84	6.3					
6.1	2.7	1.7	1.0	0.3	0.1	83	5.8					

Sand includes considerable mica

1/ Percent of the whole soil

KITCHEN CREEK LOAMY COARSE SAND
S61Calif-37-4

Description: George Borst
Date of Sampling: January 20, 1961
Collectors: Reuben Nelson, Cliff Henry, and George Borst
Location: Approximately 800 feet southwest of east 1/4 corner, sec. 3, T. 17 S., R. 6 E., in Campo Indian Reservation, San Diego County, Calif.
Elevation: 4,560 feet Relief: Moderately sloping, 5 percent.
Vegetation: Native chaparral Use: Range
Drainage: Well drained
Parent Material: Very deeply weathered granitic rocks
Climate: This soil occurs under an average annual rainfall of about 15", a mean annual temperature of 58°, a mean January temperature of 49°, and a mean July temperature of 74°.
Remarks: This soil occurs on an old, moderately sloping upland plain, which appears to be a remnant of a pre-Pleistocene land surface. Erosion is very slight, the A horizon relatively thick, and the soil very consistent in profile features. This plain is bounded abruptly by steep, eroded slopes and deeply incised barrancas which penetrate almost to the stream divides.

HORIZON		DESCRIPTION
A11	0- 4"	Dark brown (10YR 3/3) loamy coarse sand, very dark brown (10YR 2/2) when moist; weak fine crumb structure; soft, very friable, nonsticky and nonplastic; abundant very fine vertical roots; many very fine interstitial pores; micaceous; slightly acid (pH 6.5); clear smooth boundary.
A12	4-12"	Dark brown (10YR 3/3) loamy coarse sand, very dark brown (10YR 2/2) when moist; massive; soft, very friable, nonsticky and nonplastic; plentiful fine random roots; slightly acid (pH 6.5); gradual smooth boundary.
A13	12-21"	Dark brown (10YR 4/3) loamy coarse sand, dark brown (10YR 3/3) when moist; massive; slightly hard, friable, nonsticky and nonplastic; few fine random roots; slightly acid (pH 6.5); clear smooth boundary.
B1	21-28"	Light brownish gray (10YR 6/2) sandy loam, brown (10YR 5/3) when moist; massive; hard, friable, nonsticky; slightly plastic; few

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Felton loam **LOCATION** San Diego County, California

SOIL FAMILY: Ultic Argixeroll, fine-loamy, mixed, mesic.

SOIL NOS. S61Calif-37-1 **LAB. NOS.** 613 - 619

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 7	A11	7.2	14.2	9.7	24.2	12.6	20.4	11.7	36.1	11.9	8	sl
7-12	A12	8.7	14.0	8.4	23.7	12.4	19.9	12.9	35.5	11.8	10	sl
12-17	A3	7.1	14.0	8.0	23.7	12.4	21.7	13.1	37.0	11.8	8	sl
17-25	B21	6.5	10.0	6.2	21.7	10.5	22.2	22.9	32.9	13.8	4	scl
25-35	B22	3.7	7.7	5.2	22.4	12.9	25.0	23.1	38.6	14.1	2	scl
35-47	B3	5.5	8.6	6.6	29.6	15.8	20.2	13.7	44.5	10.2	2	fsl
47-60+	C1	12.8	24.2	8.4	23.0	9.8	15.4	6.4	31.2	8.1	4	lcos

pH 8C1a H_2O 1:1	pH 8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	Atterberg limits Plastic limit Pct.	Liquid limit of 0.4 mm.	Plastic index	Free Fe_2O_3 6C1b %	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
5.9	5.1		1.85	0.132	14.0	N. P.	-	N. P.	1.6			6.5
5.9	4.9		0.75	0.065	11.5				1.7			6.3
5.8	4.8		0.66	0.059	11.2				1.7			6.4
5.5	4.3		0.29	0.034		22	36	14	1.9			11.5
5.5	4.0		0.21						1.7			13.7
5.4	3.5		0.11			N. P.	-	N. P.	1.1			11.0
5.3	3.6		0.05						0.6			6.1

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 6N2b Ca	6O2c Mg	6H2a H	6P2a Na	5B1a 6Q2a K	BASE SAT. % 5C3	Sum of Cations me./100g. soil 5A3a	K				MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →							milliequivalents per liter →				
13.1	6.7	1.2	5.4	0.1	0.5	61	13.9					
10.9	5.0	1.1	3.9	0.2	0.4	63	10.6					
10.5	4.6	1.1	3.5	0.2	0.4	64	9.8					
13.9	6.1	2.7	4.5	0.2	0.4	68	13.9					
20.8	10.5	4.5	5.2	0.2	0.2	75	20.6					
20.1	11.2	4.7	4.9	0.3	0.1	77	21.2					
11.0	4.9	2.1	2.2	0.3	<0.1	77	9.5					

1/ Percent of total soil

All sands have considerable mica.

FELTON LOAM
S61Calif-37-1

Description: George Borst
Date of Sampling: January 17 1961

OIL SURVEY LABORATORY Riverside, California

SOIL TYPE Chino loam (taxadjunct) **LOCATION** Merced County, California

SOIL FAMILY: Aquic Calcixeroll, fine-loamy, mixed, thermic

SOIL NOS. S55Calif. 24-2 **LAB. NOS.** 55281-288

DEPTH INCHES	HORIZON	1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
		PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1		
0- 6	A_1	0.4	2.0	2.2	12.7	17.6	34.6	30.5	43.8	17.9	1	cl
6-16	B_1	0.5	2.2	1.8	10.8	16.0	31.3	37.4	38.7	16.6	0	cl
16-27	B_{21}	1.2	2.7	2.3	10.0	14.9	34.6	34.3	37.0	19.8	1	cl
27-35	B_{22}	2.8	3.0	2.0	9.3	15.7	34.5	32.7	39.2	17.8	1	cl
35-43	B_{31ca}	3.5	3.0	1.7	7.5	15.1	36.0	33.2	37.4	19.2	2	cl
43-52	B_{32ca}	3.6	3.3	2.0	6.7	9.7	38.8	35.9	30.1	23.1	3	cl
52-60	B_{33ca}	4.3	4.1	3.2	14.4	10.8	34.5	28.7	32.8	22.8	2	cl
60+	C	0.0	0.4	2.7	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	0	[illegible]

CHINO LOAM (taxadjunct) 1/
S55Calif-24-2

Location: Merced County, California in San Luis SCD, 565 feet north of the intersection of State Highway No. 33-152 and Delta Road, 80 feet east of Delta Road, SW1/4 SW1/4 SW1/4, sec. 15, T. 10 S., R. 11 E.
Date of Sampling: April 20, 1955.
Description by: R. C. Huff. Collectors: Ralph E. Nelson, R. C. Huff, Rudolph Ulrich.
Classification: Aquic Calcixeroll, fine-loamy, mixed, thermic.
Vegetation: Virgin areas, partially barren with native vegetation of salt grass, iodine bush, soft chess and other salt tolerant annuals and bushes. Climate: Approximate 8½ inches annual rainfall, with dry summers and cool, moist winters. Parent Material: Mixed alluvium, mainly granitic, which is underlain by stratified sediments. Relief: Slightly undulating with mounds and depressions on a basin rim. The Chino soil is located on these slight mounds. The slope is 0-2 percent. Elevation: 96 feet. Drainage: Surface drainage is slow, internal drainage is slow.
Remarks: Roots down to 26 inches. Very few roots below this depth.

HORIZON		DESCRIPTION
A_1	0- 6"	Grayish brown to gray (10YR 5/2-5/1, very dark grayish brown to very dark brown 3/2-2/2 moist) loam. Very thin (¼") platy surface layer, which grades into moderate medium angular blocky. Very hard, friable, nonplastic and nonsticky. The soil effervesces slightly with HCl. The pH is 8+. This grades abruptly and smoothly into:
[illegible]	[illegible]	[illegible]

CONSECUTIVE PEDON NUMBER C 69 55 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES MERRILL SOIL NO. S69 CALI. 36-23 SAMPLE NO. 9 291-9 298 SAN BERNARDINO AREA

SOIL FAMILY AQUIC CALCIXEROLL FINE-SILTY,MIXED,THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B-3A1 1A2A

NO.	CM.		2- .05	.05- .002	LT .002	2-1	1- .05	.5- .25	.25- .10	.1- .05	.05- .02	.02- .002	2- .2U	LT .2U	NATE 3A1A	.2- .02	OF WHOLE SOIL	75- 20 PCT.	20-5 OF	5-2 LT 75
			PERCENT OF LESS THAN 2MM																	
9 291	0- 15	AP1	24.6	51.9	23.5	0.3	0.5	0.9	5.5	17.4	25.2	26.7			11	46.4	--A	--	--	--
9 292	15- 28	AP2	19.7	56.5	23.8	0.2	0.4	0.9	4.3	13.9	29.4	27.1			11	46.2	--A	--	--	--
9 293	28- 51	A13	24.6	51.9	23.5	0.1	0.6	1.2	5.3	17.4	27.0	24.9			11	48.1	--A	--	--	--
9 294	51- 63	A14	24.4	49.6	26.0	0.2	0.4	0.9	4.9	18.0	25.1	24.5			13	46.6	--A	--	--	--
9 295	63- 79	C 1CA	26.1	46.3	27.6	0.1	0.6	1.5	7.3	16.6	22.5	23.8			16	44.0	--G	--	--	--
9 296	79-110	C 2CA	28.7	46.0	25.3	0.9	3.3	3.3	9.2	12.0	20.9	25.1			19	38.2	--G	--	--	--
9 297	110-125	C 3	37.9	46.0	16.1	0.2	1.3	3.6	16.0	16.8	22.4	23.6			9	48.9	--A	--	--	--
9 298	125-150	C 4	43.6	44.8	11.6	0.1	1.0	4.1	21.0	17.4	21.5	23.3			4	52.0	--A	--	--	--

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR-BON 6A1A	NITRO-GEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	19-2 6E1B	EXT. IRON FE 6C28	ATTERBERG LIMIT EST LL 4F2	PL 4F1	1/3 4A1F	BULK DENSITY 1/3 BAR 4A1F	DRY 4AH	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTENSBLTY LEF 4D1	LE 4D1	PH H2O 1/1 8C1A	H2O 1/10 8C1A	CACL2 8C1E
		PERCENT		PERCENT		PCT.	PCT.NO.40		GRAMS PER CC				PERCENT			PERCENT				
9 291	0- 15	3.46		26						1.07	1.17	1.00		35.9	16.1	3.0	3.0	8.0		7.7
9 292	15- 28	3.11		25					1.12			1.00			16.5			8.1		7.7
9 293	28- 51	2.16		22						1.17	1.25	1.00		32.2	14.8	2.2	2.2	8.1		7.8
9 294	51- 63	1.88		25					1.20			1.00			15.1			8.2		7.8
9 295	63- 79	1.23		33					1.67			1.00			15.2			8.3		7.8
9 296	79-110	0.64		43						1.67	1.72	1.00		18.6	12.4	1.0	1.0	8.3		7.8
9 297	110-125	0.22		18					1.67			1.00			6.0			8.2		7.9
9 298	125-150	0.20		9					1.67			1.00			5.3			8.2		7.5

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 4N2E	MG 5B4A 6D2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT.	C E C NA- OAC 5A2A	NH4- OAC 5A6A	WATER EXTRACT FROM SATURATED PASTE 8A1 EC 8A1A	CA 6N1B	MG 6O1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	B 6T1A	PH SAT. PASTE 8C1B
9 291	0- 15	30.5	10.4	0.2	1.0	42.1	100+		19.8	1.29										
9 292	15- 28	27.2	10.8	0.5	0.4	38.9	100+		19.9	1.06										

MERRILL SILT LOAM
S69Calif-36-23

Location: San Bernardino County, California; about 300 feet northwest of Eucalyptus and Central Avenue, SW¼ SE¼ SW¼ of sec. 14, T.2S., R.8W. Date of Sampling: June 10, 1969. Description by: A. A. Knecht, W. D. Nettleton, W. Z. Brock. Collectors: A. A. Knecht, W. D. Nettleton, W. Z. Brock.
Classification: Aquic Calcixeroll, fine-silty, mixed, thermic.
Vegetation: Bindweeds, forbs, annual and perennial grasses. Climate: Average annual precipitation about 16 inches. Mean temperature about 61°F. Parent Material: Mixed alluvium. Topography: Alluvial fan, nearly level, about 1 percent slope. Elevation: 600 feet. Drainage: Somewhat poorly drained; slow runoff; moderately slow to slow permeability. Area above flood plain. The water table has been lowered and wetness is not a problem. Soil Moisture: Profile was dry throughout when sampled.
Remarks: Field reaction was determined by the Truog method. All horizons were sampled and analyzed at the Riverside Soil Survey Laboratory. The A13 horizon (11-20") and C2ca horizon (31-43") were sampled for the California Division of Highways.

HORIZON	DESCRIPTION
	(0-15 cm.)
Ap1 RSL 69291	0 to 6 inches, gray (5Y 5/1) silt loam, dark olive gray (5Y 3/2) moist; weak coarse subangular blocky structure; slightly hard, friable, slightly sticky, slightly plastic; many very fine roots; many very fine interstitial pores; strongly effervescent; moderately alkaline (pH 8.0); clear smooth boundary.
	(15-28 cm.)
Ap2 RSL 69292	6 to 11 inches, gray (10YR 5/1) silt loam, very dark gray (10YR 3/1) moist; weak coarse subangular blocky structure; slightly hard, friable, slightly sticky, slightly plastic; common very fine roots; common very fine tubular pores; strongly [illegible]
A13 RSL 69293	11 to 20 inches, gray (10YR 5/1) silt loam, very dark gray (10YR 3/1) moist; weak coarse angular blocky structure; slightly hard, friable, slightly sticky, slightly plastic; common very fine roots; common very fine tubular pores; strongly effervescent; moderately alkaline (pH 8.0); gradual wavy boundary.
	(51-63 cm.)
A14 RSL 69294	20 to 25 inches, gray (10YR 6/1) silt loam, dark gray (10YR 4/1) moist; weak coarse angular blocky structure; slightly hard, friable, slightly sticky, slightly plastic; common very fine roots; common very fine tubular pores; strongly effervescent; moderately alkaline (pH 8.0); gradual wavy boundary.
	(63-79 cm.)
C1ca RSL 69295	25 to 31 inches, light gray (2.5Y 7/2) loam, grayish brown (2.5Y 5/2) moist; massive; slightly hard, friable, sticky, slightly plastic; very few roots; few very fine tubular pores; 25 percent of the mass is large, irregularly shaped extremely hard lime concretions; violently effervescent; moderately alkaline (pH 8.0); gradual wavy boundary.
	(79-110 cm.)
C2ca RSL	31 to 43 inches, light gray (2.5Y 7/2) loam, grayish brown (2.5Y 5/2) moist; massive; slightly hard, friable, sticky, plastic; few very fine tubular pores; 50

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aridic Calcixeroll, fine-silty, mixed, mesic

SOIL Calita clay loam variant SOIL Nos. S54Calif-18-1 LOCATION Lassen County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 54290 - 54296

Depth (In.)	Horizon	1B1b Total: Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Sand: Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt: 0.05–0.02	Int. III (0.02–0.002)	Int. II (0.2–0.02)	3A1 (2–0.1)			Coarse fragments: 2A2 > 2 Pct.	2 – 19	19 – 76
		Pct. of < 2 mm															Pct. of < 76 mm	
0-6	Ap	14.5	61.2	24.3	0.0	0.3	0.6	4.3	9.3	21.8	39.4		5.2			0		
6-12	A3	14.3	61.0	24.7	0.0	0.3	0.8	5.2	8.0	13.7	47.3		6.3			0		
12-20	B2	14.8	55.2	30.0	0.0	0.4	1.1	6.1	7.2	20.2	35.0		7.6			0		
20-25	B3	14.0	55.1	30.9	0.0	0.3	0.7	5.3	7.7	10.4	44.7		6.3			0		
25-42	C1ca	15.7	45.7	38.6	0.0	0.1	0.4	5.3	9.9	9.3	36.4		5.8			0		
42-53	C2	23.5	36.6	39.9	0.0	0.2	1.3	13.3	8.7	0.4	36.1		14.8			0		
53-66	C3			Incomplete dispersion - gypsum												0		

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		6E1b Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct	Pct.	4B2 15 bar Pct.				pH 8C1b Sat. Paste	8C1a 1:10
0-6	2.85	0.201	14.2		5								22.5				7.7	8.3
6-12	2.17	0.160	13.6		3								21.6				7.7	8.6
12-20	1.35	0.114	11.8		9								24.3				7.6	8.6
20-25	0.68	0.072	9.5		28								22.6				7.7	8.6
25-42	0.35				44								22.5				7.7	8.8
42-53	0.26				20								29.4				7.6	8.6
53-66	0.19				15								25.7				7.6	8.0

Depth (In.)	Extractable bases: Ca	Mg	5B1b Exch. Na 6P2a	5B1b Exch. K 6Q2a		5A2a CEC NaOAc			Water extract from saturated paste: 6N1a Ca+Mg 6O1a		6P1a Na	6Q1a K		CO_3	HCO_3	Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-6			5.4	4.8		41.4			36.7		61.0	3.3						7.30
6-12			4.9	3.9		42.1			27.8		40.0	1.9						4.95
12-20			5.1	3.5		42.4			30.5		42.5	1.5						5.75
20-25			4.6	2.3		34.7			30.4		41.0	1.3						5.66
25-42			4.7	1.6		30.5			19.2		33.7	0.6						4.50
42-53			6.9	2.0		45.8			45.1		40.0	1.0						5.50
53-66			5.7	1.8		38.8			34.5		34.2	1.1						5.31

Depth (In.)	8A Water at Saturation Pct.	5D1 Exchangeable Na Pct.		6F1a Gypsum Pct.		
0-6	63.2	13		1		
6-12	63.6	12		tr.		
12-20	67.2	12		1		
20-25	67.8	13		2		
25-42	66.8	15		tr.		
42-53	75.2	15		tr.		
53-66	73.9	15		12		

CALITA CLAY LOAM variant [1/]
S54Calif-18-1

Location: Lassen County, California, in Honey Lake Valley SCD on Paul Milton ranch, 1/4 mile south of Clinton School or 100 feet east of quarter corner between Sections 16 and 17, T. 29 N., R. 13 E.
Collectors: S. B. Johnson, J. Langdon, R. E. Nelson, S. Porter, R. C. Roberts, and L. R. Wohletz.
Date of Sampling: October 18, 1954.
Vegetation: The area where the sample was taken was planted to tall wheat grass. Some big sage along fence line. Weeds were growing on the irrigation ridges where salt had accumulated.
Climate: At Susanville, about 7½ miles west, the January average temperature is 30.7° F, the July average is 69.5° F, and the growing season is 142 days. Average annual precipitation for a 30 year period is 17.8 inches divided as follows: January 3.64, February 2.66, March 2.64, April 1.05, May 0.84, June 0.51, July 0.22, August 0.19, September 0.50, October 1.01, November 1.77, and December 2.77. The average annual precipitation at sample site is probably about 16 inches and the average snowfall is about 65 inches. Parent Material: Stratified lake-laid sediments mostly fine-texture of "Lahontan" age. Relief: Nearly level to very gently sloping basin rim or old basin floor. Leveled for irrigation. Elevation: 4050 feet. Drainage: Moderately slow drainage, both external and internal. Water table at about 4 feet. Biological activity: Many angle worms at about 2 feet and evidence of worm activity from a depth of 4 inches to 30 inches.
Classification: Aridic Calcixeroll, fine-silty, mixed, mesic.

HORIZON	DESCRIPTION
Ap RSL No. 54290	0 to 6 inches, grayish brown (10YR 5/2, very dark brown 10YR 2/2 moist) moderately plastic clay loam which is hard when dry. This layer is moderately coarse granular with a tendency to be platy. Calcareous, and pH 8.5.
A3 RSL No. 54291	6 to 12 inches, grayish brown (10YR 5/2, very dark grayish brown 10YR 3/2 moist) silty clay loam which is weak medium prismatic that breaks into moderate medium subangular blocky aggregates. This horizon is plastic when wet and very hard when dry. It has many worm holes 1/16 inch in diameter. It is calcareous and has a pH of 8.4. Gradual change to layer below.
B2 RSL No. 54292	12 to 20 inches, grayish brown (10YR 5/2, dark brown 10YR 3/3 moist) heavy silty clay loam which is extremely hard when dry and plastic when wet. It is weakly prismatic and readily breaks into moderate medium subangular blocky peds. It also has many worm holes and has a pH of 8.8. Gradual change to layer below.
B3 RSL No. 54293	20 to 25 inches, very pale brown (10YR 7/3, brown 10YR 4/3 moist) plastic silty clay loam which is faintly mottled. This horizon is hard when dry and weak fine to medium angular blocky. It has a pH of 8.8.
Clca RSL No. 54294	25 to 42 inches, light gray (10YR 7/2, brown 10YR 5/3 moist) very highly calcareous moderately plastic silty clay loam which is hard when dry. It has weak medium angular blocky aggregates with a pH of 8.6. Rather abrupt change to layer below.
C2 RSL No. 54295	42 to 53 inches, same color and consistence as layer above but coarser textured. This layer consists of brittle cemented lenses that do not slake in acid or water but readily become wet when put in a liquid. The pan is probably cemented by silica and lime and possibly sodium. pH 8.6.
C3 RSL No. 54296	53 to 66 inches, same color as layer above but much looser and only slightly plastic silt loam. This layer has many gypsum crystals and some lime. It consists mostly of brittle lenses and layers of loose friable material. pH 8.6.

[1/] The Calita series is in a fine-loamy, mixed, mesic family of Aridic Calcixerolls. This pedon is in a fine-silty family of the same subgroup.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Pachic Calcixeroll, fine-silty, mixed, thermic

SOIL Linne silty clay loam (taxadjunct) SOIL Nos. S65Calif-27-4 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65468 - 65474

Depth (cm)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (<0.002)	Size class and particle diameter (mm) 3A1 Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: (0.05-0.01)	Int. III (0.02-0.002)	Int. II (0.2-0.02)	2-0.1			Coarse fragments 1A2a: b/ > 2 Pct. of whole soil	2-19	19-76
		Pct. of <2 mm															Pct. of <76mm	
0-10	A11	19.0	50.5	30.5	0.7	0.9	1.3	5.9	10.2	30.6	19.9	44.6	8.8			G		
10-41	A12	18.8	48.0	33.2	0.8	0.9	1.3	5.5	10.3	28.0	20.0	42.0	8.5			G		
41-81	A13	19.5	46.2	34.3	1.4	0.9	1.1	5.5	10.6	26.4	19.8	40.7	8.9			G		
81-91	A14ca	18.2	46.9	34.9	0.4	1.1	1.2	5.2	10.3	25.6	21.3	39.3	7.9			G		
91-115	C1ca	31.7	46.2	22.1	3.8	5.2	4.9	7.7	10.1	16.2	30.0	30.6	21.6			G		
115-150	C2	38.1	46.1	15.8	2.8	5.7	4.8	9.9	14.9	20.2	25.9	41.4	23.2			G		
115-150	a/ shale	34.0	46.8	19.2	0.7	3.1	3.2	11.8	15.2	21.0	25.8	44.3	18.8			G		

Depth (cm)	6A1a Organic carbon c/ Pct.	6B1a Nitrogen Pct.	C/N	Carbonate as $CaCO_3$: 6E1b <2 mm Pct.	3A1a <0.002 mm Pct.	Atterberg limits: Plastic limit	Liquid limit	Plastic index	Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	3B2 Cm	Water content: 4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEf cm/cm	4D1 Extensibility COLE cm/cm	pH: 8C1b Saturated paste	8C1a H_2O 1:1	8C1c IN KCl 1:1
						Pct. of <0.4 mm												
0-10	3.90	0.330	12	1					1.42	1.67	1.00	27.6	14.5	0.056	0.056		6.9	
10-41	1.16	0.114	10	4		23	42	19	1.23	1.43	1.00	26.8	14.1	0.050	0.050		8.2	
41-81	0.79			6					1.22	1.40	1.00	27.5	15.0	0.046	0.046		8.0	
81-91	0.61			10					1.21	1.36	1.00	30.1	17.4	0.039	0.039		8.0	
91-115	0.27			31					1.07	1.08	1.00	45.2	13.7	0.004	0.004		8.5	
115-150	0.16			15		37	41	4	1.49	1.58	1.00	28.7	18.0	0.020	0.020		8.3	
a/115-150	0.18			8									21.6				8.0	

Depth (cm)	Extractable bases: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	5B1a	6H2a Ext. acidity	5A3a CEC NH_4OAc	5D2 Exchangeable Sodium	5G1 Base saturation NH_4OAc	8A Water at saturation	Water extract from saturated paste 8A1: 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K	6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1a SO_4	8A1a Electrical conductivity
	meq/100g							Pct.	Pct.	Pct.	meq/liter								mmho/cm
0-10	39.6	4.8	0.2	0.9			35.5		100+										
10-41	40.4	4.0	0.2	0.4			32.4		100+										
41-81	39.1	4.6	0.5	0.4			31.5		100+										
81-91	35.4	6.2	2.0	0.3			29.6	5	100+	64.2	8.2	2.0	10.6	0.1	0.8	1.8	17.7	0.1	2.32
91-115	28.3	4.4	3.0	0.2			20.2	8	100+	55.7	13.2	6.7	24.3	0.1	1.3	1.4	38.3	0.7	4.51
115-150	25.1	6.6	4.4	0.2			20.3	10	100+	62.4	20.1	9.8	39.8	0.1	-	1.3	63.6	3.0	7.0

Depth (cm)	Ratios to clay 8D1: CEC NH_4OAc	Ext. iron	15 bar water	d/ Clay fraction mineralogy 3A1: 7A2 X-ray e/	7A3 DTA Pct.
0-10	1.15		0.47	MT5, KK2	
10-41	0.98		0.43		
41-81	0.98		0.47		
81-91	0.93		0.54		
91-115	1.19		0.80		
115-150	1.35		1.20		
a/115-150			1.14	MT5, MI2, KK2	

a/ Sample of C2 from which carbonate coatings were removed

b/ G = ground whole. Coarse shale fragments dispersed readily

c/ 14.8 kg of organic carbon per square meter to a depth of 1 meter

d/ Mineral code
- MT = montmorillonite
- MI = mica
- KK = kaolinite

e/ Approximate weight fraction (X-ray)
- 5 = more than half
- 2 = one-twentieth to one-fifth

LINNE SILTY CLAY LOAM (taxadjunct) 1/
S65Calif-27-4

Location: Monterey County, California. About 6 miles N. of King City on Montgomery Ranch, 3800 feet NE from reservoir, 1 mile NE from main buildings, about 70 feet uphill from dirt road and 400 feet down from ridge top. Date of Sampling: May 18, 1965.

Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich. Site selection by T. D. Cook.

Classification: Pachic Calcixeroll, fine-silty, mixed, thermic. (Regosolic Chestnut).

Vegetation: Range, grass, ripgut brome, soft chess, burclover, fiddleneck; all dry. Climate: Precipitation 14 inches; mean annual temperature 60°F., mean January temperature 50°F., mean July temperature 67°F. Parent Material: Soft calcareous shale and sandstone, Plio-Pleistocene non-marine. Topography: Moderately steep rounded hills, slightly convex middle of NNE-facing slope of 26 percent. Elevation: 600 feet. Drainage: Well drained; permeability is moderately slow; runoff is medium. Soil Moisture: Slightly moist; wilting point throughout.

Remarks: pH determinations by Truog Kit. Paired sample is Linne silty clay loam, S65Calif-27-5-1 through 7. The color of the A1 horizons, A11, A12, and A13, tend a little toward 10YR 4/2 when dry; typical Linne colors have a chroma of 1. Asterisks indicate horizons sampled for California Division of Highways.

HORIZON	DESCRIPTION
A11 RSL No. 65468	0-4 inches (0-10 cm.), dark gray (10YR 4/1) light silty clay loam, very dark gray (10YR 3/1) moist; weak medium plates readily breaking to moderate medium angular blocks; very hard (dry), firm (moist), sticky and plastic (wet); abundant very fine roots; common very fine tubular pores; weakly effervescent with disseminated carbonate; platy structure at surface due to cattle trampling; gradual smooth boundary.
A12 * RSL No. 65469	4-16 inches (10-41 cm.), dark gray (10YR 4/1) light silty clay loam, very dark gray (10YR 3/1) moist; strong moderate and coarse subangular blocky structure; hard (dry), friable (moist), sticky and plastic (wet); plentiful very fine roots; many very fine interstitial and many very fine and few fine tubular pores; strongly effervescent with disseminated carbonate; clear irregular boundary.
A13 RSL No. 65470	16-32 inches (41-81 cm.), dark gray (10YR 4/1) light silty clay loam, very dark gray (10YR 3/1) moist; moderate coarse subangular blocky structure; hard (dry), friable (moist), sticky and plastic (wet); plentiful very fine roots; many very fine interstitial, common very fine and few fine tubular pores; strongly effervescent with disseminated carbonate and many fine filaments of carbonate toward bottom of horizon; gradual irregular boundary.
A14ca RSL No. 65471	32-36 inches (81-91 cm.), dark grayish brown (10YR 4/2) light silty clay loam, very dark grayish brown (10YR 3/2) moist; weak medium angular blocky structure; hard (dry), friable (moist), sticky and plastic (wet); few very fine roots; common very fine interstitial and few very fine tubular pores; violently effervescent with disseminated carbonate, fine carbonate filaments cover 65 percent of ped faces; abrupt wavy boundary.
C1ca RSL No. 65472	36-45 inches (91-115 cm.), very pale brown (10YR 7/3) calcareous shale and sandstone with many white, thick to thin coats of carbonate on rock joints; soft (dry); carbonate appears to have accumulated from above; gradual wavy boundary.
C2 RSL No. 65473 & 65474	45-60 inches (115-150 cm.), similar to above but with few thin carbonate coats on rock joints.

1/ The Linne series is in a fine-loamy, mixed, thermic family of Calcic Pachic Haploxerolls. This pedon is in the fine-silty family and has a calcic horizon, and is calcareous in all parts of all horizons above the calcic horizon.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Posa silt loam **LOCATION** Madera County, California

SOIL FAMILY: Orthidic Durixeroll, fine-loamy, mixed, thermic

SOIL NOS. S58Calif-20-1 **LAB. NOS.** 58251 - 58258

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	3A1 0.02-0.002	2A2 > 2	TEXTURAL CLASS
0-4	A11	0.3	1.7	2.1	7.2	8.3	51.8	28.6	35.6	29.2	0	sil-cl sicl
4-12	A12	0.4	3.4	4.7	14.4	15.5	47.1	14.5	53.0	19.1	0	l
12-19	C1sfca	1.5	4.3	3.4	9.1	11.6	45.3	24.8	37.6	25.2	0	l
19-22	C2sica	4.2	5.4	3.4	9.5	13.6	49.9	14.0	43.1	26.8	0	l
22-30	C3sica	13.9	18.0	6.7	10.0	7.0	31.6	12.8	22.8	21.5	0	cosl
30-37	C4ca	6.0	8.5	4.9	13.0	15.4	42.3	9.9	42.0	24.9	0	l
37-47	C5	4.4	23.5	4.4	11.6	3.9	44.7	7.5	29.7	24.7	0	l
47-62	C6	0.1	2.4	4.2	31.3	28.5	31.6	1.9	73.6	10.1	0	fsl

pH 8C1b SATURATED		8C1a	ORGANIC MATTER 6A1a ORGANIC	6B1a NITRO		6I1a CO_3	8A1a Elec. Conduc.	6E1b $CaCO_3$ equiv.	6J1a HCO_3	MOISTURE TENSIONS 1/10	1/3	4B2 15

POZO SILT LOAM
S58Calif-20-1

Location: Madera County, California. 0.3 mile southeast of intersection of power line and dike and 200 feet east of dike in NW1/4, Sec. 8, T. 11 S., R. 14 E. Date of Sampling: October 29, 1958. Description by: R. Ulrich. Collectors: R. E. Nelson, Jess Glouser, Gene Novak, and R. Ulrich. Classification: Orthidic Durixeroll, fine-loamy, mixed, thermic. Vegetation: Dominantly lippia and saltgrass with spikeweed, alkali weed, vinegar weed (turpentine weed), wild rye, and some cocklebur. In range pasture use. Climate: About 9 inches rainfall. Mean annual temperature about 63° F. Parent Material: Largely water transported, sorted, and deposited sediments from granitic sources in Sierra Nevada to the eastward. Position is basin or basin rim by University of California physiographic terminology. Associated soils largely Fresno, El Peco, and Rossi. Topography: Essentially a level plain, less than 1 percent slope, with very gently undulating or hummocky (gilgai) microrelief. Elevation: 120 feet as given by Rose Farm USGS Topographic Quadrangle

sheet. Drainage: Imperfect at present time, possibly poorly drained sometime in the past before widespread pump irrigation and surface diversion from canals and reservoirs higher on the alluvial fans. No water table above 8 feet by auger sampling, but soil was moist below the hardpan and dry above. Internal drainage very slow to slow due to hardpan. Surface runoff very slow.

DESCRIPTION

A11 RSL No. 58251 — 0-4 inches, gray (10YR 5/1) silt loam, very dark gray (10YR 3/1) moist; weak medium and fine granular; hard, friable, slightly plastic, nonsticky; abundant coarse saltgrass stolons and abundant secondary fine root distribution pattern throughout horizon; many extremely fine and very fine interstitial pores; few worm casts; noncalcareous to dilute acid; reaction not determined in field, see lab data; clear wavy lower boundary.

A12 RSL No. 58252 — 4-12 inches, similar colored silt loam; very weak very fine granular moist drying essentially massive; hard, friable, slightly plastic, nonsticky; few coarse saltgrass stolons and plentiful fine roots; many extremely fine and very fine tubular pores, many extremely fine and very fine interstitial pores; few worm casts; noncalcareous to dilute acid; abrupt, slight wavy lower boundary.

C1sica RSL No. 58253 — 12-19 inches, white (10YR 8/1) heavy loam, light brownish gray (2.5Y 6/2) moist; massive; hard, friable, very slightly plastic, nonsticky; plentiful fine roots; common very fine and few fine tubular pores; moderately calcareous; clear smooth lower boundary.

C2sica RSL No. 58254 — 19-22 inches, similar colored loam; massive to weak medium platy or laminated presumably due to stratification; hard, friable, very slightly plastic, nonsticky; few fine roots with some platelets or laminae having root mat concentrations; few extremely fine and very fine tubular pores; moderately calcareous; abrupt smooth lower boundary.

C3sicam RSL No. 58255 — 22-30 inches, light gray (5Y 7/2) strongly cemented lime-silica hardpan, olive gray (5Y 5/2) moist, with a very thin white (5Y 8/1) capping (light gray (5Y 7/1) moist, on the upper surface and alternating in very thin seams throughout the horizon; upper surface of hardpan continuously covered with a root mat and secondary root mats of a less continuous nature occur along cracks

OIL SURVEY LABORATORY Riverside, California

OIL TYPE Crouch loamy coarse sand (taxadjunct) **LOCATION** San Diego County, California

SOIL FAMILY: Typic Haploxeroll, coarse-loamy, mixed, mesic

SOIL NOS. S61Calif-37-5 **LAB. NOS.** 6132 - 6140

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1	2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	
0- 3	A11	13.8	21.5	12.4	25.3	9.8	11.6	5.6	30.6	5.5	15	lcos
3-11	A12	13.8	20.8	10.8	25.2	10.1	12.4	6.9	31.5	6.1	14	lcos
11-18	A13	13.3	18.3	9.4	23.6	10.8	16.3	8.3	32.8	8.5	16	cosl
18-22	B1	14.7	19.6	9.2	21.8	9.2	17.6	7.9	30.1	9.5	18	cosl
22-30	B21t	11.5	18.8	8.8	22.3	10.6	19.3	8.7	33.2	10.3	15	cosl
30-38	B22t	13.8	19.0	9.6	21.4	9.1	18.9	8.2	30.1	10.5	16	cosl
38-46	B3	13.1	19.7	9.6	22.0	9.0	19.4	7.2	30.7	10.8	17	cosl
46-62	C1	12.2	30.7	10.2	20.5	8.1	13.2	5.1	26.3	6.3	29	lcos
62-68	C2	7.5	24.4	10.6	23.6	9.7	17.6	6.6	31.7	9.1	26	cosl-lcos

CROUCH LOAMY COARSE SAND (taxadjunct) 1/
S61Calif-37-5

Description: George Borst
Date of Sampling: January 23, 1961
Collectors: Reuben Nelson, Lothair Grant, and George Borst
Location: Approximately 1,500 feet west-northwest of southeast corner, sec. 27, T. 16 S., R. 6 E., in Manzanita Indian Reservation, San Diego County, Calif.
Elevation: 4,480 feet Relief: Moderately sloping, 6 percent.
Vegetation: Native chaparral Use: Range
Drainage: Well drained, moderately rapid subsoil permeability
Parent Material: Very deeply weathered granitic rock
Climate: This soil occurs under an average annual rainfall of about 15", a

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haploxeroll, coarse-loamy, mixed, thermic

SOIL San Andreas sandy loam SOIL Nos. S65Calif-27-6 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65598

Depth (In.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int III (0.02-0.002)	Int II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	* Coarse fragments 1A2a > 2 Pct.	2-19	19-76
		Pct. of < 2 mm															Pct of < 76mm	
0-3	A1***															28	26	2 **

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content: Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility LEf Pct.	4D1 Extensibility LE Pct.	pH: 8C1a H_2O 1:1		
0-3	1.76	0.100	18															

Depth (In.)	Extractable bases 5B4a: 6N2a Ca	6O2d Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A6a NH_4OAc	5A3a Sum		6G1e KCl Ext. Al^{+++}						Base saturation: Pct.	Pct.
	meq/100 g																

Depth (In.)	Clay mineralogy (<0.002 mm.) 3A1: Mont.	Verm.	Mica	Kaolin.	
	X-ray				

* From characterization sample (5Kg) (except as noted).
** 2-19 mm. from characterization sample, 19-76 mm. from field volume estimate.
*** Composite A horizon sample.

SAN ANDREAS SANDY LOAM
S65Calif-27-6

Location: Monterey County, California, about 10 miles NE of King City on Montgomery Ranch, 300 feet due west from Pinalito VABM, about 75 feet down from ridge top. (Site #1 North Slope). Date of Sampling: May 18, 1965.

Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, and T. D. Cook.

Classification: Typic Haploxeroll, coarse-loamy, mixed, thermic. (minimal) Noncalcic Brown.

Vegetation: Blue oak, chamise, pine blue grass, wild buckwheat, red brome, annual fescue, wild oats, melic, digger pine. Climate: Precipitation is 14 inches; mean annual temperature 60°F., January mean 58°F., July mean 67°F. Parent Material: Granodiorite. Topography: Steep and very steep slopes, locally slightly concave, 50 percent North facing slope, smooth, 75 feet below ridge crest. Elevation: 1500 feet. Drainage: Somewhat excessive, moderately rapid permeability, medium runoff. Soil Moisture: Very slightly moist, wilting or below.

Remarks: Composite sample of A1 by orchard type auger to depth of 8 inches. Composite sample included individual samples (borings) from a plot with 4 rows 12 feet apart. Each row was sampled 6 times at 6-foot intervals. The paired site for this location on south slope S65Calif-27-7-1. Other samples in this study are south

CONSECUTIVE PEDON NUMBER C 69 27 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES SAN ANDREAS TAXADJUNCT SOIL NO. S69 CALI. 31-22 SAMPLE NO. 9 131-9 136 PLACER COUNTY AREA

SOIL FAMILY TYPIC HAPLOXEROLL COARSE-LOAMY,MIXED,THERMIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2- .05	SILT .05- .002	CLAY LT .002	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 SANDS VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	SILTS CSI .05- .02	FSI .02- .002	CLAYS CO. 2- .2U	FINE LT .2U	CARBONATE 3A1A	1A2A INT. II .2- .02	COARSE FRAGMENTS PCT. OF WHOLE SOIL I 75- 20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75		
9 131	0- 13	A11	74.3	18.6	7.1	13.3	21.3	13.8	18.6	7.3	7.9	10.7				24.2	5W	--	--	5
9 132	13- 38	A12	73.7	18.6	7.7	12.0	20.0	14.1	19.4	8.2	7.5	11.1				25.5	6W	--	--	6
9 133	38- 61	B2	72.7	19.2	8.1	13.2	19.3	12.6	18.6	9.0	7.5	11.7				26.2	10W	--	--	10
9 134	61- 74	B3	72.3	18.1	9.6	13.6	20.1	13.0	18.1	7.5	7.1	11.0				23.8	10W	--	--	10
9 135	74- 94	C 1	72.0	14.6	13.4	17.3	22.1	11.2	15.7	5.7	5.5	9.1				19.0	10W	--	--	10
9 136	94-120	C 2	83.2	8.1	8.7	24.9	27.3	12.1	14.6	4.3	3.2	4.9				14.4	19W	--	--	19

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A	NITROGEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	19-2 6E1B	EXT. IRON FE 6C2B	ATTERBERG LIMIT LL 4F2	PL 4F1	BULK DENSITY EST 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4AH	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTENSBLTY LEF 4D1	LE 4D1	PH H2O 1/1 8C1A	H2O 1/10 8C1A	CACL2 8C1E
		-PERCENT--		-PERCENT-		PCT.	PCT.	NO.40	GRAMS PER CC				-----PERCENT---			-PERCENT--				
9 131	0- 13	1.90	0.118							1.34	1.35	0.97		4.4	4.1	0.2	0.2	6.2		5.5
9 132	13- 38	0.61	0.043							1.63	1.63	0.96		7.5	3.3			5.9		5.2
9 133	38- 61	0.19								1.67	1.67	0.94		8.0	3.0			5.9		5.2
9 134	61- 74								1.73			0.94			3.7			5.7		5.0
9 135	74- 94									1.79	1.82	0.93		10.2	5.5	0.6	0.5	5.4		4.9
9 136	94-120									1.97	2.02	0.85		9.2	4.4	0.8	0.7	5.3		4.7

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6Q2D	NA 5B4A 6P2A	K 6Q2A	SUM	BASE SAT.	C E C NA- OAC 5A2A	NH4- OAC 5A6A	WATER EXTRACT FROM SATURATED PASTE 8A1 EC 8A1A	CA 6N1B	MG 6O1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	B 6T1A	PH SAT. PASTE 8C1B
		MEQ/100G					PCT.	-MEQ/100G		MMHO	MEQ/1000 ML								PPM	
9 131	0- 13	6.6	0.6	0.1	0.3	7.6	79		9.6											
9 132	13- 38	4.4	0.7	0.1	0.2	5.4	78		6.9											
9 133	38- 61	2.3	0.7	0.1	0.2	3.3	75		4.4											
9 134	61- 74	2.4	0.8	0.1	0.2	3.5	83		4.2											
9 135	74- 94	3.9	0.7	0.1	0.2	4.9	82		6.0											
9 136	94-120	5.5	0.8	0.1	0.1	6.5	90		7.2											

SAMPLE NO.	DEPTH CM.	WATER AT SAT. 8A PCT.	EXCHANGBLE NA 5B4B MEQ.	NA 5D2 PCT.	SAR 5E	GYPSUM AMT. REQ. 6F1B PERCENT	6P3A	CLAY MINERALOGY LESS THAN 0.002 MM X RAY I 7A2E	II 7A2E	III 7A2E	DTA IV 7A2E	PCT. 7A3	LT. .0002 MM X RAY I 7A2E	II 7A2E	III 7A2E	MINERAL CODE	RELATIVE AMOUNT
9 131	0- 13		0.1	1				MI 3	KK 3							MI MICA	1 TRACE
9 132	13- 38		0.1	1												KK KAOLINIT	2 SMALL
9 133	38- 61		0.1	2				MI 3	KK 3								3 MODERATE
9 134	61- 74		0.1	2													4 ABUNDANT
9 135	74- 94		0.1	2													5 DOMINANT
9 136	94-120		0.1	1													6 INDETER.

SAMPLE PREPARATION CODES
W=DATA DETERMINED ON WHOLE SOIL AND RECALCULATED TO 2 MM BASIS. GRAVEL FROM SLAKED SUBSAMPLE.

SAN ANDREAS COARSE SANDY LOAM (TAXADJUNCT) 1/
S69Calif-31-22

Location: Placer County, California. From Sierra College Blvd., 1.5 miles east of Douglas Blvd. Pit location approximately 80 feet north and directly across from Ken L. Services, 2,700 feet west and 80 feet north of the SE corner of sec. 3, T.10N., R.7E. Date of Sampling: May 9, 1969.

Description by: L. A. Brittan, T. M. Ryan and G. M. Kennedy. Collectors: K. W. Flach, W. D. Nettleton, G. M. Kennedy, T. M. Ryan, L. A. Brittan.

Classification: Typic Haploxeroll, coarse-loamy, mixed, thermic.

Vegetation: Blue oak, soft chess, ripgut, and wild oats. Climate: Annual average precipitation 22 inches; mean annual air temperature 62°F. Parent Material: Weathered granodiorite. Topography: Low foothill, gently rolling, 8 percent with west aspect. Elevation: 310 feet. Drainage: Well drained, slow to medium runoff and moderately rapid permeability. Soil Moisture: Dry to 24 inches, slightly moist below.

Remarks: A stone line 2 to 3 inches thick was overlying the C1 horizon and composed mainly of quartz fragments. Roots were matted between C1 and C2 horizons. The C1 horizon was noticeably easier to dig than the C2 horizon. Perhaps clay films make this material harder to dig compared to the decomposed granodiorite in Vista soils.

HORIZON	DESCRIPTION
	(0-13 cm.)
A11 RSL 69131	0 to 5 inches, grayish brown (10YR 5/2) coarse sandy loam, very dark brown (10YR 2/2) moist; moderate medium granular structure; soft, very friable, nonsticky, nonplastic; many very fine and fine vertical roots; many very fine tubular and interstitial pores; slightly acid (pH 6.5); clear smooth boundary.
	(13-38 cm.)
A12 RSL 69132	5 to 15 inches, grayish brown (2.5Y 5/2) coarse sandy loam, dark brown (10YR 3/3) moist; massive; slightly hard, very friable, nonsticky, nonplastic; many very fine and common fine and medium vertical roots; common very fine tubular and interstitial and few fine tubular pores; medium acid (pH 6.0); clear smooth boundary.
	(38-61 cm.)
B2 RSL 69133	15 to 24 inches, pale brown (10YR 6/3) coarse sandy loam, brown and dark brown (10YR 4/3) moist; massive; hard, very friable, nonsticky, nonplastic; common very fine, medium and coarse random roots; common very fine tubular and interstitial, and few fine tubular pores; neutral (pH 7.0); clear smooth boundary.
	(61-74 cm.)
B3 RSL 69134	24 to 29 inches, very pale brown (10YR 7/3) coarse sandy loam, brown (10YR 5/3) moist; massive; hard, very friable, nonsticky, nonplastic; few fine and medium horizontal roots; common very fine tubular and interstitial, and few fine tubular pores; few thin clay films line pores and bridge mineral grains; slightly acid (pH 6.5); abrupt wavy boundary.
	(74-94 cm.)
C1 RSL 69135	29 to 37 inches, very pale brown matrix (10YR 7/4) with variegated mineral grains of weathered granodiorite, reddish yellow matrix (7.5YR 6/6) moist; rock fabric structure; hard, friable, nonsticky, nonplastic; few fine and medium horizontal roots; common very fine tubular and interstitial pores; many moderately thick clay films line pores and bridge mineral grains; slightly acid (pH 6.5); gradual wavy boundary.
	(94-120 cm.)
C2 RSL 69136	37 to 47 inches, variegated weathered granodiorite; rock fabric structure; hard, friable, nonsticky, nonplastic; very few fine and coarse random roots; few very fine tubular and interstitial pores; many moderately thick clay films line pores and bridge mineral grains; slightly acid (pH 6.5); gradual wavy boundary.

1/ The San Andreas series is in a coarse-loamy, mixed, thermic family of Typic Haploxerolls. This pedon is in the same family and subgroup but has more coarse sand than is typical of the series.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haploxeroll, coarse-loamy, mixed, thermic

SOIL Sheridan loam (taxadjunct) SOIL Nos. S65Calif-27-20 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65559 - 65564

Depth	Horizon	1B1b Total Sand	Total Silt	Total Clay	Sand Very	Sand Coarse	Sand Medium	Sand Fine	Sand Very fine	Silt Int. III-	Silt Int. II (0.2-0.02)	Silt (2-0.1)	3A1a Carbon-	3A1a Non-carbon-	* Coarse fragments 1A2a >2	2-19	19-76

SHERIDAN LOAM (taxadjunct) 1/
S65Calif-27-20

Location: Monterey County, California, NE corner, NW¼, NW¼, Sec. 21, T18S, R1E, about 3.5 miles on old Coast Road from Bixby Bridge on Hwy. 1. 1 mile up Rancho Calera Road

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haploxeroll, coarse-loamy, mixed, thermic

SOIL Sheridan loam (taxadjunct) SOIL Nos. S65Calif-27-23 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65575 - 65579

Depth (In.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Size class and particle diameter (mm) 3A1 Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	* Coarse fragments 1A2a > 2 Pct.	2-19	19-76	
		Pct. of < 2 mm															Pct. of < 76mm		
0-7	A11	73.0	16.3	10.7	18.7	22.0	9.5	15.1	7.7	6.7	9.6	22.0	65.3	0	11	11	11	0	**
7-18	A12	71.7	17.1	11.2	15.1	21.0	10.6	16.8	8.2	7.1	10.0	23.9	63.5	0	11	15	15	0	**
18-35	A13	71.1	17.2	11.7	14.3	20.6	9.6	17.6	9.0	7.1	10.1	25.0	62.1	0	12	16	16	0	**
35-42	C1	71.3	15.6	13.1	14.2	23.2	9.6	16.7	7.6	7.4	8.2	23.5	63.7	0	13	16	16	0	**
42-60+	C2	78.4	12.9	8.7	15.4	26.2	10.6	18.3	7.9	7.4	5.5	24.8	70.5	0	9	13	13	0	**

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1	
0-7	0.97	0.077	13			1.63	1.64		0.93		10.3	4.3	0.002	0.002	6.3	
7-18	0.52	0.043	12			1.49	1.54		0.91		12.6	5.0	0.013	0.011	6.5	
18-35	0.39					1.55	1.58		0.90		11.6	5.5	0.006	0.005	6.6	
35-42	0.22					1.70	1.78		0.89		14.4	7.7	0.015	0.013	6.5	
42-60+	0.17					1.77	1.83		0.91		12.5	5.2	0.012	0.011	6.7	

Depth (In.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1a NH_4OAc	5A2a NaOAc		6G1d KCl Ext. Al^{+++}		Atterberg limits: Plastic limit	Liquid limit	Plastic index			Base saturation 5C1 NH_4OAc Pct	Pct.
	meq/100 g											Pct. of <0.4 mm						
0-7	8.8	2.0	0.3	0.1	11.2		11.7										96	
7-18	9.8	2.7	0.4	0.1	13.0			16.5					-	N.P.			95	
18-35	10.4	3.4	0.4	0.1	14.3		14.4										99	
35-42	14.2	5.8	0.5	0.1	18.1		19.2					20	30	10			94	
42-60+	8.6	3.4	0.4	0.1	12.5		11.6										100+	

Depth (In.)	Clay mineralogy (<0.002 mm) 7A: Mont.	Verm.	Mica	Kaolin	
	X-ray				

* From characterization sample (5Kg) (except as noted).

** 2-19 mm. from characterization sample, 19-76 mm. from field volume estimate.

SHERIDAN FINE GRAVELLY LOAM (taxadjunct) 1/
S65Calif-27-23

Location: Monterey County, California, center NE¼ SE¼ SE¼ Sec. 30, T15S, R6E. About 3.5 miles up Johnson Canyon Road from Armstrong Ranch main gate. (16" rainfall, inland, grass, south slope). Date of Sampling: May 26, 1965.

Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich; site selection by T. D. Cook.

Classification: Typic Haploxeroll, coarse-loamy, mixed, thermic. Regosol (Brunizemic).

Vegetation: Range grass, soft chess, annual fescue, ripgut brome, popcorn flower, lupine. Climate: Precipitation is 14 to 20 inches. Mean annual temperature is 60°F., mean January 50°F., mean July 66°F. Parent Material: Weathered diorite (gruss). Topography: Slightly convex moderately steep slope on south exposure, about 20 percent and about 100 feet below ridgecrest. Elevation: 2850 feet. Drainage: Well drained, moderately permeable, medium runoff. Soil Moisture: Dry (All), moist below.

Remarks: The closely related soil about 300 feet away on the North slope is S65Calif-

mile away. Other related soils are S65Calif-27-26, North inland and S65Calif-27-20, S65Calif-27-21, and S65Calif-27-22 near the coast. Asterisks indicate horizons sampled for California Division of Highways.

HORIZON	DESCRIPTION
All RSL 65575	0 to 7 inches, dark grayish brown (10YR 4/2) fine gravelly loam, very dark grayish brown (10YR 3/2) moist; weak coarse subangular blocky structure; slightly hard, very friable, slightly sticky and slightly plastic; plentiful

very fine roots; many very fine interstitial and few fine tubular pores; slightly acid (pH 6.2); surface 1/2 inch soft and much darkened by organic matter, horizon somewhat compacted by trampling, 17 percent fine gravel by volume and 40 percent coarse sand; gradual smooth boundary.

A12 7 to 18 inches, dark grayish brown (10YR 4/2) fine gravelly light loam, very

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haploxeroll, coarse-loamy, mixed, thermic
SOIL Sheridan loam (taxadjunct) SOIL Nos. S65Calif-27-25 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65586 - 65591

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	* Coarse fragments 1A2a > 2 Pct.	2 - 19 (Pct. of < 76mm)	19-76 (Pct. of < 76mm)	
		Pct. of < 2 mm																	
0-5	A11															15	15	0	**
5-15	A12															19	19	0	**
15-26	A13	70.0	19.3	10.7	15.6	19.6	9.3	17.2	8.3	9.0	10.3	26.2	61.7	0	11	21	21	0	**
26-41	C1															19	19	0	**
41-53	C2															26	26	0	**
53-72+	C3															26	26	0	**

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	Bulk density: 4A1f 1/3 bar g/cc	Bulk density: 4A1h Oven dry g/cc	Bulk density: g/cc	Water content: Pct.	Water content: 4B1c 1/3 bar Pct.	Water content: 4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1a H_2O 1:1		
0-5	1.10	0.094	12											6.4		
5-15	0.53	0.040	13											6.4		
15-26	0.38										6.3			6.5		
26-41	0.16													6.4		
41-53	0.10													6.7		
53-72+	0.06													6.5		

Depth (In.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1a NH_4OAc	5A3a Sum	6G1d KCl Ext. Al^{+++}	Base saturation: 5C1 NH_4OAc Pct.	Pct.
	meq/100 g										
0-5	8.3	1.4	0.3	0.2	10.2					96	
5-15	8.8	1.7	0.3	0.2	11.0		11.7			94	
15-26	10.6	2.4	0.4	0.1	13.5		14.2			95	
26-41	18.1	5.6	0.4	0.2	24.3		25.5			95	
41-53	19.4	6.4	0.4	0.1	26.3		28.0			94	
53-72+	18.1	6.7	0.5	0.1	25.4		26.3			97	

Depth (In.)	Clay mineralogy (<0.002 mm.) 7A: Mont.	Verm.	Mica	Kaolin	
	X-ray				

* From characterization sample (5Kg)(except as noted).
** 2-19 mm. from characterization sample, 19-76 mm. from field volume estimate.

SHERIDAN FINE GRAVELLY SANDY LOAM (taxadjunct) 1/
S65Calif-27-25

Location: Monterey County, California, SW¼ NE¼ NE¼ Sec. 29, T15S, R6E, 4.5 miles up Johnson Canyon Road from Armstrong Ranch main gate, 600 feet N. at dirt road, about 150 feet below ridge top. (Inland, grass, south slope). Date of Sampling: May 26,1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich; site selection by T. D. Cook.
Classification: Typic Haploxeroll, coarse-loamy, mixed, thermic. Regosol (brunizemic).
Vegetation: Range, grass, soft chess, lupine, ripgut brome, popcorn flower.
Climate: Precipitation is 14 to 20 inches; mean annual temperature 60°F., mean January 50°F., mean July 66°F. Parent Material: Weathered diorite (grus). Topography: Slightly convex, moderately steep slope on south exposure, about 27 percent and about 150 feet below ridge top. Elevation: 3100 feet. Drainage: Well drained, moderately rapid permeability, medium runoff. Soil Moisture: All and A12 are dry. Moist below.
Remarks: All Sheridan soils sampled in this study are rather similar in texture, though individual pits may vary from light loam to coarse sandy loam. Mean soil temperature is assumed to be more than 59°F., though coastal sites may be slightly cooler. The matched profile to this site is S65Calif-27-23 and is about a mile away on a south slope. Profile S65Calif-27-26, a few hundred feet away on a north slope; S65Calif-27-24 is also on a north slope. Other Sheridan profiles under grass and brush under 35-45 inches of rainfall near the coast are numbers 20, 21 and 22.

HORIZON	DESCRIPTION
A11 RSL 65586	0 to 5 inches, dark grayish brown (10YR 4/2) fine gravelly heavy sandy loam, very dark brown (10YR 2/2) moist; weak coarse subangular blocky structure; slightly hard, very friable, slightly sticky and slightly plastic; plentiful very fine roots; common very fine interstitial and few fine tubular pores; slightly acid (pH 6.5); loose layer at top of strong fine granular structure with upper 1/2 inch mixed organic and mineral soil; gradual smooth boundary.
A12 RSL 65587	5 to 15 inches, dark grayish brown (10YR 4/2) fine gravelly heavy sandy loam, very dark brown (10YR 2/2) moist; moderate medium granular and some medium subangular blocky structure; slightly hard, very friable, slightly sticky and slightly plastic; plentiful very fine roots; common very fine interstitial and few very fine, common fine and medium tubular pores; medium acid (pH 6.0); diffuse smooth boundary.
A13 RSL 65588	15 to 26 inches, dark grayish brown (10YR 4/2 toward 4/1) fine gravelly heavy sandy loam, very dark brown (10YR 2/2) moist; strong medium granular structure; slightly hard, very friable, slightly sticky and slightly plastic; plentiful very fine roots; many very fine interstitial pores; slightly acid (pH 6.5); considerable rodent activity just above C1; clear irregular boundary.
C1 RSL 65589	26 to 41 inches, brown (7.5YR 5/4) fine gravelly heavy sandy loam, brown and dark reddish brown (7.5YR 4/4, 5YR 3/4) moist; massive; shows definite rock structure; difficult to cut and break with hand tools; hard, friable, sticky and slightly plastic; few very fine roots; common very fine interstitial and common very fine tubular pores; common thin clay films in pores and continuous colloidal stains; medium acid (pH 6.0); clear smooth boundary.
C2 RSL 65590	41 to 53 inches, reddish brown (5YR 4/4 moist or dry) with 20 percent darker mineral grains; massive; rock structure, difficult to cut or break with hand tools; very hard, firm, slightly sticky and nonplastic; very few very fine roots; few very fine interstitial pores; common colloidal stains; slightly acid (pH 6.5); few large rock joints have thin clay films; gradual wavy boundary.
C3 RSL 65591	53 to 72 inches, brown (7.5YR 5/4) fine gravelly sandy loam, brown and reddish brown (7.5YR 4/4, 5YR 4/4) moist; massive; difficult to break with hand tools; very hard, firm; no roots; slightly acid (pH 6.5).

1/ The Sheridan series is in a coarse-loamy, mixed, thermic family of Pachic Haploxerolls. This pedon has a mollic epipedon less than 20 inches thick.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haploxeroll, coarse-loamy, mixed, thermic
SOIL Sheridan loam (taxadjunct) SOIL Nos. S65Calif-27-26 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65592 - 65597

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand Very coarse (2-1)	Sand Coarse (1-0.5)	Sand Medium (0.5-0.25)	Sand Fine (0.25-0.1)	Sand Very fine (0.1-0.05)	Silt (0.05-0.02)	Silt Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	* Coarse fragments 1A2a > 2 Pct.	2 - 19 (Pct. of < 76 mm)	19-76 (Pct. of < 76 mm)	
		Pct. of < 2 mm																	
0-6	A11															12	12	0	**
6-20	A12															14	14	0	**
20-34	A13	76.0	16.4	7.6	15.0	22.1	11.1	19.5	8.3	7.3	9.1	25.5	67.7	0	8	16	16	0	**
34-47	A14															22	22	0	**
47-54	C1															19	19	0	**
54-68+	C2															19	19	0	**

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct	Bulk density 4A1f 1/3 bar g/cc	Bulk density 4A1h Oven dry g/cc	g/cc	Water content Pct.	Water content 4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1b Saturated Paste	pH 8C1a H_2O 1:1	pH 8C1a H_2O 1:10
0-6	1.01	0.077	13												6.3	
6-20	0.54	0.042	13												6.2	
20-34	0.45										5.0				6.4	
34-47	0.42														6.4	
47-54	0.37														6.3	
54-68+	0.27														6.2	

Depth (In.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	5B1a 6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 8A1 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-6	6.1	1.2	0.2	0.2	7.7			8.6										
6-20	8.8	1.3	0.3	0.1	10.5			11.3										
20-34	9.7	1.9	0.3	0.1	12.0			12.4										
34-47	13.8	3.4	0.4	0.1	17.7			15.8	0.7	0.2	1.2	0.1		-	0.9	0.4		0.23
47-54	6.3	1.6	0.4	0.1	8.4				0.5	0.4	1.2	0.1		-	0.9	0.4		0.19
54-68+	5.8	1.5	0.4	0.3	8.0			6.5	0.6	0.2	0.8	tr.		-	1.1	0.5		0.22

Depth (In.)	8A Water at Saturation Pct.	5D2 Exchangeable Na Pct	5B1b Exch. Na meq./100g.	Gypsum Pct.	5C1 Base Sat. NH_4OAc Pct.	Clay mineralogy (<0.002 mm.) 7A Mont.	Verm.	Mica	Kaolin	
						X-ray				
0-6					90					
6-20					93					
20-34					97					
34-47	31.9		0.4		100+					
47-54	30.8		0.3		100+					
54-68+	29.7	6	0.4		100+					

* From characterization sample (5Kg) (except as noted).

** 2-19 mm. from characterization sample, 19-76 mm. from field volume estimate.

SHERIDAN SANDY LOAM (taxadjunct) 1/
S65Calif-27-26

Location: Monterey County, California. SW corner NE¼ NE¼ Sec. 29, T15S, R6E, 4.5 miles up Johnson Canyon Road from Armstrong Ranch main gate, 600 feet north of dirt road. (16" rainfall, inland, grass, north slope). Date of Sampling: May 26, 1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich: site selection by T. D. Cook.
Classification: Typic Haploxeroll, coarse-loamy, mixed, thermic. Regosol (brunizemic). (See temperature data).
Vegetation: Range, grass, soft chess, ripgut brome, fiddleneck and scattered weeds.
Climate: Precipitation is 14 to 20 inches. Mean annual temperature is 60°F., mean January 50°F., mean July 66°F. Parent Material: Weathered diorite (gruss).
Topography: Steep, smooth, north-facing slope of 34 percent, about 50 feet below top of rounded ridge crest. Elevation: 3100 feet. Drainage: Well drained, moderately rapid permeability, medium runoff. Soil Moisture: All is dry; moist below.

Remarks: The closely related paired profile for this sample is about a mile away, S65Calif-27-24; a south exposure a few hundred feet away is S65Calif-27-25 and the one a mile away is S65Calif-27-23. Other Sheridan samples are from near the coast under 35 to 45 inches of rainfall and are numbers 20, 21 and 22.

HORIZON	DESCRIPTION
A11 RSL 65592	0 to 6 inches, dark grayish brown (10YR 4/2) heavy sandy loam, very dark brown (10YR 2/2) moist; moderate fine and medium subangular blocky structure; only very slightly compacted by cattle; slightly hard, very friable, slightly sticky and slightly plastic; abundant very fine roots; many very fine interstitial and few fine tubular pores; medium acid (pH 6.0); gradual smooth boundary.
A12 RSL 65593	6 to 20 inches, dark grayish brown (10YR 4/2) sandy loam, very dark brown (10YR 2/2) moist; moderate fine subangular blocky structure; slightly hard, very friable, slightly sticky and slightly plastic; plentiful very fine roots; many very fine interstitial and common fine tubular pores; medium acid (pH 6.0); diffuse smooth boundary.
A13 RSL 65594	20 to 34 inches, dark grayish brown (10YR 4/2) sandy loam, very dark brown (10YR 2/2 toward 3/2) moist; strong medium granular structure; slightly hard, very friable, slightly sticky and slightly plastic; plentiful very fine roots; many very fine interstitial and few fine tubular pores; slightly acid (pH 6.3); clear irregular boundary.
A14 RSL 65595	34 to 47 inches, brown (10YR 5/3) light sandy loam, darker brown (10YR 4/3) moist; weak fine subangular blocky structure; slightly hard, very friable, nonsticky and nonplastic; very few very fine roots; many very fine interstitial pores; considerable rodent activity and mixing, 20 to 30 percent, upper A soil material; clear wavy boundary.
C1 RSL 65596	47 to 54 inches, brown (7.5YR 5/4) coarse sand, strong brown and brown (7.5YR 5/6 and 4/4) moist; massive; has rock structure and is easily cut with hand tools; hard, friable, nonsticky and nonplastic; no roots; common very fine interstitial pores; few colloidal stains and few thin clay films in pores; slightly acid (pH 6.5); gradual irregular boundary.
C2 RSL 65597	54 to 68 inches, black and white clean decomposed diorite, easily cut with hand tools.

1/ The Sheridan series is in a coarse-loamy, mixed, thermic family of Pachic Haploxerolls. This pedon has a mollic epipedon less than 20 inches thick.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Sweeney silty clay loam (taxadjunct) **LOCATION** San Mateo County, California

SOIL FAMILY: Typic Haploxeroll, fine-loamy, mixed, mesic

SOIL NOS. S55-Calif.-41-10 **LAB. NOS.** 55373-55379

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	3A1 0.02-0.002	2A2 > 2 2/	TEXTURAL CLASS
0- 7	A1	0.9	4.5	6.4	23.8	14.4	25.1	24.9	41.9	13.2	1	scl
7-16	B2	0.6	3.6	5.3	22.6	13.7	25.3	28.9	40.2	13.7	1	scl
16-22	B3	0.8	7.1	7.8	26.1	15.3	23.9	19.0	42.9	13.2	0	fsl
22-33	C1	1.4	9.9	9.5	27.8	14.2	24.5	12.7	42.4	13.5	0	fsl
33-45	C21	3.1	15.7	11.8	27.6	13.9	19.5	8.4	40.4	9.6	2	fsl
45-57	C22	3.0	18.1	13.5	28.9	13.0	17.8	5.7	39.7	8.0	0	sl
57-65	C23	1.9	16.3	14.6	30.3	13.6	18.2	5.1	41.0	8.1	0	sl

8C1b pH SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	Free Fe_2O_3 6C1b %	GYPSUM me./100g. SOIL	4A3a Bulk Density g/cc		4B2 Soil water 15 atmos. Pct.
6.4		6.9	1.48	0.112	13.2		0.4	1.1		1.33		12.8
6.3		6.9	1.12	0.096	11.7		0.5	1.2		1.48		14.0
6.7		7.2	0.40	0.042	9.5		0.4	1.0				14.0
6.7		7.3	0.19				0.4	1.1				14.0
6.9		7.4	0.08				0.3	0.7		1.65		11.6
7.0		7.2	0.05				0.3	0.7				11.0
7.2		7.3	0.09				0.3	0.7				10.2

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	5B1b Exch. Na 6P2a	5B1b Exch. K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg	6J1a 1/ HCO_3	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
47.7	21.0	19.8	6.6	1.1	0.2	86	0.8	Tr.	2.8	2.4	0.4	46.0
52.8	23.1	20.0	6.3	0.9	0.2	88	0.8	0.1	3.3	2.2	0.3	45.6
56.2	25.8	23.1	6.5	0.9	0.2	89	0.7	Tr.	2.5	1.5	0	41.4
56.4	28.0	24.4	5.5	0.9	0.1	91	0.6	Tr.	2.4	1.2	0	40.1
53.5	26.8	22.6	6.6	1.1	0.1	89	0.6	Tr.	1.4	0.9	0	39.5
52.5	26.7	22.2	6.1	1.2	0.1	89	0.6	Tr.	4.2	0.9	0	39.1
47.7	24.3	21.1	4.9	1.2	0.1	91	0.7	Tr.	2.0	0.7	0	37.5

1/ No $CO_3^{=}$ found

2/ Percent of the whole soil

SWEENEY SILTY CLAY LOAM (taxadjunct) 1/
S55Calif-41-10

Location: San Mateo County, California. In the San Mateo County SCD, west side of Highway No. 5 (Skyline Highway), 4.5 miles south of Sky Londa. See Sheet 24, Soil Survey of San Mateo Area, California, May 1961. Date of Sampling: May 17, 1955
Description by: Ralph E. Nelson. Collectors: Reuben E. Nelson, Ralph E. Nelson, and Rudolph Ulrich.
Classification: Typic Haploxeroll, fine-loamy, mixed, mesic.
Vegetation: Annual grasses; soft chess, wild oat, lupine. Parent Material: Basaltic material classified as diabase. Relief: Hilly to mountainous with gently to moderately steep rolling hills. The slope at the sample site location was about 25 percent, plain, toward the southeast. Elevation: 2,100 feet. Drainage: Surface drainage is rapid, internal drainage is moderate. Climate: 35-40 inches of rainfall.
Remarks: Many worms in surface 7-inch and occasional to few worms to a depth of 16 inches, krotovinas very abundant to a depth of 22 inches.

HORIZON		DESCRIPTION
A_1	0-7"	Dark grayish brown (10YR 4/2; very dark grayish brown to very dark brown, 10YR 2.5/2, moist) light silty clay loam. Moderate medium to fine granular structure. Hard, friable, slightly plastic and nonsticky. There are many roots and worms, also krotovinas. The reaction is 7.0. Grades clearly and smoothly into:
B_2	7-16"	Dark grayish brown (10YR 4/2; very dark brown, 10YR 2/2, moist)

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haploxeroll, fine-loamy, mixed, thermic
SOIL San Andreas sandy clay loam (taxadjunct) SOIL Nos. S65Calif-27-28 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65533 - 65537

Depth (In.)	Horizon	1B1b Total Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Sand: Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt: 0.05–0.02	Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	* Coarse fragments 1A2a > 2	2 – 19	19 – 76
		Pct. of < 2 mm														Pct.	Pct. of < 76mm	
0-1	A11	64.7	16.1	19.2	17.4	17.8	8.5	13.6	7.4	8.2	7.9	22.7	57.3	0	19	12	12	0
1-8	A12	64.4	16.2	19.4	17.0	17.4	8.1	14.3	7.6	8.0	8.2	22.7	56.8	0	19	10	10	0
8-17	B2	64.4	14.6	21.0	19.4	19.0	7.9	11.9	6.2	6.7	7.9	18.8	58.2	0	21	9	9	0
17-21	B3	65.8	14.8	19.4	18.1	19.3	8.8	13.4	6.2	6.3	8.5	18.8	59.6	0	19	10	10	0
21-36+	C	83.1	9.9	7.0	21.6	28.6	11.3	16.1	5.5	5.0	4.9	17.3	77.6	0	7	13	13	0

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content: Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1a H_2O 1:1	
0-1	1.65	0.135	12		-							9.7			6.6	
1-8	0.88	0.067	13		-	1.67	1.72		1.00		11.4	7.5	0.011	0.011	6.6	
8-17	0.30				-	1.74	1.82		1.00		12.7	9.0	0.014	0.014	6.7	
17-21	0.20				-							9.1			6.5	
21-36+	0.05				-	1.99	1.99		1.00		7.7	5.1	0.002	0.002	6.3	

Depth (In.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1a NH_4OAc	5A3a Sum	6G1d KCl Ext. Al^{+++}	Base saturation: 5C1 NH_4OAc Pct.	Pct.
	meq/100 g										
0-1	15.1	5.4	0.2	0.5	21.2		21.8			97	
1-8	15.0	6.2	0.2	0.2	21.6		22.5			96	
8-17	20.2	8.0	0.3	0.1	28.6		28.3			100+	
17-21	18.2	7.3	0.4	0.1	26.0		25.0			100+	
21-36+	13.1	5.3	0.4	0.1	18.9		17.7			100+	

Depth (In.)	Clay mineralogy (≤0.002 mm) 7A: Mont.	Verm.	Mica	Kaolin
	X-ray			
0-1	xxxx			
1-8				
8-17	xxxx			t
17-21				
21-36+	xxxx	x		

* From characterization sample (5Kg) (except as noted).

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

SAN ANDREAS SANDY CLAY LOAM (taxadjunct) 1/
S65Calif-27-28

Location: Monterey County, California, 400 feet west of S¼ corner of SE¼, Sec. 18, T17S, R7E, 6 miles on road to Pinnacles from Soledad. Date of Sampling: May 27, 1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton and B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich; site selection by T. D. Cook.
Classification: Typic Haploxeroll, fine-loamy, mixed, thermic. Noncalcic Brown(medial).
Vegetation: California sage, deer vetch, wild oats, red brome, some soft chess. Climate: Precipitation 10 inches; mean annual temperature 65°F., mean January temperature 55°F., mean July temperature 70°F. Parent Material: Weathered diorite (gruss). Topography: Steep smooth 30 percent south-facing slope. Elevation: 1500 feet. Drainage: Well drained, permeability moderately slow to slow, runoff medium. Soil Moisture: Dry throughout.
Remarks: The paired sample for this site is S65Calif-27-27. Classification assumes mollic epipedon for series as a whole; mollic marginal this site because of weak structure.

HORIZON	DESCRIPTION
A11 RSL 65533	0 to 1 inch, dark brown (10YR 3/3 moist or dry) sandy clay loam; weak coarse platy structure readily breaking to moderate coarse granular; slightly hard, friable, sticky and plastic; plentiful very fine roots; common very fine interstitial pores; slightly acid (pH 6.5); thin layer of fine gravel covers 1/2 of surface; clear smooth boundary.
A12 RSL 65534	1 to 8 inches, dark brown (7.5YR 3/2) sandy clay loam, very dark brown (7.5YR 2/2) moist; weak coarse prismatic structure; very hard, firm, sticky and plastic; plentiful very fine roots mostly on ped exterior; common very fine interstitial and few fine tubular pores; neutral (pH 7.0); vertical joints darkened by organic material; clear wavy boundary.
B2 RSL 65535	8 to 17 inches, dark reddish brown (5YR 3/4 ped interiors) sandy clay loam, dark reddish brown (5YR 3/4m ped interior), dark brown (7.5YR 3/2 ped exteriors, 3/4 rubbed) moist; weak coarse prismatic structure; extremely hard, firm, very sticky and plastic; plentiful very fine roots mostly on ped exteriors; common very fine interstitial pores; continuous colloidal stems, common thin clay stains in pores and on ped faces; neutral (pH 7.0); clear wavy boundary.
B3 RSL 65536	17 to 21 inches, brown (7.5YR 5/4) light sandy clay loam, dark brown (7.5YR 3/4) moist; massive; very hard, firm, sticky and plastic; very few very fine roots; many very fine interstitial pores; few thin clay films in bridges and common colloidal stains; slightly acid (pH 6.5); B3 is an indistinct transition between B2 and C; clear smooth boundary.
C RSL 65537	21 to 36 inches, clean decomposed diorite with common colloidal stains upper part, rock structure, can be dug with hand tools with difficulty, becomes lithic contact at 36 inches.

1/ The San Andreas series is in a coarse-loamy, mixed, thermic family of Typic Haploxerolls. This pedon is in a fine-loamy (more than 18 percent clay) family of the same subgroup.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haploxeroll, fine, montmorillonitic, thermic
SOIL Rincon clay loam (taxadjunct) SOIL Nos. S64Calif-57-17 LOCATION Yolo County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64488 - 64494

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	Coarse fragments 1A2a > 2	2 - 19	19 - 76
		Pct. of < 2 mm														Pct.	Pct. of < 76 mm	
0-8	Ap	13.6	51.0	35.4	0.4	1.6	2.4	5.1	4.1	15.0	36.0	22.0	9.5			0		
8-17	B1	13.0	44.7	42.3	0.3	1.1	1.6	4.9	5.1	14.0	30.7	22.2	7.9			0		
17-34	B21t	20.0	45.5	34.5	0.7	1.2	2.1	7.5	8.5	16.8	28.7	30.2	11.5			0		
34-45	B22t	18.4	46.0	35.6	0.4	0.9	1.7	6.9	8.5	15.6	30.4	28.7	9.9			0		
45-54	B31	19.7	47.6	32.7	0.8	0.8	1.5	7.6	9.0	17.8	29.8	32.0	10.7			0		
54-74	B32	34.3	37.9	27.8	1.8	2.5	5.4	13.4	11.2	16.2	21.7	34.7	23.1			0		
74-84+	C1	25.3	43.4	31.3	1.2	2.1	4.2	9.7	8.1	16.0	27.4	29.4	17.2			0		

Depth (in.)	6A1a Organic carbon	6B1a Nitrogen	C/N	6C2a Ext. Iron as Fe	6E1b Carbonate as $CaCO_3$		Atterberg limits: Plastic limit	Liquid limit	Plastic index		Water content:	4B1c 1/3 bar	4B2 15 bar	4D1 Extensibility COLE	4D1 Extensibility COLE	pH: 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
	Pct.	Pct.		Pct.	Pct.		Pct. of < 0.4 mm.				Pct.	Pct.	Pct.	in./in.	in./in.			
0-8	0.87	0.088	10		-		19	29	10				13.1				6.2	6.7
8-17	0.50	0.069	7		-								16.4				6.6	7.1
17-34	0.35	0.048	7		-		19	37	18				15.5				7.4	7.8
34-45	0.30				-								14.3				7.9	8.3
45-54	0.19				tr.								13.0				8.1	8.7
54-74	0.17				1								11.2				8.2	8.8
74-84+	0.17				2		16	29	13				12.5				8.2	8.9

Depth (in.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 6A1: 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-8	11.5	9.5	0.2	0.9	22.1	4.1	26.9											0.33
8-17	14.9	11.5	0.2	0.6	27.2	3.2	31.8											0.13
17-34	14.2	9.1	0.4	0.6	24.3		24.6											0.18
34-45	15.9	9.0	0.4	0.5	25.8		27.4											0.25
45-54	17.5	8.4	0.5	0.5	26.9		23.0											0.31
54-74	18.0	7.2	0.4	0.3	25.9		20.8											0.33
74-84+	24.8	8.1	0.5	0.3	33.7		24.2											0.32

Depth (in.)	8A Water at Saturation	Exch. bases 5B3b Ca	5B3b Mg	5B1b Na	5D2 Exchangeable Na	5C2 Base Sat.	Gypsum	Clay mineralogy (<0.002 mm.) 13A1: Mont.	Verm.	Mica	Kaolin	
	Pct.	meq./100 g.			Pct.	Pct.	Pct.	X-ray 7A2				
0-8	42.2				1	82						
8-17	45.0				1	86		xxxx	x	-	xx	
17-34	41.2				2	99						
34-45	46.9				1	94		xxxx	-	-	xx	
45-54	46.1				2	-						
54-74	42.5				2	-						
74-84+	45.3				2	-		xxxx	-	-	xx	

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

RINCON CLAY LOAM (taxadjunct) 1/
S64Calif-57-17

Location: Yolo County, California. A pit located 100 feet south of the northwest corner of NE¼ NE¼, Sec. 26, T.11N., R.2W., located about 9.5 miles west of Zamora. Photo ABB-7K-160.
Date of Sampling: September 29, 1964.
Description by: G. J. Staidl. Collectors: R. E. Nelson. B. R. Brasher. G. M. Kennedy. W. F.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE: Landlow silty clay loam **LOCATION** Merced County, California

SOIL FAMILY: Aquic Haploxeroll, fine, montmorillonitic, thermic

SOIL NOS. S55Calif. 24-1 **LAB. NOS.** 55273-280

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0- 4	A_{1p}	0.3	0.8	1.8	9.8	8.2	46.1	33.0	31.2	29.4	0	cl
4-10	B_{21}	0.1	0.5	2.0	11.1	9.3	36.8	40.2	30.4	23.3	0	c
10-17	B_{22}	0.0	0.4	1.9	11.0	10.1	38.1	38.5	30.0	25.8	0	cl
17-21	B_{31}	0.1	0.4	2.0	10.9	10.6	39.1	36.9	32.0	25.4	0	cl
21-32	B_{32}	0.3	0.8	2.4	10.1	10.4	46.4	29.6	34.8	29.0	0	cl
32-48	B_{33}	0.0	0.6	1.1	4.4	4.9	48.0	41.0	21.6	34.5	0	sic
48-60	C_1	0.1	0.4	0.8	15.9	12.9	44.7	25.2	44.0	26.5	0	l
60-65	C_2	0.2	0.7	1.6	10.8	8.1	54.5	24.1	32.3	37.9	0	sil

8C1b pH SATURATED PASTE 1:1	8C1a pH 1:5	8C1a pH 1:10	6A1a ORGANIC MATTER ORGANIC CARBON %	6B1a NITROGEN %	C/N	4A3a Bulk Density g/cc	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
7.5		8.1	1.35	0.115	11.7	1.54	2.7		0			12.9
8.1		8.4	0.74	0.059	12.5	1.65	2.4		0			15.0
8.3		9.2	0.62	0.053	11.7	1.68	2.9	<0.5	0			15.7
8.4		9.0	0.39	0.032	12.2	1.71	3.9	<0.5	0			15.1
8.3		9.1	0.15	0.018		1.76	7.1	<0.5	0			13.3
8.0		8.7	0.14	0.020		1.75	14.5	<0.5	0			14.6
8.2		9.5	0.09			1.75	14.1	<1	0			11.2
8.4		9.6	0.09			1.58	16.2	1	0			10.9

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	5D1 EXCH. Na %	SATURATION EXTRACT SOLUBLES 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg	6I1a 6J1a HCO_3 + CO_3	6K1a Cl	8A1 8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
29.4	13.4	11.8		2.4	0.8	8.2	20.0	0.3	9.1	8.2	12.6	48.3
33.9	11.0	12.4		6.3	0.7	18.6	20.2	0.2	2.2	5.8	13.4	58.0
30.3	8.6	10.8		9.4	0.6	31.0	26.2	0.2	1.2	6.4	12.9	77.6
27.6	8.0	9.9		10.6	0.5	38.4	34.2	0.2	1.4	5.9	18.5	76.7
22.5	10.2	7.7		9.3	0.5	41.3	62.5	0.4	4.2	3.9	39.9	54.0
27.8	11.9	8.2		9.2	0.7	33.1	128.7	0.2	34.9	2.0	53.2	56.9
18.3	10.7	6.3		9.3	0.5	50.8	132.5	0.2	14.0	2.7	88.3	45.3
17.3	11.4	7.5		8.4	0.5	48.6	145.0	0.2	18.3	2.7	114.3	42.3

LANDLOW SILTY CLAY LOAM
S55Calif-24-1

Location: Merced County, California. In the Los Banos SCD on the Los Banos Outlying Nursery in the middle of Check No. 15 from the south field fence, 100 feet into the field east of the west fence, SW1/4 SE1/4 SW1/4 sec. 35, T. 9 S., R. 10 E.
Date of Sampling: April 19, 1955.
Description by: R. C. Huff. Collectors: R. C. Huff, R. E. Nelson, George Edmunson, Rudolph Ulrich.
Classification: Aquic Haploxeroll, fine, montmorillonitic, thermic.
Vegetation: The area where the sample was taken is part of the Los Banos Outlying Nursery and is planted to a mixture of Goars tall fescue and birdsfoot trefoil.
Climate: Rainfall approximately 8 inches, characterized by hot, dry summers and relatively cool, moist winters. Parent Material: Fine-textured sedimentary alluvium, which originated mainly from sandstone and shale rocks from the eastern side of the Coast Range Mountains. Relief: Nearly level to level (0-1 percent) very gently sloping basin rim; very slightly concave. Leveled for irrigation. Elevation: 95 feet. Drainage: Surface drainage is very slow, internal drainage, slow. The water table normally fluctuates within 10 feet of the surface.
Remarks: Some evidence of worm activity.

HORIZON		DESCRIPTION
A_{1p}	0- 4"	Grayish brown (10YR 5/2, very dark grayish brown 3/2 moist) silty clay loam; moderate fine and medium subangular blocky. Hard, firm, slightly plastic and nonsticky. Noncalcareous. pH of 8.0. The lower boundary is clear and smooth.
B_{21}	4-10"	Grayish brown (10YR 5/2, very dark brown 2/2 moist) clay. Moderate fine angular blocky. Very hard, firm, plastic and slightly sticky. Noncalcareous. pH of 8.0. The lower boundary is gradual and smooth.
B_{22}	10-17"	Grayish brown (10YR 5/2, very dark grayish brown 3/2 moist, with darker faces 10YR 2/2) slightly coarser-textured clay than the horizon above. Very hard, firm, plastic and slightly sticky. Very slight evidence of being calcareous. pH 8.0. This grades clearly and with a wavy boundary into:
B_{31}	17-21"	Grayish brown (2.5Y 5/2, very dark grayish brown 3/2 moist) silty clay loam. Weak, fine subangular blocky. Hard, firm, slightly plastic and nonsticky. There are a very few very fine (less than 1 mm.) manganese shot scattered throughout this horizon. The soil is very slightly calcareous. pH is 8.0. The lower boundary is clear and smooth.
B_{32}	21-32"	Gray brown (2.5Y 5/2, dark grayish brown 4/2 moist) slightly coarser-textured silty clay loam than the horizon above. Weak medium subangular blocky. Hard, firm, slightly less plastic

OIL SURVEY LABORATORY Riverside, California

SOIL TYPE Landlow silty clay (taxadjunct) **LOCATION** Riverside County, California

SOIL FAMILY: Aquic Haploxeroll, fine, montmorillonitic, thermic.

SOIL NOS. S57Calif-33-6 **LAB. NOS.** 57495 - 57500

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2	
0-6	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	1.8	[illegible]	[illegible]	[illegible]	[illegible]	0	[illegible]

LANDLOW SILTY CLAY (taxadjunct) 1/
S57Calif-33-6

Sampled by: George Borst, R. E. Nelson, Cliff Henry and Arnold Knecht, June 4, 1957. Description by: Fred Eggers.

Location: 200' N of the S 1/4 corner of Sec. 33, T4S, R3W. On the Rynsberger Ranch in the triangle between Ellis Road, highway 395 and the San Jacinto River, about 1 mile east of Ferris, Riverside County, California.

Elevation: 1413 feet. Precipitation: 12 inches.

Classification: Aquic Haploxeroll, fine, montmorillonitic, thermic.

Vegetation: Salt grass and annual grasses and weeds.
Parent Material: Mixed alluvium, chiefly granitic. Physiography: Flood plain.
Drainage: Slow.

HORIZON	DESCRIPTION
S57Calif-33-6-1 A1 0-6"	Dark gray (10YR 4/1 dry, very dark gray 3/1 moist) silty clay; moderately defined fine granular; hard when dry, firm when moist; sticky and very plastic when wet; moderately slow permeability; many plant roots; pH 8.2; slight effervescence; occasional small pores; clear smooth lower boundary.
S57Calif-33-6-2 B1 6-10"	Gray (10YR 5/1 dry, very dark gray 3/1 moist) silty clay; strongly defined medium subangular blocky with a slight tendency to prismatic structure; very hard when dry, very firm when moist; sticky and very plastic when wet; moderately slow permeability; many plant roots; occasional small pores; pH 8.2; slight effervescence; gradual smooth lower boundary.
S57Calif-33-6-3 B21 10-24"	Gray (10YR 6/1-5/1 dry, dark gray 4/1 moist) clay; strongly defined medium subangular blocky; definitely prismatic on drying; many prominent clay skins, extremely hard when dry, very firm when moist; very sticky and very plastic when wet; slow permeability; occasional plant roots; occasional small pores; pH 8.4; strong effervescence; gradual smooth lower boundary.
S57Calif-33-6-4 B22 24-30"	Gray (10YR 6/1-5/1 dry, dark gray 4/1 moist) clay; strongly defined medium subangular blocky; definitely prismatic; on drying prominent clay skins; extremely hard when dry, very firm when moist; very sticky and very plastic when wet; slow permeability; few roots; pH 8.4; calcium carbonate and calcium sulphate in seams; gradual smooth lower boundary.
S57Calif-33-6-5 C1 30-42"	Gray (10YR 6/1-5/1 dry, dark gray 4/1 moist) silty clay; blocky; very hard when dry, very firm when moist; sticky and plastic when wet; moderately slow permeability; no plant roots; pH 8.4; many seams of gypsum and some line concretions; abrupt wavy lower boundary.
S57Calif-33-6-6 C2 42-60"+	Gray (10YR 6/1-5/1 dry, dark gray 4/1 moist) silty clay, blocky; very hard when dry, very firm when moist; sticky and plastic when wet; moderately slow permeability; no plant roots; pH 8.4; many seams of gypsum and some line concretions.

1/ This pedon lacks mottles with chromas of 2 or less but is presumed to be saturated with water for 90 days or more within 1 meter in most years.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aridic Haploxeroll, coarse-loamy, mixed, mesic

SOIL Calpine sandy loam SOIL Nos. S61Calif-32-19 LOCATION Sierra Valley, Plumas County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 61730 - 61735

Depth (in.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)			Coarse fragments 2A2: > 2 Pct.	2 - 19	19 -76
		Pct. of < 2 mm															Pct. of < 76 mm	
0-3	A11	79.5	11.5	9.0	9.5	27.5	16.9	21.0	4.6	4.2	18.5	7.3	65.2			4		
3-13	A12	77.8	11.8	10.4	6.1	20.3	17.2	27.0	7.2	4.4	25.5	7.4	56.7			4		
13-21	A3	78.7	10.9	10.4	5.8	21.1	17.8	27.4	6.6	4.1	24.3	6.8	58.5			4		
21-30	B1	76.2	12.3	11.5	4.7	19.5	17.6	27.7	6.7	4.3	24.6	8.0	55.9			4		
30-46	B2	81.2	11.0	7.8	7.0	21.9	19.9	26.8	5.6	4.2	21.5	6.8	63.9			8		
46-59	C1	84.3	9.7	6.0	2.1	9.2	20.7	44.2	8.1	3.7	29.3	6.0	58.7			1		
59-81+	C2g	Not sampled																

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		6E1b Carbonate as $CaCO_3$ Pct.	6C2a Ext. Iron as Fe Pct.	Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	4B2 15 bar Pct.				pH 8C1b Sat. Paste	8C1a 1:10
0-3	2.30	0.144	16		-	0.8							6.4				5.8	6.2
3-13	0.98	0.076	13		-	0.9							5.8				6.0	6.3
13-21	0.63	0.048	13		-	0.9							5.2				6.1	6.4
21-30	0.41	0.033	12		-	1.0							5.8				6.3	6.6
30-46	0.15				-	0.8							5.3				6.5	6.6
46-59	0.08				-	0.6							5.3				6.6	6.6
59-81+																		

Depth (in.)	Extractable bases 5B1a: 6N2c Ca	6O2c Mg	6P2a Na	6Q2a K	6H2a Ext. Acidity	Cat. Exch. Cap. 5A2a NaOAc		Water extract from saturated paste: Ca	Mg	Na	K		CO_3	HCO_3	Cl	SO_4	Electrical conductivity
	meq/100 g							meq/liter									mmho/cm

CALPINE SANDY LOAM

S61Calif-32-19

Location: Sierra Valley, Plumas County, California. Center of NE1/4 SE1/4 section 27, T. 23 N., R. 14 E. Approximately 3/4 mile southwest of Beckwourth just west of Highway A-23 (Beckwourth-Calpine road).
Date of Sampling: September 19, 1961.
Collectors: Grant Kennedy, Reuben Nelson, John Rogers, Paul Nazar, James McLaughlin, Eugene Begg.
Vegetation: The vegetation is silver sage, big sage, dryland sedges, and bitter brush. Climate: Estimated annual precipitation 15 inches. Nearest weather station (Portola): Mean annual temperature 45.1° F.; January average 28.7° F.; July average 62.6° F.; mean annual precipitation 20.62 inches.
Parent Material: Mixed but dominantly granitic alluvium. Relief: Slightly undulating low terrace with a slope less than 2 percent. Elevation: 4880 feet. Drainage: Moderately well drained, runoff very slow and permeability is rapid in the upper part of the profile and moderate below.
Remarks: This is a Brunizem with a weakly prismatic B2 horizon. The C2g horizon was not sampled. The C2g was stratified sand and coarse sand. A water table was found at 8 feet. Samples were taken for the California Division of Highways and the Bureau of Public Roads from the A12 horizon (3-13 inches), the B2 horizon (30-46 inches) and the C1 horizon (46-59 inches).
Classification: Aridic Haploxeroll, coarse-loamy, mixed, mesic.

HORIZON	DESCRIPTION
A11 RSL No. 61730	0 to 3 inches, dark grayish brown (10YR 4/2) sandy loam (near loamy sand), very dark brown (10YR 2/2) when moist; moderate very fine platy structure; loose, very friable, nonsticky and nonplastic; abundant very fine, plentiful fine and few medium roots; many very fine tubular and interstitial pores; strongly acid (pH 5.5); clear, smooth boundary.
A12 RSL No. 61731	3 to 13 inches, dark grayish brown (10YR 4/2) sandy loam, very dark brown (10YR 2/3) when moist; weak coarse platy structure; loose, very friable, slightly sticky and slightly plastic; abundant [illegible] plentiful fine and few medium roots; many very fine tubular and interstitial pores; strongly acid (pH 5.5); gradual, smooth boundary.
A3 RSL No. 61732	13 to 21 inches, dark grayish brown (10YR 4/2) sandy loam, very dark brown (10YR 2/3) when moist; weak medium to coarse subangular blocky structure; soft, very friable, nonsticky and nonplastic; abundant very fine, plentiful fine and few medium roots; common very fine tubular and interstitial, few fine and medium tubular pores; few thin clay films in pores and bridges; strongly acid (pH 5.5); clear, smooth boundary.
B1 RSL No. 61733	21 to 30 inches, brown (10YR 5/3) sandy loam, dark yellowish brown (10YR 3/4) when moist; weak coarse prismatic structure; hard, firm, nonsticky and nonplastic; plentiful very fine and few fine to medium roots; common very fine tubular and interstitial and few fine to medium tubular pores; common thin clay films in pores and bridges; medium acid (pH 6.0); gradual, smooth boundary.
B2 RSL No. 61734	30 to 46 inches, variegated light yellowish brown (10YR 6/4) and yellow (10YR 7/6) sandy clay loam (near sandy loam), yellowish brown (10YR 5/4) when moist; weak coarse prismatic structure; hard, firm, slightly sticky and slightly plastic; few very fine, fine, and medium roots; common very fine tubular and interstitial and few fine to medium tubular pores; common thin clay films in pores and bridges; medium acid (pH 6.0); gradual, smooth boundary.
C1 RSL No. 61735	46 to 59 inches, light yellowish brown (10YR 6/4) loamy fine sand, dark yellowish brown (10YR 4/4) when moist; massive; slightly hard, firm, nonsticky and nonplastic; few fine, very fine and medium roots; many very fine interstitial and few fine to medium tubular pores; medium acid (pH 6.0); clear, wavy boundary.
C2g RSL No. 61736	59 to 81 inches +, three of the dominant colors are light yellowish brown (10YR 6/4), dark reddish brown (5YR 3/2) and yellow (10YR 7/8), moist colors are reddish yellow (7.5YR 6/6), yellowish red (5YR 5/8) and dark reddish brown (2.5YR 3/4); manganese stains are apparent; texture varies in stratified layers from coarse sand to sand; single grained; loose, nonsticky and nonplastic; very few very fine, fine and medium roots; many very fine interstitial and few very fine tubular pores; medium acid reaction (pH 6.0).

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Aridic Haploxeroll, coarse-loamy, mixed, mesic

SOIL Calpine sandy loam SOIL Nos. S61Calif-46-5 LOCATION Sierra Valley, Sierra County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 61724 - 61729

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.2)			Coarse fragments 2A2 > 2 Pct.	2 - 19	19 - 76
		Size class and particle diameter (mm) 3A1																
		Pct. of < 2 mm															Pct. of < 76 mm	
0-4	A11	77.2	13.7	9.1	5.5	16.2	15.5	29.8	10.2	6.7	7.0	33.3	50.6			2		
4-11	A12	76.5	12.1	11.4	6.0	14.7	14.9	29.2	11.7	6.4	5.7	34.2	48.7			2		
11-17	A3	75.2	12.3	12.5	3.9	12.7	15.5	30.8	12.3	6.3	6.0	36.1	45.4			2		
17-27	B1	74.6	11.7	13.7	5.9	15.2	14.9	28.1	10.5	5.8	5.9	32.3	48.1			2		
27-41	B2	71.7	14.9	13.4	2.8	11.9	12.1	30.2	14.7	7.5	7.4	40.4	38.8			3		
41-58	C1	80.4	10.5	9.1	4.2	21.3	19.4	28.2	7.3	4.6	5.9	25.2	59.8			4		
58-76+	IIC2	Not sampled																

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		6E1b Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	4B2 15 bar Pct.			pH 8C1b Sat. Paste	8C1a 1:10
0-4	3.56	0.251	14		-								10.1			6.0	6.3
4-11	0.93	0.073	13		-								6.1			6.2	6.5
11-17	0.56	0.054	10		-								6.1			6.2	6.5
17-27	0.27				-								6.7			6.1	6.5
27-41	0.12				-								7.6			6.3	6.7
41-58	0.06				-								6.6			6.8	6.9
58-76+																	

Depth (in.)	Extractable bases 6N2c Ca	6O2c Mg	6P2a Na	5B1a 6Q2a K	6H2a Ext. Acidity	Cat. Exch. Cap. 5A2a NaOAc			Water extract from saturated paste Ca	Mg	Na	K		CO_3	HCO_3	Cl	SO_4	Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-4	9.5	2.5	0.1	0.7	7.6	19.3												
4-11	6.2	1.9	0.1	0.6	4.7	10.5												
11-17	5.8	2.2	0.1	0.3	3.8	13.3												
17-27	5.8	3.0	0.2	0.2	3.5	13.3												
27-41	7.6	4.7	0.2	0.1	2.5	14.8												
41-58	7.0	4.9	0.4	0.1	1.4	14.3												
58-76+																		

Depth (in.)	Water at Saturation Pct.	5C3 Base Sat. (Sum) Pct.	Clay Mineralogy (<0.002 mm.) Montmorillonite	Vermiculite	Mica	Kaolinite	Kaolinite % DTA
			X-ray				
0-4		63	-	t	xx	x	15
4-11		65					
11-17		69					
17-27		72					
27-41		83	xx	x	-	x	15
41-58		89	xxx	-		x	8
58-76+							

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

CALPINE SANDY LOAM

S61Calif-46-5

Location: Sierra Valley, Sierra County, California. Southeast corner of the NW1/4SW1/4 of section 11, T. 21 N., R. 14 E. 3/4 mile north of Terral Farm Headquarters. Date of Sampling: September 18, 1961.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Soquel loam (taxadjunct) **LOCATION** San Luis Obispo Co.

SOIL FAMILY: Cumulic Haploxeroll, fine-loamy, mixed, mesic

SOIL NOS. S55Ca1. 40-3 **LAB. NOS.** 55459-464

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	3A1 2A2 > 2 1/	TEXTURAL CLASS
0- 3	A_{p1}	0.5	1.2	3.9	25.0	17.4	33.4	18.6	50.6	17.3	1	1
3-8½	A_{p2}	0.4	1.2	3.9	24.6	17.4	34.3	18.2	51.3	17.2	1	1
8½-15	C_1	0.3	1.2	4.0	25.3	17.4	34.1	17.7	51.5	17.3	2	1
15-24	C_2	0.0	0.6	2.4	19.6	19.5	40.3	17.6	55.6	18.0	0	1
24-40	C_3	0.1	0.2	1.2	11.6	16.7	46.7	23.5	58.5	13.5	0	1
40-50	A_{1b}	0.0	0.1	0.3	5.7	12.2	51.6	30.1	37.6	31.1	0	sicl

pH 8C1b SATURATED PASTE 1:1	1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC x 10 MILLIMHOS PER CM @25°C	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	MOISTURE TENSIONS 4B2 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS.
7.2		7.7	1.15	0.092	12.5		1.9					9.8
7.2		8.2	1.12	0.087	12.9		0.4					10.5
7.4		8.3	1.02	0.078	13.1		0.4					10.3
7.3		8.1	0.83	0.068	12.2		0.5					10.9
7.4		8.2	1.24	0.089	13.9		0.4					13.3
7.4		8.2	1.45	0.115	12.6		0.4					20.2

5A2a CATION EXCHANGE CAPACITY (← Na)	EXTRACTABLE CATIONS 5B 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg		6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
26.6	14.1	10.7	2.6	0.5	0.4	91	5.00	0.3	14.3			36.8
27.1	13.5	10.3	2.6	0.4	0.4	90	1.7	0.1	2.5			35.0
26.3	14.0	10.6	2.7	0.4	0.4	90	1.6	0.1	2.2			35.9
31.4	14.1	12.4	2.9	0.4	0.3	90	1.6	0.1	3.3			40.6
35.8	15.7	16.9	3.2	0.1	0.4	91	0.4	0.04	3.9			48.0
37.5	19.9	18.3	3.4	0.3	0.4	92	0.5	0.1	3.5			58.2

1/ Percent of the whole soil

SOQUEL LOAM (taxadjunct) 1/
S55Calif-40-3

Location: About 5/8 mile W. of farm buildings on George Mello farm, San Luis Obispo County, California.

Collectors: W. M. Johnson, Rudolph Ulrich, Lewis Leifer, W. G. Harper, L. R. Wohletz, J. M. Williams. Samples taken September 8, 1955. Description by W. M. Johnson.
Classification: Cumulic Haploxeroll, fine-loamy, mixed, mesic.
Elevation: About 40 feet.

Vegetation: Cultivated field. Planted to sugar beets at time samples were taken. Irrigated.

Relief: Gently sloping, low terrace-like surface. Slope about 1½ percent, plane, toward the east.

Drainage: Well drained. Runoff is medium; permeability is moderate.

Parent material: Loamy, noncalcareous alluvium from mixed sources.

Remarks: Fertilization history of the field was not obtained. Sugar beets looked very healthy. Clods on the soil surface in many places had a thin, patchy white efflorescence that suggested soluble salts.

HORIZON		DESCRIPTION
A_{p1}	0-3"	Very dark grayish brown to grayish brown (10YR 5/2; 3/2 moist) loam. Cloddy from tillage. Weak coarse and medium granular. Very hard, friable, plastic, slightly sticky. Many roots. pH 7.5. Abrupt, smooth lower boundary.
A_{p2}	3-8½"	Very dark grayish brown (10YR 3/2, moist) loam. Weak medium and coarse granular. Friable, slightly plastic, nonsticky. Very many worm casts and burrows. Roots numerous. pH 7.5. Gradual lower boundary.
C_1	8½-15"	Finely, indistinctly mottled very dark gray and very dark grayish brown (10YR 3/1 and 3/2, moist) loam. Massive. Friable, slightly plastic, nonsticky. Compact in places. Common worm burrows and casts. pH 7.5. Common roots. Gradual lower boundary.
C_2	15-24"	Mottled very dark gray and olive brown (10YR 3/1 and 2.5Y 4/4, moist) loam with occasional specks of strong brown. Weak, medium subangular blocky with many worm casts and burrows. Roots common. pH 7.5. Gradual lower boundary.
C_3	24-40"	Very dark gray (10YR 3/1, moist) loam finely mottled with dark grayish brown and very dark grayish brown. Massive with very many worm casts and burrows. Friable, plastic, slightly sticky. Roots common. pH 7.5. Gradual lower boundary.
A_{1b}	40-50"+	Black (10YR 2/1, moist) silty clay loam, finely mottled with very dark grayish brown. Moderate medium granular. Friable, plastic, slightly sticky. Many worm casts and burrows. Roots common. pH 7.5. Bits of carbonized plant material are scattered throughout. Probably surface of old marsh.

1/ The Soquel series is in a fine-loamy, mixed, mesic family of Pachic Haploxerolls. This pedon has an organic carbon content that decreases irregularly with depth and remains above 0.3 percent at 50 inches.

U S DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Pachic Haploxeroll, coarse-loamy, mixed, thermic

SOIL Sheridan loam SOIL Nos. S65Calif-27-21 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65565 - 65569

[redacted]

SHERIDAN LOAM
S65Calif-27-21

Location: Monterey County, California. SE corner SW¼ SW¼, Sec. 16, T18S, R1E. About 3.5 miles from Bixby Bridge (on Hwy. #1) on old Coast Road, 1 mile up Rancho Calera Road from Old Coast Road, about 20 feet down from ridge top. (Coastal grass site). Date of Sampling: May 25, 1965.

Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich; site selection by T. D. Cook.

Classification: Pachic Haploxeroll, coarse-loamy, mixed, thermic. Regosol (brunizemic).

Vegetation: Range grass, red brome, soft chess, lupine (brush side slopes). Climate: Precipitation 35 to 45 inches; mean annual soil temperature at 20 inches is assumed to be greater than 59° F. Parent Material: Weathered diorite (gruss). Topography: Convex smooth ridgetop on W. and slightly N. side of ridge, 45 percent west-facing slope. Elevation: 1400 feet. Drainage: Well drained, moderately permeable, medium runoff. Soil Moisture: Dry to 8 inches, moist below.

Remarks: Field pH by Truog Kit. Closely related soil on brush covered south slope a few hundred feet away is S65Calif-27-20-1 through 6. The closely paired site is S65Calif-27-22-1 through 5. Other closely related but not paired to this site are S65Calif-27-23, S65Calif-27-24, S65Calif-27-25, S65Calif-27-26. Gopher activity makes up 5 to 10 percent of profile. Asterisks indicate horizons sampled for California Division of Highways.

HORIZON	DESCRIPTION
A11 RSL 65565	0 to 6 inches, dark grayish brown (10YR 4/2) loam, very dark brown (10YR 2/2) moist; weak fine and medium granular structure; slightly hard, very friable, slightly sticky and slightly plastic; plentiful very fine roots; common very fine interstitial and few very fine and fine tubular pores; medium acid (pH 6.0); gradual smooth boundary.
*A12 RSL 65566	6 to 15 inches, dark grayish brown (10YR 4/2) loam, very dark brown (10YR 2/2) moist; moderate fine and medium granular structure; slightly hard, very friable, slightly sticky and slightly plastic; plentiful very fine roots; many very fine interstitial and few very fine, fine and medium tubular pores; slightly acid (pH 6.3); diffuse smooth boundary.
A13 RSL 65567	15 to 30 inches, dark grayish brown (10YR 4/2) loam, very dark brown (10YR 2/2) moist; dark grayish brown (10YR 4/2) moist, rubbed; moderate medium and coarse granular structure; slightly hard, very friable, slightly sticky and slightly plastic; plentiful very fine roots; many very fine interstitial and few very fine, fine and medium tubular pores; slightly acid (pH 6.5); few lumps, 5 to 10 percent C1 material in lower part of horizon; abrupt irregular boundary.
C1 RSL 65568	30 to 41 inches, strong brown (7.5YR 5/6) dry and moist, coarse sandy loam, brown (7.5YR 5/4 when rubbed) moist; massive; weathered diorite readily cut by hand tools; hard, friable, slightly sticky and slightly plastic; few very fine roots; common very fine interstitial pores; common thin clay films and very few

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Pachic Haploxeroll, coarse-loamy, mixed, thermic
SOIL Sheridan loam SOIL Nos. S65Calif-27-22 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65570 - 65574

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand Very coarse (2-1)	Sand Coarse (1-0.5)	Sand Medium (0.5-0.25)	Sand Fine (0.25-0.1)	Sand Very fine (0.1-0.05)	Silt 0.05-0.02	Silt Int. III (0.02-0.002)	Silt Int. II (0.2-0.02)	Silt (2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	* Coarse fragments > 2 Pct.	1A2a 2-19 Pct. of < 76mm	19-76 Pct. of < 76mm	
		Pct. of < 2 mm																	
0-6	A11															10	10	0	**
6-17	A12															14	14	0	**
17-32	A13	57.8	30.6	11.6	5.9	13.3	9.1	17.6	11.9	13.2	17.4	35.1	45.9	0	12	13	13	0	**
32-43	C1															14	14	0	**
43-60+	C2															16	16	0	**

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1		
0-6	1.76	0.139	13											5.9		
6-17	0.89	0.072	12											6.1		
17-32	0.66										10.5			6.3		
32-43	0.20													6.7		
43-60+	0.10													6.7		

Depth (In.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	5B1a Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A1a NH_4OAc	5A3a Sum	6G1d KCl Ext. Al^{+++}	Base saturation 5C1 NH_4OAc Pct	Pct
	meq/100 g										
0-6	15.8	8.3	0.3	0.3	24.7	10.2	28.2			88	
6-17	15.8	9.6	0.3	0.1	25.8	3.6	29.6			87	
17-32	15.9	10.6	0.4	0.1	27.0		29.5			92	
32-43	15.8	11.4	0.6	0.1	27.9		30.5			92	
43-60+	17.0	13.3	0.6	0.1	31.0		31.6			98	

Depth	Clay mineralogy (<0.002 mm)7A
	[illegible]

* From characterization sample (5Kg)(except as noted).
** 2-19 mm. from characterization sample, 19-76 mm. from field volume

SHERIDAN LOAM
S65Calif-27-22

Old Coast Road, 1200 feet SSE from Ranch house, about 100 feet down from ridge top. (Coastal grass site). Date of Sampling: May 25, 1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich; site selection by T. D. Cook.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Pachic Haploxeroll, coarse-loamy, mixed, thermic.

SOIL Sheridan loam SOIL Nos. S65Calif-27-24 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65580 - 65585

Depth (in.)	Horizon	1B1b Total: Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Sand: Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt: (0.05–0.02)	Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	* Coarse fragments 1A2a > 2 Pct.	2 – 19	19 – 76	
		Pct. of < 2 mm															Pct. of < 76mm		
0-8	A11	71.3	17.7	11.0	16.5	22.0	9.9	16.1	6.8	7.3	10.4	21.7	64.5	0	11	9	9	0	**
8-18	A12	71.6	17.1	11.3	17.3	21.9	9.4	15.7	7.3	6.7	10.4	22.0	64.3	0	11	12	12	0	**
18-28	A13	69.5	18.5	12.0	12.9	20.9	9.7	17.1	8.9	7.9	10.6	25.3	60.6	0	12	16	16	0	**
28-39	A14	70.0	18.6	11.4	11.8	20.4	9.6	18.6	9.6	7.9	10.7	26.6	60.4	0	11	11	11	0	**
39-43	C1	87.7	8.4	3.9	26.9	30.9	9.4	15.1	5.4	4.7	3.7	17.1	82.3	0	4	16	16	0	**
43-57+	C2	89.2	8.7	2.1	25.1	30.8	10.6	16.4	6.3	5.3	3.4	19.4	82.9	0	2	25	25	0	**

Depth (in.)	6A1a Organic carbon Pct	6B1a Nitrogen Pct	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct		Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content: Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1a H_2O 1:1		
0-8	1.68	0.129	13				1.51	1.54		0.95		12.0	5.0	0.005	0.005	6.4		
8-18	0.73	0.062	12				1.44	1.45		0.93		13.2	5.1	0.003	0.003	6.4		
18-28	0.52						1.44	1.49		0.91		13.6	5.8	0.010	0.009	6.5		
28-39	0.37						1.55	1.59		0.93		13.3	6.1	0.007	0.007	6.6		
39-43	0.15						1.95	1.99		0.88		7.5	2.7	0.007	0.006	6.5		
43-57+	0.08						2.07	2.07		0.79		6.7	2.3	0.001	0.000	6.5		

Depth (in.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1a NH_4OAc	5A3a Sum		6G1d KCl Ext. Al^{+++}				5D2 Exch. Na		Base saturation: 5C1 NH_4OAc	
	meq/100 g													Pct.		Pct	Pct.
0-8	9.4	1.7	0.3	0.2	11.6		12.7									91	
8-18	10.4	2.1	0.4	0.1	13.0		13.7									95	
18-28	13.1	3.0	0.4	0.1	16.6		16.9									98	
28-39	14.2	3.8	0.5	0.1	18.6		18.0									100+	
39-43	6.4	1.9	0.4	0.1	8.8		8.1							5		100+	
43-57+	5.0	1.4	0.4	0.1	6.9		5.8							7		100+	

Depth (in.)	Clay mineralogy (<0.002 mm) 7A: Mont.	Verm.	Mica	Kaolin	
	X-ray				

* From characterization sample (5Kg) (except as noted).

** 2-19 mm. from characterization sample, 19-76 mm. from field volume estimate.

SHERIDAN FINE GRAVELLY LOAM
S65Calif-27-24

Location: Monterey County, California, about 100 feet below top of ridge, 250 feet west of E¼ corner SE¼ Sec. 30, T15S, R6E, about 3.5 miles up Johnson Canyon Road from Armstrong Ranch main gate. (16" rainfall, inland, grass, north slope).
Date of Sampling: May 26, 1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich; site selection by T. D. Cook.
Classification: Pachic Haploxeroll, coarse-loamy, mixed, thermic.
Regosol (brunizemic).
Vegetation: Range, grass, annual fescue, soft chess, ripgut. Climate: Precipitation is 14 to 20 inches; mean annual temperature is 60°F., mean January is 50°F., and mean July is 66°F. Parent Material: Weathered diorite (gruss). Topography:

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Pachic Haploxeroll, coarse-silty, mixed, thermic

SOIL Salinas silty clay loam (taxadjunct) SOIL Nos. S65Calif-27-3 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65518 - 65525

Depth (in.)	Horizon	1B1b Total: Sand (2–0.05)	Total: Silt (0.05–0.002)	Total: Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	* Coarse fragments 1A2a: > 2 (Pct)	2 – 19 (Pct. of < 76 mm)	19 -76 (Pct. of < 76 mm)	
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm																	
0-5	Ap1	31.7	47.6	20.7	0.2	0.8	1.3	12.4	17.0	23.8	23.8	49.5	14.7	0	21	0	0	0	**
5-15	A12	30.1	49.9	20.0	0.1	0.7	1.4	11.8	16.1	24.7	25.2	49.1	14.0	0	20	0	0	0	**
15-23	A13	30.1	52.1	17.8	0.1	0.8	1.5	11.8	15.9	26.9	25.2	51.0	14.2	0	18	0	0	0	**
23-39	C1	33.6	50.7	15.7	0.2	0.8	1.6	13.1	17.9	27.0	23.7	54.3	15.7	0	16	0	0	0	**
39-59	C2	49.5	42.0	8.5	0.1	0.4	0.5	7.3	41.2	31.7	10.3	78.7	8.3	0	9	0	0	0	**
59-64	C3	27.3	62.3	10.4	0.0	0.2	0.2	2.2	24.7	44.8	17.5	71.2	2.6	0	10	0	0	0	**
64-71	C4	25.9	59.3	14.8	0.1	0.2	0.4	7.2	18.0	30.6	28.7	54.4	7.9	0	15	0	0	0	**
71-82+	C5	69.2	25.2	5.6	0.0	0.2	0.3	18.7	50.0	19.0	6.2	86.4	19.2	0	6	0	0	0	**

Depth (in.)	6A1a Organic carbon (Pct.)	6B1a Nitrogen (Pct.)	C/N	6C2a Ext. Iron as Fe (Pct.)	6E1b Carbonate as $CaCO_3$ (Pct.)		Bulk density: 4A1f 1/3 bar (g/cc)	Bulk density: 4A1h Oven dry (g/cc)	(g/cc)	5C1 Cm	Water content (Pct.)	4B1c 1/3 bar (Pct.)	4B2 15 bar (Pct.)	4D1 Extensibility COLEF (in./in.)	4D1 Extensibility COLE (in./in.)	pH: 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-5	2.74	0.252	11	0.9	-		1.27	1.40		0.83		25.4	18.3	0.032	0.032		7.9	
5-15	1.32	0.126	10	1.0	-		1.39	1.47		1.00		22.7	12.0	0.019	0.019		8.0	
15-23	0.77			0.9	-		1.30	1.38		1.00		23.8	10.6	0.020	0.020		8.0	
23-39	0.52			0.9	tr.		1.29	1.37		1.00		21.4	9.6	0.021	0.021		8.0	
39-59	0.23			0.7	1		1.34	1.36		1.00		14.4	6.4	0.004	0.004		8.2	
59-64	0.23			0.8	2		1.37	1.39		1.00		20.9	6.8	0.004	0.004		8.3	
[illegible]	[illegible]			[illegible]	[illegible]		[illegible]	[illegible]		[illegible]		[illegible]	[illegible]	[illegible]	[illegible]		[illegible]	

SALINAS SILTY CLAY LOAM (taxadjunct) 1/
S65Calif-27-3

Location: Monterey County, California. 150 feet SSW from NE corner of field, 0.7 mile along dirt road in a southwesterly direction from Metz Road, 400 feet N of the entrance to the Nello Solari Ranch on Metz Road about 5.5 miles north of King City.
Date of Sampling: May 18, 1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich; site selection by T. D. Cook.
Classification: Pachic Haploxeroll, coarse-silty, mixed, thermic. Brunizem (minimal).
Vegetation: Irrigated alfalfa. 20 tons per acre. barnyard manure applied 1963-64 season.

[redacted]

July mean 65°F. Parent Material: Moderately fine textured alluvium from dominantly sedimentary formations. Topography: Nearly level (<1 percent), high flood plain not subject to new deposition. Elevation: 260 feet. Drainage: Moderately well drained, moderately slow permeability, medium runoff. Soil Moisture: Moist throughout, about field capacity.
Remarks: Field pH by Truog Kit; paired sample S65Calif-27-11-1 through 7.

HORIZON	DESCRIPTION
Ap RSL 65518	0 to 5 inches, dark gray (10YR 4/1 toward 4/2) silty clay loam, very dark gray (10YR 3/1) moist; moderate medium subangular blocky structure; hard, firm, sticky and plastic; plentiful very fine and medium roots; few very fine interstitial, many very fine and few fine tubular pores; moderately alkaline (pH 8.0), noncalcareous; gradual smooth boundary.
A12 RSL 65519	5 to 15 inches, dark gray (10YR 4/1 toward 4/2) silty clay loam, very dark gray (10YR 3/1) moist; strong medium subangular blocky structure; hard, firm, sticky and plastic;plentiful very fine and medium roots; common very fine interstitial and common fine and few medium tubular pores; moderately alkaline (pH 8.0), noncalcareous; slight compaction of horizon due to tillage; gradual smooth boundary.
A13 RSL 65520	15 to 23 inches, grayish brown (2.5Y 5/2) silty clay loam, very dark grayish brown (10YR 3/2) moist; moderate medium subangular blocky structure; hard, firm, sticky and plastic; few fine and medium, plentiful very fine roots; common very fine interstitial and common fine and medium tubular pores; moderately alkaline (pH 8.0), noncalcareous, lower portion of profile slightly effervescent with disseminated lime; horizon has some very diffuse mixture of A and C; gradual smooth boundary.
C1 RSL 65521	23 to 39 inches, grayish brown (2.5Y 5/2) silty loam, very dark grayish brown (2.5Y 3/2) moist; weak medium subangular blocky structure; slightly hard, friable, sticky and plastic; few very fine and plentiful medium roots; many very fine interstitial and many very fine and common tubular pores; moderately alkaline (pH 8.0), slightly effervescent with disseminated lime; some large rodent holes filled with A1 material; clear smooth boundary.
C2 RSL 65522	39 to 59 inches, light brownish gray (2.5Y 6/2) very fine sandy loam, olive brown (2.5Y 4/3) moist; massive; soft, very friable, slightly sticky and slightly plastic; few very fine and few medium roots; many very fine interstitial and few fine tubular pores; moderately alkaline (pH 8.0), slightly effervescent with dis-

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Pachic Haploxeroll, fine-loamy, mixed, mesic

SOIL: Linne silty clay — SOIL Nos. S62Calif-42-11 — LOCATION: Santa Barbara County, California

SOIL SURVEY LABORATORY: Riverside, California — LAB. Nos. 62130 - 62135

Depth (in.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)			Coarse fragments 2A2: > 2 Pct	2 - 19	19 -76
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm															Pct. of < 76 mm	
0-9	Ap	36.1	31.4	32.5	0.2	1.1	3.0	19.3	12.5	15.1	16.3	39.8	23.6			0		
9-14	A12	42.7	27.3	30.0	0.2	1.1	3.1	23.7	14.6	13.6	13.7	43.6	28.1			0		
14-29	A13	43.8	26.7	29.5	0.2	1.0	2.9	24.3	15.4	13.3	13.4	45.1	28.4			0		
29-32	AC	47.2	24.7	28.1	0.3	1.0	2.6	26.1	17.2	11.7	13.0	47.0	30.0			0		
32-36	C1ca	57.2	24.6	18.2	1.0	2.8	2.7	30.6	20.1	10.2	14.4	53.3	37.1			0		
36-51+	C2	55.6	30.7	13.7	2.1	3.4	2.2	21.1	26.8	17.1	13.6	60.3	28.8			0		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	3A1a Carbonate in <0.002 mm. Pct.	6E1b Carbonate as $CaCO_3$ Pct.	6C2a Ext. Iron as Fe Pct.	Bulk density: 4A1f 1/3 bar clods g/cc	4A1h Oven-dry clods g/cc	g/cc	Water content: Pct.	4B1c 1/3 bar clods Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.		pH: 8C1b Sat. Paste H_2O	8C1a 1:10 H_2O
0-9	1.90	0.175	11	tr.	2	0.5	1.19	1.42			29.5	18.6	0.059		7.3	8.1
9-14	0.97	0.100	10	1	4	0.5	1.26	1.45			26.3	16.3	0.047		7.5	8.4
14-29	0.72	0.072	10	3	5	0.3	1.21	1.43			29.4	16.3	0.056		7.6	8.6
29-32	0.56	0.060	9	9	16	0.1	1.32	1.42			27.0	16.3	0.025		7.8	8.7
32-36	0.35			6	27	0.1	1.38	1.40			24.7	7.7	0.007		7.8	8.8
36-51+	0.23			5	24	0.1	1.55	1.58			12.4	7.7	0.007		8.1	9.0

Depth (in.)	Extractable bases 5B1a: 6N2b Ca	6O2c Mg	6P2a Na	6Q2a K	6H2a Ext. Acidity	Cation Exch. Capacity: 5A2a NaOAc			Water extract from saturated paste 8A1: 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-9	44.0	2.0	0.3	0.4	0.2	39.8			7.1	0.7	1.0	0.1		-	7.0	0.3		0.69
9-14	41.2	1.6	0.3	0.3	-	34.0			4.2	0.6	1.0	Tr.		-	4.2	0.1		0.45
14-29	38.3	1.5	0.3	0.2	-	31.6			3.5	0.4	1.0	0.1		-	3.7	Tr.		0.40
29-32	32.3	1.6	0.4	0.1	-	24.0			3.5	0.6	1.5	0.1		-	4.1	Tr.		0.46
32-36	25.3	1.2	0.4	0.1	-	14.1			2.6	0.4	1.6	Tr.		-	3.3	0.2		0.38
36-51+	23.0	1.4	0.5	0.1	-	12.9			1.9	0.3	2.1	Tr.		-	2.9	0.3		0.37

Depth (in.)	8A Water at Saturation Pct.	Exchangeable Na Pct.	5B1b Exch. Na meq/100g.	Gypsum Pct.		
0-9	59.1	1	0.3			
9-14	53.4	1	0.3			
14-29	54.9	1	0.3			
29-32	56.3	1	0.3			
32-36	42.7	2	0.3			
36-51+	35.2	3	0.4			

LINNE CLAY LOAM
S62Calif-42-11

Location: Santa Barbara County, California. Latitude 34° 48' 62" N., Longitude 120° 21' 06" W; equivalent to center of Sec. 4, T8N, R33W; from just north of Solomon Summit 1½ mi. south on Union Oil Company road and 750 feet east; 11 miles southeast of Santa Maria; field sheet 2B108. Date of Sampling: April 20, 1962.
Description by: L. C. Leifer. Collectors: R. E. Nelson, Lothair Grant, G. E. Shipman, C. J. Hilliker, L. C. Leifer.
Classification: **Pachic Haploxeroll, fine-loamy, mixed, mesic.**
Vegetation: Heavy growth wild oats, bur clover, a little annual lupine and ripgut brome. Use: Range, but was cultivated 10 to 15 years ago; little evidence of tillage in field, but fairly sharp break in soil structure on pit walls. Climate: Precipitation 15 inches. Mean annual temperature 57°F., July 61°F., and January 50°F. Parent Material: Soft marly sediments. Topography: Round smooth hills, locally slightly convex smooth, 22 percent northeast slope near crest of ridge. Elevation: 850 feet. Drainage: Well drained. Runoff is medium. Permeability is moderately slow. Soil Moisture: Moderately moist at sampling time. All horizons approaching wilting point.
Remarks: Field pH determinations by Hellige-Truog triplex indicator. Paired site not sampled.

HORIZON	DESCRIPTION
Ap RSL 62130	0 to 9 inches, very dark gray (10YR 3/1) clay loam, black (10YR 2/1) when moist; moderate coarse subangular blocky structure, 1/6 to 1/3 of horizon strong fine granular between larger peds, larger peds break mostly to granules; very hard, friable, sticky and plastic; plentiful very fine roots; many very fine, few fine interstitial pores; slightly effervescent disseminated lime, moderately alkaline (pH 8.0); clear smooth boundary.
A12 RSL 62131	9 to 14 inches, dark gray (10YR 4/1) clay loam, black (10YR 2/1) when moist; weak coarse prismatic structure; very hard, friable, sticky and plastic; plentiful very fine roots; many very fine, common fine interstitial pores and few fine tubular pores; strongly effervescent disseminated lime and some fine lime filaments especially on faces of larger peds, moderately alkaline (pH 8.0); gradual wavy boundary.
A13 RSL 62132	14 to 29 inches, gray (10YR 5/1) clay loam, very dark gray (10YR 3/1) when moist; weak coarse prismatic structure, breaks to medium and coarse angular blocky with much medium granular; very hard, friable, sticky and plastic; few very fine roots; many very fine interstitial and few fine and medium tubular pores; violently effervescent disseminated lime and many fine lime filaments especially on faces of larger peds, moderately alkaline (pH 8.0); gradual smooth boundary.
AC RSL 62133	29 to 32 inches, gray and light gray (10YR 5/1 and 7/1) sandy clay loam, gray and light grayish brown (10YR 5/1 and 6/2) when moist; massive; tends to break to weak coarse and medium angular blocky with some coarse granules; very hard, friable, sticky and plastic; very few very fine roots; many very fine interstitial and few fine tubular pores; violently effervescent disseminated lime and many distinct lime filaments, moderately alkaline (pH 8.0); clear irregular boundary.
Clca RSL 62134	32 to 36 inches, white (N8/ and 10/) fine sandy loam, very pale brown and white (10YR 7/2 and 8/2) when moist; massive; extremely hard, firm, sticky and plastic; no roots; common very fine interstitial and few very fine tubular pores; this horizon somewhat discontinuous and variable in thickness, also broken by squirrel holes; violently effervescent, moderately alkaline (pH 8.0); clear irregular boundary.
C2 RSL 62135	36 to 51 inches +, white (N 8/) mudstone, light gray and pale yellow (2.5Y 7/2 and 8/4) when moist; massive; very hard, firm, slightly sticky and slightly plastic; common very fine interstitial and very few fine tubular pores, strongly effervescent, moderately alkaline (pH 8.0).

NOTE: About 1/4 of Ap made up of krotovinas 2 to 3 inches across; 1/3 of A12 and A13 is krotovinas filled with strong fine and medium granular soil. Roots tend to concentrate in krotovinas and along weak vertical joints. Several krotovinas gray and dark gray (10YR 5/1 and 4/1m) 2 to 3 inches in diameter go several feet into C and also cross the AC and Clca horizon. A few white fragments of C 1/16 to 1/8 inches scattered through all of A horizons.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Sweeney silty clay loam (taxadjunct) **LOCATION** San Mateo County, California

SOIL FAMILY: Pachic Haploxeroll, fine-loamy, mixed, mesic

SOIL NOS. S55-Calif.-41-9 **LAB. NOS.** 55366-55372

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 2/	TEXTURAL CLASS
0- 6	A1	1.2	4.1	4.5	14.5	12.0	38.3	25.4	38.3	21.6	1	l
6-17	B2	1.5	4.3	4.2	13.8	12.1	37.6	26.5	39.2	19.7	2	l
17-23	B3	2.2	6.3	5.6	15.9	12.5	35.0	22.5	38.0	19.9	5	l
23-30	C1	2.2	9.2	8.4	22.0	15.3	28.6	14.3	43.0	14.9	1	fsl
30-48	C21	1.9	9.6	8.2	21.3	15.1	30.1	13.8	44.2	14.6	1	fsl
48-64	C22	8.3	21.5	11.6	22.5	12.6	17.1	6.4	35.6	7.6	0	cosl
64-70	C23	4.7	18.9	13.3	25.1	11.1	21.7	5.2	38.1	9.7	0	fsl

pH 8C1b SATURATED PASTE 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C.	Free Fe_2O_3 6C1b %	GYPSUM me./100g. SOIL	4A3a Bulk Density g./cc.		4B2 Soil Water 15 atmos. Pct.
6.4		6.9	2.14	0.177	12.1		0.7	1.9		1.12		14.3
6.5		6.9	1.50	0.130	11.5		0.5	1.9		1.32		15.0
6.5		7.0	0.86	0.077	11.2		0.3	1.7				14.9
6.6		7.2	0.26				0.4	1.5				14.7
6.9		6.8	0.24				0.3	1.5		1.49		14.2
6.8		7.2	0.07				0.3	1.0				11.9
6.3		7.3	0.07				0.3	0.9				12.5

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6N1a 6O1a Ca+Mg	6J1a 1/ HCO_3	6K1a Cl	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil						milliequivalents per liter					
51.5	24.7	20.1	8.6	0.7	0.3	84	1.1	0.1	6.3	5.2	0.4	49.3
56.3	25.4	21.2	7.7	1.2	0.2	86	1.0	0.1	2.2	2.4	0.5	47.8
60.8	25.5	23.1	7.0	1.0	0.2	88	0.8	0.1	1.1	2.2	0	45.1
62.4	28.7	24.2	6.9	0.7	0.1	89	0.8	Tr.	2.2	1.3	0	43.2
60.9	25.3	25.4	6.8	1.1	0.1	88	0.8	Tr.	1.3	1.0	0	42.2
66.2	26.6	24.1	7.8	1.3	0.1	87	0.7	Tr.	1.6	0.8	0	41.0
68.9	33.8	23.6	8.8	0.9	0.7	87	0.7	Tr.	1.0	0.7	0	43.9

SWEENEY SILTY CLAY LOAM (taxadjunct) 1/
S55Calif-41-9

Location: San Mateo County, California in the San Mateo County SCD, west side of Highway 5 (Skyline Highway), 3.65 miles south of Sky Londa, 100 feet north of U. S. Coast and Geodetic survey benchmark No. 0.387 dated 1936. Date of Sampling: May 17, 1955.

Description by: Ralph E. Nelson. Collectors: Reuben E. Nelson, Ralph E. Nelson, and Rudolph Ulrich.

Classification: Pachic Haploxeroll, fine-loamy, mixed, mesic.

Vegetation: Annual grasses consisting of wild oat, soft chess, bur clover, some California lupine. Climate: 35-40 inches rainfall. Parent Material: Basaltic material classified as diabase. Relief: Hilly to mountainous with gently to moderately steep rolling hills. The slope at the sampling location was about 6 percent, slightly concave toward the southeast. Elevation: 2,100 feet. Drainage: Surface drainage is medium, internal drainage is moderate.

Remarks: Many worms in surface 6 inch and occasional to few worms to a depth of 17 inches.

HORIZON		DESCRIPTION
A_1	0- 6"	Dark grayish brown (10YR 4/2; very dark grayish brown to very dark brown, 10YR 2.5/2, moist) light silty clay loam. Moderate medium to fine granular structure. Hard, friable, slightly plastic and nonsticky. This horizon contains many roots and many worm casts. The reaction is 7.0. The lower boundary is smooth and clear.
B_2	6-17"	Dark grayish brown (10YR 4/2; very dark grayish brown and very dark brown, 10YR 2/2-3/2, moist) silty clay loam, slightly finer textured than horizon above. Weak medium subangular blocky. Hard, firm to friable, slightly plastic and nonsticky. There are thin continuous clay skins on the vertical faces of the structural aggregates. There are many roots and a few worm casts. The reaction is 7.0. This horizon grades clearly and with a wavy boundary into:
B_3	17-23"	Indistinctly mottled dark brown and dark grayish brown (10Y 4/3-4/2; dark brown and very dark grayish brown, 10YR 3/3-3/2, moist) clay loam. Weak medium and fine subangular blocky. Hard, firm to friable, plastic and slightly sticky. There are thin patchy clay skins on the darker vertical faces of the structural aggregates. There are numerous roots, a few worm casts, and some krotovinas. The reaction is 7.0. The lower boundary is clear but tonguelike.
C_1	23-30"	Mottled yellowish brown and dark brown (10YR 5/4-3/3; dark yellowish brown and very dark grayish brown, 10YR 4/4-3/2, moist) light clay loam. Very weak medium subangular blocky. Hard, firm, slightly plastic and slightly sticky. There are some thin patchy clay skins on vertical surfaces and dark metallic segregations which glisten in the sunlight. Very few roots. The reaction is 7.0. Grades with a gradual and wavy lower boundary into:
C_{21}	30-48"	Mottled yellowish brown and dark brown (10YR 5/4-3/3; dark yellowish brown and very dark grayish brown, 10YR 4/4-3/2, moist)

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Pachic Haploxeroll, fine-loamy, mixed, thermic
SOIL Salinas silty clay loam SOIL Nos. S65Calif-27-11 LOCATION Monterey County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65526 - 65532

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	* Coarse fragments 1A2a > 2 Pct.	2 - 19 Pct. of < 76 mm	19 - 76 Pct. of < 76 mm	
		Pct. of < 2 mm (Size class and particle diameter (mm) 3A1)																	
0-5	Ap1	43.2	32.2	24.6	0.4	1.0	1.7	19.4	20.7	14.9	17.3	50.2	22.5	0	25	0	0	0	**
5-13	Ap2	43.7	31.7	24.6	0.5	0.8	1.6	19.3	21.5	14.6	17.1	50.9	22.2	0	25	0	0	0	**
13-23	A13	32.8	40.9	26.3	0.1	0.1	0.3	12.5	19.8	18.1	22.8	48.4	13.0	0	26	0	0	0	**
23-33	A14	42.9	36.7	20.4	0.1	0.3	0.4	17.0	25.1	18.6	18.1	58.1	17.8	0	20	0	0	0	**
33-40	A15	63.4	25.1	11.5	0.2	0.3	0.7	26.7	35.5	16.8	8.3	74.6	27.9	0	12	0	0	0	**
40-49	C1	70.1	21.0	8.9	0.2	0.3	0.7	29.0	39.9	15.6	5.4	79.8	30.2	0	9	0	0	0	**
49-75+	C2	61.3	29.0	9.7	0.1	0.2	0.3	15.2	45.5	21.9	7.1	80.6	15.8	0	10	0	0	0	**

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Atterberg limits: Plastic limit Pct. of <0.4 mm	Liquid limit	Plastic index		Water content: Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-5	1.36	0.129	11		-								11.8				7.8	
5-13	1.26	0.116	11		-		19	33	14				12.7				7.8	
13-23	1.17				-								14.0				7.7	
23-33	0.84				-								11.2				7.8	
33-40	0.41				-								6.9				8.0	
40-49	0.30				tr.								5.4				8.3	
49-75+	0.15				2			-	N.P.				6.0				8.3	

Depth (in.)	Extractable bases (meq/100 g): 6N2a Ca	6O2a Mg	6P2a Na	5B1a 6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste (meq/liter): 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity mmho/cm
0-5	17.3	7.7	1.5	1.4	27.9			24.5										
5-13	17.4	7.7	1.4	1.4	27.9			25.5										
13-23	18.7	9.0	1.8	0.8	30.3			28.4										
23-33	16.0	7.7	1.6	0.5	25.8			24.9										
33-40	10.2	4.9	4.0	0.8	19.9		18.2		2.2	1.6	6.4	0.2		0.2	2.6	1.5		1.06
40-49	14.3	4.0	0.9	0.3	19.5			11.7										
49-75+	18.7	4.6	0.9	0.1	24.3			12.3										

Depth (in.)	8A Water at Saturation Pct.	5D2 Exchangeable Na Pct.	5B1b Exch. Na meq./100g.	Gypsum Pct.	5C1 Base Sat. NH_4OAc Pct.	Clay mineralogy (<0.002 mm.) 7A: Mont.	Verm.	Mica	Kaolin.	
						X-ray				
0-5		6			100+					
5-13		5			100+					
13-23		6			100+					
23-33		6			100+					
33-40	37.4	21	3.2		100+					
40-49		8			100+					
49-75+		7			100+					

* From characterization sample (5Kg) (except as noted).
** 2-19 mm. from characterization sample, 19-76 mm. from field volume estimate.

SALINAS SILTY CLAY LOAM
S65Calif-27-11

Location: Monterey County, California, 1.3 mile south of Chualar (town) underpass on Highway 101, 1100 feet southwest on paved road, 600 feet southeast on Farm Road, about 30 feet northeast into field. Date of Sampling: May 20, 1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich. Site selection by T. D. Cook.
Classification: **Pachic Haploxeroll, fine-loamy, mixed, thermic. (Brunizem).**
Vegetation: Truck crops, onions just harvested. Climate: Precipitation 12 inches;

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Pachic Haploxeroll, fine, montmorillonitic, thermic

SOIL Rincon clay loam (taxadjunct) SOIL Nos. S64Calif-57-16 LOCATION Yolo County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64480 - 64487

		1B1b	Size class and particle diameter (mm) 3A1											
		Total	Sand					Silt		Int. II		3A1a	3A1a Non-	Coarse fragments 1A2a

RINCON CLAY LOAM (taxadjunct) 1/
S64Calif-57-16

Location: Yolo County, California. A pit located 1/8 mile west of County Road 95B and 3/8 mile north of County Road 18A, T.10N., R.1E. (section lines not surveyed), 5 miles northwest of Woodland. Photo ABB-47T-16. Date of Sampling: September 29, 1964.
Description by: C. R. Goudey. Collectors: R. E. Nelson, B. R. Brasher, G. M. Kennedy.

[redacted]

Classification: Pachic Haploxeroll, fine, montmorillonitic, thermic.
Vegetation: Fallow (safflower was prior crop). Climate: 15.5 inches average annual precipitation and mean annual temperature of 61°F. Parent Material: Alluvium from sedimentary rocks. Topography: Alluvial fan; plane; less than one percent slope; near toe. Elevation: 80 feet

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Pachic Ultic Haploxeroll, clayey skeletal, mintmorillonitic, thermic

SOIL Santa Lucia shaly clay loam (taxadjunct) SOIL Nos. S62Calif-42-10 LOCATION Santa Barbara County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 62126 - 62129

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002) 1/	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)			Coarse fragments 2A2 > 2 Pct.	2-19	19-76
		Pct. of < 2 mm															Pct. of < 19 mm	
0-8	A11	26.8	35.3	37.9	5.8	8.0	3.8	5.8	3.4	8.0	27.3	14.2	23.4				25	
8-17	A12	23.9	35.8	40.3	4.1	6.3	3.4	5.8	4.3	8.5	27.3	15.8	19.6				19	
17-24	A13	23.2	34.0	42.8	4.7	5.8	3.2	5.6	3.9	8.0	26.0	14.9	19.3				26	
24-36+	R		Rock															

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N		Carbonate as $CaCO_3$ Pct.		Bulk density g/cc	g/cc	g/cc		Water content Pct.	Pct.	4B2 15 bar Pct.			pH 8C1a H_2O 1:1	8C1c 1 N KCl 1:1	(1:1)
0-8	9.43	0.743	13										36.7			6.0	5.2	
8-17	6.54	0.494	13										37.0			5.9	5.2	
17-24	5.55	0.443	13										36.2			5.8	5.0	
24-36+		Rock																

Depth (In.)	Extractable bases 5B1a: 6N2b Ca	602c Mg	6P2a Na	6Q2a K		6H2a Ext. Acidity	Cation Exch. Capacity 5A3a Sum									Base saturation 5C3 Sum Pct.	Pct.
	meq/100 g																
0-8	29.6	9.2	0.5	5.6		24.1	70.0									66	
8-17	30.0	10.9	0.9	2.8		24.5	69.1									64	
17-24	25.1	12.4	1.0	1.9		26.5	66.9									60	
24-36+		Rock															

Depth (In.)	3A1 Clay fraction mineralogy: 7A2 X-ray 2/	
0-8		
8-17	MM 3	
17-24		
24-36+	AM 6	

1/ Shale fragments of sand and silt size break down in particle size distribution analysis. The soil has the chemical properties of a material of higher clay content and the mechanical properties of one of lower clay content.

2/ Mineral code:
MM = montmorillonite
AM = amorphous
Approximate weight fraction (X-ray)
3 = one-fifth to one-third
6 = indeterminate

SANTA LUCIA SHALY CLAY LOAM (taxadjunct) 1/
S62Calif-42-10

Location: Santa Barbara County, California. NW¼ of NE¼ Sec. 5, T.9N, R.35W. Very near watershed divide in Casmalia Hills, reached by trail road east southeast out of Corralitos Ranch headquarters; 5 airline miles south and slightly west of Guadalupe. Date of Sampling: April 19, 1964.

Description by: L. C. Leifer. Collectors: R. E. Nelson, Lothair Grant, G. E. Shipman, C. J. Hilliker, L. C. Leifer.

Classification: Pachic Ultic Haploxeroll, clayey skeletal, montmorillonitic, thermic.

Vegetation: Oak-grass used for range; ripgut brome, foxtail barley, alfilaria, a few thistle, scattered coast live oak with 35 percent canopy; sample site 35 and 50 feet from trees and no tree roots in pit. Climate: Precipitation 16 inches. Mean annual temperature 57°F., January 49°F., July 63°F. Parent Material: Diatomaceous shale, Monterey. Topography: Generally steep hills, locally plane 18 percent north northwest. Elevation: 1500 feet. Drainage: Well drained. Runoff is medium. Permeability is moderate. Soil Moisture: Nearly dry in upper 8 inches, slightly moist below.

Remarks: Field determinations of pH by Hellige-Truog triplex indicator. Paired site sample numbers are Santa Lucia S62Calif-42-9-1 through 4.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Pachic Ultic Haploxeroll, fine-loamy, mixed, frigid

SOIL Tournquist loam SOIL Nos. S66Calif-25-1 LOCATION Modoc County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66327 - 66333

		1B1b	Size class and particle diameter (mm) 3A1

TOURNQUIST LOAM
S66Calif-25-1

Location: Modoc County, California. NW¼ NW¼ Sec. 33, T48N, R11E, 402 feet east of line tag between Sections 32 and 33. Date of Sampling: May 20, 1966.
Description by: J. S. Cahoon. Collectors: J. E. McClelland, J. P. Tribe, K. Flach, D. R. Taylor, G. H. Simonson, R. W. Mayko, J. S. Cahoon.
Classification: Pachic Ultic Haploxeroll, fine-loamy, mixed, frigid.
Vegetation: Timber and grasses. Ponderosa pine, white fir, squaw carpet, serviceberry, strawberry, bitterbrush, balsamroot, pine grass. Climate: Estimated as follows: average annual precipitation is 18 inches; mean annual air temperature 44°F., mean summer air temperature is 57°F. Deep snow covers the ground in winter. Parent Material: Reddish mudflow consisting of rounded andesitic boulders and cobbles in loamy matrix capped by a thin mantle of higher ash content is suggested. Topography: Tableland. Slope gradient is 1 percent. Elevation: 5450 feet. Drainage: Well drained, permeability is moderate, runoff is slow. Soil Moisture: Moist throughout.
Remarks: BPR samples were taken of the A horizon, 0 to 14 inches, and the B2 horizon, 14 to 34 inches. Field determination of pH was made with bromthymol blue indicator. About 10 percent volume of fine rounded concretions or "shot" occurred throughout. Coarse fragment volume commonly is 30 to 40 percent in the B2 of Tournquist soils. The A horizons of these soils appear to have low bulk density and a higher ash content compared to underlying horizons. Soil temperature at 20 inches was 49°F. at time of sampling.

HORIZON	DESCRIPTION
O1	2 to 0 inches, fresh and partly decomposed litter of pine needles and leaves.
A1 RSL 66327	0 to 4 inches, reddish brown (5YR 4/3) loam, dark reddish brown (5YR 3/2) moist; strong very fine granular structure; soft to slightly hard, very friable, slightly sticky and slightly plastic; many fine roots; many very fine interstitial pores; slightly acid (pH 6.5); clear smooth boundary.
A3 RSL 66328	4 to 14 inches, reddish brown (5YR 4/3) loam, dark reddish brown (5YR 3/3) moist; moderate very fine granular structure; soft to slightly hard, very friable, slightly sticky and slightly plastic; many fine roots; many very fine interstitial pores; slightly acid (pH 6.4); clear smooth boundary.
IIB21 RSL 66329	14 to 21 inches, reddish brown (5YR 4/3) gravelly loam, dark reddish brown (5YR 3/3) moist; moderate very fine granular and weak medium subangular blocky structure; slightly hard, friable, slightly sticky and plastic; many fine roots; common very fine tubular pores; stone line of hard andesite or basalt at top of this horizon; slightly acid (pH 6.4); gradual smooth boundary.
IIB22 RSL	21 to 34 inches, reddish brown (5YR 4/4) gravelly clay loam, dark reddish brown (5YR 3/4) moist; moderate fine subangular blocky structure; slightly hard,

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Crouch loamy coarse sand (taxadjunct) **LOCATION** San Diego County, California

SOIL FAMILY: Ultic Haploxeroll, sandy, mixed, mesic.

SOIL NOS. S60Calif-37-5 **LAB. NOS.** 60418 - 60424

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1	2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	

CROUCH LOAMY COARSE SAND (taxadjunct) 1/
S60Calif-37-5

Description: George Borst
Date of Sampling: November 10, 1960
Collectors: Reuben Nelson, Lothair Grant, Fred C. Eggers, John Wachtell, and George Borst
Location: Approximately 4200 feet south of northeast corner, sec. 23, T. 11 S., R. 4 E., southeast of Montezuma Valley Prison Camp, San Diego County, Calif.
Elevation: 4,000 feet Relief: Moderately sloping, 6 percent.
Vegetation: Native chaparral Use: Range.
Drainage: Somewhat excessively drained
Parent Material: Deeply weathered granitic rock
Climate: This soil occurs under about 17" annual rainfall. The mean annual temperature is about 57°, the mean January temperature about 44°, and the mean July temperature about 73°.
Remarks: This soil has a thicker A horizon and a more prominent B horizon than that sampled as S60Calif37-3. This site is probably more nearly modal for the series.
Classification: Ultic Haploxeroll, sandy, mixed, mesic.

HORIZON		DESCRIPTION
A11	0- 2"	Grayish brown (10YR 5/2) loamy coarse sand, very dark grayish brown (10YR 3/2) when moist; weak very fine crumb structure; soft, very friable, nonsticky and nonplastic; abundant very fine random roots; slightly acid (pH 6.5); micaceous; clear smooth boundary.
A12	2- 7"	Grayish brown (10YR 5/2) loamy coarse sand, very dark grayish brown (10YR 3/2) when moist; weak fine crumb structure; soft, friable, nonsticky and nonplastic; abundant fine random roots; neutral (pH 7.2); micaceous; clear wavy boundary, with tongues of this horizon penetrating the underlying material.
AC	7-10"	Brown (10YR 5/3) loamy coarse sand, dark brown (10YR 3/3) when moist; massive; slightly hard, friable, nonsticky and nonplastic; plentiful fine to medium random roots; neutral (pH 7.2); micaceous; clear wavy boundary.
C1	10-20"	Brown (10YR 5/3) loamy coarse sand, containing very pale brown (10YR 7/3) and dark gray (10YR 4/1) mineral particles, dark brown (10YR 4/3), pale brown (10YR 6/3), and very dark gray (10YR 3/1) when moist; massive; hard, friable, nonsticky and nonplastic; few medium to coarse angular roots; common fine random tubular pores; few thin clay films in pores and on mineral grains; slightly acid (pH 6.5); micaceous; diffuse smooth boundary.
C2	20-29"	Brown (10YR 5/3) loamy coarse sand, containing very pale brown (10YR 7/3) and dark gray mineral particles, dark brown (10YR 4/3), pale brown (10YR 6/3), and very dark gray (10YR 3/1) when moist; massive; hard, friable, nonsticky and nonplastic; few medium to coarse random roots; common fine random tubular pores; few thin clay films in pores and on mineral grains; slightly acid (pH 6.3); micaceous; gradual smooth boundary.
C3	29-37"	Brown (10YR 5/3) loamy coarse sand, containing very pale brown (10YR 7/3) and dark gray (10YR 4/1) mineral particles, dark brown

OIL SURVEY LABORATORY Riverside, California

SOIL TYPE Sheephead stony fine sandy loam **LOCATION** San Diego County, California

SOIL FAMILY: Ultic Haploxeroll, loamy, mixed, mesic, shallow

SOIL NOS. S60Calif-37-7 **LAB. NOS.** 60431 - 60436

DEPTH INCHES	HORIZON	1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	3A1 0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
		PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)										
0- 4	A11	5.7	11.6	8.3	32.7	17.5	17.9	6.3	48.6	8.7		fs1-ls
4- 8	A12	3.9	12.0	7.7	31.0	17.2	21.7	6.5	49.7	10.6		fs1
8-24	R &C	1.7	6.4	7.7	36.0	24.5	19.0	4.7	62.8	5.9		fs1
24-40	R &C	3.7	10.6	9.6	31.9	19.4	17.3	7.5	51.9	6.2		fs1
40-51	R &C	1.6	7.2	7.3	31.9	21.6	24.3	6.1	58.2	9.3		fs1
51-57+	C		not		determined							

pH 8C1a H_2O 1:1	pH 8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC x10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	Free Fe_2O_3 6C1b %	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.2	5.4		4.09	0.161	25.4				1.6			6.7
6.7	5.3		0.83	0.044	18.9				1.5			4.5
6.6	4.9		0.26	0.013					1.7			3.9
6.2	4.5		0.12						1.7			4.2
6.3	4.2		0.15						2.0			5.9
		not		determined					1.5			

5A2a CATION EXCHANGE CAPACITY (Na)	6N2b Ca	EXTRACTABLE CATIONS 6O2c Mg	6H2a H	6P2a Na	5B1a 6Q2a K	BASE SAT. % 5C3	Sum of Cations me./100 g. soil 5A3a	g. K				MOISTURE AT SATURATION %
		milliequivalents per 100g. soil →						milliequivalents per liter →				
15.5	10.1	1.3	5.5	0.1	0.5	69	17.5					
9.6	5.2	1.2	2.5	0.1	0.3	73	9.3					
6.7	3.1	1.0	1.8	0.1	0.2	71	6.2					
7.3	3.5	1.2	2.2	0.1	0.2	69	7.2					
11.9	6.3	2.2	3.0	0.2	0.1	75	11.8					
9.9	5.2	2.1	2.3	0.2	<0.1	77	9.8					

1/ The coarse fragments were discarded when the soils were sampled.

SHEEPHEAD STONY FINE SANDY LOAM
S60Calif-37-7

Location: Approximately 800 feet south-southeast of entrance to the Lucky S Ranch gate on Sunrise Highway in SW1/4 of Section 7, T. 14 S., R. 5 E., San Diego County, California. Date of Sampling: November 15, 1960.
Description by: George Borst. Collectors: Reuben Nelson, Fred C. Eggers, Cliff Henry, and George Borst.
Classification: Ultic Haploxeroll, loamy, mixed, mesic, shallow.
Vegetation: Native chaparral. Climate: This soil occurs under an average annual rainfall of about 30 inches, a mean annual temperature of 53° F., a mean January temperature of 39° F., and a mean July temperature of 70° F. Parent Material: Deeply weathered micaceous schist. Use: Range. Relief: Steep, 40 percent. Elevation: 5,000 feet. Drainage: Somewhat excessively drained. Reaction: pH was determined in the field with a Truog (Hellige) soil reaction kit. Remarks: The weathered schist is thinly laminated, highly micaceous, with a dip of about 76°. The R&C horizon was subdivided for laboratory analysis at 24 inches and 40 inches.

HORIZON	
A11 RSL No. 60431	0 to 4 inches, dark grayish brown (10YR 4/2) fine sandy loam, dark brown (10YR 3/3) when moist; weak very fine crumb structure; soft, very friable, nonsticky and nonplastic; abundant very fine random roots; slightly acid (pH 6.5); highly micaceous; clear smooth boundary.
A12 RSL No. 60432	4 to 8 inches, brown (10YR 5/3) highly micaceous fine sandy loam, dark brown (10YR 3/3) moist; weak fine crumb structure; soft, friable, nonsticky and nonplastic; abundant fine roots; common fine tubular pores; slightly acid (pH 6.2); clear, irregular boundary.
R&C RSL Nos. 60433 60434 60435	8 to 51 inches, fractured foliar micaceous schist and light yellowish brown (10YR 6/4) angular gravelly or stony fine sandy loam, yellowish brown (10YR 5/4) moist; massive; soft, friable, nonsticky and nonplastic; common medium to coarse roots; medium acid (pH 6.0); diffuse, irregular boundary.
R RSL No. 60436	51 to 57 inches +, pale yellow (2.5Y 7/4) deeply weathered foliar schist; yellowish brown (10YR 5/4) when moist, with yellowish red (5YR 4/8) colloidal coatings on the rock fragments; few coarse random roots; medium acid (pH 6.0).

SOIL SURVEY LABORATORY Riverside, California

OIL TYPE Crouch loamy sand **LOCATION** San Diego County, California

SOIL FAMILY: Ultic Haploxeroll, coarse-loamy, mixed, mesic

SOIL NOS. S61Calif-37-3 **LAB. NOS.** 6117 - 6123

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 5	A11	14.2	19.4	9.4	19.5	8.7	21.7	7.1	29.3	12.3	42	cosl
5-12	A12	16.6	22.2	10.0	19.4	6.6	18.7	6.5	25.2	11.1	41	cosl-lcos
12-24	AC	14.2	21.4	10.3	19.0	11.4	17.8	5.9	29.0	9.8	45	lcos-cosl
24-34	C1	15.1	21.8	10.0	20.9	8.4	18.6	5.2	27.9	10.8	34	" "
34-42	C2	9.5	18.9	9.9	22.6	9.2	23.4	6.5	32.1	13.7	18	cosl
42-55	C3	16.6	29.9	9.2	19.1	6.1	13.1	6.0	22.7	7.1	36	lcos
55-61+	C4	18.2	32.6	10.2	18.4	5.2	9.5	5.9	19.6	4.9	59	lcos

pH 8C1a H_2O 1:1	pH 8C1c KCl 1:1	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	Free Fe_2O_3 6C1b %	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.9	6.1	2.58	0.122	21.1				0.8			7.7
6.7	5.7	1.13	0.059	19.2				0.7			6.2
6.5	5.3	0.63	0.032					0.8			5.4
6.4	4.8	0.22	0.015					0.9			5.4
6.4	4.9	0.23						0.8			5.8
6.4	5.0	0.13						0.7			5.6
6.4	4.9	0.14						0.7			5.4

5A2a CATION EXCHANGE CAPACITY (+ Na)	EXTRACTABLE CATIONS 6N2b Ca	6O2c Mg	6H2a H	6P2a Na	6Q2a K	BASE SAT. % 5C3	Sum of Cations me./100g. soil 5A3a	g. K	MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						milliequivalents per liter →		
20.5	14.9	1.7	5.4	0.2	1.0	77	23.2		
15.4	9.1	1.5	5.0	0.2	1.0	70	16.8		
14.9	8.2	0.5	3.6	0.2	0.6	73	13.1		
13.0	8.8	0.5	2.3	0.2	0.2	81	12.0		
12.2	8.8	0.5	2.3	0.2	0.3	81	12.1		
8.9	5.2	0.5	1.6	0.2	0.3	79	7.8		
8.3	4.4	0.4	1.8	0.2	0.3	75	7.1		

1/ Percent of the whole soil

CROUCH LOAMY SAND
861Calif-37-3

Description: George Borst
Date of Sampling: January 18, 1961
Collectors: Reuben Nelson, Cliff Henry, and George Borst
Location: About 1/2 mile southwest of Matagual Ranch headquarters on the south side of reservation road, in NW1/4 sec. 35, T. 11 S., R. 3 E. (projected), Santa Ysabel Indian Reservation, San Diego County, California.
Elevation: 4,500 feet
Relief: Very steep, 60 percent
Use: Forestry
Vegetation: Mixed coniferous and deciduous forest
Drainage: Somewhat excessively drained
Parent Material: Deeply weathered granitic rock
Climate: This soil occurs under an average annual rainfall of about 38", a mean annual temperature of 53°, a mean January temperature of 39°, and a mean July temperature of 70°.
Remarks: A deep layer of forest duff, approximately 2 inches in thickness, occurs at the surface
Classification: Ultic Haploxeroll, coarse-loamy, mixed, mesic.
Reaction: pH was determined in the field with a Truog (Hellige) soil reaction kit

HORIZON		DESCRIPTION
A11	0- 5"	Dark grayish brown (10YR 4/2) loamy sand, dark brown (10YR 3/3) [illegible] nonsticky and nonplastic; abundant very fine vertical roots; many very fine interstitial pores; slightly acid (pH 6.5); clear smooth boundary.
A12	5-12"	Brown (10YR 5/3) loamy sand, dark brown (10YR 3/3) when moist; weak fine granular structure; soft, very fine, nonsticky and nonplastic; few medium random roots; many very fine interstitial pores; slightly acid (pH 6.2); gradual smooth boundary.
AC	12-24"	Pale brown (10YR 6/3) loamy sand, dark brown (10YR 4/3) when moist; massive; soft, very friable, nonsticky and nonplastic; plentiful medium to coarse random roots; medium acid (pH 6.0); gradual smooth boundary.
C1	24-34"	Pale yellow (2.5Y 8/4) loamy sand, light olive brown (2.5Y 5/4) when moist; massive; soft, very friable, nonsticky and nonplastic; few medium to coarse random roots; medium acid (pH 6.0); abrupt smooth boundary.
C2	34-42"	Pale yellow (2.5Y 8/4) sandy loam, light yellowish brown (2.5Y 6/4) when moist; massive; hard, firm, nonsticky and nonplastic; few medium to coarse random roots; common medium random interstitial pores; medium acid (pH 6.0); abrupt smooth boundary.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Crouch sandy loam **LOCATION** San Diego County, California

SOIL FAMILY: Ultic Haploxeroll, coarse-loamy, mixed, mesic.

SOIL NOS. S61Calif-37-6 **LAB. NOS.** 6141 - 6147

DEPTH INCHES	HORIZON	1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	2/ CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
		PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)										
0- 3	A11	11.7	23.3	8.8	14.7	7.1	23.0	11.4	25.2	13.4	20	cosl
3-13	A12	10.0	21.9	8.9	15.8	7.8	23.5	12.1	26.8	13.5	18	cosl
13-29	A3	12.2	24.6	8.4	14.2	7.4	22.6	10.6	25.4	12.7	17	cosl
29-42	B1	8.1	19.1	9.9	18.4	10.2	24.2	11.1	32.3	13.2	8	cosl
42-56	B2	11.6	21.7	9.0	16.3	9.2	21.6	10.6	29.3	11.4	7	cosl
56-70	B31	27.2	24.1	8.7	13.8	6.7	11.9	7.6	21.0	5.8	17	lcos
70-88	B32	19.3	23.3	11.8	20.9	6.6	11.1	7.0	25.5	4.4	17	lcos

pH 8C1a H_2O 1:1	8C1c KCl 1:1		ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	Free Fe_2O_3 6C1b %	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
6.4	5.2		2.25	0.148	15.2				1.8			8.0
6.0	4.8		1.31	0.103	12.7				2.0			8.1
6.2	4.4		0.48	0.040	12.0				1.8			8.3
6.1	4.0		0.35	0.029					2.0			10.9
6.0	3.8		0.15						1.9			11.8
6.4	3.7		0.08						1.6			7.8
6.3	3.7		0.05						1.9			6.9

5A2a CATION EXCHANGE CAPACITY (Na)	EXTRACTABLE CATIONS 6N2b Ca	6O2c Mg	6H2a H	6P2a Na	5B1a 6Q2a K	BASE SAT. % 5C3	Sum of Cations me./100 g. soil 5A3a	Sum of Cations me./100g. clay	Sum of Cations me./100g. clay + fine silt
	milliequivalents per 100g. soil →								
19.8	11.1	2.4	6.4	0.2	0.6	69	20.7	182	83
16.5	9.5	2.3	6.5	0.2	0.5	66	19.0	157	74
19.3	11.0	3.2	5.3	0.3	0.3	74	20.1	190	86

CROUCH SANDY LOAM
S61Calif-37-6

Description: George Borst
Date of Sampling: January 25, 1961
Collectors: Reuben Nelson, Lothair Grant, and George Borst
Location: Approximately 500 feet south of north 1/4 corner, sec. 29, T. 12 S., R. 4 E., about 2 miles north-northeast of Julian, **San Diego County, California.**
Elevation: 4,500 feet Relief: Steep, 40 percent
Vegetation: Forest, mostly oak and cedar Use: Woodland
Drainage: Well drained, moderately slow subsoil permeability
Parent Material: Deeply weathered granitic rock
Climate: This soil occurs under an average annual rainfall of about 38", a mean annual temperature of 53°, a mean January temperature of 39°, and a mean July temperature of 70°.
Remarks: The A horizon of this soil is relatively thick, and erosion relatively slight, as compared with most soils in this series in San Diego County.
Classification: **Ultic Haploxeroll, coarse-loamy, mixed, mesic.**

HORIZON		DESCRIPTION
A11	0- 3"	Very dark grayish brown (10YR 3/2) sandy loam, very dark brown (10YR 2/2) when moist; weak very fine crumb structures; soft, very friable, nonsticky and slightly plastic; abundant very fine random roots; many very fine interstitial pores; slightly acid (pH 6.5); abrupt smooth boundary.
A12	3-13"	Dark grayish brown (10YR 4/2) sandy loam, very dark grayish brown (10YR 3/2) when moist; massive; slightly hard, friable, nonsticky and slightly plastic; plentiful fine random roots; common fine random tubular pores; slightly acid (pH 6.5); clear smooth boundary.
A3	13-29"	Dark yellowish brown (10YR 4/4) sandy loam, containing burrows filled with soil of the overlying horizon, dark yellowish brown (10YR 3/4) when moist; massive; slightly hard, friable, slightly sticky and plastic; plentiful medium random roots; common fine random pores; slightly acid (pH 6.5); clear wavy boundary.
B1	29-42"	Yellowish brown (10YR 5/4) sandy loam, dark yellowish brown (10YR 4/4) when moist; massive; hard, firm, slightly sticky and plastic; few medium random roots; few fine random tubular pores; colloid in bridges between mineral grains; slightly acid (pH 6.5); clear wavy boundary.
B2	42-56"	Yellowish brown (10YR 5/4) sandy clay loam, dark yellowish brown (10YR 4/4) when moist; massive; hard, firm, sticky and plastic; few coarse random roots; common medium random tubular pores; many thick clay films on fractures planes; slightly acid (pH 6.5); gradual smooth boundary.
B31	56-70"	Yellowish brown (10YR 5/4) sandy clay loam, containing very dark grayish brown (10YR 3/2) and very pale brown (10YR 8/4) mineral particles, dark yellowish brown (10YR 4/4) when moist; massive; hard, firm, sticky and plastic; few coarse random roots; common medium random tubular pores; many thick clay films on fractured planes; slightly acid (pH 6.5); gradual smooth boundary.
B32	70-88"+	Yellowish brown (10YR 5/4) coarse sandy loam, containing very dark grayish brown (10YR 3/2) and very pale brown (10YR 8/4) mineral [illegible]

OIL SURVEY LABORATORY Riverside, California

SOIL TYPE Crouch sandy loam **LOCATION** San Diego County, California

SOIL FAMILY: Ultic Haploxeroll, coarse-loamy, mixed, mesic.

SOIL NOS. S61Calif-37-7 **LAB. NOS.** 6148 - 6155

DEPTH INCHES	HORIZON	1B1b VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	2/ CLAY < 0.002	0.2-0.02	3A1 0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
		PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)										
0-4	A11	7.1	19.9	11.0	21.8	8.4	22.4	9.4	31.7	12.0	3	cosl
4-19	A12	8.3	21.5	11.1	20.6	8.1	20.3	10.1	29.4	11.1	2	cosl
19-28	A3	7.8	19.7	10.6	21.6	8.2	20.7	11.4	30.2	11.3	2	cosl
28-38	B1	9.1	23.8	10.8	19.6	7.0	18.8	10.9	26.9	10.3	3	cosl
38-50	B21	5.4	21.6	12.0	23.4	8.2	19.2	10.2	30.6	10.2	2	cosl
50-61	B22	6.9	22.1	12.0	22.9	8.0	17.5	10.6	29.8	8.9	1	cosl
61-73	B23	4.5	26.1	11.2	22.5	8.9	16.7	10.1	30.4	9.1	1	cosl
73-91+	B3	4.4	19.1	9.7	24.7	13.5	21.6	7.0	40.8	10.2	4	sl
pH 8C1a	8C1c	ORGANIC MATTER 6A1a	6B1a	EST%	ELECTRICAL CONDUC-		Free Fe O	MOISTURE TENSIONS 4B2				

CROUCH SANDY LOAM
S61Calif-37-7

Description: George Borst
Date of Sampling: January 25, 1961
Collectors: Reuben Nelson, Lothair Grant, and George Borst
Location: Approximately 900 feet northeast of south 1/4 corner, sec. 19, T. 12 S., R. 4 E., about 2 miles north of Julian, San Diego County, Calif.
Elevation: 4,100 feet Relief: Steep, 35 percent. Use: Woodland - range
Vegetation: Open forest, with scattered oak and manzanita
Drainage: Well drained, moderately slow subsoil permeability
Parent Material: Deeply weathered granitic rock
Climate: This soil occurs under an average annual precipitation of about 38". The mean annual temperature is 53°, the mean January temperature is 39°, and the mean July temperature 70°.
Remarks: The A horizon of this soil is relatively thick, and erosion relatively slight, as compared with most soils in this series in San Diego County.
Classification: **Ultic Haploxeroll, coarse-loamy, mixed, mesic.**

HORIZON	DESCRIPTION
	friable; nonsticky and nonplastic; abundant very fine vertical roots; many very fine interstitial pores; highly micaceous; slightly acid (pH 6.5); abrupt smooth boundary.
A12 4-19"	Dark brown (10YR 4/3) sandy loam, very dark grayish brown (10YR 3/2) when moist; massive; slightly hard, friable, slightly sticky and

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Ultic Haploxeroll, coarse-loamy, mixed, mesic
SOIL Stoner loam (taxadjunct) SOIL Nos. S66Calif-47-9 LOCATION Siskiyou County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66393 - 66400

Depth (in.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand Very coarse (2-1)	Sand Coarse (1-0.5)	Sand Medium (0.5-0.25)	Sand Fine (0.25-0.1)	Sand Very fine (0.1-0.05)	Silt 0.05-0.02	Silt Int. III (0.02-0.002)	Silt Int. II (0.2-0.02)	Silt (2-0.1)		3A1a Non-carbonate clay	1/ Coarse fragments 1A2a > 2 Pct.	2-19	19-76
		Pct. of < 2 mm														Pct.	Pct. of < 76mm	
0-3	A11	64.5	27.2	3.3	22.0	17.6	7.1	10.9	6.9	10.4	16.8	23.2	57.6		8	51	48	3
3-6	A12	62.4	28.6	9.0	18.9	15.9	7.3	12.1	8.2	11.1	17.5	26.0	54.2		9	50	47	3
6-11	A3	64.2	26.8	9.0	20.1	16.4	7.8	11.9	8.0	9.9	16.9	24.5	56.2		9	47	45	2
11-24	B1t	63.2	27.4	9.4	19.7	16.5	7.3	11.7	8.0	10.1	17.3	24.6	55.2		9	41	39	2
24-40	B21t	65.0	23.9	11.1	21.8	17.7	7.1	10.8	7.6	9.2	14.7	22.3	57.4		11	36	35	1
40-46	B22t	64.4	24.0	11.6	18.5	17.9	8.0	12.3	7.7	9.5	14.5	23.7	56.7		12	38	37	1
46-58+	B23t	65.1	24.1	10.8	19.6	17.0	7.9	12.9	7.7	9.8	14.3	24.3	57.4		11	32	34	8
0-6	A1 2/															43	42	1

Depth (in.)	6A1a Organic carbon Pct	6B1a Nitrogen Pct	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 C'	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1		
0-3	2.77	0.100	28	2.4									6.5			6.3		
3-6	1.93	0.078	25	2.3			1.41	1.42		0.65		10.8	6.4	0.004	0.003	6.4		
6-11	0.84	0.042	20	2.5			1.52	1.54		0.66		11.0	5.8	0.004	0.003	6.2		
11-24	0.20			2.4			1.76	1.79		0.68		12.2	6.1	0.007	0.005	6.2		
24-40	0.17			2.4			1.71	1.75		0.73		13.0	6.6	0.009	0.006	6.1		
40-46	0.11			2.5			1.76	1.80		0.71		12.6	6.9	0.009	0.006	6.1		
46-58+	0.13			2.6			1.73	1.78		0.68		11.5	6.9	0.010	0.006	6.2		
0-6	1.33	0.063	21	2.3									6.1			6.5		

Depth (in.)	Extractable bases 5B1a 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A1a NH_4OAc	5A3a Sum		6G1d KCl Ext. Al^{+++}						Base saturation 5C1 NH_4OAc	5C3 Sum
	meq/100 g															Pct.	Pct.
0-3	11.8	2.0	0.1	0.8	14.7	7.5	19.5	22.2								75	66
3-6	9.9	2.1	0.1	0.6	12.7	6.1	16.3	18.8								78	68
6-11	7.1	2.3	0.1	0.5	10.0	4.1	13.2	14.1								76	71
11-24	5.7	3.1	0.1	0.4	9.3	3.7	12.0	13.0								78	72
24-40	6.9	4.7	0.1	0.4	12.1	3.4	14.9	15.5								81	78
40-46	9.4	6.0	0.1	0.3	15.8	4.2	18.6	20.0								85	79
46-58+	10.7	6.9	0.1	0.2	17.9	3.6	20.2	21.5								89	83
0-6	8.9	2.2	0.1	0.7	11.9	4.5	13.3	16.4								89	73

Depth (in.)	Clay mineralogy (<0.002 mm) 7A Mont.	Verm.	Illite	Kaolin	Hydrous Mica
	X-ray				
0-3					
3-6					
6-11			xxx		
11-24					
24-40	t	x	xxx	xx	x
40-46					
46-58+		xx	xx	x	x
0-6					

1/ From characterization sample (5 kg).

2/ Composite of 20 subsamples taken in a 15x12' grid adjacent to the master pedon.

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

STONER GRAVELLY SANDY LOAM (taxadjunct) 1/
S66Calif-47-9

Location: Siskiyou County, California. Pit located in the upper reaches of McConnaughy Gulch in Scott Valley and on the Denny Ranch. Follow the McConnaughy Gulch Road from the east road around Scott Valley to the locked gate. Take Bee Hive Gulch Road for about $\frac{1}{4}$ mile. Site is south of road about 100 yards and located 1100 feet east and 1500 feet north of SW corner of Sec. 27. T42N. R8W. Date of Sampling: May 25. 1966.

Classification: Ultic Haploxeroll, coarse-loamy, mixed, mesic.

Vegetation: Dominant vegetation is ponderosa pine and white oak. Very little grass or shrub understory. Well stocked pine stand which is growing moderately slow. Area had been logged and possibly burned. Tree canopy is about 55 percent. Climate: Average annual precipitation is approximately 22 inches, with some in the form of snow. Most precipitation comes in winter and early spring months. Summers are warm and dry. Average January temperature is 31°F., the average July temperature 70°F., and the mean annual temperature 50°F. Parent Material: The soil has formed in gravelly alluvium from mixed rock sources. The parent rock is mainly metamorphic including ultramafic, greenstone, schists, etc. Parent alluvium is often somewhat stratified at depths greater than 36 inches. Parent alluvium was deposited in fan, bottom and terrace positions along drainages from the mountains. The soil contains cobbles, stones and gravel, but most coarse fragments are fine to medium gravels;20 to 60 percent of the soil volume may be gravels. The amount of material > 2mm varies widely within short distances. Topography: Site is located on a single convex slope with a northwest aspect and a 13 percent gradient. Physiographic position is a terrace. Elevation: 3520 feet. Drainage: Well drained, moderate permeability, medium runoff. Soil Moisture: About 10 percent of field capacity throughout the profile.

Remarks: Reactions determined by Truog soil reaction test. Paired profile is S66Calif-47-10. This profile is a representative of the upper limits of the precipitation zone in which these soils occur in this soil survey area. It has a mollic epipedon and a cambic horizon. There are soils in the area with thicker A horizons. Most of the Stoner soils are cultivated. Surface soil has been lost by erosion. This profile and the paired profile have not been disturbed as far as could be ascertained.

HORIZON	DESCRIPTION
O1 & O2	$\frac{1}{2}$ to 0 inch, new and partially decomposed organic debris. Abrupt smooth boundary.
A11 RSL 66393	0 to 3 inches, dark grayish brown (10YR 4/2) gravelly sandy loam, very dark grayish brown (10YR 3/2) moist; weak medium to fine platy breaking to moderate fine granular structure; slightly hard, very friable, slightly sticky and slightly plastic; many fine and very fine roots; many random interstitial and few fine tubular pores; medium acid (pH 6.0); 40 percent gravels (included in sample); abrupt smooth boundary.
A12 RSL 66394	3 to 6 inches, brown (10YR 5/3) gravelly sandy loam, dark brown (10YR 3/3)moist;moderate medium subangular blocky structure; slightly hard, friable, slightly sticky and slightly plastic; many fine and medium roots; few fine and medium tubular pores; few thin clay films in pores and on ped faces; medium acid (pH 6.0); 40 percent fine and medium gravels and these were included in the sample; clear smooth boundary.
A3	6 to 11 inches, brown (10YR 5/3) gravelly sandy loam, dark brown (10YR 3/3) moist;

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE San Andreas loam (taxadjunct) **LOCATION** Amador County, California

SOIL FAMILY: Ultic Haploxeroll, coarse-loamy, mixed, thermic.

SOIL NOS. S59Calif-3-17 **LAB. NOS.** 59340 - 59344

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1	2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2 1/	
0-9	A_1	7.8	13.5	6.0	12.5	10.8	39.4	10.0	38.8	19.1	2	1-fsl
9-16	A_3	8.7	13.6	5.8	12.2	11.0	38.7	10.0	36.0	21.3	2	1-fsl
16-27	B_2	8.0	12.0	5.7	12.3	11.4	39.1	11.5	36.9	21.3	7	1

SAN ANDREAS LOAM (taxadjunct) 1/
S59Calif-3-17

Collectors: Grant Kennedy, Reuben Nelson, Bob Sketchley, and John Rogers.
Date of Sampling: June 19, 1959. Description by: Grant Kennedy.
Location: On H. Farnham Ranch, 1/2 mile from the Junction of Votaw and Shenandoah Roads, on Votaw Road, in the SE1/4 of the SE1/4 of Sec. 31, T8N,R11E., Amador County, California.
Elevation: 1,450 feet.
Relief: Near the crest of a hill in the gently rolling uplands on a slightly convex slope of 5 percent to the NW.
Drainage: Well drained, slow runoff, moderate permeability.
Parent Material: Deeply weathered granodiorite.
Biological activity: Earthworms and burrowing rodents are common in these soils.
Classification: Ultic Haploxeroll, coarse-loamy, mixed, thermic.
Climate: Mean annual rainfall 32".
[illegible]

filaree.
Remarks: The soil is a weakly developed Brunizem. A few rock outcrops occur in the area. Oriented ring samples were obtained in all horizons and Bureau of Public Roads samples in the A_1, B_2, and D_r horizons.

HORIZON		DESCRIPTION
A_1	0-9"	Brown (10YR 5/3) loam near fine sandy loam, dark brown (10YR 3/3) when moist; massive; slightly hard when dry, friable when moist, nonsticky and slightly plastic when wet; abundant very fine roots; many very fine and fine pores, few medium and coarse; medium acid, pH 5.8; clear, smooth lower boundary.
A_3	9-16"	Brown (10YR 5/3) loam near fine sandy loam, brown (10YR 4/3) when moist; massive; slightly hard when dry, friable when moist, slightly sticky and slightly plastic when wet; abundant very fine, common fine, occasional medium and coarse roots; many very fine and fine, few medium and coarse pores; medium acid, pH 5.7; clear, smooth lower boundary.
B_2	16-27"	Brown (7.5YR 5/4) loam, brown (7.5YR 4/4) when moist; massive; hard when dry, friable when moist, slightly sticky and plastic when wet; abundant very fine, few fine, occasional medium and coarse roots; many very fine and fine, few medium and coarse pores; few thin, discontinuous clay films, colloid mainly in bridges between sand grains; medium acid, pH 5.8; clear, wavy lower boundary.
B_3	27-32"	Mixed light brown, strong brown and pink (7.5YR 6/4, 4/6, and 7/4) loam, reddish brown and brown (5YR 4/4 and 7.5YR 5/4) when moist; massive with granitic appearance in place; hard when dry

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE San Andreas loam (taxadjunct) **LOCATION** Amador County, California

SOIL FAMILY: Ultic Haploxeroll, coarse-loamy, mixed, thermic.

SOIL NOS. S59Calif-3-18 **LAB. NOS.** 59345 - 59348

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0-7	A_p	9.2	11.5	5.5	12.6	11.7	40.3	9.2	40.2	19.4	9	1-fsl
7-16	A_3	9.0	11.9	5.5	12.0	11.6	40.4	9.6	39.9	19.7	15	1-fsl
16-28	B_2	7.8	11.2	5.1	11.9	12.3	41.9	9.8	41.2	20.5	19	1
28-37+	D_r	Weathered granodiorite										

pH 8C1a H_2O 1:1	pH 8C1c KCl 1:1	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	"Free" Fe_2O_3 6C1b %	ELECTRICAL CONDUCTIVITY EC x 10³ MILLIMHOS PER CM @25°C.	$CaCO_3$ equivalent %	Bulk Density 4A3a g/cc	MOISTURE TENSIONS 4B1a 1/10 ATMOS. %	4B1a 1/3 ATMOS. %	4B2 15 ATMOS. %
6.0	4.9	1.15	0.096	12.0	1.6			1.37	25.7	18.3	4.8
5.9	4.7	0.51	0.053	9.6	1.9			1.55			5.1
6.0	4.6	0.35	0.040	8.8	1.9			1.60	24.3	16.8	4.7
								1.80			

5A2a CATION EXCHANGE CAPACITY (-(Na)	EXTRACTABLE CATIONS 5B1a Ca 6N2b	5B1a Mg 6O2c	6H2a H	Na 6P2a	K 6Q2a	BASE SAT. % 5C3	SATURATION EXTRACT SOLUBLE Na	K				MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
11.5	6.3	1.1	3.7	<0.1	0.3	68						
11.4	6.3	1.3	3.7	<0.1	0.2	68						
11.2	7.0	1.2	2.4	<0.1	0.2	78						

1/ Percent of total soil

SAN ANDREAS LOAM (taxadjunct) 1/
S59Calif-3-18

Collectors: Grant Kennedy, Reuben Nelson, Bob Sketchley, John Rogers.
Date of Sampling: June 20, 1959. Description by: Grant Kennedy.
Location: Amador County, California, on the Fererro Ranch, 1/2 mile N. of the junction of Shenandoah Road and Shenandoah School Road, in NE1/4 of SW1/4 of Sec. 36, T8N, R10E.
Elevation: 1,240 feet.
Relief: Gently rolling uplands on a plain slope of 5 percent to the northeast.
Drainage: Well drained, slow runoff, moderate permeability.
Parent Material: Quartz diorite or granodiorite.
Biological activity: Worms and burrowing rodents are present in these soils.
Climate: Mean annual rainfall, 32".
Vegetation: Soft chess, ripgut, filaree, brodela, Digger pine, blue oak, live oak.
Remarks: Minimal Brunizem; a few rock outcrops are present. Oriented ring samples were obtained for all horizons and BPR samples in all except the A_3 horizon.
Classification: Ultic Haploxeroll, coarse-loamy, mixed, thermic.

HORIZON		DESCRIPTION
A_p	0-7"	Brown (10YR 5/3) loam near fine sandy loam, dark brown (10YR 3/3) when moist; massive; slightly hard when dry, friable when moist, nonsticky and nonplastic when wet; abundant very fine roots in upper 2 inches, common very fine and fine roots below; many very fine, few fine pores; medium acid, pH 6.0; clear, smooth lower boundary.
A_3	7-16"	Yellowish brown (10YR 5/5) loam near fine sandy loam, dark brown (10YR 3/3) when moist; massive; slightly hard when dry, friable when moist, slightly sticky and slightly plastic when wet; common very fine and fine roots; many very fine, few fine pores; few thin discontinuous clay films in pores; medium acid, pH 5.9; gradual smooth lower boundary.
B_2	16-28"	Brown (10YR 5/3) gravelly loam, dark brown (7.5YR 4/4) when moist; massive; very hard when dry, friable when moist, slightly sticky and slightly plastic when wet; common very fine, few fine roots; many very fine, few fine pores; common thin discontinuous clay films in pores; colloid mainly in bridges between sand grains; medium acid, pH 6.0; abrupt, irregular lower boundary.
D_r	28-37"+	Mixed strong brown, very pale brown, or white decomposing granodiorite with a few very fine roots and moderately thick clay films along fracture planes and an occasional quartz dike up to 6 inches thick. The weathered rock is easily displaced with a spade but the fragments are very hard when dry.

1/ The San Andreas series is in a coarse-loamy, mixed, thermic family of Typic Haploxerolls. This pedon is a taxadjunct because the base saturation of the argillic horizon is less than 75 percent.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Vertic Haploxeroll, fine, montmorillonitic, thermic
SOIL Zaca silty clay loam SOIL Nos. S65Calif-27-5 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65475 - 65481

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	1/ Coarse fragments 1A2a > 2 Pct.	2-19	19-76
		Pct. of < 2 mm															Pct. of < 76mm	
0-4	A11	27.0	30.9	42.1	2.4	4.0	4.2	10.3	6.1	11.3	19.6	22.9	20.9	0	42			
4-16	A12	27.0	29.4	43.6	2.0	3.8	3.9	10.0	7.3	10.2	19.2	23.4	19.7	1	43			
16-23	A13	26.2	30.0	43.8	1.9	3.7	3.9	9.5	7.2	11.2	18.8	24.1	19.0	2	42			
23-40	A14ca	26.3	30.8	42.9	1.9	3.9	4.0	9.5	7.0	11.5	19.3	24.4	19.3	5	38			
40-46	C1ca	27.4	37.5	35.1	1.7	3.1	3.1	9.6	9.9	13.9	23.6	30.1	17.5	4	31			
46-58	C2	39.0	35.9	25.1	2.7	4.9	6.0	15.3	10.1	14.5	21.4	33.5	28.9	1	24			
58-70+	C3	7.6	28.9	63.5	0.3	1.0	0.7	2.2	3.4	5.6	23.3	10.5	4.2	0	64			

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.	Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEP in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1	
0-4	2.73	0.229	12		4						18.5			7.9	
4-16	1.05	0.100	11		6						18.4			8.2	
16-23	0.77				8						18.5			8.3	
23-40	0.47				13						18.0			8.2	
40-46	0.22				13						15.8			8.2	
46-58	0.09				9						12.7			8.5	
58-70+	0.16				1						25.2			8.2	

Depth (In.)	Extractable bases 5B1a 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A1a NH_4OAc	5A3a Sum	6G1d KCl Ext. Al^{+++}	Base saturation 5C1 NH_4OAc Pct.	Pct.
	meq/100 g										
0-4	41.0	5.6	0.1	1.8	48.5		38.8			100+	
4-16	40.1	5.2	0.3	1.0	46.6		34.6			100+	
16-23	39.2	6.8	0.2	0.8	47.0		35.3			100+	
23-40	36.1	6.9	0.3	0.6	43.9		32.2			100+	
40-46	30.0	9.6	0.5	0.5	40.6		27.8			100+	
46-58	23.7	10.2	0.9	0.4	35.2		23.1			100+	
58-70+											

Depth (In.)	Clay mineralogy (<0.002 mm) 7A Mont.	Verm.	Mica	KaoHn.
	X-ray			
0-4	xxxx			x
4-16				
16-23				
23-40				
40-46				
46-58				
58-70+	xxxx			x

1/ Apparent coarse fragments dispersed readily. Hence the characterization sample was ground whole and all chemical and physical data are on a whole soil basis.

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

ZACA SILTY CLAY LOAM
S65Calif-27-5

Location: Monterey County, California. About 11 miles NNE of King City or 6 miles NE of main ranch headquarters, Montgomery Ranch, about 1.4 mile (air) N. from Pinalito VABM, 1400 feet NE of a reservoir. Date of Sampling: May 18, 1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton, B. R. Brasher, C. J. Weisel, R. D. Waggoner, F. J. Pecarich. Site selection by T. D. Cook.
Classification: Vertic Haploxeroll, fine, montmorillonitic, thermic. (Regosolic Chestnut).
Vegetation: Range, annual bromes, bur clover, wild oats, foxtail, fiddleneck and little stipa. Climate: Precipitation 14 inches; mean annual temperature 60°F., January mean 50°F., July mean 67°F. Parent Material: Soft calcareous sandstone and shale, Plio-Pleistocene non-marine. Topography: Round, rolling hills locally slightly convex, 13 percent North-facing slope near crest, upper 1/3. Elevation: 1300 feet. Drainage: Well drained, permeability moderately slow, runoff medium. Soil Moisture: Slightly moist, about wilting point throughout.
Remarks: Field pH by Truog Kit. Paired sample is Linne silty clay loam, S65Calif-27-4-1 through 6. Surface is somewhat compacted to a depth of 4 inches by cattle trampling.

[redacted]

HORIZON	DESCRIPTION
A11 RSL 65475	0 to 4 inches, dark gray (10YR 4/1) heavy silty clay loam, very dark gray (10YR 3/1) moist; moderate medium and coarse subangular blocky structure; friable, very sticky and plastic; many plentiful fine and very fine roots; few very fine interstitial and few very fine tubular pores; moderately alkaline (pH 8.0), strongly effervescent with disseminated lime; gradual smooth boundary.
A12 RSL 65476	4 to 16 inches, dark gray (10YR 4/1) heavy silty clay loam, very dark gray (10YR 3/1) moist; moderate medium subangular blocky structure; very hard, friable, very sticky and plastic; plentiful very fine roots; common very fine interstitial and common very fine tubular pores; moderately alkaline (pH 8.0),

OIL SURVEY LABORATORY Riverside, California

SOIL TYPE Waukena loam variant **LOCATION** Merced County, Calif.

SOIL FAMILY: Typic Natrixeroll, fine-loamy, mixed, thermic

SOIL NOS. S55Calif. 24-3 **LAB. NOS.** 55289-296

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1 VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	2A2 > 2 1/	TEXTURAL CLASS
0- 6	A_1	0.3	0.9	1.4	10.2	19.7	48.5	19.0	52.0	24.5	0	l
6-14	B_1	0.2	0.9	1.3	10.2	20.1	44.8	22.5	50.7	22.7	0	l
14-23	B_{21}	0.2	0.8	1.1	8.8	21.9	38.1	29.1	48.3	19.0	0	cl
23-31	B_{22}	0.2	0.7	1.2	8.2	19.6	37.4	32.7	45.1	18.7	0	cl
31-38	B_{31ca}	0.3	0.9	0.9	7.4	18.4	38.3	33.8	43.5	19.5	0	cl
38-43	B_{32ca}	2.3	1.8	1.0	7.6	16.6	41.6	29.1	39.9	24.7	0	cl
43-57	B_{33ca}	3.6	3.9	2.1	8.5	20.2	40.9	20.8	48.1	19.9	2	l
57-68	C	0.8	0.6	0.4	7.8	31.0	45.0	14.4	60.0	23.3	4	l

8C1b pH SATURATED PASTE 1:1	8C1a pH 1:5	8C1a pH 1:10	6A1a ORGANIC MATTER ORGANIC CARBON %	6B1a NITROGEN %	C/N	4A3a Bulk Density g./cc.	8A1a Elec. Conduc. EC×10³ MILLIMHOS PER CM @25°C	6E1b $CaCO_3$ equivalent %	6F1a GYPSUM me./100g. SOIL	4B2 MOISTURE TENSIONS 1/10 ATMOS. %	4B2 1/3 ATMOS. %	4B2 15 ATMOS. %
7.9		8.6	1.95	0.192	10.2	1.37	1.0	2	0			10.8
8.2		9.1	1.22	0.112	10.9	1.33	3.2	2	0			12.5
8.6		9.8	0.47	0.049	9.6	1.45	6.9	6	0			17.8
8.9		9.9	0.30	0.033	9.1	1.44	8.7	7	0			18.8
9.0		10.0	0.23	0.027		1.40	9.2	6	0			20.2
9.0		9.9	0.09			1.65	11.4	30	0			12.8
8.9		10.0	0.08			1.65	8.9	15	0			10.7
8.5		9.8	0.04			1.75	8.6	1	0			8.01

5A2a CATION EXCHANGE CAPACITY (+(Na)	EXTRACTABLE CATIONS 5B 5B1a Ca 6N2b	5B1a Mg 6Q2c	6H2a H	Exch. 5B1b Na 6P2a	Exch. 5B1b K 6Q2a	5D1 EXCH. Na %	SATURATION EXTRACT SOLUBLE 8A1 6P1a Na	6Q1a K	6W1a 6O1a Ca+Mg	6I1a 6J1a HCO_3+CO_3+	6K1a	8A MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						← milliequivalents per liter →					
24.6	23.4	8.8		1.0	1.4	4.1	5.0	0.1	4.7	7.6	0.8	44.3
23.8	20.5	9.3		4.6	1.3	19.3	28.7	0.5	3.3	8.8	19.7	42.1
22.3	17.4	7.4		11.0	0.9	49.3	65.0	0.1	1.7	6.4	46.3	75.3
23.4	16.2	6.9		3.2	0.7	13.7	84.0	0.2	1.2	5.9	56.6	92.6
26.7	15.0	6.6		5.8	0.7	21.7	88.0	0.2	0.9	6.6	60.5	102.3
15.8	13.5	7.5		10.5	0.3	66.5	101.0	0.04	1.6	4.4	77.8	65.4
15.8	13.7	4.7		10.6	0.3	67.1	85.0	0.1	1.4	4.5	60.0	64.0
16.3	8.4	6.7		7.1	0.2	43.6	80.0	0.1	5.9	3.9	62.9	2.0

1/ Percent of the whole soil

WAUKENA LOAM variant 1/
S55Calif-24-3

Location: Merced County, California. In San Luis SCD, southwest corner of Delta Road and Henry Miller Road, 100 feet south and 100 feet west of corner. NE1/4 NE1/4 NE1/4 sec. 4, T. 10 S., R. 11 E. Date of Sampling: April 21, 1955.
Description by: R. C. Huff. Collectors: Ralph E. Nelson, Richard C. Huff, Rudolph Ulrich.
Classification: Typic Natrixeroll, fine-loamy, mixed, thermic.
Vegetation: Probably has been cultivated in the past. Cover now consists of native annual grasses and occasional salt-tolerant species. Climate: Approximately 8½ inches rainfall with dry summers and cool, moist winters. Parent Material: Mixed alluvium, mainly granitic, which is underlain by stratified sediments. Relief: Slightly undulating upper edge of basin rim. Slope 0-2 percent. Elevation: 100 feet. Drainage: Surface drainage is slow, internal drainage is slow.
Remarks: Many roots in surface foot, few roots below 31 inches.

HORIZON		DESCRIPTION
A_1	0- 6"	Gray brown (10YR 5/2, very dark brown 2/2 moist) loam. Moderate medium subangular blocky. Hard, friable, nonsticky and nonplastic. The soil reacts slightly to HCl. The pH is above 8. This grades abruptly and smoothly into:
B_1	6-14"	Gray brown (10YR 5/2, very dark brown 2/2 moist) light clay loam. Moderate medium and fine subangular blocky. Very hard, firm, slightly plastic and nonsticky. The soil reacts slightly to HCl. The pH is above 8. This grades clearly and with a smooth lower boundary into:
B_{21}	14-23"	Light brownish gray to grayish brown (2.5Y 6/2 and 5/2, very dark grayish brown 4/2 moist, with darker (2.5Y 3/2) faces) clay. Moderate medium prismatic and moderate medium angular blocky. Very hard, very firm, plastic and slightly sticky. The soil reacts slightly with HCl. The pH is above 8. This grades clearly and with a wavy lower boundary into:
B_{22}	23-31"	Light brownish gray (2.5Y 6/2, dark grayish brown 4/2 moist) clay. Moderate fine subangular blocky. Very hard, very firm, plastic and slightly sticky. The soil reacts to HCl. There are conspicuous seams of lime throughout this horizon. The pH is above 8. Grades abruptly and with a wavy lower boundary into:
B_{31ca}	31-38"	Pale olive (5Y 6/3, olive 4/3 moist) clay. Moderate medium and fine subangular blocky. Very hard, very firm, plastic and slightly sticky. The soil effervesces with HCl. The pH is above 8. This grades clearly and smoothly into:
B_{32ca}	38-43"	Pale olive and white (5Y 6/3 and 8/2, olive and light gray 4/3 and 7/2 moist) clay. Weak fine subangular blocky. Very hard, very firm, plastic and slightly sticky. The soil effervesces with HCl. The pH is above 8. This grades abruptly and smoothly into:
B_{33ca}	43-57"	Pale olive and white (5Y 6/3 and 10YR 8/1, olive 5Y 4/3 moist) light clay. Weak medium subangular blocky. Very hard, firm, plastic and slightly sticky. The soil reacts violently to HCl and has many pinkish lime concretions, 5 to 10 mm. in diameter. The pH is above 8. This grades abruptly and smoothly into:
C	57-68"	Pale yellow (5Y 7/3, olive 5/3 moist) variable textured, stratified sediments. The texture ranges from sandy loam to clay loam. Massive. Hard, firm, slightly plastic and slightly sticky. There are many iron and manganese streaks throughout this horizon. The pH is above 8.

1/ The Waukena series is in a fine-loamy, mixed, thermic family of Typic Natrixeralfs. This pedon has a mollic epipedon.

SOIL SURVEY LABORATORY Lincoln, Nebraska

SOIL TYPE Aiken loam **LOCATION** Amador County, Calif rnia

SOIL FAMILY: Xeric Haplohumult, clayey, kaolinitic, mesic.

SOIL NOS. S59Calif-3-12 **LAB. NOS.** 11356 - 11363

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	3A1 2A2* > 2	TEXTURAL CLASS
0- 9	A1	5.6a	7.5a	5.5a	11.3a	12.0a	37.0	21.1	31.2	24.2	9	1
9-20	A3	4.9a	5.4a	4.4a	10.7a	12.4a	39.4	22.8	33.7	24.5	10	1
20-29	B11t	2.5a	3.9a	3.5a	9.1a	11.5a	40.5	29.0	31.4	26.1	10	cl
29-40	B12t	2.4a	2.9a	2.5a	6.3a	8.6a	38.5	38.8	25.7	25.3	11	cl
40-54	B21t	2.0c	3.3c	3.2c	8.0c	8.9c	35.2	39.4	23.3	25.6	7	cl
54-69	B22t	1.4b	3.5b	3.3b	7.8b	7.7b	32.0	44.3	21.1	23.2	20	c
69-92	B23t	1.1c	2.4c	2.4b	6.9b	8.0b	35.7	43.5	22.9	25.0	23	c
92-100	B3t	1.2b	2.4b	2.5b	6.9b	8.4b	36.4	42.2	22.7	26.4	29	c

pH 8C1a 1:1	8C1a 1:5	8C1a 1:10	ORGANIC MATTER 6A1a ORGANIC CARBON %	6B1a NITROGEN %	C/N	EST% SALT (BUREAU CUP)	ELECTRICAL CONDUCTIVITY EC×10³ MILLIMHOS PER CM @25°C.	6C1a Free iron as % Fe_2O_3	4A3a Bulk Density g/cc	MOISTURE TENSIONS 1/10 ATMOS. %	1/3 ATMOS. %	4B2 15 ATMOS. %
5.9	6.1	6.2	4.98	0.179	28			5.0	0.98			18.5
6.0	6.4	6.4	2.11	0.090	23			5.2	0.92			19.1
5.9	6.0	6.2	1.02	0.052	20			5.5	0.99			19.2
6.0	6.2	6.4	0.48					6.2				20.1
5.3	5.6	5.8	0.09					6.2	1.33			23.2
5.4	5.8	6.0	0.14					6.7	1.35			23.0
6.0	6.4	6.6	0.18					6.4	1.36			21.1
5.2	5.5	5.6	0.09					6.3	1.26			24.2

5A3a1/ CATION EXCHANGE CAPACITY (Sum)	EXTRACTABLE CATIONS 6N2b Ca	6Q2b Mg	6H2a H	6P2a Na	6Q2a K	BASE SAT. % 5C3	Clay mineralogy (<0.002 mm.) Mont.	Verm. X-ray	Mixed layer Mica-Verm. 7A2	Mica	% Kaolin DTA 7A3	% Gibb. DTA 7A3
	milliequivalents per 100g. soil											
34.4	6.2	1.5	25.8	<0.1	0.9	25		xx			20	17
26.0	5.8	0.8	18.6	<0.1	0.8	28						
19.4	4.5	1.0	13.2	<0.1	0.7	32		xx			22	20
16.3	3.2	1.9	10.4	<0.1	0.8	36						
9.9	0.5	0.4	8.7	0.1	0.2	12		x			45	10
10.4	1.2	0.6	7.8	0.1	0.7	25						
13.0	2.2	1.8	8.0	<0.1	1.0	38						
9.4	0.4	<0.1	8.8	<0.1	0.2	6					55	tr.

a. Many (>50 percent) Fe? nodules
b. Few (5 - 25 percent) Fe? nodules
c. Common (25 - 50 percent) nodules
* 2 mm. - 3/4" fraction
1/ In acid soils CEC (NH_4OAc) is almost always lower than either CEC (Na) or CEC (sum).
based on CEC (sum)

AIKEN LOAM
S59Calif-3-12

Location: On the Allen Ranch on the road east of Fiddletown to Highway 88, 100 yards west of the cattle chute and 100 feet south of the road in the SW corner of the NE1/4 of sec. 33, T. 8 N., R. 13 E., Amador County, Calif.
Date of Sampling: June 19, 1959 Description by: Grant Kennedy.
Collectors: Grant Kennedy, Reuben Nelson, Bob Sketchley
Vegetation: Ponderosa pine, incense cedar, sweet birch, bracken fern
Climate: Mean annual rainfall - 43 inches
Parent Material: Andesitic tuff-breccia.
Relief: Broad, gently sloping ridgetop with a plane or slightly concave slope of 8 percent, facing north
Elevation: 3,685 feet
Drainage: Well drained; slow runoff; moderate over moderately slow permeability
Remarks: This is a Reddish-Brown Lateritic soil with a good stand of timber. Samples were taken for the Bureau of Public Roads of horizons I, 6, and 8 (A1, B22t, and B3t). Soil was moist when described.
Classification: Xeric Haplohumult, clayey, kaolinitic, mesic.

HORIZON		DESCRIPTION
O1	4- 0"	Partially decomposed forest litter.
A1 LSL No. 11356	0- 9"	Dark brown (7.5YR 4/4) loam, darker brown (7.5YR 3/3) when moist; strong very fine and fine granular structure; soft, very friable; abundant very fine and fine, few medium and coarse roots; many very fine, fine, medium, few coarse pores; many very fine, few fine shot*; medium acid (pH 6.0); gradual smooth boundary.
A3 LSL No. 11357	9-20"	Reddish brown (5YR 4/4) loam, dark reddish brown (5YR 3/3) when moist; moderate very fine and fine granular structure; otherwise similar to overlying horizon except slightly plastic when wet; clear smooth boundary.
B11t LSL No. 11358	20-29"	Yellowish red (5YR 5/6) clay loam, yellowish red (5YR 3/6) when moist; massive; slightly hard, friable, slightly plastic and slightly sticky; abundant very fine, fine and medium, few coarse roots; many very fine, fine and medium pores; thin continuous clay films, mainly in pores; common very fine, few fine shot*; medium acid (pH 6.0); gradual smooth boundary.
B12t LSL No. 11359	29-40"	Yellowish red (5YR 5/6) heavy clay loam, dark red (2.5YR 3/6) when moist; massive; slightly hard, firm, plastic and slightly sticky; plentiful very fine and fine, few medium and coarse roots; many very fine, fine, and medium pores; thin continuous clay films in pores and bridges; common very fine, few fine shot*; medium acid (pH 6.0); gradual smooth boundary.
B21t LSL No. 11360	40-54"	Yellowish red (5YR 5/6) clay, dark red (2.5YR 3/6) when moist; massive; hard, firm, plastic and sticky; occasional very fine to coarse root; many very fine, few fine pores; moderately thick continuous clay films in pores and bridges; common very fine shot*; medium acid (pH 6.0); diffuse smooth boundary.
B22t LSL No. 11361	54-69"	Similar to horizon above except strongly acid (pH 5.5); clear irregular boundary.
B23t LSL No. 11362	69-92"	Light brown (7.5YR 6/4) clay, reddish brown (5YR 4/4) when moist; massive; hard, firm, plastic and slightly sticky; occasional roots; common very fine, few fine pores; thin, continuous clay films in pores and bridges; very strongly acid (pH 5.0); gradual smooth boundary.
B3t LSL No. 11363	92-100"	Light brown (7.5YR 6/4) clay loam, reddish brown (5YR 4/4) when moist; massive; hard, very firm, slightly plastic and slightly sticky; occasional root; common very fine and fine pores; few thin continuous clay films in pores and along fracture planes; very strongly acid (pH 4.8).

* Shot are magnetic when dry. Below 69 inches shot are hard to discern but about half the soil mass is attracted by a magnet.

CONSECUTIVE PEDON NUMBER C 68118 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES SITES SOIL NO. S68 CALI. 29-31 SAMPLE NO. 8 883-8 891 NEVADA COUNTY AREA

SOIL FAMILY XERIC HAPLOHUMULT CLAYEY,KAOLINITIC,MESIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL SAND 2-.05	TOTAL SILT .05-.002	TOTAL CLAY LT .002	VCS 2-1	CS 1-.5	MS .5-.25	FS .25-.10	VFS .1-.05	CSI .05-.02	FSI .02-.002	CO. 2-.2U	FINE LT .2U	CARBO NATE 3A1	INT. II .2-.02	COARSE FRAGMENTS PCT. OF WHOLE SOIL 1A2A	I 75-20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75			
8 883	0- 18	A11	22.0	45.2	32.8	2.7	5.3	3.0	5.9	5.1	14.0	31.2				22.5	--A	--	--	--
8 884	18- 53	A12	20.3	45.1	34.6	1.8	4.1	3.1	5.9	5.4	14.6	30.5				23.5	6A	--	4	2
8 885	53- 79	A3	20.8	43.4	35.8	1.5	3.7	3.0	6.0	6.6	13.2	30.2				23.2	6A	--	6Z	N
8 886	79-100	B21T	18.8	38.5	42.7	0.9	3.0	2.9	5.8	6.2	11.6	26.9				21.0	3A	--	3Z	N
8 887	100-130	B22T	19.7	34.9	45.4	1.5	3.1	2.8	6.2	6.1	11.7	23.2				21.6	TRA	--	TR	TR
8 888	130-165	B23T	24.1	32.8	43.1	1.3	3.8	3.6	7.9	7.5	10.8	22.0				23.2	2A	--	2Z	N
8 889	165-190	B31	27.3	31.4	41.3	1.0	3.9	4.4	9.8	8.2	10.6	20.8				24.6	--A	--	--	--
8 890	190-215	B32	34.0	29.9	36.1	1.9	6.8	5.7	10.9	8.7	10.2	19.7				25.4	--A	--	--	--
8 891	0- 18	A11															3A	--	3Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR-BON 6A1A	NITRO-GEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	2-20 6E1B	SESQUIOXIDES DI-CIT FE 6C2B	DI-CIT AL 6G7A	EXT PYROP FE 6C5A	EXT PYROP AL 6G5A	ATTERBERG L L 4F2	P L 4F1	I P 4A1F	T EST. 1/3 BAR 4A1F	BULK DENSITY 1/3 BAR 4A1H	DRY 3B2	CM	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1	LE 4D1
		PERCENT		PERCENT		PERCENT				PCT NO 40 GRAMS PER CC							PERCENT			PERCENT	
8 883	0- 18	6.00	0.249			9.2	1.4			63	47		0.90	0.94	1.00		33.7	23.5	1.5	1.5
8 884	18- 53	1.72	0.085			9.8	1.2					0.95			0.98			21.1		
8 885	53- 79	1.07				9.9	1.1			46	33	1.00			0.98			20.9		
8 886	79-100	0.51				9.9	1.0					1.00			0.99			20.3		
8 887	100-130	0.38				10.5	0.9			56	20	1.10			1.00			21.3		
8 888	130-165	0.30				12.6	1.1			59	47		1.17	1.24	0.99		41.7	24.3	2.0	1.9
8 889	165-190	0.22				12.6	1.1					1.17			1.00			26.2		
8 890	190-215					12.4	1.2			59	49		1.17	1.22	1.00		40.9	24.8	1.4	1.4
8 891	0- 18	7.44	0.284									0.90			0.99			23.9		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACI-DITY 6H2A	KCL EXT. AL 6G1E	C E C SUM 5A3A	C E C NH4-OAC 5A6A	BASE SUM+ ACI-DITY	SATURATION SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 883	0- 18	9.9	2.7	0.1	1.0	13.7	22.1		35.8	31.8	38		43		5.9	5.3
8 884	18- 53	3.4	1.5	0.1	1.0	6.0	14.0		20.0	17.4	30		34		5.9	5.1
8 885	53- 79	2.8	1.8	0.1	0.6	5.3			5.3	17.7	100+		30		5.7	5.1
8 886	79-100	2.1	2.3	TR	0.6	5.0		0.3	5.0	13.3	100+	94	38		5.5	4.9
8 887	100-130	2.8	2.0	0.1	0.5	5.4		0.3	5.4	11.2	100+	95	48		5.4	4.9
8 888	130-165	2.9	1.8	0.1	0.4	5.2		0.2	5.2	13.0	100+	96	40		5.2	4.7
8 889	165-190	3.1	2.1	0.1	0.4	5.7		0.2	5.7	12.4	100+	97	46		5.2	4.6
8 890	190-215	2.9	2.2	TR	0.4	5.5		0.7	5.5	18.9	100+	89	29		5.1	4.5
8 891	0- 18														5.9	5.3

C L A Y M I N E R A L O G Y

SAMPLE NO	DEPTH CM.	LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	L.T. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III
8 883	0- 18	KK 3	VR 1	MI 1						
8 884	18- 53									
8 885	53- 79	KK 3	VR 1							
8 886	79-100									
8 887	100-130									
8 888	130-165	KK 3								
8 889	165-190									
8 890	190-215	KK 3								
8 891	0- 18									

MINERAL CODE
KK KAOLINIT
VR VERMICLT
MI MICA

RELATIVE AMOUNT
1 TRACE
2 SMALL
3 MODERATE
4 ABUNDANT
5 DOMINANT
6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.

SITES LOAM
S68Calif-29-31

Location: Nevada County, California; 600 feet east of center of sec. 4, T. 15 N., R. 9 E., 150 feet west of Colfax Highway, 40 feet south of Dalmatian Way, 5½ miles southeast of Grass Valley, California. Date of Sampling: September 5, 1968.
Description by: G. Staidl and L. A. Brittan. Collectors: L. A. Brittan, G. Staidl, T. M. Ryan, C. B. Goudey, G. M. Kennedy.
Classification: Xeric Haplohumult, clayey, kaolinitic, mesic.
Vegetation: Not cultivated. Mixed coniferous forest, ponderosa pine, sugar pine, Douglas fir, incense cedar, black oak, bracken fern, bear clover. Climate: Average annual precipitation is 55 inches. Mean annual temperature is about 56°F. Parent Material: Metabasic rock and diabase. Topography: Mountainous upland, moderately steep, convex slope 25 percent, NNE aspect near toe of slope. Elevation: 2,640 feet. Drainage: Well drained; runoff is medium; permeability is moderate over slow. Soil Moisture: Essentially dry to 31 inches. Very slightly moist 31 to 85 inches.
Remarks: Erosion at this site appeared to be slight. pH taken by Hellige method. Companion profile is S68Calif-29-32. The B21t horizon contains 3 percent by weight fragments 3 to 4 inches in diameter. The B22t horizon contains 25 percent by weight fragments 3 to 7 inches in diameter. The A3, B21t, B22t and B32 horizons were sampled for California Division of Highways.

HORIZON	DESCRIPTION
O1	4 to 2½ inches, litter, twigs, leaves.
O2	2½ to 0 inches, partially decomposed organic matter.
	(0-18 cm.)
A11 RSL 68883	0 to 7 inches, reddish brown (5YR 4/3) loam, dark reddish brown (5YR 3/3) moist; strong fine subangular blocky parting to strong fine and medium crumb structure; slightly hard, friable, slightly sticky, plastic; many very fine and fine random and many medium and coarse horizontal and oblique roots; many very fine interstitial pores; slightly acid (pH 6.3); clear smooth boundary.
	(18-53 cm.)
A12 RSL 68884	7 to 21 inches, yellowish red (5YR 4/6) heavy loam, dark red (2.5YR 3/6) moist; moderate fine and medium subangular blocky parting to strong fine and medium crumb structure; slightly hard, friable, slightly sticky, plastic; many very fine and fine random and many medium and coarse horizontal and oblique roots; many very fine interstitial pores; medium acid (pH 6.0); clear smooth boundary.
	(53-79 cm.)
A3 RSL 68885	21 to 31 inches, yellowish red (5YR 4/6) clay loam, dark red (2.5YR 3/6) moist; moderate medium subangular blocky parting to strong fine crumb structure; slightly hard, friable, sticky, plastic; many very fine and fine random and many medium and coarse horizontal and oblique roots; many very fine interstitial pores; medium acid (pH 6.0); clear wavy boundary.
	(79-100 cm.)
B21t RSL 68886	31 to 40 inches, red (2.5YR 4/6) clay loam, dark red (2.5YR 3/6) moist; moderate medium subangular blocky structure; slightly hard, friable, sticky, plastic; common very fine and fine and few medium and coarse oblique roots; common very fine tubular and interstitial pores; few thin clay films on faces of peds and lining pores; 5 percent by volume rock fragments 1 to 4 inches in diameter; medium acid (pH 5.8); clear smooth boundary.
	(100-130 cm.)
B22t RSL 68887	40 to 52 inches, red (2.5YR 4/6) clay, dark red (2.5YR 3/6) moist; moderate medium and coarse subangular blocky structure; hard, firm, sticky, plastic; common very fine and fine, few medium and coarse oblique roots; common very fine tubular pores; many thin clay films on faces of peds and many moderately thick clay films line pores; 2 to 5 percent by volume rock fragments 1 to 4 inches in diameter; strongly acid (pH 5.5); clear wavy boundary.
	(130-165 cm.)
B23t RSL 68888	52 to 64 inches, red (2.5YR 5/6) clay, red (2.5YR 4/6) moist; moderate medium and coarse subangular blocky structure; hard, firm, sticky, plastic; few very fine random and very few medium vertical roots; common very fine tubular pores; many thin clay films on faces of peds and many moderately thick clay films line pores; 10 to 15 percent by volume rock fragments 2 to 7 inches in diameter; strongly acid (pH 5.5); gradual smooth boundary.
	(165-190 cm.)
B31 RSL 68889	64 to 75 inches, red (2.5YR 5/6) clay, red (2.5YR 4/6) moist; weak coarse subangular blocky structure; slightly hard, friable, slightly sticky, plastic; few very fine random roots; many very fine tubular pores; many thin clay films, red (2.5YR 4/8 moist) on faces of peds and many moderately thick clay films line pores; strongly acid (pH 5.5); gradual wavy boundary.
	(190-215 cm.)
B32 RSL 68890	75 to 85 inches, yellowish red (5YR 5/8) clay loam, red (2.5YR 4/6) moist; massive; slightly hard, friable, slightly sticky, plastic; very few very fine random roots; many very fine tubular pores; many thin red (2.5YR 4/8 dry and moist) clay films line pores; strongly acid (pH 5.5).

CONSECUTIVE PEDON NUMBER C 68119 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES SITES SOIL NO. S68 CALI. 29-32 SAMPLE NO. 8 892-8 901 NEVADA COUNTY AREA

SOIL FAMILY XERIC HAPLOHUMULT CLAYEY,KAOLINITIC,MESIC RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	TOTAL FRACTION SAND 2-.05	SILT .05-.002	CLAY LT .002	SANDS VCS 2-1	CS 1-.5	MS .5-.25	FS .25-.10	VFS .1-.05	SILTS CSI .05-.02	FSI .02-.002	CLAYS CO. 2-.2U	FINE LT .2U	CARBONATE 3A1A	INT. II .2-.02	COARSE FRAGMENTS PCT. OF WHOLE SOIL I 75-20	II 20-5	III 5-2
			PERCENT OF LESS THAN 2MM														PCT. OF LT 75		
8 892	0- 18	A1	23.2	39.4	37.4	1.6	4.2	3.3	7.4	6.7	13.7	25.7	30.2	7.2		24.9	13A --	13Z	N
8 893	18- 36	B1	19.3	35.3	45.4	0.9	2.8	2.9	6.6	6.1	11.4	23.9				21.5	5A --	5Z	N
8 894	36- 46	B21T	17.8	31.1	51.1	1.3	2.8	2.6	5.9	5.2	9.6	21.5				18.4	5A --	5Z	N
8 895	46- 84	B22T	18.6	30.8	50.6	0.9	2.8	2.8	6.2	5.9	10.1	20.7	36.3	14.3		19.8	6* --	1Z	N
8 896	84-120	B31	22.1	28.2	49.7	0.5	2.7	3.9	8.6	6.4	9.0	19.2				20.3	7* --	--	--
8 897	120-155	B32	26.8	32.8	40.4	0.6	3.4	4.5	10.2	8.1	10.2	22.6				24.0	6* --	--	--
8 898	155-185	C 1	26.7	34.7	38.6	0.4	2.4	4.3	11.3	8.3	10.7	24.0				25.4	--A --	--	--
8 899	185-205	C 2	32.7	33.2	34.1	1.3	6.0	6.5	11.6	7.3	11.6	21.6				25.3	--A --	--	--
8 900	205-235	C 3	36.0	33.2	30.8	1.6	7.6	7.6	12.6	6.6	10.3	22.9				23.4	--A --	--	--
8 901	0- 18	A1															3A --	3Z	N

SAMPLE NO.	DEPTH CM.	ORG.MATTER CARBON 6A1A -PERCENT-	NITROGEN 681A -PERCENT-	CARBONATE AS CACO3 6E1B LT 2	6E1B 2-20	SESQUIOXIDES DI-CIT FE 6C2B	EXT AL 6G7A	PYROP EXT FE 6C5A	AL 6G5A	ATTERBERG LIMIT LL 4F2 PCT.NO.40	PL 4F1	BULK DENSITY EST. 1/3 BAR 4A1F GRAMS PER CC	1/3 BAR 4A1H	DRY 3B2	CM	WATER CONTENT 1/10 BAR 4B1C -PERCENT-	1/3 BAR 4B1C	15 BAR 4B2	EXTNSBLTY LEF 4D1 -PERCENT-	LE 4D1
8 892	0- 18	6.69	0.260			8.3	1.5	0.4	0.7	60	44	0.94	0.97	0.95		37.9	23.9	1.1	1.0
8 893	18- 36	2.16	0.103			9.7	1.2					1.00	1.07	0.98		40.2	24.1	2.3	2.2
8 894	36- 46	1.39				11.1	1.3			53	35	1.06	1.14	0.98		40.6	25.5	2.5	2.4
8 895	46- 84	0.84				10.5	1.3					1.11	1.20	0.98		42.1	26.7	2.6	2.6
8 896	84-120	0.85				13.2	1.3					1.08	1.15	0.97		49.1	28.7	2.1	2.0
8 897	120-155	0.42				12.6	1.2					1.18	1.25	0.97		42.4	27.7	1.9	1.9
8 898	155-185	0.37				12.9	1.3			65	52	1.01	1.05	1.00		54.2	27.4	1.3	1.3
8 899	185-205	0.20				12.6	1.5					1.05	1.10	1.00		49.7	27.0	1.6	1.6
8 900	205-235					12.5	1.4	0.1	0.2	62	54	1.00	1.05	1.00		54.5	25.8	1.6	1.6
8 901	0- 18	7.43	0.276								0.94			0.99			22.4		

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6D2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	EXT. ACIDITY 6H2A	KCL EXT. AL 6G1E	CEC SUM 5A3A	NH4-OAC 5A6A	BASE SATURATION SUM+ ACI-	SUM+ AL	NH4-OAC 5C1	PH NAF 2 MIN 8C1D	H2O 1/1 8C1A	CA-CL2 8C1E
		MEQ/100G									PERCENT					
8 892	0- 18	12.1	3.6	TR	1.1	16.8	29.4		46.2	33.9	36		50		5.8	5.3
8 893	18- 36	3.4	3.2	TR	0.7	7.3	10.9	0.3	18.2	20.2	40	96	36		5.4	4.8
8 894	36- 46	3.1	3.7	TR	1.0	7.8	12.6	0.2	20.4	17.9	38	98	44		5.2	4.8
8 895	46- 84	2.7	3.2	TR	1.0	6.9		0.2	6.9	20.6	100+	97	34		5.3	4.8
8 896	84-120	2.4	2.0	TR	0.8	5.2	16.7	0.5	21.9	17.2	24	91	31		5.2	4.6
8 897	120-155	1.3	1.3	TR	0.6	3.2	17.7	1.4	20.9	14.6	15	70	22		5.2	4.5
8 898	155-185	1.0	1.0	0.1	0.5	2.6	17.5	2.9	20.1	14.8	13	47	18		5.2	4.4
8 899	185-205	0.6	0.6	TR	0.4	1.6	17.7	2.7	19.3	20.2	9	38	8		5.2	4.4
8 900	205-235	0.6	0.6	0.1	0.3	1.6	18.7	2.6	20.3	16.4	8	38	10		5.2	4.4
8 901	0- 18														5.8	5.3

CLAY MINERALOGY

SAMPLE NO	DEPTH CM.	LESS THAN 0.002 MM X RAY 7A2E I	7A2E II	7A2E III	7A2E IV	DTA 7A3 -PERCENT-	7A3	CLT. 0.0002 MM X RAY 7A2E I	7A2E II	7A2E III	MINERAL CODE	RELATIVE AMOUNT
8 892	0- 18	KK 3									KK KAOLINIT	1 TRACE
8 893	18- 36											2 SMALL
8 894	36- 46	KK 3										3 MODERATE
8 895	46- 84											4 ABUNDANT
8 896	84-120											5 DOMINANT
8 897	120-155											6 INDETER.
8 898	155-185	KK 3										
8 899	185-205											
8 900	205-235	KK 3										
8 901	0- 18											

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.
N=NOT DETERMINED.
Z=NOT ALL FRACTIONS SEPERATED. ADJACENT FRACTIONS INCLUDED HERE.
*=THE 75-250MM FRACTION IS INCLUDED.

SITES LOAM
S68Calif-29-32

Location: Nevada County, California; 2,300 feet south and 1,000 feet west of NE corner of sec. 4, T. 15 N., R.9E., 5 miles ESE of Grass Valley, California; 1,350 feet due north of You Bet-Red Dog Road. Date of Sampling: September 5, 1968.

Description by: L. A. Brittan and G. Staidl. Collectors: L. A. Brittan, G. Staidl, C. B. Goudey, T. M. Ryan, G. M. Kennedy.

Classification: Xeric Haplohumult, clayey, kaolinitic, mesic.

Vegetation: Mixed coniferous forest, ponderosa pine, incense cedar, black oak, bear clover. Not cultivated. Climate: Mean annual precipitation is 56 inches. Mean annual temperature is 56°F. Parent Material: Diabase and metabasic rock. Topography: Strongly sloping, concave slope, 14 percent with NE aspect near crest of slope. Elevation: 2,720 feet. Drainage: Well drained; runoff is medium; permeability is moderate over moderately slow. Soil Moisture: Dry to 14 inches. Very slightly moist 14 to 81 inches.

Remarks: pH taken by Hellige method. Depth to the B2t horizon varies from 14 to 40 inches. This pedon is almost one full pH unit more acid than S68Calif-29-31, and there are more rock fragments in the C1 and C2 horizons.

HORIZON	DESCRIPTION
	Colors are for dry soil unless otherwise indicated.
O1	2-0 inches (5-0 cm.), undecomposed oak leaves and pine needles.
A1 RSL No. 68892	0-7 inches (0-18 cm.), reddish brown (5YR 4/4) heavy clay loam, dark reddish brown (5YR 3/4) moist; moderate medium subangular blocks breaking to strong fine and medium crumbs; slightly hard (dry), friable (moist), slightly sticky and plastic (wet); many very fine and fine random and many medium and coarse horizontal and oblique roots; many very fine interstitial, few very fine tubular pores; 5 percent gravel by volume; clear wavy boundary.
B1t RSL No. 68893	7-14 inches (18-36 cm.), yellowish red (5YR 4/6) light clay, dark red (2.5YR 3/6) moist; weak medium subangular blocks breaking to weak fine crumbs; slightly hard (dry), friable (moist), sticky and plastic (wet); common very fine and fine random and many medium and coarse horizontal and oblique roots; many very fine interstitial and common very fine tubular pores; 2 percent gravel by volume; clear wavy boundary.
B21t RSL No. 68894	14-18 inches (36-46 cm.), red (2.5YR 4/6) light clay, dark red (2.5YR 3/6) moist; weak medium subangular blocks breaking to weak fine crumbs; slightly hard (dry), friable (moist), sticky and very plastic (wet); common very fine and fine random and many medium and coarse horizontal roots; many very fine interstitial and common very fine tubular pores; many thin clay skins line pores; 2 percent gravel by volume; clear wavy boundary.
B22t RSL No. 68895	18-33 inches (46-84 cm.), red (2.5YR 4/6) light clay, dark red (2.5YR 3/6) moist; weak medium and coarse angular blocky structure; hard (dry), friable (moist), sticky and plastic (wet); few very fine and fine vertical and common medium and coarse horizontal and oblique roots; common very fine interstitial and few very fine and fine tubular pores; many thin clay skins on faces of peds and continuous moderately thick clay skins line pores; less than 1 percent gravel, 2 percent weathered cobbles by volume; gradual wavy boundary.
B31t RSL No. 68896	33-48 inches (84-120 cm.), yellowish red (5YR 5/8) light clay, red (2.5YR 4/6) moist; weak medium and coarse angular blocky structure; hard (dry), friable (moist), sticky and plastic (wet); few very fine and fine vertical and common medium horizontal and oblique roots; common very fine interstitial and few very fine and fine tubular pores; many thin clay skins on faces of peds and continuous moderately thick clay skins line pores; 3 percent cobbles by volume; gradual wavy boundary.
B32t RSL No. 68897	48-61 inches (120-155 cm.), yellowish red (5YR 5/8) clay loam, red (2.5YR 5/8) moist; weak medium and coarse angular blocky structure; hard (dry), friable (moist), sticky and plastic (wet); few very fine and fine vertical and common medium horizontal and oblique roots; few very fine tubular and interstitial pores; many thin clay skins on faces of peds and continuous moderately thick clay skins line pores; 3 percent cobbles by volume; clear wavy boundary.
C1 RSL No. 68898	61-72 inches (155-185 cm.), yellowish red (5YR 5/8) loam, red (2.5YR 4/6) moist; massive; slightly hard (dry), friable (moist), slightly sticky and plastic (wet); few very fine, fine and medium vertical roots; few very fine tubular pores; continuous moderately thick red (2.5YR 4/6) dry, 4/8 moist) clay skins line pores; 20 percent by volume of soft weathered rock fragments 1 to 8 cm. in diameter, gradual wavy boundary.
C2 RSL No. 68899	72-81 inches (185-205 cm.), yellowish red (5YR 5/8) light loam, red (2.5YR 4/6) moist; massive; slightly hard (dry), friable (moist), slightly sticky and slightly plastic (wet); few very fine, fine and medium vertical roots; few very fine tubular pores; many moderately thick red (2.5YR 4/8 dry and moist) clay skins line pores; 50 to 60 percent by volume weathered rock fragments greater than 5 cm. in diameter; gradual wavy boundary.
C3 RSL No. 68900	81-92 inches (205-235 cm.), yellowish red (5YR 5/8) light loam with many medium distinct reddish yellow (7.5YR 6/8) and common fine light gray (10YR 7/2) mottles, red (2.5YR 4/6), strong brown (7.5YR 5/8) and light gray (10YR 7/2) moist; massive; slightly hard (dry), friable (moist), slightly sticky and slightly plastic (wet); many moderately thick red (2.5YR 4/8 dry and moist) clay skins line pores and joint planes in the saprolite; 60 percent rock fragments greater than 5 cm. in diameter.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Goldridge fine sandy loam **LOCATION** Sonoma County, California

SOIL FAMILY: Typic Haploxerult, fine-loamy, mixed, mesic.

SOIL NOS. S61Calif-49-5 **LAB. NOS.** 61414 - 61422

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2	

GOLDRIDGE FINE SANDY LOAM
S61Calif-49-5

Location: Sonoma County, California. 300 feet northeast of Springhill School - 4 tree rows east and 4 tree rows north - (LeBallister Ranch) northeast corner of SE1/4, section 4, T. 6 N., R. 9 W.
Date of Sampling: June 22, 1961.
Collectors: Reuben Nelson, Leo C. Klameth, V. C. Miller, G. Lambert, J. H. Kashiwagi, F. Graham.
Classification: **Typic Haploxerult, fine-loamy, mixed, mesic.**
Vegetation: Natural mixed conifers and hardwoods. Use: Apple orchard. Climate: Subhumid mesothermal. Annual precipitation 40 inches, mean January temperature 45.6°F., mean July temperature 65.8°F., mean annual temperature 56.8°F. Parent Material: Soft sedimentary rock - Merced formation. Elevation: About 262 feet. Relief: Gently sloping upland on a 5 percent south facing slope. Drainage: Moderately well drained, medium runoff, moderate to moderately slow permeability. Biological Activity: Worm casts found 17-30 inches.
Remarks: Dead roots found throughout B31t, B32t, B33 horizons on fracture planes, but no live roots below 50 inches.

HORIZON	DESCRIPTION
A11 * RSL No. 61414	0 to 7 inches, light brownish gray (10YR 6/2) fine sandy loam, brown (10YR 4/3) when moist; massive; soft when dry, very friable when moist, nonsticky and nonplastic when wet; abundant very fine and fine roots; many very fine and fine tubular and interstitial pores; slightly acid, pH 6.5; abrupt, smooth boundary.
A12	7 to 17 inches, pale brown (10YR 6/3) fine sandy loam, dark brown (10YR 4/3) when moist;

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Goldridge fin sandy loam **LOCATION** Sonoma County, California

SOIL FAMILY: Typic Haploxerult, fine-loamy, mixed, mesic.

SOIL NOS. S61Calif-49-6 **LAB. NOS.** 61423 - 61430

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	3A1 0.2-0.02	0.02-0.002	2A2 > 2	TEXTURAL CLASS
0- 7	Ap	0.2	0.7	1.1	49.0	18.6	20.9	9.5	72.5	11.4	1	fsl
7-20	A1	0.1	0.4	0.6	46.4	19.3	21.0	12.2	71.4	11.9	0	fsl
20-24	A2	0.0	0.3	0.5	46.5	18.9	21.3	12.5	71.1	12.4	0	fsl
24-28	B1	0.1	0.3	0.4	43.9	16.5	20.9	17.9	64.8	13.5	0	fsl
28-41	B21t	0.0	0.2	0.4	37.8	11.1	17.1	33.4	50.5	12.7	0	scl
41-57	B22t	0.0	0.0	0.2	38.5	12.9	19.5	28.9	54.2	14.9	0	scl
57-73	B31	0.0	0.2	0.2	38.3	16.6	23.1	21.6	60.9	16.0	0	scl
73+	B32	0.0	0.2	0.2	38.1	18.1	25.3	18.1	62.9	17.4	0	fsl

8C1a pH H_2O 1:1	8C1c pH KCl 1:1		6A1a ORGANIC MATTER ORGANIC CARBON %	6B1a NITROGEN %	C/N	Bulk Density 4A3a g/cc		Ext. Iron 6C1b % Fe				4B2 Soil Water 15 atmos. Pct.
5.0	4.3		1.48	0.109	14			0.3				4.2
5.2	4.0		0.52	0.048	11	1.43		0.3				4.7
5.1	4.0		0.20	0.023		1.56		0.3				4.5
5.0	3.9		0.18	0.024		1.6		0.4				6.5
4.6	3.7		0.20	0.030		1.65		1.0				14.0
4.5	3.6		0.06			1.60		0.8				13.5
4.5	3.6		0.03			1.52		0.7				11.8
4.5	3.6		0.01			1.50		0.6				11.6

5A2a CATION EXCHANGE CAPACITY (+ Na)	EXTRACTABLE CATIONS 6N2b Ca	6O2c Mg	6H2a H	6P2a Na	6Q2a K	5B1a BASE SAT. % 5C3	Sum of Cations 5A3a					MOISTURE AT SATURATION %
	milliequivalents per 100g. soil →						meq/100g.					
9.8	2.0	0.5	6.8	0.2	0.6	33	10.1					
7.4	1.3	0.3	5.2	0.1	0.3	28	7.2					
5.7	1.3	0.5	3.5	0.1	0.2	38	5.6					
7.3	1.7	0.9	3.4	0.1	0.2	46	6.3					

GOLDRIDGE FINE SANDY LOAM
S61Calif-49-6

Location: Sonoma County, California. Stanley Ross Ranch, northeast corner of NW1/4, SW1/4, section 11, T. 6 N., R. 9 W., 2 rows in from Elphic Road and 2 rows from young orchard.
Date of Sampling: June 22, 1961.
Collectors: Reuben Nelson, Leo C. Klameth, V. C. Miller, G. Lambert, J. H. Kashiwagi, F. Graham.
Classification: [illegible]

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Haploxerult, fine-loamy, mixed, mesic

SOIL Josephine gravelly silt loam SOIL Nos. S60Calif-3-24 LOCATION Amador County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 60128 - 60133

		Size class and particle diameter (mm) 3A1						Coarse fragments 1A2a
		Total	Sand	Silt				

JOSEPHINE GRAVELLY SILT LOAM
S60Calif-3-24

Location: Amador County, California. 1/2 mile west of Shake Ridge Road, crossing over Rancheria Creek, 150 feet north of Shake Ridge Road, 3.8 miles west of junction of Charleston, Volcano Road and Shake Ridge Road. 0.35 mile north of the SE corner of sec. 24, T. 7 N., R. 11 E.
Date of Sampling: May 4, 1960.
Description by: Grant Kennedy. Collectors: Grant Kennedy, Reuben Nelson, Robert Sketchley, Clifford Henry, Paul Nazar, Jack McClelland.
Classification: Typic Haploxerult, fine-loamy, mixed, mesic.
Vegetation: Ponderosa pine (Pinus ponderosa), sugar pine (Pinus lambertiana), black oak (Quercus velutina), mountain misery (Chamaebatia foliolosa), manzanita (Arctostaphylos sp.), toyon (Photinia arbutifolia).
Climate: Mean annual rainfall 33 inches; average temperatures (Tiger Creek Station): January - 40° F.; July - 70° F.; mean annual - 55° F. Parent Material: Metasedimentary rocks of the Calaveras formation (schists with vertical cleavage). Topography: Steep uplands on a north-facing slope of 25 percent.
Elevation: 2,050 feet. Drainage: Well drained; moderate runoff; moderate permeability.
Remarks: Field estimates of the volume of gravel for the B2t, B3t, and C1 horizons are given in the profile description.

HORIZON	DESCRIPTION
	Colors are for the dr · soil unless otherwise indicated.
O1	1-0 inch (3-0 cm.), organic layer consisting of fresh needles, litter, and decomposing litter.
A1 RSL No. 60128	0-2 inches (0-5 cm.), brown (10YR 4/3) gravelly loam, dark brown (7.5YR 3/2) when moist; weak fine subangular blocky structure; slightly hard (dry), friable (moist), slightly sticky and slightly plastic (wet); abundant very fine, few fine and medium roots; many very fine, few fine interstitial and tubular pores; abrupt smooth boundary.
A3 RSL No. 60129	2-9 inches (5-23 cm.), brown (7.5YR 5/4) gravelly light loam, dark brown (7.5YR 4/4) when moist; slightly hard (dry), friable (moist), slightly sticky and slightly plastic (wet); abundant very fine, few fine and medium roots; many very fine tubular and interstitial pores; few fine and medium tubular pores; clear smooth boundary.
B1t RSL No. 60130	9-22 inches (23-56 cm.), brown (7.5YR 5/5) light clay loam, reddish brown (5YR 4/5) when moist; moderate medium subangular blocky structure; hard (dry), friable (moist), sticky and plastic (wet); abundant very fine, few fine to very coarse roots; pores as above; thin, nearly continuous clay skins in pores; gradual irregular boundary because of presence of a few partially disintegrated stones and bedrock.
B2t RSL No. 60131	22-37 inches (56-94 cm.), brown (7.5YR 5/5) heavy silty clay loam, strong brown (7.5YR 4/6) when moist; moderate medium subangular blocky structure; very hard (dry), firm (moist), sticky and plastic (wet); few fine to coarse roots; many very fine interstitial and tubular pores, few fine and medium tubular pores; moderately thick continuous clay skins on ped faces and in many pores; about 25 percent by volume of weathered soft rocks; gradual irregular boundary.
B3t RSL No. 60132	37-47 inches (94-120 cm.), yellowish brown (10YR 5/6) silty clay loam, dark yellowish brown (10YR 4/6) when moist; moderate medium subangular blocky structure; very hard (dry), firm (moist), slightly sticky and slightly plastic (wet); few very fine to large roots; many very fine interstitial pores, few very fine, common fine tubular pores; clay skins as above but thin; about 35 percent by volume of weathered soft rocks; clear irregular boundary.
C1 RSL No. 60133	47-68 inches (120-175 cm.), brownish yellow, yellowish red, and gray (10YR 6/6, 5YR 5/6, and N 6/) decomposing rock, brownish yellow, red, and dark gray (10YR 6/6, 2.5YR 4/6, and N 4/) when moist; massive; extremely hard (dry), very firm (moist); very few roots; few very fine tubular, common fine interstitial pores; moderately thick, continuous clay skins along fracture faces.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Mariposa silt loam (taxadjunct) **LOCATION** Amador County, California

SOIL FAMILY: Typic Haploxerult, fine-loamy, mixed, mesic

SOIL NOS. S60Calif-3-25 **LAB. NOS.** 60108 - 60113

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1	2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	≥ 3 1/	
0- 1	A11	3.9	5.7	4.4	10.4	11.5	55.5	8.6	43.5	29.9	20	sil
1- 4	A12	3.2	5.2	4.1	8.9	11.6	55.7	11.3	41.3	31.7	12	sil
4- 8	B1t	2.0	3.8	3.0	6.7	9.2	56.1	19.2	33.8	35.8	8	sil
8-15	B2t	0.9	3.8	2.6	6.3	8.1	58.2	20.1	32.2	38.2	25	sil
15-23	[illegible]	1.0	3.2	2.1	5.5	7.2	64.6	16.4	31.1	44.2	0	sil

Date of sampling: May 5, 1960.
Location: 0.1 mile NW of the center of the SE1/4 sec. 34, T. 8 N., R. 12 E., 25 feet W of Fiddletown-Silver Lake Road; 3.5 miles W of Fiddletown-Silver Lake

OIL SURVEY LABORATORY Riverside, California

OIL TYPE Red Bluff gravelly loam (taxadjunct) **LOCATION** Amador County, California

SOIL FAMILY: Typic Haploxerult, clayey, mixed, thermic

s59Calif-3-9 LAB. NOS. 59314 - 59319

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	3A1 0.02-0.002	2A2 >2 1/	TEXTURAL CLASS
0-3	A_{11}	4.3	12.3	10.5	17.5	7.2	32.2	16.0	32.4	16.4	18	1-fsl
3-7	A_{12}	3.9	11.3	11.1	18.5	7.9	29.5	17.8	34.0	14.0	17	fsl
7-11	B_1	3.0	10.3	10.2	16.9	7.0	27.1	25.5	30.7	13.1	10	fsl
11-19	B_{21}	2.2	5.5	4.5	7.9	3.9	17.3	58.7	15.6	10.1	3	c
19-27	B_{22}	1.3	4.4	3.9	7.2	3.6	17.2	62.4	15.3	9.7	2	c
27-40	D	3.8	6.6	6.4	11.1	5.1	19.9	47.1	20.2	11.2	26	c

pH		ORGANIC MATTER		6C1b	ELECTRICAL CONDUC-			MOISTURE TENSIONS
[illegible]	[illegible]	[illegible]	[illegible]					[illegible]

RED BLUFF GRAVELLY LOAM (taxad.junct) 1/
S59Calif-3-9

Collectors: Grant Kennedy, Reuben Nelson, Bob Sketchley.
Date of Sampling: June 16, 1959. Description by: Grant Kennedy.
Location: Amador County, California on Meiss Road, 1/2 mile from intersection with Highway 104, 20 feet east of road, SW1/4 of SW1/4 of Sec. 9, T6N, R9E.
Elevation: 250 feet.
Relief: On an old terrace with a plane slope of 8 percent, facing west.
Drainage: Well drained; moderate runoff; moderate over slow permeability.
Parent Material: Mixed gravels of uncertain age resting unconformably on the Ione formation.
Climate: Rainfall about 18 inches.
Vegetation: Blue oak with wild oats and soft chess.
Remarks: This is a medial developed Noncalcic Brown soil with about 10 percent by volume of cobbles and gravels in the B horizons. The B_{22} may be formed in the Ione formation as well as the D and probably should be B_{22u}.
Classification: Typic Haploxerult, clayey, mixed, thermic.

HORIZON		DESCRIPTION
A_{11}	0-3"	Reddish brown (5YR 4/3) gravelly loam, near gravelly fine sandy loam, dark reddish brown (5YR 3/3) when moist; massive; hard when dry, friable when moist, nonsticky and nonplastic when wet; abundant very fine roots; many very fine, common fine, few medium pores; slightly acid, pH 6.5; abrupt, smooth lower boundary.
A_{12}	3-7"	Reddish brown (5YR 4/5) gravelly fine sandy loam, near gravelly loam, dark reddish brown (5YR 3/4) when moist; massive; hard when dry, friable when moist, slightly plastic and slightly sticky when wet; abundant very fine and fine, few medium and coarse roots, many very fine, common fine, few medium and coarse pores; occasional thin discontinuous clay film; slightly acid, pH 6.1; clear, smooth lower boundary.
B_1	7-11"	Yellowish red (5YR 4/6) loam near clay loam, dark reddish brown (2.5Y 3/5) when moist; massive, hard when dry, friable when moist, slightly sticky and slightly plastic when wet; plentiful very fine and fine, few medium and coarse roots; many very fine, common fine, few medium and coarse pores; thin continuous clay films in pores; very strongly acid, pH 5.3; clear, smooth lower boundary.
B_{21}	11-19"	Red (2.5YR 4/6) clay, dark red (10R 3/6) when moist; massive; hard when dry, firm when moist, sticky and plastic when wet; few very fine, fine and medium roots; common very fine, few fine, medium and coarse pores; moderately thick, continuous clay films in pores and in bridges; extremely acid, pH 4.0; gradual, smooth lower boundary.
B_{22}	19-27"	Weak red (10R 4/4) clay with common streaks of pale brown (10YR 6/3), dark red (10R 3/6) and brown (10YR 5/3) when moist; massive; very hard when dry, very firm when moist, sticky and plastic when wet; few fine and medium roots; common very fine, few fine, medium and coarse pores; thick continuous clay films in pores and bridges; extremely acid, pH 3.6; gradual, smooth lower boundary.
D	27-40"	Pale brown (10YR 6/3) gravelly clay with dark red (10R 3/6) coatings, brown (10YR 5/3) with dark red (10R 3/6) when moist; massive; very hard when dry, extremely firm when moist,

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Chromoxerert, fine, montmorillonitic, mesic

SOIL Montague clay SOIL Nos. S66Calif-47-7 LOCATION Siskiyou County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 66371 - 66375

Depth (in.)	Horizon	1B1b Total Sand (2–0.05)	Total Silt (0.05–0.002)	Total Clay (< 0.002)	Sand Very coarse (2–1)	Sand Coarse (1–0.5)	Sand Medium (0.5–0.25)	Sand Fine (0.25–0.1)	Sand Very fine (0.1–0.05)	Silt (0.05–0.02)	Silt Int. III (0.02–0.002)	Silt Int. II (0.2–0.02)	Silt (2–0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	1/ Coarse fragments > 2 Pct.	1A2a 2 - 19	1A2a 19 - 76
		Size class and particle diameter (mm) 3A1																
		Pct. of < 2 mm															Pct. of < 76mm	
0-4	Ap	23.2	29.8	47.0	1.5	2.8	3.5	8.3	7.1	10.0	19.8	21.6	16.1		47	7	4	3
4-16	A12	23.1	27.9	49.0	1.6	2.9	4.0	8.1	6.5	9.4	18.5	20.3	16.6		49	6	5	1
16-24	A13	22.3	28.1	49.6	2.0	3.0	3.6	7.7	6.0	9.9	18.2	20.2	16.3		50	4	4	-
24-30	C1cam	55.1	28.2	16.7	14.2	13.5	7.3	11.7	8.4	10.7	17.5	25.5	46.7	2	15			
30-36	C2cam	61.7	25.7	12.6	19.8	15.4	7.4	11.5	7.6	10.3	15.4	23.9	54.1	3	10			

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	Bulk density 4A1h Oven dry g/cc	Bulk density g/cc	3B2 C'	Water content Pct.	Water content 4B1c 1/3 bar Pct.	Water content 4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1a H_2O 1:1		
0-4	0.63	0.059	11	0.6			1.16	1.57		0.97		38.2	23.3	0.107	0.103	6.6		
4-16	0.51	0.050	10	0.5			1.30	1.75		0.97		34.0	27.2	0.104	0.100	6.7		
16-24	0.42	0.046	9	0.6	tr.		1.28	1.70		0.98		32.4	26.5	0.098	0.096	7.2		
24-30	0.53			0.1	31		1.12	1.18		1.00		40.5	26.5	0.019	0.019	8.1		
30-36	0.56			0.1	40		1.26	1.27		1.00		31.8	23.8	0.002	0.002	8.3		

Depth (in.)	Extractable bases 5B1a 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A1a NH_4OAc	5A3a Sum		6G1d KCl Ext. Al^{+++}						Base saturation 5C1 NH_4OAc Pct	5C3 Sum Pct.
	meq/100 g																
0-4	17.8	19.8	0.2	2.6	40.4	2.9	40.1	43.3								100	93
4-16	18.0	17.0	0.2	1.0	36.2	2.5	41.8	38.7								87	93
16-24	18.0	17.5	0.4	1.2	37.1	1.9	41.5	39.0								89	95
24-30	20.0	13.9	0.3	0.7		-	27.7	34.9								100+	100
30-36	18.7	10.6	0.3	0.6		-	22.1	30.2								100+	100

Depth (in.)		Clay mineralogy (<0.002 mm) 7A Mont.	Verm.	Mica	Kaolin	
		X-ray				
0-4		xxxx				
4-16						
16-24						
24-30		xxxx				
30-36						

1/ From characterization sample (5 kg.). Samples 66374 and 66375 were crushed and >2 mm. fractions were not determined.

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

MONTAGUE CLAY
S66Calif-47-7

Location: Siskiyou County, California. A pit located just north of where the Montague ditch crosses the County Airport Road on the Lipert Ranch. Pit site located between power poles #30 and 7/25, 47 paces north and 48 paces west from pole #30. Section 29, T46N, R5W. Date of Sampling: May 26, 1966.

Description by: S. B. Johnson. Collectors: K. W. Flach, S. B. Johnson and D. Lucas.

Classification: Typic Chromoxerert, fine, montmorillonitic, mesic.

Vegetation: Cultivated. Summer fallow at time of description and sampling. Climate: The average annual precipitation is approximately 13 inches with some in the form of snow. The mean January temperature is 34°F., the mean July temperature is 73°F., and the mean annual temperature is 52°F. Parent Material: Alluvium from basic igneous rock of the Western Cascade Series formation. The alluvium was deposited over semi-consolidated tuff and a hardpan formed in the surface layers of the tuff. The dominant mineral is montmorillonitic clay. Topography: The site is located on a nearby level valley bottom. There is very faint undulation of the land surface. Slope 1 percent or less. Elevation: 2680 feet plus or minus 10 feet. Drainage: Well drained, slow permeability, hardpan very slow, runoff slow to ponded. Soil Moisture: 0 to 4 inches about 70 percent of field capacity and 4 to 24 inches at field capacity.

Remarks: Reactions were determined by the Truog soil reaction test. This was the only Montague soil sampled, therefore, there is no paired sample. This soil varies mainly in depth and color. Soil depth to the duripan ranges from 20 to 60 inches with an average depth of about 24 to 26 inches. Soil colors are normally in the 10YR hues and range from black to dark brown (2/1, 2/2, 3/2, 3/3) moist and very dark gray to brown (3/1, 4/1, 4/2, 4/3, 5/3) dry. Slickensides and pressure faces are common in the lower horizons and middle horizons. Depth of cracking ranges from 8 to 20 inches. Under irrigation the soil produces good alfalfa hay and seed, plus excellent irrigated pastures. Coarse fragments are negligible in most areas of the Montague soil. Gravel occurs throughout the soil in places and cobbles and stones may be found near rock outcroppings. A gravelly phase hasn't been mapped to date. The pan ranges in thickness from 6 inches to a foot or two and is strongly cemented to nearly indurated.

HORIZON	DESCRIPTION
Ap RSL 66371	0 to 4 inches, dark gray (10YR 4/1) clay, very dark gray (10YR 3/1) moist; strong coarse angular blocky breaking to strong medium angular blocky structure; very hard, firm, very sticky, very plastic; few fine and very fine roots; few fine tubular pores; neutral (pH 6.6); abrupt smooth boundary.
A12 RSL	4 to 16 inches, brown (10YR 5/3) rubbed, very dark grayish brown (10YR 3/2) ped faces, clay, dark brown (10YR 3/3) moist and rubbed, with very dark grayish brown

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Chromoxerert, fine, montmorillonitic, thermic.

SOIL Altamont clay SOIL Nos. S58Calif-1-3 LOCATION Alameda County, California

SOIL SURVEY LABORATORY Beltsville, Maryland LAB. Nos. 591013 - 591018

Depth	Horizon	1B1b Total Sand	Total Silt	Total Clay	Size class and particle diameter (mm) 3A1 Sand Very	Sand Coarse	Sand Medium	Sand Fine	Sand Very fine	Silt	Silt Int III	Silt Int. II		3A1a Carbon	3A1a Non-carbon	Coarse fragments 1A2a 2A2 >2	2-19	19-76

ALTAMONT CLAY
S58Calif-1-3

Location: 750 feet east and 2250 feet south of northwest corner of Section 19 in SE1/4, SW1/4, NW1/4, Section 19, R. 3 E., T. 2 S., Dyer Ranch, Alameda County, California.
Date of Sampling: November 12, 1958.
Description by: R. Ulrich. Collectors: R. E. Nelson, Larry Welch, Robert Mitchel, Lynn Whittig, R. Ulrich.
Classification: Typic Chromoxerert, fine, montmorillonitic, thermic.
Vegetation: Annual bromes (soft chess and ripgut mostly), mouse barley, bur clover, mustard. Present use is annual grass range; may have been cultivated sometime in past. Climate: Rainfall is 13-14 inches (estimated); mean annual soil temperature about 60° F. Distinct summer-dry, winter-moist Mediterranean type climate. Parent Material: Interbedded calcareous fine grained sandstone and shale. Lime content is variable, but it usually is weak or slightly calcareous although thin seams of limestone and other strongly calcareous sediments are often present. Slope: 25 percent (estimated), west facing, plane surface. Elevation: About 1100 feet. Drainage: Well drained with slow surface runoff when dry and deeply cracked and medium runoff when moist and expanded during the rainy season. Internal drainage is variable ranging from very rapid when dry and deeply cracked during the dry season and for some interval after the start of the rainy season to slow internal drainage when moist and essentially massive following expansion during the rainy season.

HORIZON	DESCRIPTION
A11 BSL No. 591013	0 to 7 inches, dark brown (10YR 4/2.5) clay, darker brown (10YR 3/2.5) moist, drying 10YR 5/2 or lighter in the very thin immediate surface crust; strong coarse prismatic and strong coarse and medium blocky; very hard to extremely hard, very firm to extremely firm, very plastic, sticky; numerous fine roots which tend to concentrate along structure planes and facings, roots appear flattened or compressed; few extremely fine and very fine tubular pores; noncalcareous to dilute acid; reaction not determined in field, see lab data; structure facings have very dark brown to black carbonaceous films and mats on them in places; clear smooth lower boundary.
A12 BSL No. 591014	7 to 19 inches, dark brown (10YR 3.5/3) clay, darker brown (10YR 3/3) moist, moderate coarse prismatic and moderate coarse blocky; very hard, very firm, very plastic, sticky; few fine roots, mostly along structural planes; few extremely fine and very fine tubular pores; noncalcareous but may have a few whitish lime films on surface of peds in lower portion; many slickensides; clear smooth lower boundary.
A13 BSL No. 591015	19 to 28 inches, similar colored clay with few fine whitish soft to slightly hard lime segregations and films; moderate coarse prismatic and moderate coarse and medium blocky; very hard, very firm, very plastic, sticky; few fine roots; concentrated mainly along structure facings; few extremely fine and very fine tubular pores; slightly calcareous, the lime mostly disseminated; many slickensides; clear smooth lower boundary.
A14ca BSL No.	28 to 37 inches, finely mottled dark brown and dark yellowish brown (10YR 4/3 and 4/4) clay, darker brown and darker yellowish brown (10YR 3/3 and 4/3) moist. with few fine whitish

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Chromoxerert, fine, montmorillonitic, thermic.

SOIL Altamont clay SOIL Nos. S58Calif-1-4 LOCATION Alameda County, California

SOIL SURVEY LABORATORY Beltsville, Maryland LAB. Nos. 591019 - 591024

ALTAMONT CLAY
S58Calif-1-4

Location: 700 feet east and 2600 feet south of northwest corner of Section 24 in SE1/4, NW 1/4, Section 24, R. 3 E., T. 2 S., Flynn Ranch, Alameda County, California. On the north side of Altamont Pass Road. Date of Sampling: November 12, 1958.
Description by: R. Ulrich. Collectors: R. E. Nelson, Larry Welch, Robert Mitchel, R. Ulrich.
Classification: Typic Chromoxerert, fine, montmorillonitic, thermic.
Vegetation: Annual bromes (soft chess and ripgut mostly), wild oats, mouse barley, turkey mullein, and tar weed. Present use is annual grass range; may have been cultivated sometime in the past. Climate: Rainfall is 11-12 inches (estimated); mean annual soil temperature about 60° F. Distinct summer-dry, winter-moist Mediterranean type climate. Parent Material: Interbedded calcareous fine grained sandstone and shale. Lime content is mainly weak or slight although thin seams of limestone and other strongly calcareous sediments are often present. Slope: 6-7 percent (estimated), south facing, slightly concave. Elevation: About 450 feet. Drainage: Well drained with slow surface runoff when dry and deeply cracked and slow to medium runoff when moist and expanded during the rainy season. Internal drainage is variable varying from very rapid when dry and deeply cracked during the dry season and for some interval after the start of the rainy season to slow internal drainage when moist and essentially massive following expansion during the rainy season.

HORIZON	DESCRIPTION
A11 BSL No. 591019	0 to 7 inches, dark brown (10YR 3.5/3) clay, darker brown (10YR 3/3) moist, drying to 10YR 5/2 or lighter in the very thin immediate surface crust; strong coarse prismatic and strong coarse and medium angular blocky; very hard, very firm, very plastic, sticky; numerous fine roots tending to concentrate along structure planes and facings, roots appear flattened or compressed; few extremely fine and very fine tubular pores; noncalcareous to dilute acid; reaction not determined in field, see lab data; structure facings have very dark brown to black carbonaceous films and mats on them in places; clear smooth lower boundary.
A12 BSL No. 591020	7 to 18 inches, similar colored clay; strong coarse prismatic and strong coarse angular blocky; very hard, very firm, very plastic, sticky; few fine roots, mostly along structure planes and flattened or compressed; few extremely fine and very fine tubular pores; noncalcareous in most of mass with few white films or small soft white lime segregations in lowest portion of horizon; many slickensides especially in lower portion; clear smooth lower boundary.
A13 BSL No. 591021	18 to 26 inches, similar colored clay with some dark yellowish brown (10YR 3.5/4) mottles, darker yellowish brown (10YR 3/4) moist, and common fine soft to hardened whitish lime segregations and lime films; moderate coarse prismatic and moderate medium blocky; very hard, very firm, very plastic, sticky; few fine roots, mostly along structure faces; few extremely fine and very fine tubular pores; matrix is slightly calcareous, with the lime segregations, nodules, and films strongly calcareous; many slickensides; clear wavy lower boundary.
A14ca BSL No. 591022	26 to 37 inches, dark yellowish brown (10YR 4/4) clay, darker yellowish brown (10YR 3/4) moist, with fine mottlings of 10YR 3.5/3 dry and 3/3 moist and many fine soft to hardened whitish lime segregations, nodules, and films; moderately coarse prismatic and moderate medium blocky; very hard, very firm, very plastic, sticky; few fine roots, mostly along structure planes; few extremely fine and very fine tubular pores; matrix is slightly calcareous, with the lime segregations strongly calcareous; many slickensides; clear smooth lower boundary.
ACca BSL No. 591023	37 to 47 inches, yellowish brown (10YR 5/4) heavy silty clay loam, dark yellowish brown (10YR 4/4) moist, with a few fine soft whitish lime segregations and nodules; weak medium and fine subangular blocky; hard, firm, plastic, slightly sticky; very few fine roots; few extremely fine and very fine tubular pores; matrix is very slightly to slightly calcareous with strong reaction in lime segregations; few slickensides; abrupt smooth lower boundary.
Dr BSL No. 591024	47 inches +, pale olive (5Y 6/3 and 6/4) fine grained sandstone, olive (5Y 5/3 and 5/4) moist, and pale yellow (5Y 8/3 and 8/4) arenaceous limestone,(5Y 7/3 and 7/4) moist, in thin interbedded strata, the sandstone is very weakly calcareous, and the limestone is

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Chromoxerert, fine, montmorillonitic, thermic
SOIL Canay clay (crusty surface) SOIL Nos. S67Calif-33-1 LOCATION Riverside, County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 6716 - 6724

Depth (cm.)	Horizon	1B1b Total: Sand (2-0.05)	Silt (0.05-0.002)	Clay (< 0.002)	Sand: Very coarse (2-1)	Coarse (1-0.5)	Medium (0.5-0.25)	Fine (0.25-0.1)	Very fine (0.1-0.05)	Silt: 0.05-0.02	Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non carbonate clay	1/ Coarse fragments 1A2a > 2	2 - 19	19 - 76
		Size class and particle diameter (mm) 3A1																
		Pct. of < 2 mm														Pct.	Pct. of < 76 mm	
0-5	A11	12.8	40.3	46.9	1.5	2.0	1.7	3.2	4.4	23.1	17.2	29.3	8.4	0	47	3	1	2
5-18	A12	12.7	37.9	49.4	1.5	2.0	1.4	2.9	4.9	20.6	17.3	27.2	7.8	0	49	3	2	1
18-36	A13	13.3	38.5	48.2	1.8	2.0	1.3	2.6	5.6	20.6	17.9	27.7	7.7	0	48	5	4	1
36-58	A14	12.7	37.8	49.5	1.3	1.8	1.2	2.7	5.7	19.8	18.0	27.1	7.0	0	50	4	4	-
58-79	A15	12.1	38.4	49.5	1.4	1.9	1.2	2.4	5.2	20.4	18.0	27.0	6.9	0	50	5	4	1
79-110	A16	12.2	38.0	49.8	1.3	1.7	1.1	2.6	5.5	19.0	19.0	25.9	6.7	0	50	2	2	-
110-140	AC	11.1	40.2	48.7	1.0	1.7	1.2	2.9	4.3	16.3	23.9	22.2	6.8	0	49	2	2	-
140-160	C1	51.7	29.5	18.8	1.2	5.6	9.6	23.7	11.6	12.8	16.7	35.9	40.1	0	19	1	1	-
160-225+	C2	69.1	22.9	8.0	0.5	3.5	8.8	40.4	15.9	10.9	12.0	48.7	53.2	0	8	-	-	-

Depth (cm.)	6A1a Organic carbon	6B1a Nitrogen	C/N	6C2b Ext. Iron as Fe	6E1b Carbonate as $CaCO_3$	Bulk density: 4A1f 1/3 bar	4A1h Oven dry		3B2 Cm	Water content	4B1c 1/3 bar	4B2 15 bar	4D1 Extensibility LEf	4D1 Extensibility LE	pH: 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
	Pct.	Pct.		Pct.	Pct.	g/cc	g/cc	g/cc		Pct.	Pct	Pct.	Pct	Pct.			
0-5	1.20	0.114	11		-							19.4			7.2		7.9
5-18	0.80	0.074	11		-		1.35	1.83		0.98		29.7	19.4	10.7	10.4	7.3	8.3
18-36	0.57	0.048	12		tr.		1.40	1.84		0.97		27.9	19.7	9.5	9.2	7.7	8.9
36-58	0.50	0.043	12		tr.		1.45	1.95		0.98		27.9	20.2	10.4	10.1	7.7	8.8
58-79	0.50	0.040	12		1		1.45	1.89		0.97		28.3	20.8	9.2	9.0	7.7	8.9
79-110	0.39	0.034	11		1		1.45	1.92		0.99		29.0	21.6	9.8	9.7	7.6	8.8
110-140	0.28	0.026	11		1		1.34	1.81		0.99		34.2	25.0	10.5	10.4	7.6	8.7
140-160	0.17				-		1.40	1.68		0.99		27.5	21.1	6.3	6.2	7.3	8.5
160-225+	0.15				-		1.43	1.62		1.00		28.5	16.0	4.2	4.2	7.1	8.0

Depth (cm.)	Extractable bases 5B1a: 6N2e Ca	6O2d Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation exch. Capacity: 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 8A1: 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1b SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-5	29.7	12.4	0.5	2.4				42.3										
5-18	31.0	11.6	0.9	1.3				43.3										
18-36	40.7	10.4	1.4	1.0				42.9										
36-58	45.9	10.8	2.3	1.0				44.3	3.1	1.3	4.2	0.1		-	2.9	3.2	2.7	0.87
58-79	41.0	10.2	3.4	1.0				44.0	4.3	1.1	7.5	0.2		-	2.8	9.0	3.5	1.52
79-110	43.2	10.4	4.3	0.9				45.6	8.5	2.7	13.0	0.2		-	2.9	17.5	6.3	2.59
110-140	44.7	11.4	5.2	0.8				52.2	10.5	4.0	15.6	0.8		-	2.3	19.3	8.3	2.94
140-160	39.4	11.0	4.6	0.5				52.5	8.0	2.5	14.8	0.5		-	2.1	17.7	6.9*	2.66
160-225+	31.0	8.5	4.0	0.4				42.7	4.7	1.2	11.4	0.2		-	1.3	13.7	4.1	2.08

Depth (cm.)	8A Water at Saturation	Exch. bases: 5B3b Ca	5B3b Mg	5B1b Na	5D2 Exchangeable Na	SAR	Gypsum	Clay mineralogy (<0.002 mm) 3A1: Mont.	Verm.	Mica 7A2	Kaolin.
	Pct.	meq./100 g.			Pct.		Pct.	X-ray			
0-5								xxxx		tr.	tr.
5-18											
18-36											
36-58	75.4										
58-79	75.3							xxxx			tr.
79-110	76.4										
110-140	85.8										
140-160	66.1										
160-225+	58.3							xxxx			tr.

1/ Coarse fragments from characterization sample (5 kg.).

* Indicator destroyed indicating nitrites.

CAPAY CLAY (Crusty surface)
S67Calif-33-1

Location: Riverside County, California, Section 19, T. 4 S., R. 5 W., SBBM on aerial photo AXM 13K-2.
Date of Sampling: May 5, 1967. Description by: B. R. Brasher.
Classification: Typic Chromoxerert, fine, montmorillonitic, thermic.
Topography: Plane slope, 5 percent, midslope position, aspect ENE.
Collectors: F. F. Peterson, K. W. Flach, W. D. Nettleton and B. R. Brasher.

HORIZON	DESCRIPTION
A11 RSL No. 6716	0 to 5 cm., grayish brown (10YR 5/2) clay, very dark grayish brown (10YR 3/2) moist; moderate, medium subangular blocky structure, common, inclined pressure faces; hard, friable, sticky, plastic; plentiful fine and very fine roots; many fine interstitial exped pores and many fine vesicular inped pores; clear, smooth boundary.
A12 RSL No. 6717	5 to 18 cm., dark grayish brown (10YR 4/2) clay, dark brown (10YR 3/3) moist; moderate, coarse, prismatic, breaking to moderate coarse subangular blocky, common, inclined pressure faces; very hard, firm, sticky, plastic; plentiful fine and very fine roots; common very fine inped pores; clear smooth boundary.
A13 RSL No. 6718	18 to 36 cm., dark grayish brown (10YR 4/2) clay, dark brown (10YR 3/2) moist; moderate, coarse, prismatic breaking to moderate medium, subangular blocky, common, inclined pressure faces; very hard, firm, sticky, plastic; plentiful fine and very fine roots; common very fine inped pores; clear, smooth boundary.
A14 RSL No. 6719	36 to 58 cm., grayish brown (10YR 4/2) clay, very dark grayish brown (10YR 3/2) moist; strong, medium, angular blocky, common intersecting slickensides from 1 to 10 cm. long on ped surfaces; very hard, very firm, sticky, plastic; plentiful fine and very fine mostly exped roots; common fine and very fine inped pores; arbitrary boundary for sampling purposes.
A15 RSL No. 6720	58 to 79 cm., dark grayish brown (10YR 4/2) clay, dark brown (10YR 3/2) moist; strong, medium, angular blocky, common intersecting slickensides from 1 to 10 cm. long on ped surfaces; very hard, very firm, sticky, plastic; plentiful very fine roots; common fine and very fine inped pores; gradual wavy boundary.
A16	79 to 110 cm., dark grayish brown (10YR 4/2) clay, with common medium, distinct olive gray

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Chromoxerert, fine, montmorillonitic, thermic

SOIL Capay clay (smooth surface) taxadjunct SOIL Nos. S67Calif-33-2 LOCATION Riverside County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 6725 - 6735

Depth (cm)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (<0.002)	Sand Very coarse (2-1)	Sand Coarse (1-0.5)	Sand Medium (0.5-0.25)	Sand Fine (0.25-0.1)	Sand Very fine (0.1-0.05)	Silt (0.05-0.02)	Silt Int. III (0.02-0.002)	Silt Int. II (0.2-0.02)	Silt 2-0.2			Coarse fragments 1A2a > 2 Pct. of whole soil	2-19 Pct. of <76mm	19-76 Pct. of <76mm
		Pct. of <2 mm																
0-10	A11	22.9	50.0	27.1	2.4	3.4	2.4	5.2	9.5	32.9	17.1	45.2	13.4			23	11	12
10-25	A12	19.2	43.5	37.3	1.9	3.2	2.1	4.0	8.0	27.2	16.3	37.3	11.2			31	7	24
25-36	A13	20.1	39.7	40.2	2.2	3.5	2.0	4.1	8.3	23.4	16.3	33.9	11.8			10	9	1
36-61	A14	18.4	41.4	40.2	2.2	2.9	1.8	4.0	7.5	23.8	17.6	33.5	10.9			5	5	-
61-84	C1ca	19.4	39.1	41.5	2.6	2.8	1.9	4.1	8.0	21.3	17.8	31.7	11.4			6	6	-
84-135	C2	15.0	40.6	44.4	1.4	2.0	1.4	3.1	7.1	19.5	21.1	28.5	7.9			2	2	-
135-150	C3	8.4	44.3	47.3	0.9	1.6	0.8	1.9	3.2	18.9	25.4	23.3	5.2			-	-	-
150-170	C4	7.6	45.0	47.4	0.5	0.9	0.6	1.9	3.7	14.2	30.8	19.1	3.9			-	-	-
170-185	C5	20.8	44.3	34.9	0.7	1.7	2.6	8.7	7.1	12.3	32.0	24.7	13.7			1	1	-
185-225	IIC6	66.1	23.2	10.7	4.0	8.2	13.9	29.9	10.1	8.8	14.4	33.6	56.0			4	4	-
225-260	IIIC7	2.1	57.4	40.5	0.1	0.3	0.3	0.7	0.7	4.1	53.3	5.3	1.4			-	-	-

Depth (cm)	6A1a Organic carbon a/ Pct.	6B1a Nitrogen Pct.	C/N	6C2b Ext. iron as Fe Pct.	Carbonate as $CaCO_3$ 6E1b <2 mm Pct.	Carbonate as $CaCO_3$ 3A1a <0.002 mm Pct.	Bulk density 4A1f 1/3 bar g/cc	Bulk density 4A1h Oven dry g/cc	Bulk density g/cc	3B2 Cm	Water content 4B1c 1/3 bar Pct.	Water content 4B2 15 bar Pct.	Water content 4C1 1/3 to 15 bar cm/cm	4D1 Extensibility COLEf cm/cm	4D1 Extensibility COLE cm/cm	pH 8C1b Saturated paste	pH 8C1c IN KCl 1:1	pH 8C1a H_2O 1:10
0-10	1.85	0.171	11		-		1.32	1.55		0.87	22.1	12.2		0.055	0.047	7.1		7.8
10-25	0.78	0.080	10		-		1.46	1.85		0.80	26.2	15.3		0.082	0.064	6.8		7.8
25-36	0.59	0.052	11		tr.		1.55	1.91		0.94	23.3	16.7		0.072	0.067	7.5		8.9
36-61	0.59	0.036	16		4		1.54	1.93		0.97	24.2	16.2		0.078	0.077	7.9		9.3
61-84	0.40	0.028	14		5		1.56	1.91		0.96	24.0	17.4		0.070	0.067	8.0		9.3
84-135	0.32				4		1.52	1.87		0.99	25.3	19.9		0.072	0.071	7.8		9.0
135-150	0.23				2		1.44	1.82		1.00	28.8	23.5		0.081	0.081	7.6		8.7
150-170	0.22				1		1.28	1.76		1.00	38.7	29.3		0.112	0.112	7.6		8.7
170-185	0.13				tr.		1.22	1.64		1.00	39.7	26.3		0.104	0.104	7.4		8.7
185-225	0.10				-		1.53	1.69		0.98	20.6	13.9		0.034	0.033	7.2		8.2
225-260	0.13				-		1.26	1.82		1.00	40.9	37.1		0.130	0.130	7.1		8.1

Depth (cm)	Extractable bases 6N2e Ca	6O2d Mg	6P2a Na	5B1a 6Q2a K	6H2a Ext. acidity	5A1a CEC NH_4OAc	5D2 Exchangeable Sodium Pct.	5E Sodium adsorption ratio	8A Water at satur., ratio Pct.	Water extract from saturated paste 8A1 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K	6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1b SO_4	8A1a Electrical conductivity mmho/cm
	meq/100 g									meq/liter								
0-10	16.8	5.6	0.4	1.9		24.6												
10-25	20.2	8.6	0.8	1.2		30.8												
25-36	28.6	8.4	1.6	0.8		32.2												
36-61	29.8	7.8	2.8	0.6		30.6		6	63.6	1.0	0.4	4.6	0.1	-	3.8	1.4	1.7	0.68
61-84	31.5	7.8	4.0	0.6		30.7		8	70.9	1.3	0.4	6.9	0.1	-	3.1	3.3	3.9	1.01
84-135	34.2	9.1	5.9	0.6		36.7		9	80.2	6.0	2.1	17.4	0.2	-	2.0	20.4	10.3	3.17
135-150	42.1	12.8	8.1	0.8		47.9		7	88.9	23.5	8.8	27.2	0.5	-	1.8	23.9	44.2	5.51
150-170	51.2	14.1	8.6	0.8		53.0		7	93.3	26.2	9.6	30.8	0.3	-	1.9	23.6	48.8	5.65
170-185	47.5	15.2	8.8	0.9		58.8		7	85.4	26.2	9.7	30.8	0.3	-	1.7	21.6	52.9	5.77
185-225	22.3	9.1	5.0	0.3		35.7		9	51.2	4.1	1.7	16.1	0.1	-	1.3	13.0	11.3	2.53
225-260	42.2	17.0	10.8	1.1		66.2		9	110.7	7.0	3.4	19.9	0.2	-	0.9	16.8	15.3 b/	3.17

Depth (cm)	Ratios to clay 8D1 CEC NH_4OAc	Ext. iron	15 bar water	Salt Pct.	6F1a Gypsum Pct.	c/ Clay fraction mineralogy 3A1 7A2 X-ray d/	7A3 DTA Pct.
0-10	0.91		0.45			MT4, MI3, KK1	
10-25	0.83		0.41				
25-36	0.80		0.42				
36-61	0.76		0.42			MT5, MI2, KK1	
61-84	0.74		0.42				
84-135	0.83		0.45			MT5, MI2, KK1	
135-150	1.01		0.50		tr.		
150-170	1.12		0.62		tr.		
170-185	1.68		0.75		tr.		
185-225	3.34		1.30				
225-260	1.63		0.92			MT5, KK1	

a/ 8.8 kg of organic carbon per square meter to a depth of 1 meter

b/ Indicator destroyed indicating nitrites

c/ Mineral code
- MT = montmorillonite
- MI = mica
- KK = kaolinite

d/ Approximate weight fraction (X-ray)
- 5 = more than half
- 4 = one-third to half
- 3 = one-fifth to one-third
- 2 = one-twentieth to one-fifth
- 1 = less than one-twentieth

CAPAY CLAY (Smooth surface) (taxadjunct) 1/
S67Calif-33-2

Location: Riverside County, California. Sec. 19, T. 4 S., R. 5 W., SBBM on aerial photo AXM 13K-2.
Date of Sampling: May 5, 1967.
Description by: B. R. Brasher. Collectors: B. R. Brasher, F. F. Peterson, K. W. Flach, and W. D. Nettleton.
Classification: Typic Chromoxerert, fine, montmorillonitic, thermic.
Vegetation: Red brome, soft chess, sawtooth goldenbush, bunchgrass, wild mustard, needlegrass, and occasional bunches of wild buckwheat. Climate: Average annual precipitation 12 inches; average annual temperature 64° F. Parent Material: Soft marine sandstone and shale. Topography: East northeast-facing, 5 percent slope, rolling uplands. Elevation: 1,770 feet. Drainage: Well drained. Remarks: Moist when described. A crack, 1 cm. in width, begins in the C1 and extends downward to near the bottom of the C2. Many 1-2 cm.^{2} ped surfaces are either slickensides or pressure faces. Rocks may be concentrated on the surface (1 to 5 cm. in diameter) especially when this soil occurs adjacent to the crusty surface Capay. This pedon is located about 200 feet north of S67Calif-33-1.

HORIZON	DESCRIPTION
A11 RSL No. 6725	0-4 in., (0-10 cm.), dark brown (10YR 4/3) light clay loam, dark brown (10YR 3/2) moist; moderate, medium, subangular blocky structure; hard (dry), friable (moist), slightly sticky and slightly plastic (wet); plentiful fine and very fine roots; common fine and many very fine pores; 13 percent gravel by volume; clear, smooth boundary.
A12 RSL No. 6726	4-10 in., (10-25 cm.), dark brown (10YR 4/3) gravelly heavy silty clay loam, very dark grayish brown (10YR 3/2) moist; moderate medium prisms breaking to moderate medium and fine subangular blocks; few pressure faces; slightly hard (dry), friable (moist), sticky and plastic (wet); few fine and plentiful very fine roots; common very fine and micro pores; 20 percent gravel by volume; clear, smooth boundary.
A13 RSL No. 6727	10-14 in., (25-36 cm.), dark brown (10YR 4/3) light clay, dark reddish brown (5YR 3/2) moist; moderate, medium, angular blocky structure; few pressure faces; slightly hard (dry), friable (moist), sticky and plastic (wet); plentiful fine and micro roots; common very fine and micro pores; 6 percent gravel by volume; gradual, wavy boundary.
A14 RSL No. 6728	14-24 in., (36-61 cm.), dark yellowish brown (10YR 4/4) silty clay, dark reddish brown (5YR 3/3) moist; weak medium, prisms breaking to moderate, medium, angular blocks; few horizontal and vertical nonintersecting slickensides; hard (dry), firm (moist), sticky and plastic (wet); plentiful fine and micro roots; common very fine and micro pores; strongly effervescent; 3 percent gravel by volume; gradual, wavy boundary.
C1ca RSL No.	24-33 in., (61-84 cm.), dark brown (10YR 4/3) light clay, with common, fine distinct dark brown (7.5YR 3/2) mottles, dark brown (7.5YR 3/2) and dark reddish brown (5YR 3/4) moist; moderate, coarse, prism breaking to

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Chromoxerert, fine, montmorillonitic, thermic

SOIL: Capay clay SOIL Nos. S64Calif-48-2 LOCATION: Solano County, California

SOIL SURVEY LABORATORY: Riverside, California LAB. Nos. 64418 - 64426

Depth (in.)	Horizon	1B1b Total Sand (2–0.05)	Total Silt (0.05–0.002)	Total Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	Int II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	Coarse fragments 1A2a > 2	2 – 19	19 – 76
		Pct. of < 2 mm														Pct.	Pct. of < 76 mm	
0-5	Ap	4.3	42.0	53.7	0.1	0.2	0.4	1.2	2.4	11.7	30.3	14.9	1.9			0		
5-21	A12	3.2	42.0	54.8	0.1	0.1	0.2	0.9	1.9	10.2	31.8	12.7	1.3			0		
21-32	A13	3.8	45.5	50.7	0.3	0.3	0.2	0.9	2.1	11.5	34.0	14.2	1.7			0		
32-40	AC	3.5	51.3	45.2	tr.	0.1	0.1	1.0	2.3	11.9	39.4	14.9	1.2			0		
40-50	C1	7.0	55.1	37.9	tr.	0.1	0.1	3.2	3.6	15.5	39.6	21.3	3.4			0		
50-62	C2	7.8	54.0	38.2	tr.	tr.	0.2	2.8	4.8	14.7	39.3	21.4	3.0			0		
62-81	C3	25.9	38.1	36.0	0.3	2.1	6.1	11.1	6.3	12.0	26.1	23.7	19.6			0		
81-88	C4	59.1	16.8	24.1	0.1	2.8	21.7	29.2	5.3	6.3	10.5	21.6	53.8			0		
88-102	IIC5	82.3	4.3	13.4	tr.	3.2	44.7	32.0	2.4	1.9	2.4	12.2	79.9			0		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEf in./in.	4D1 Extensibility COLE in./in.	pH 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-5	1.09	0.103	11	1.9	-								19.5				7.1	7.7
5-21	0.79	0.078	10	1.9	tr.		1.51	1.95		1.00		26.0	19.7	0.090	0.090		7.9	8.6
21-32	0.52	0.057	9	2.1	1		1.49	1.89		1.00		25.9	20.4	0.083	0.083		8.4	9.1
32-40	0.30			2.1	tr.								19.0				8.3	9.1
40-50	0.24			2.0	tr.		1.52	1.78		1.00		24.9	19.1	0.054	0.054		8.3	9.0
50-62	0.23			1.9	-								18.3				8.2	8.9
62-81	0.11			1.4	1		1.62	1.81		1.00		21.5	16.6	0.038	0.038		8.2	9.2
81-88	0.09			1.0	1								10.7				8.3	8.9
88-102	0.07			0.6	-								6.3				8.1	8.1

Depth (in.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1a SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-5	19.7	19.8	1.2	0.7	41.4		43.0		1.1	1.0	2.6	0.1		-	1.8	0.8		0.48
5-21	25.4	17.3	1.6	0.5	44.8		42.4		0.8	0.8	3.6	0.1		-	3.2	0.2		0.50
21-32	25.6	18.2	3.6	0.4	47.8		42.5		0.8	0.6	9.3	0.1		-	3.4	0.6		0.78
32-40	23.3	14.8	4.4	0.4	42.9		40.1		2.8	0.6	8.5	0.2		-	3.4	0.6	5.9	1.08
40-50	19.8	13.4	5.1	0.3	38.6		37.6		1.1	1.1	11.0	0.2		-	3.6	0.6	7.6	1.22
50-62	18.8	13.0	5.9	0.3	38.0		36.9		1.9	2.1	15.2	0.3		-	2.7	0.4	15.6	1.79
62-81	18.5	11.2	5.2	0.3	35.2		33.3		3.0	2.0	17.9	0.3		-	2.2	0.6	19.8	2.18
81-88	16.3	7.8	3.3	0.3	27.7		22.6		1.2	1.6	15.9	0.3		-	2.6	0.9	16.3	1.94
88-102	5.4	4.1	1.7	0.2	11.4		12.5		2.6	0.6	8.5	0.4		-	1.6	0.6	7.9	1.04

Depth (in.)	8A Water at Saturation Pct.	Exch. bases 5B3b Ca	5B3b Mg	5B1b Na	5D2 Exchangeable Na Pct.	5E SAR	Gypsum Pct.	Atterberg limits Plastic limit	Liquid limit	Plastic index
		meq./100 g.						Pct. of <0.4 mm.		
0-5	53.9			1.1	3	3				
5-21	53.7			1.4	3	4		28	49	21
21-32	55.0			3.1	7	11				
32-40	54.7			4.0	10	7				
40-50	59.8			4.4	12	10				
50-62	56.4			5.0	14	11				
62-81	52.3			4.3	13	11		14	45	31
81-88	43.1			2.6	12	13				
88-102	37.0			1.4	11	7				

CAPAY CLAY
S64Cal-48-2

Location: A pit in milo field 2½ miles NE of Elmira on Hawkins Road, 200 feet east and 50 feet north of the SW corner of Sec. 16, T6N, R1E, MDB&M, Solano County, California. Date of Sampling: October 6, 1964.

Description by: Grant Kennedy, Wells Andrews, Sam Johnson. Collectors: Reuben Nelson, Benny

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Chromoxerert, fine, montmorillonitic, thermic
SOIL Capay clay SOIL Nos. S64Calif-57-13 LOCATION Yolo County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64413 - 64417

Depth (in.)	Horizon	1B1b Total: Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Sand: Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt: 0.05–0.02	Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	Coarse fragments 1A2a: > 2 Pct.	2 – 19	19 – 76
		Size class and particle diameter (mm) 3A1 — Pct. of < 2 mm															Pct. of < 76 mm	
0-11	Ap	8.0	46.4	45.6	0.2	0.4	0.8	3.1	3.5	12.6	33.8	18.0	4.5			0		
11-18	A12	9.4	47.7	42.9	0.1	0.4	0.8	3.4	4.7	12.4	35.3	19.3	4.7			0		
18-36	A13	7.3	41.0	51.7	tr.	0.2	0.7	2.7	3.7	10.1	30.9	15.6	3.6			0		
36-49	ACca	6.4	42.8	50.8	tr.	0.2	0.5	2.3	3.4	9.9	32.9	14.9	3.0			0		
49-65+	C1	5.0	49.0	46.0	tr.	0.2	0.4	1.8	2.6	12.1	36.9	15.9	2.4			0		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content: Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLE in./in.	4D1 Extensibility COLE in./in.	pH: 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-11	1.16	0.102	11	1.7	-		1.45	1.82		1.0		24.3	16.9	0.078	0.078		7.2	7.8
11-18	0.59	0.066	9	1.8	-		1.53	1.87		1.0		23.4	17.2	0.070	0.070		7.0	7.8
18-36	0.53	0.057	9	2.1	-		1.52	1.94		1.0		25.6	19.1	0.084	0.084		7.4	8.2
36-49	0.43			2.0	tr.		1.57	1.92		1.0		24.6	19.1	0.070	0.070		8.0	8.5
49-65+	0.31			2.0	tr.		1.55	1.89		1.0		24.6	18.8	0.068	0.068		8.2	8.8

Depth (in.)	Extractable bases 5B1a: 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 8A1: 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-11	13.5	19.6	1.0	0.8	34.0		33.9		0.6	1.6	2.8	0.2		-	1.5	1.2		0.44
11-18	13.4	17.7	1.2	0.4	32.7		33.4		0.3	0.8	2.1	0.1		-	0.5	1.1		0.33
18-36	17.2	20.0	2.1	0.4	39.7		42.2		0.5	0.5	2.4	0.1		-	1.0	0.7		0.34
36-49	18.5	20.0	2.9	0.5	41.9		39.8		0.6	0.6	4.8	0.1		-	2.3	0.5		0.67
49-65+	17.2	17.0	3.6	0.4	38.2		38.0		0.5	0.6	5.8	0.1		-	2.4	1.0		0.72

Depth (in.)	8A Water at Saturation Pct.	Atterberg limits (Pct. of <0.4 mm): Plastic limit	Liquid limit	Plastic index	5D2 Exchangeable Na Pct.	5E SAR	5B1b Exchangeable Na meq/100 g	Clay mineralogy (<0.002 mm.) 3A1 X-ray 7A2: Mont.	Verm.	Mica	Kaolin.	
0-11	49.8	24	41	17	2	3	0.8					
11-18	44.0				3	3	1.1	xxxx	-	-	xx	
18-36	50.2	24	47	23	5	3	2.0					
36-49	53.4				7	6	2.7					
49-65+	50.6	22	51	29	9	8	3.3	xxxx	-	-	xx	

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

CAPAY CLAY
S64Cal-57-13

Location: Yolo County, California. A pit located 100 feet south of the NE corner of NW ¼ SW ¼ of Sec. 2, T.9N., R.1E., 10 miles west of Woodland. Photo ABB-5K-171. Date of Sampling: September 28, 1964.
Description by: C. B. Goudey. Collectors: R. E. Nelson, B. R. Brasher, G. M. Kennedy, W. F. Andrews, L. A. Bates. G. S. Staidl, C . B. Goudey.
Classification: Typic Chromoxerert, fine, montmorillonitic, thermic. 1/
Vegetation: Sudan grass. Climate: 16 inches average annual precipitation and mean annual temperature of 61°F. Parent Material: Alluvium from sedimentary rocks. Topography: Basin rim; plane; less than one percent; field was leveled (very slight cuts and fills). Elevation: 140 feet. Drainage: Runoff is slow; Permeability is slow. Soil Moisture: Profile moist throughout below 5 inches.
Remarks: (1) Truog and Thymol blue field pH determination; (2) Paired Sample No. 64 Cal. 48-2-1 through 9 ; (3) Diagnostic features- clay content greater than 30 percent, intersecting slickensides, and a Ca horizon; (4) Soil temperature-Ap-32.5°C; A12-30°C; A13-28°C; ACca-25°C; C1-23°C; (5) Mottling in A horizon is due to previous rice culture; (6) B.P.R. samples taken from the Ap, A13 and C horizons.

HORIZON	DESCRIPTION
Ap RSL 64413	0 to 11 inches, grayish brown (2.5Y 5/2) clay with common fine faint mottles of brown (7.5YR 5/4),very dark grayish brown (2.5Y 3/2) with common fine distinct mottles of brown to dark brown (7.5YR 4/4) when moist; moderate very coarse prismatic structure; hard, very firm, sticky and very plastic; plentiful micro to very fine and few fine random roots; common micro vesicular and interstitial pores; moderately alkaline (pH 8.0); clear smooth boundary.
A12 RSL 64414	11 to 18 inches, grayish brown (2.5Y 5/2) clay with common fine distinct mottles of yellowish brown (10YR 5/6), very dark grayish brown (2.5Y 3/2) with common fine distinct mottles of brown to dark brown (10YR 4/3) moist; moderate very coarse prismatic structure; very hard, very firm, sticky and very plastic; plentiful micro to very fine random roots; common micro tubular pores; mildly alkaline (pH 7.5); gradual wavy boundary.
A13 RSL 64415	18 to 36 inches, dark grayish brown (2.5Y 4/2) clay, very dark grayish brown (2.5Y 3/2) when moist; moderate very coarse prismatic structure; very hard, very firm, sticky and very plastic; few micro to very fine random roots; common micro tubular pores; many prominent intersecting slickensides; moderately alkaline (pH 8.0); clear wavy boundary.
ACca RSL 64416	36 to 49 inches, grayish brown (10YR 5/2) clay, dark grayish brown (2.5Y 4/2) with very dark grayish brown (2.5Y 3/2) when moist (probably due to organic staining along cracks); massive; very hard, very firm, sticky and very plastic; few micro to very fine random roots; common micro tubular pores; many prominent intersecting slickensides; moderately alkaline (pH 8.2), strongly effervescent, lime occurs in concretions; gradual wavy boundary.
C1 RSL 64417	49 to 64 inches, pale brown (10YR 6/3) clay, brown to dark brown (10YR 4/3); massive; very hard, very firm, sticky and plastic; few micro random roots; common micro tubular pores; many prominent intersecting slickensides; moderately alkaline (pH 8.3); effervescent disseminated lime.

1/ Mottles in this pedon resulted from rice culture and are disregarded in the classification of the pedon.

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Clear Lake clay **LOCATION** Sonoma County, California

SOIL FAMILY: Typic Pelloxerert, fine, montmorillonitic, thermic.

SOIL NOS. S61Calif-49-9 **LAB. NOS.** 61446 - 61453

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2	

CLEAR LAKE CLAY
S61Calif-49-9

Location: Sonoma County, California. Matteri Ranch. NW 1/4, SW1/4, section 26, T. 5 N., R. 7 W., projected. From the junction of East Washington Street with Ely Road, 4000 feet due southeast on Ely Road, thence 880 feet due southwest along Homestead Road and into pastured field, 30 feet northwest of field boundary fence (on southerly side of field). (Site spotted on photo No. CSH-1BB-163). Date of Sampling: June 26, 1961.
Collectors: Reuben Nelson, Leo C. Klameth, V. C. Miller, G. Lambert, J. H. Kashiwagi, F. Graham.
Classification: **Typic Pelloxerert, fine, montmorillonitic, thermic.**

Vegetation: Annual grasses and forbs (before cultivation). Use: Oat hay. Climate: About 25 inches rainfall, average temperature 57.5° F., January temperature 46.7° F., July temperature 66.3° F. Parent Material: Mixed alluvium. Elevation: About 43 feet. Relief: Flat basin, on 1 percent slope. Drainage: Poorly drained, slow runoff, very slow permeability. Biological Activity: No worm or rodent activity apparent at this time of year.

HORIZON	DESCRIPTION
Ap1 RSL No. 61446	0 to 1 inches, dark gray (N 4/) clay, very dark gray (10YR 3/1) when moist; very fine and fine granular structure; very hard when dry, extremely firm when moist, sticky and very plastic when wet; abundant very fine and fine roots; medium acid, pH 6.0; abrupt, wavy boundary.
Ap2 * RSL No. 61447	1 to 7 inches, very dark gray (N 3/) clay, black (10YR 2/1) with dark grayish brown (2.5Y 4/2) blotches when moist; massive; very hard when dry, extremely firm when moist, sticky and very plastic when wet; abundant very fine and fine roots; many very fine and fine tubular and interstitial pores; medium acid, pH 6.0; clear, wavy boundary.
A11g RSL No. 61448	7 to 25 inches, very dark gray (N 3/) clay with common fine distinct mottles of gray (N 6/) when dry; black (N 2/) with a few scattered blotches of dark grayish brown when moist; hard when dry, extremely firm when moist, sticky and very plastic when wet; abundant very fine and fine roots; common very fine tubular pores; neutral, pH 7.0; few slickensides; gradual, smooth boundary; few segregated lime blotches.
A12g RSL No. 61449	25 to 40 inches, dark gray (N 4/) clay with common fine distinct light gray (2.5Y 7/2) mottles when dry, black (N 2/) with common fine distinct light gray mottles of lime; massive; extremely hard when dry, extremely firm when moist, sticky and very plastic when wet; plentiful very fine roots; common very fine and fine tubular concretions; evidence of vertical cracking from surface down, streaks of lighter color in old cracks; moderately alkaline, pH 8.0; gradual, smooth boundary; lime is segregated.
ACg RSL No. 61450	40 to 48 inches, gray (N 5/) clay with many medium distinct light gray (N 7/) lime blotches, very dark gray (10YR 3/1) when moist; massive; extremely hard when dry, extremely firm when moist, sticky and very plastic when wet; few fine and very fine roots; common very fine and fine tubular pores; slickensides; segregated lime blotches; fragments of black A horizon churned in throughout; mildly alkaline, pH 7.5 - 8.0; gradual, wavy boundary.
C1g RSL No. 61451	48 to 54 inches, light olive gray (5Y 6/2) clay with many medium distinct white (5Y 8/1) lime blotches, olive gray (5Y 4/2) when moist; massive; very hard when dry, very firm when moist; sticky and very plastic when wet; few very fine and fine roots; common very fine and fine interstitial and tubular pores; slickensides; segregated lime blotches oriented horizontally; moderately alkaline, pH 8.0; clear, wavy boundary.
C2g *	54 to 77 inches, light olive gray (5Y 6/2) clay, dark grayish brown (2.5Y 4/2) [illegible]

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Clear Lake clay **LOCATION** Sonoma County, California

SOIL FAMILY: Typic Pelloxerert, fine, montmorillonitic, thermic.

SOIL NOS. S61Calif-49-10 **LAB. NOS.** 61454 - 61462

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE	COARSE	MEDIUM	FINE	VERY FINE						

CLEAR LAKE CLAY
S61Calif-49-10

Location: Sonoma County, California. Dangers Ranch - Ely Road, 1/2 mile southeast of junction with Corona Road. NW1/4, NW1/4, section 21, T. 5 N., R. 7 W., projected. Date of Sampling: June 27, 1961.
Collectors: Reuben Nelson, Leo C. Klameth, V. C. Miller, G. Lambert, J. H. Kashiwagi, F. Graham.
Classification: **Typic Pelloxerert, fine, montmorillonitic, thermic.**

Vegetation: Annual grasses, forbs and scattered oak (before cultivation). Use: Oat hay. Climate: About 25 inches rainfall, average temperature 57.5° F., January temperature 46.7° F., July temperature 66.3° F. Parent Material: Mixed alluvium. Elevation: About 39 feet. Relief: Flat basin on 1 percent slope. Drainage: Poorly drained, slow runoff, very slow permeability. Biological Activity: No worm or rodent activitiy apparent at this time of the year.

HORIZON	DESCRIPTION
Ap1 RSL No. 61454	0 to 2 inches, dark gray (N 4/) clay, very dark gray (10YR 3/1) when moist; fine and medium granular structure; extremely hard when dry, very firm when moist, sticky and plastic when wet; abundant very fine and fine roots; medium acid, pH 6.0; clear, wavy boundary.
Ap2 * RSL No. 61455	2 to 8 inches, dark gray (N 4/) clay, very dark gray (10YR 3/1) when moist; massive breaking to coarse subangular blocks; very hard when dry, very firm when moist, sticky and plastic when wet; abundant very fine and fine roots; many fine and micro pores; medium acid, pH 6.0; clear, wavy boundary.
A11g RSL No. 61456	8 to 25 inches, dark gray (N 4/) clay, black (10YR 2/1) when moist, massive; extremely hard when dry, extremely firm when moist, sticky and plastic when wet; abundant very fine and fine roots; common very fine and fine tubular pores; slickensides; medium acid, pH 6.0; gradual, smooth boundary.
A12g RSL No. 61457	25 to 39 inches, dark gray (N 4/) clay with common fine distinct light gray (2.5Y 7/2) lime spots when dry; black (10YR 2/1) when moist, massive; extremely hard when dry, extremely firm when moist, sticky and plastic when wet; abundant very fine and fine roots; common very fine tubular pores; slickensides; quarter inch vertical cracks; part of this horizon has lime blotches; moderately alkaline, pH 8.0; gradual, smooth boundary.
AC[illegible]	39 to 46 inches, dark gray (N 4/) and gray (N 5/) clay with common fine distinct [illegible]

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Diablo clay (taxadjunct) LOCATION San Luis Obispo Co.

SOIL FAMILY: Typic Pelloxerert, fine, montmorillonitic, thermic.

SOIL NOS. S55Cal. 40-2 LAB. NOS. 55455-458

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE SAND 2-1	COARSE SAND 1-0.5	MEDIUM SAND 0.5-0.25	FINE SAND 0.25-0.10	VERY FINE SAND 0.10-0.05	SILT 0.05-0.002	CLAY < 0.002	0.2-0.02	0.02-0.002	> 2	
0- 1	Ap1	2.2	1.6	1.0	4.3	2.3	31.2	57.4	13.1	23.2	5	c
1- 4	Ap2	2.7	1.2	0.9	4.1	2.4	30.3	58.4	13.6	21.9	4	c

DIABLO CLAY (taxadjunct) 1/
S55Calif-40-2

Location: About 1.31 miles N.N.E. of farm buildings on George Mello farm, San Luis Obispo County, California. On aerial photograph AXH-4F-148.

Collectors: W. M. Johnson, Rudolph Ulrich, Lewis Leifer, W. G. Harper, L. R. Wohletz, J. M. Williams. Samples taken September 7, 1955. Description by W. M. Johnson.

Classification: Typic Pelloxerert, fine, montmorillonitic, thermic.

Elevation: About 330 feet.

Vegetation: Cultivated field. In small-grain stubble at time of sampling.

Relief: Hilly, eroded, residual upland. Slope about 5 percent, plane, toward the north.

Drainage: Well drained. Runoff is medium; permeability is slow.

Parent material: Weathered shale of the San Luis Obispo formation.

Remarks: Classified as Grumusol. Depth to bedrock varies from about 24 to about 46 inches in this field. When dry, the soil surface has cracks in a polygonal pattern, 12 to 18 inches in diameter, with cracks about 1½ inches wide and 10 to 20 inches deep. pH's were measured in the field with Truog Triplex indicator.

Sample numbers: S55Cal-40-2-1 through 4.

HORIZON		DESCRIPTION
Ap1	0-1"	Very dark gray to dark gray (10YR 3.5/1, dry; 3/1, moist) clay. Strong coarse, medium and fine granular. Extremely hard, very firm, very plastic, very sticky. pH 6.5. Abrupt, smooth boundary.
Ap2	1-4"	Very dark gray (10YR 3/1. moist) clay. Moderate medium and

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Pelloxerert, fine, montmorillonitic, thermic

SOIL Willows clay SOIL Nos. S64Calif-57-14 LOCATION Yolo County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64548 - 64556

Depth (in.)	Horizon	1B1b Total: Sand (2–0.05)	Total: Silt (0.05–0.002)	Total: Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Noncarbonate clay	Coarse fragments 1A2a: > 2	2 – 19	19 – 76
		Pct. of < 2 mm														Pct.	Pct. of < 76 mm	
0-9	Ap	2.2	42.1	55.7	tr.	0.3	0.3	0.6	1.0	8.1	34.0	9.4	1.2			0		
9-12	A12	2.1	41.4	56.5	0.1	0.3	0.2	0.6	0.9	6.9	34.5	8.1	1.2			0		
12-19	A13	2.2	42.0	55.8	0.1	0.3	0.3	0.7	0.8	5.8	36.2	7.0	1.4			0		
19-30	AC	1.6	50.0	48.4	0.2	0.3	0.2	0.3	0.6	4.0	46.0	4.8	1.0			0		
30-40	C1	4.3	55.0	40.7	0.6	1.1	0.5	0.8	1.3	5.8	49.2	7.5	3.0			0		
40-55	C2	7.9	58.2	33.9	0.6	0.9	0.4	0.8	5.2	15.5	42.7	21.2	2.7			0		
55-66	C3	4.7	51.3	44.0	0.5	1.0	0.6	0.9	1.7	3.6	47.7	5.8	3.0			0		
66-74	C4	2.3	39.9	57.8	0.1	0.4	0.4	0.6	0.8	2.1	37.8	3.2	1.5			0		
74-82	C5	7.8	48.3	43.9	0.2	0.3	0.4	2.7	4.2	8.0	40.3	14.1	3.6			0		

Depth (in.)	6A1a Organic carbon	6B1a Nitrogen	C/N	6C2a Ext. Iron as Fe	6E1b Carbonate as $CaCO_3$	6E1c Clay size carbonate	Bulk density: 4A1f 1/3 bar	Bulk density: 4A1h Oven dry		3B2 Cm	Water content	Water content: 4B1c 1/3 bar	4B2 15 bar	4D1 Extensibility COLEf	4D1 Extensibility COLE	pH: 8C1b Saturated Paste	pH: 8C1a H_2O 1:1	pH: 8C1a H_2O 1:10
	Pct.	Pct.		Pct.	Pct.	Pct.	g/cc	g/cc	g/cc		Pct.	Pct.	Pct.	in./in.	in./in.			
0-9	0.55	0.066	8	1.6	tr.		1.59	2.08		1.00		24.5	18.6	0.095	0.095		8.3	9.0
9-12	0.74	0.065	11	1.6	tr.		1.58	2.06		1.00		25.6	18.7	0.092	0.092		8.3	9.2
12-19	0.63	0.062	10	1.4	2		1.54	2.02		1.00		26.7	19.3	0.094	0.094		8.6	9.6
19-30	0.36	0.049	7	1.6	2		1.41	1.81		1.00		32.5	25.6	0.087	0.087		8.9	9.9
30-40	0.28			1.7	7	tr.	1.43	1.72		1.00		31.5	22.8	0.063	0.063		9.1	9.9
40-55	0.26			1.6	4		1.51	1.61		1.00		24.3	22.8	0.022	0.022		9.0	9.9
55-66	0.30			1.8	5	tr.	1.37	1.61		1.00		32.9	24.1	0.054	0.054		8.9	9.9
66-74	0.38			1.5	1		1.34	1.69		1.00		36.1	25.8	0.081	0.081		8.7	9.7
74-82	0.23			1.6	tr.								20.8				8.6	9.5

Depth (in.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	5B1a 6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 8A1: 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	6L1a SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-9	16.7	19.0	4.9	0.7	41.3		36.6		2.1	2.6	13.8	0.2		-	3.2	4.0	12.5	1.98
9-12	16.9	20.0	6.3	0.8	44.0		38.0		1.0	1.6	12.8	tr.		-	3.0	3.1	9.9	1.67
12-19	19.8	19.5	9.4	0.8	49.5		34.6		0.6	1.0	18.6	0.1		-	5.2	3.7	11.3	2.05
19-30	9.4	16.5	14.8	0.4	41.1		30.4		0.4	0.3	19.4	0.1		1.0	4.4	3.9	10.4	2.06
30-40	6.7	13.4	16.9	0.5	37.5		26.7		0.5	0.6	25.6	0.1		1.8	4.3	5.7	15.8	1.78
40-55	7.2	12.8	14.0	0.2	34.2		25.3		0.5	0.7	30.0	0.2		1.6	3.9	7.6	19.6	3.29
55-66	8.0	15.6	19.0	0.3	42.9		32.4		0.6	1.0	31.0	0.1		0.6	4.2	9.2	18.7	3.26
66-74	11.6	19.6	19.3	0.4	50.9		38.6		0.4	0.8	25.1	0.1		-	4.1	7.9	15.1	2.81
74-82	7.6	15.5	14.5	0.4	38.0		31.9		1.0	0.2	23.7	0.1		-	3.6	7.1	14.0	2.58

8A Water | Exch. bases 5B3b | 5B2b | 5B1b | 5D2 Gypsum | 5E | Clay mineralogy (<0.002 mm. 5A1)

WILLOWS CLAY
S64Calif-57-14

Location: Yolo County, California. A pit located about 150 feet northeast of the southwest corner of Sec. 32, T.12N., R.1E. Area located about 3 miles north of Zamora. Photo ABB-42T-23. Date of Sampling: September 30, 1964.
Description by: George J. Staidl. Collectors: R. E. Nelson, B. R. Brasher, G. M. Kennedy, W. F. Andrews, L. A. Bates, G. J. Staidl, C. B. Goudey.
Classification: Typic Pelloxerert, fine, montmorillonitic, thermic.
Vegetation: Fallow at the time of sampling. Climate: Average annual precipitation is 15 inches; mean annual temperature 62°F. Parent Material: Fine textured alluvium from sedimentary rocks. Topography: Basin; plane; less than one percent slope. Elevation: 28 feet above sea level. Drainage: Runoff is slow; permeability is slow; drainage class is poorly drained altered to somewhat poorly drained; ground water table was at 66 inches. Soil Moisture: The profile was moist below 4 inches.
Remarks: (1) Thymol blue used in field pH determinations. (2) Paired Sample No. 64 Cal. 57-15. (3) Diagnostic horizon - Mollic epipedon. (4) More than 30 percent clay; soil too wet to observe slickensides. (5) Soil temperature at about 10"-27°C; 16"-22.5°C.; 25"-22.0°C.; 35"-21.5°C.; 48"-21.0°C.; 60"-21.0°C.; 70"-20.5°C.; 78"-20.5°C. (6) Soil temperature taken in C5 horizon or at 78 inches was taken in the water. (7) The A12 horizon 9-12 inches is probably a plow pan. (8) The A13 (12-19 inches) and AC (19-30 inches) horizons have some fine manganese shot.

HORIZON	DESCRIPTION
Ap RSL 64557	0 to 9 inches, gray (5Y 5/1) clay with common fine distinct mottles of brown to dark brown (7.5YR 4/4), very dark gray (5Y 3/1) and olive gray (5Y 4/2) with common fine distinct mottles of brown to dark brown (10YR 4/3) moist; moderate coarse prismatic structure; hard, extremely firm, sticky and very plastic; very few, very fine and fine random roots; common micro tubular pores; moderately alkaline (pH 8.3); clear wavy boundary.
A12 RSL 64558	9 to 12 inches, gray (5Y 5/1) clay with few, fine distinct yellowish brown (10YR 5/8) mottles, very dark gray (5Y 3/1) and olive gray (5Y 4/2) with few fine distinct brown to dark brown (10YR 4/3) mottles moist; moderate coarse prismatic structure; hard, extremely firm, sticky and very plastic; very few, very fine to fine random roots; common micro tubular pores; very slightly effervescent disseminated lime; strongly alkaline (pH 8.6); clear wavy boundary.
A13 RSL 64559	12 to 19 inches, gray (5Y 5/1) clay with few, fine distinct yellowish brown (10YR 5/8) mottles, olive gray (5Y 4/2) with few, fine distinct brown to dark brown (10YR 4/3) mottles moist; massive; extremely hard, very firm, sticky and very plastic; very few, very fine to fine random roots; common micro tubular pores; strongly effervescent disseminated and segregated lime in soft masses; strongly alkaline (pH 8.8); gradual wavy boundary.
AC RSL 64560	19 to 30 inches, light olive gray (5Y 6/2) clay, with common fine distinct yellowish brown (10YR 5/6) mottles, dark grayish brown (2.5Y 4/2) with common fine distinct dark yellowish brown (10YR 4/4) mottles moist; massive; extremely hard, very firm, sticky and plastic; very few, very fine to fine random roots; common micro and few very fine tubular pores; strongly effervescent disseminated lime, strongly alkaline (pH 8.8); gradual wavy boundary.
C1 RSL 64561	30 to 40 inches, mottled light olive gray (5Y 6/2), light yellowish brown (2.5Y 6/4), and yellowish brown (10YR 5/6) clay, mottled olive gray (5Y 5/2), olive (5Y 4/3), and dark grayish brown (2.5Y 4/2) moist; massive; very hard, very firm, sticky and very plastic; common micro and few very fine tubular pores; strongly effervescent disseminated lime; strongly alkaline (pH 8.8); diffuse boundary.
C2 RSL 64562	40 to 55 inches, mottled olive gray (5Y 5/2), light yellowish brown (2.5Y 6/4) and strong brown (7.5YR 5/8) clay, mottled olive (5Y 4/4), dark grayish brown (2.5Y 4/2) and brown to dark brown (10YR 4/3) moist; massive; hard, firm, sticky and very plastic; common micro and few very fine tubular pores; strongly effervescent disseminated lime; strongly alkaline (pH 8.8); gradual irregular boundary.
C3 RSL 64563	55 to 66 inches, mottled pale olive (5Y 6/4), light brownish gray (2.5Y 6/2) and yellowish brown (10YR 5/6) clay, mottled dark grayish brown (2.5Y 4/2) and brown to dark brown (7.5YR 4/4) moist; massive; hard, firm, sticky and very plastic; many micro and few fine tubular pores; strongly effervescent disseminated lime; strongly alkaline

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Typic Pelloxerert, very-fine, montmorillonitic, thermic

SOIL Willows clay (taxadjunct) SOIL Nos. S64Calif-57-15 LOCATION Yolo County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 64557 - 64563

Depth (in.)	Horizon	1B1b Total: Sand (2–0.05)	Silt (0.05–0.002)	Clay (< 0.002)	Sand: Very coarse (2–1)	Coarse (1–0.5)	Medium (0.5–0.25)	Fine (0.25–0.1)	Very fine (0.1–0.05)	Silt: 0.05–0.02	Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	Coarse fragments 1A2a: > 2 Pct.	2 – 19	19 – 76
		Pct. of < 2 mm														Pct.	Pct. of < 76 mm	
0-4	Ap	1.0	36.3	62.7	0.1	0.1	0.1	0.3	0.4	2.8	33.5	3.5	0.6			0		
4-13	A12	1.0	34.3	64.7	tr.	0.1	0.1	0.4	0.4	2.1	32.2	2.7	0.6			0		
13-28	A13	1.6	35.2	63.2	tr.	0.2	0.3	0.6	0.5	2.1	33.1	2.9	1.1			0		
28-38	A14	1.9	36.5	61.6	tr.	0.3	0.3	0.6	0.7	2.6	33.9	3.6	1.2			0		
38-48	AC	1.8	38.2	60.0	0.1	0.3	0.3	0.5	0.6	3.4	34.8	4.3	1.2			0		
48-61	C1	2.2	40.1	57.7	0.1	0.4	0.3	0.6	0.8	4.9	35.2	6.0	1.4			0		
61-72	C2	3.3	41.0	55.7	0.5	0.6	0.4	0.6	1.2	6.4	34.6	7.9	2.1			0		

Depth (in.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Bulk density: 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm		Water content: 4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLE in./in.	4D1 Extensibility COLE in./in.	pH: 8C1b Saturated Paste	8C1a H_2O 1:1	8C1a H_2O 1:10
0-4	1.92	0.162	12	1.9	-								21.3				6.6	7.1
4-13	1.34	0.121	11	1.9	-		1.44	1.90		1.00		28.5	21.5	0.099	0.099		6.7	7.6
13-28	0.86	0.077	11	1.7	1		1.54	2.00		1.00		26.6	21.0	0.091	0.091		8.4	9.5
28-38	0.73	0.067	11	1.7	1		1.45	1.94		1.00		28.5	21.3	0.103	0.103		8.7	9.7
38-48	0.61	0.059	10	1.8	1		1.45	1.93		1.00		27.8	21.7	0.100	0.100		8.5	9.7
48-61	0.48			1.6	2		1.49	1.94		1.00		27.8	20.1	0.092	0.092		8.8	9.7
61-72	0.40			1.6	2		1.53	1.97		1.00		25.5	19.3	0.089	0.089		8.6	9.6

Depth (in.)	Extractable bases 6N2a Ca	6O2a Mg	6P2a Na	6Q2a K	5B1a Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A2a NaOAc	5A1a NH_4OAc	Water extract from saturated paste 6N1b Ca	6O1b Mg	6P1a Na	6Q1a K		6T1a CO_3	6J1a HCO_3	6K1a Cl	6L1a SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-4	17.1	24.6	1.1	1.4	44.2		47.3		2.6	4.7	3.4	0.4		-	1.4	3.0	5.8	1.04
4-13	17.2	25.0	1.2	0.9	44.3		46.9		0.9	1.4	2.6	0.1		-	1.0	0.8	2.7	0.50
13-28	16.9	28.7	6.0	0.5	52.1		42.4		0.3	0.9	9.9	0.1		-	6.1	0.5	2.8	0.96
28-38	15.0	27.7	8.9	0.5	52.1		40.1		0.4	0.8	13.1	tr.		0.6	5.2	1.4	5.7*	1.34
38-48	14.8	27.1	9.4	0.5	51.8		38.2		0.6	0.7	13.0	0.1		0.8	5.1	1.0	6.0*	1.35
48-61	12.4	26.2	7.5	0.5	46.6		37.6		0.5	0.6	13.4	tr.		-	5.1	2.8	5.2*	1.36
61-72	12.5	25.7	8.2	0.5	46.9		35.4		0.4	0.6	13.8	tr.		-	4.2	2.3	6.6	1.46

Depth (in.)	8A Water at Saturation Pct.	Atterberg limits: Plastic limit	Liquid limit	Plastic index	5D2 Exchangeable Na Pct.	5E SAR	5B1b Exchangeable Na meq/100 g	Clay mineralogy (<0.002 mm.) 7A2: Mont.	Verm.	Mica	Kaolin.	
		Pct. of <0.4 mm.						X-ray				
0-4	73.9	30	68	38	2	2	0.8					
4-13	72.7				2	2	1.0					
13-28	71.4	24	65	41	12	13	5.3	xxxx	-	tr.	xx	
28-38	86.3				19	17	7.8					
38-48	89.9				21	16	8.2					
48-61	80.9				17	18	6.4					
61-72	82.4	21	66	45	20	19	7.1	xxxx		tr.	xx	

* Nitrites detected

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

WILLOWS CLAY (taxadjunct) 1/
S64Calif-57-15

Location: Yolo County, California. A pit located in the center of southwest ¼ of Sec. 31. T.10N., R.2E., 3 1/4 miles east of Woodland. Photo ABB-41T-68. Date of Sampling: October 2, 1964.

Description by: C. B. Goudey Collectors: R. E. Nelson, B. R. Brasher, G. M. Kennedy, W. F. Andrews, L. A. Bates, G. J. Staidl, C. B. Goudey.

Classification: Typic Pelloxerert, very-fine, montmorillonitic, thermic.

Vegetation: Safflower stubble. Climate: 17.3 inches average annual precipitation and mean annual temperature of 61°F. Parent Material: Mixed fine alluvium. Topography: Basin; plane; less than one percent. Elevation: 24 feet. Drainage: Runoff-slow; permeability-slow: altered drainage class (Soil formed under poor drainage, presently imperfectly drained). Soil Moisture: Profile was moist throughout below 13 inches.

Remarks: (1) Truog and Thymol Blue field pH determinations. (2) Paired profile 64 Cal. 57-14. (3) Diagnostic features -- Mollic epipedon; natric horizon; profile more than 30 percent clay; many prominent intersecting slickensides. (4) Soil temperature-Ap-19°C, A12-20°C, A13-20°C, A14-19.7°C, AC-19.5°C, C1-19°C, C2-19°C. (5) BPR samples taken from the Ap, A13, and C2 horizons. (6) Very few pores in upper portion of A12 horizon. It has the appearance of a plow pan. (7) The C1 and C2 horizons were too moist to show structure in place but break out in the structure described.

HORIZON	DESCRIPTION
Ap RSL 64548	0 to 4 inches, gray (5Y 5/1) clay with many fine distinct strong brown (7.5YR 5/6) mottles, very dark gray (5Y 3/1) with many fine distinct yellowish brown 10YR 5/6) mottles when moist; granular structure; extremely hard, very firm, sticky and very plastic; plentiful micro and very fine roots; many micro tubular pores; neutral (pH 7.0); abrupt smooth boundary.
A12 RSL 64549	4 to 13 inches, gray (5Y 5/1) clay with many fine distinct strong brown (7.5YR 5/6) mottles, very dark gray (5Y 3/1) with many fine distinct yellowish brown (10YR 5/6) mottles when moist: strong very coarse prismatic structure; extremely hard. very firm, sticky and very plastic; abundant micro and very fine exped and plentiful micro inped roots; few micro tubular pores: many prominent intersecting slickensides: mildly alkaline (pH 7.5); clear smooth boundary.
A13 RSL 64550	13 to 28 inches, olive gray (5Y 4/2) clay, very dark gray (5Y 3/1) when moist; strong very coarse prismatic structure; very hard, very firm, sticky and very plastic; plentiful micro roots; many micro and common very fine tubular pores; many prominent intersecting slickensides; slightly effervescent segregated lime in soft masses: strongly alkaline (pH 8.8); diffuse boundary.
RSL 64551	coarse prismatic structure; very hard, very firm, sticky and very plastic; few micro roots; many micro and few fine tubular pores; many prominent intersecting slickensides; slightly effervescent disseminated lime; strongly alkaline (pH 9.0); clear smooth

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Chromic Pelloxerert, fine, montmorillonitic, mesic

SOIL Pit silty clay (taxadjunct) SOIL Nos. S54Calif-18-3 LOCATION Lassen County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 54305 - 54310

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)	3A1		2A2 Coarse fragments > 2 Pct.	2 - 19	19-76
		Pct. of < 2 mm															Pct. of < 76 mm	
0-6	A1p	7.6	44.6	47.8	0.6	1.2	0.8	2.7	2.3	6.8	37.8		5.3			0		
6-21	A12	5.5	32.4	62.1	0.0	0.4	0.3	1.8	3.0	8.4	24.0		2.5			0		
21-36	B2	6.5	29.0	64.5	0.6	0.8	1.0	2.2	1.9	15.2	13.8		4.6			0		
36-44	B3	25.8	30.6	43.6	0.0	3.1	4.0	7.3	11.4	12.3	18.3		14.4			0		
44-60	C1	21.7	39.4	38.9	6.7	3.3	1.9	3.3	6.5	13.8	25.6		15.2			0		
60-94	C2	47.4	41.6	11.0	0.1	0.5	0.9	17.7	28.2	19.1	22.5		19.2			0		

Depth (In.)	6A1a Organic carbon Pct	6B1a Nitrogen Pct.	C/N	6E1b Carbonate as $CaCO_3$ Pct.	Bulk density g/cc	g/cc	g/cc	Water content Pct.	Pct.	4B2 15 bar Pct.			pH 8C1b Sat. Paste	8C1a 1:10
0-6	2.84	0.195	14.6	-						31.3			7.8	8.3
6-21	1.15	0.071	16.2	1						32.6			8.4	9.2
21-36	0.82	0.056	14.6	2						38.5			8.5	9.3
36-44	0.18			6						25.4			7.2	9.2
44-60	0.12			2						25.3			8.0	9.2
60-94	0.05			1						15.2			7.9	8.6

Depth (In.)	Extractable bases: 5B1a Ca 6N2b	5B1a Mg 6O2c	5B1b Exch. Na 6P2a	5B1b Exch. K 6Q2a	Sum of bases	5A2a CEC NaOAc			Water extract from saturated paste: 6N1a 6O1a Ca + Mg		6P1a Na	6Q1a K		CO_3	HCO_3	8A1 Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0-6	20.4	18.4	20.3	3.2	62.3	62.6			46.1		235.0	2.8						16.30
6-21	22.5	18.0	18.3	1.2	60.0	57.2			0.7		35.5	0.4						3.63
21-36	23.0	27.2	26.0	1.2	77.4	66.6			0.6		14.0	0.1						1.38
36-44	21.9	22.6	11.1	0.7	56.3	44.9			0.4		10.6	0.1						1.05
44-60	34.9	22.5	8.1	0.7	66.2	49.8			0.3		8.7	0.1						0.94
60-94	23.3	13.2	2.0	0.4	38.9	33.1			1.6		4.2	0.1						0.65

Depth (In.)	8A Water at Saturation Pct.	5D1 Exchangeable Na Pct.		6F1a Gypsum Pct.		
0-6	96.7	32		tr.		
6-21	158.3	32		-		
21-36	219.6	39		-		
36-44	74.9	25		-		
44-60	70.4	16		-		
60-94	45.3	6		-		

PIT SILTY CLAY (taxadjunct) 1/
S54Calif-18-3

Location: Lassen County, California, in Honey Lake Valley SCD on John Theodore ranch, 1/4 mile N of SW corner, section 29, T. 29 N., R. 15 E., and 75 feet east of road, about 5 miles E of Wendel.
Date of Sampling: October 19, 1954.
Collectors: S. B. Johnson, R. E. Nelson, S. Porter, R. C. Roberts and L. R. Wohletz.
Classification: Chromic Pelloxerert, fine, montmorillonitic, mesic.
Vegetation: The area where sample was taken had been plowed or disked and was in grasses and weeds such as sweet clover, saltgrass, rye grass, and brassia. Virgin area across the road was a wet meadow and in grasses with some big greasewood and rabbitbrush. Climate: At Susanville, 18 miles west of the sample site and 200 feet higher elevation, the January average temperature is 30.7°F, the July average is 69.5° F, and the growing season is 142 days. Average annual precipitation for a 30-year period is 17.8 inches divided as follows, January 3.64, February 2.66, March 2.64, April 1.05, May 0.84, June 0.51, July 0.22, August 0.19, September 0.50, October 1.01, November 1.79 and December 2.77. The average annual precipitation at the sample site is probably 11 to 13 inches. The average snowfall is slightly less than at Susanville, which is 71 inches. Parent Material: Stratified lake-laid sediments mostly fine-texture of "Lahontan" age. Relief: Nearly level basin floor with many winding old abandoned stream channels or old slough ways. Elevation: 4000 feet. Drainage: Slow external drainage, and slow drainage through the soil after it becomes saturated; water table at 3 to 5 feet, and the site may be flooded at times. Biological activity: Little or no evidence of macroscopic activity.

HORIZON	DESCRIPTION
A1p RSL No. 54305	0 to 6 inches, dark gray (10YR 4/1, very dark gray 10YR 3/1 moist) silty clay which is plastic and sticky when wet, firm when moist and hard when dry. This layer is light weight and appears to be high in organic matter and breaks into medium subangular blocky units. This layer has considerable visible salt and a thin layer covers the surface in places. Saltgrass roots are very numerous in this horizon. It has a pH of 8.6 and rather abruptly changes to layer below.
A12 RSL No. 54306	6 to 21 inches, dark gray (10YR 4/1, very dark gray 10YR 3/1 moist) moderate medium prismatic heavy clay which is very plastic and very sticky when wet and very hard when dry. The prisms readily break to fine angular blocky aggregates. This layer has about 10 percent of the surface area occupied by hard light brownish gray (10YR 6/2, grayish brown 10YR 5/2 moist) lime concretions about 1/8 inch across. When dry this layer has cracks about 1 inch wide every 4 to 8 inches. The pH is 8.9, and this layer changes abruptly to layer below.
B2 RSL No. 54307	21 to 36 inches, very dark gray (10YR 3/1, black 2/1 moist) very plastic sticky clay that has lots of silt. This layer is extremely hard when dry and the structure is weak very coarse prismatic that very readily breaks into strong fine angular blocky aggregates. The prisms have very shiny polished sides like slickensides. This layer has clay flows around the structural aggregates, and about 8 percent of the surface area is occupied by very hard grayish brown (10YR 5/2, dark grayish brown 10YR 4/2 moist) irregular shaped lime concretions. The pH is 8.9. When dry this layer has wide cracks about 8 inches apart. It gradually changes to layer below.
B3 RSL No. 54308	36 to 44 inches, grayish brown (10YR 5/2, very dark grayish brown 2.5Y 3/2 moist) strong medium angular blocky silty clay loam which breaks very readily into finer angular aggregates. All aggregates are colloidal coated. This layer is not nearly as hard as layer above and is less plastic. It has a pH of 9.0 and roots are quite numerous from the surface to the bottom of this layer. It has some lime concretions and considerable material from layer above that occurs as mottlings.
C1 RSL No. 54309	44 to 60 inches, light brownish gray (10YR 6/2, dark grayish brown 2.5Y 4/2 moist) weak medium angular blocky silty clay loam which is plastic when wet and friable when moist and slightly calcareous. The pH is 8.4.
C2 RSL No. 54310	60 to 94 inches, saturated stratified loam, silt loam, and sandy loam, all noncalcareous and dark grayish brown (10YR 4/2) when wet. This horizon is loose and friable and only slightly plastic. It has some mica and is noncalcareous. pH 8.2.

1/ The Pit series is in a fine, montmorillonitic, mesic family of Typic Chromoxererts. This pedon has gray colors throughout the upper 30 cm.

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Chromic Pellixerert, fine, montmorillonitic, thermic
SOIL Bosanko clay SOIL Nos. S64Calif-37-8 LOCATION San Diego County, California
SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 6450 - 6455

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	3A1 Int. II (0.2-0.02)	(2-0.1)	* Clay <.002	** Clay <.002	1/ Coarse fragments 1A2a ≥ 2 Pct.	2 - 19 (Pct. of < 76 mm)	19-76 (Pct. of < 76 mm)
		Pct. of < 2 mm																
0- 4	Ap	38.5	25.5	36.0	2.4	6.6	6.4	13.9	9.1	11.7	13.8	28.9	29.4	35.4	35.2	1	1	0
4-18	A11	33.3	28.8	37.9	2.4	6.2	5.9	11.5	7.3	15.7	13.1	29.6	26.0	39.1	38.4	2	2	0
18-22	A12	35.1	26.6	38.3	2.5	6.2	5.3	12.5	8.6	15.9	10.7	32.0	26.5	34.4	34.2	1	1	0
22-28	ACca	50.7	22.9	26.4	1.6	6.6	8.1	20.4	14.0	11.2	11.7	36.9	36.7	24.5	24.3	1	1	0
28-45	C1	79.6	11.1	9.3	7.0	19.6	14.3	29.3	9.4	6.6	4.5	25.8	70.2	6.7	5.9	12	12	0
45-57+	C2	81.5	11.3	7.2	5.3	19.3	14.2	32.0	10.7	6.8	4.5	29.0	70.8	5.4	4.5	17	17	0

Depth (In.)	6A1a * Organic carbon Pct.	6B1a* Nitrogen Pct.	C/N	* 6C2a Ext. Iron as Fe Pct.	6E1b ** Carbonate as $CaCO_3$ Pct.	Bulk density 4A1f 1/3 bar g/cc	Bulk density 4A1h Oven dry g/cc	g/cc	3B2 C'	Water content Pct.	Water content 4B1c 1/3 bar Pct.	Water content 4B2* 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH 8C1b Saturated Paste	pH 8C1a H_2O 1:1	pH 8C1a H_2O 1:10
0- 4	1.59	0.131	12	0.6	-							15.6			5.5	5.9	
4-18	0.54	0.057	9	0.6	-	1.50	1.75		0.99		25.0	17.0	0.053	0.052	6.7	7.1	
18-22	0.44	0.045	10	0.5	1	1.50	1.81		0.99		25.7	17.3	0.065	0.064	7.8	8.2	
22-28	0.28			0.5	1	1.53	1.71		0.99		23.7	17.2	0.038	0.037	7.8	8.4	
28-45	0.01			0.5	-	1.59	1.74		0.92		19.4	15.4	0.031	0.028	8.0	8.7	
45-57+	-			0.5	-	1.73	1.86		0.92		15.5	14.3	0.025	0.022	8.2	8.7	

Depth (In.)	Extractable bases 6N2a* Ca	6O2a* Mg	6P2a* Na	5B1a 6Q2a* K	Sum* of bases	6H2a* Ext. Acidity	Cation Exch. Capacity 5A2a* NaOAc	5A3a* Sum	Water extract from saturated paste 8A1 ** 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0- 4	17.2	12.5	0.6	1.0	31.3	4.5	34.7	35.8	2.1	2.1	2.3	0.3		-	3.6	2.4		0.65
4-18	23.2	14.6	1.4	0.3	39.5	1.5	41.7	41.0	1.2	0.7	3.0	tr.		-	3.4	1.4		0.52
18-22	31.9	14.7	2.5	0.3	49.4	-	44.7	49.4	0.7	0.8	4.7	tr.		-	4.5	0.4		0.55
22-28	36.4	17.2	3.9	0.2	57.7	-	53.3	57.7	0.6	0.8	5.5	tr.		-	4.8	0.5		0.62
28-45	38.9	23.3	7.6	0.1	69.9	1.4	65.7	71.3	0.4	0.4	5.8	tr.		-	3.5	0.7		0.63
45-57+	42.1	20.3	6.6	0.1	69.1	0.2	64.0	69.3	0.7	0.3	6.1	0.1		-	3.8	0.8		0.70

Depth (In.)	8A Water at Saturation Pct.	5D2 Exchangeable Na Pct.	5F SAR	Gypsum Pct.	Base Sat. Pct.	Clay mineralogy (<0.002 mm) 7A Relative abundance based on height of diagnostic X-ray peak 7A2: Mont.	Verm.	Intev.	Mica	Kaol.	% Kaolinite (DTA)	*** % Mont. & Verm.	Charge density meq x10³/m²	7D1 Ethylene glycol retention m²/gram
						X-ray								
0- 4	54.5	2	2		87									
4-18	61.0	3	3		96	xxxx	-	t	-	x	-	25	2.04	200
18-22	58.0	5	5		100									
22-28	51.9	7	7		100									
28-45	49.0	11	9		98									
45-57+	48.7	10	9		100	xxxx	x	t	-	-	-	34	2.03	262

1/, *, and **: Same as for Bonsall 37-5

*** Mehra and Jackson 1959 Constancy of the sum of mica unit cell potassium surface and interlayer sorption surface in vermiculite-illite clays. Soil Sci. Soc. Amer. Proc. 23:101-105.

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

BOSANKO CLAY
S64Calif-37-8

Location: San Diego County, California. A soil profile pit located 120 feet east of the center line of Camp Pendleton Road and 150 feet north, 200 feet west of the southeast 1/4 corner of the northeast 1/4 of section 33, T. 10 S., R. 4 W. Date of Sampling: March 11, 1964.
Description By: Roy H. Bowman and Gerald Kester. Collectors: Reuben E. Nelson, George Borst, Gerald Kester, Lothair Grant, Roy H. Bowman and Donald Markewich.
Classification: Chromic Pelloxerert, fine, montmorillonitic, thermic.
Vegetation: Wild oats, mustard. Climate: Annual precipitation is 13 inches and mean annual temperature is 59° F. Parent Material: Tonalite grus. Topography: West-facing, smooth 4 percent slopes on rolling upland. Elevation: 145 feet above sea level. Drainage: Well drained; medium to rapid runoff; slowly permeable. Soil Moisture: Slightly moist.

HORIZON	DESCRIPTION
Ap RSL No. 6450	0 to 4 inches, dark gray (10YR 4/1) clay, very dark gray (10YR 3/1) moist; moderate coarse to fine granular structure; very hard, firm, sticky and plastic; abundant very fine and fine roots; many fine tubular and many fine to medium interstitial pores; slightly acid (pH 6.2); abrupt smooth boundary.
A11 RSL No. 6451	4 to 18 inches, dark gray (10YR 4/1) clay, very dark gray (10YR 3/1) moist; moderate medium angular blocky structure; extremely hard, very firm, sticky and plastic; few very fine and fine roots; common very fine and fine tubular pores; few slickensides; slightly acid (pH 6.5); abrupt wavy boundary.
A12 RSL No. 6452	18 to 22 inches, grayish brown (10YR 5/2) clay, very dark grayish brown (10YR 3/2) moist; moderate medium angular blocky structure; extremely hard, very firm, sticky and plastic; very few, very fine roots; few very fine to fine tubular pores; few slickensides; few small soft masses of lime; mildly alkaline (pH 7.8), slightly effervescent; clear wavy boundary.
ACca RSL No. 6453	22 to 28 inches, grayish brown (10YR 5/2) clay, very dark grayish brown (10YR 3/2) moist; moderate medium angular blocky structure; very hard, very firm, sticky and plastic; very few very fine roots; very few very fine tubular pores; moderately alkaline (pH 8.0), slightly to strongly effervescent; many medium to small soft

SCS-420
10-64

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Chromic Pelloxerert, fine, montmorillonitic, thermic

SOIL Bosanko clay SOIL Nos. S64Calif-37-9 LOCATION San Diego County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 6456 - 6460

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Int. II (0.2-0.02)	(2-0.1)			1/ Coarse fragments 1A2a > 2 Pct.	2 - 19 (Pct. of < 76 mm)	19 - 76 (Pct. of < 76 mm)
		Size class and particle diameter (mm) 3A1																
		Pct. of < 2 mm																
0- 5	Ap	50.5	28.9	20.6	4.9	10.8	8.5	15.6	10.7	13.8	15.1	33.1	39.8			4	4	0
5-18	A11	39.0	24.0	37.0	3.9	8.5	6.3	12.0	8.3	10.8	13.2	25.6	30.7			9	9	0
18-23	A12	43.4	19.4	37.2	7.3	10.7	6.7	11.7	7.0	9.2	10.2	22.6	36.4			11	11	0
23-30	ACca	61.5	11.7	26.8	13.1	20.4	9.2	13.8	5.0	3.9	7.8	15.4	56.5			14	14	0
30-48+	C1	75.8	9.7	14.5	16.5	26.7	10.5	16.4	5.7	2.8	6.9	16.4	70.1			22	22	0

Depth (In.)	6A1a Organic carbon Pct.	6B1a Nitrogen Pct.	C/N	6C2a Ext. Iron as Fe Pct.	6E1b Carbonate as $CaCO_3$ Pct.		Exch. Bases: Na meq /100 g	Exch. Bases: K meq /100 g	g/cc	3B2 C'	Water content: Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility COLEF in./in.	4D1 Extensibility COLE in./in.	pH: 8C1b Saturated Paste	pH: 8C1a H_2O 1:1	pH: 8C1a H_2O 1:10
0- 5	0.86	0.075	11	0.5			0.5	0.4					8.9			5.7	6.0	
5-18	0.34			0.6			2.1	0.2					16.0			6.6	7.0	
18-23	0.32			0.5			3.3	0.2					16.7			7.4	7.7	
23-30	0.12			0.4	1		5.3	0.1					16.7			7.6	8.0	
30-48+	0.02			0.6	<1		4.9	0.1					12.9			7.6	8.0	

Depth (In.)	Extractable bases: 6N2a Ca	6O2a Mg	6P2a Na	5B1a 6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A2a NaOAc	5A3a Sum	Water extract from saturated paste 8A1: 6N1a Ca	6O1a Mg	6P1a Na	6Q1a K		6I1a CO_3	6J1a HCO_3	6K1a Cl	SO_4	8A1a Electrical conductivity
	meq/100 g								meq/liter									mmho/cm
0- 5	7.4	6.9	0.6	0.4		3.3	18.6	18.5	0.9	1.2	2.4	0.1		-	2.2	1.5		0.46
5-18	18.5	10.4	2.4	0.2		2.0	33.4	33.2	1.3	1.2	6.2	0.1		-	2.6	5.9		0.95
18-23	22.3	10.1	4.2	0.2		1.1	38.4	37.0	3.1	4.2	14.2	0.1		-	2.0	18.4		2.33
23-30	31.2	12.0	6.6	0.1		0.8	46.0	49.4	8.4	8.7	24.6	0.1		-	2.0	35.6		4.08
30-48+	22.0	9.5	6.2	0.1		1.5	38.2	39.3	10.2	11.8	29.4	0.1		-	1.3	48.9		5.33

Depth (In.)	8A Water at Saturation Pct	5D2 Exchangeable Na Pct.	5F SAR	Base Sat. Cations Pct.	Base Sat. NaOAc CEC Pct.	Clay mineralogy (<0.002 mm) 7A: Relative abundance based on height of diagnostic X-ray peak 7A2: Mont.	Verm.	Int.	Mica	Kaol.
						X-ray				
0- 5	36.5	3	3	82	82					
5-18	50.8	6	5	94	93	xxxx	xx	-	-	t
18-23	59.2	9	7	97	93					
23-30	56.2	12	8	98	100+					
30-48+	44.9	13	9	96	99	xxx	xxx	-	-	t

1/ From characterization sample.

- = looked for but not found
t = trace
x = small
xx = moderate
xxx = abundant
xxxx = dominant

BOSANKO CLAY
S64Calif-37-9

Location: San Diego County, California. A soil profile pit located 3700 feet north of Haverford Road and 140 feet west of Lilac Road in the SW1/4 of the NE1/4 of assumed section 3, T. 13 S.

CONSECUTIVE PEDON NUMBER C 69 57 C H A R A C T E R I Z A T I O N D A T A

SOIL SERIES CROPLEY SOIL NO. S69 CALI. 36-26 SAMPLE NO. 9 302-9 303 SAN BERNARDINO AREA

SOIL FAMILY CHROMIC PELLOXERERT FINE,MONTMORILLONITIC,THERM RIVERSIDE SOIL SURVEY LABORATORY

SAMPLE NO.	DEPTH CM.	HORZ.	SIZE, CLASS AND PARTICLE DIAMETER (MM) 1B1B,3A1 TOTAL FRACTION 2- .05	SILT .05- .002	CLAY LT .002	VCS 2-1	CS 1- .05	MS .5- .25	FS .25- .10	VFS .1- .05	CSI .05- .02	FSI .02- .002	CO. 2- .2U	FINE LT .2U	CARBO NATE 3A1A	1A2A INT. II .2- .02	PCT. OF WHOLE SOIL	COARSE FRAGMENTS- I 75-	II 20-5	III 5-2 PCT. OF LT 75
			PERCENT OF LESS THAN 2MM																	
9 302	0- 20	AP	26.0	42.5	31.5	1.4	3.9	2.9	8.0	9.8	21.2	21.3				36.1	--A	--	--	--
9 303	58- 86	AC2	29.3	34.8	35.9	0.8	3.3	3.7	10.4	11.1	16.1	18.7				33.8	--A	--	--	--

SAMPLE NO.	DEPTH CM.	ORG.MATTER CAR- BON 6A1A -PERCENT--	NITRO GEN 6B1A	CARBONATE AS CACO3 LT 2 6E1B	19-2 6E1B	EXT. IRON FE 6C2B -PERCENT-	ATTERBERG L I M I T LL 4F2 PCT.	PL 4F1 PCT.NO.40	BULK DENSITY EST 1/3 BAR 4A1F	1/3 BAR 4A1F	DRY 4AH	CM 3B2	WATER CONTENT 1/10 BAR 4B1C	1/3 BAR 4B1C	15 BAR 4B2	EXTENSBLTY LEF 4D1	LE 4D1	PH H2O 1/1 8C1A	H2O 1/10 8C1A	CACL2 8C1E
									GRAMS PER CC				----PERCENT----			-PERCENT--				
9 302	0- 20	2.69	0.242							1.15	1.48	1.00		33.2	17.5	8.8	8.8	7.2		6.9
9 303	58- 86	0.76	0.078							1.27	1.59	1.00		34.1	18.5	7.8	7.8	7.5		7.0

SAMPLE NO.	DEPTH CM.	EXTRACTABLE BASES CA 5B4A 6N2E	MG 5B4A 6O2D	NA 5B4A 6P2A	K 5B4A 6Q2A	SUM	BASE SAT. PCT.	C E C NA- DAC 5A2A	NH4- OAC 5A6A	EC 8A1A	WATER EXTRACT FROM SATURATED PASTE CA 6N1B	MG 6O1B	NA 6P1A	K 6Q1A	CO3 6I1A	HCO3 6J1A	CL 6K1A	SO4 6L1B	B 6T1A	PH SAT. PASTE 8C1B
		-MEQ/100G						-MEQ/100G		MMHO	-MEQ/1000 ML								PPM	
9 302	0- 20	25.1	6.6	0.3	2.0	34.0	100+		30.9											
9 303	58- 86	27.1	7.0	1.4	0.6	36.1	100+		33.9											

SAMPLE NO.	DEPTH CM.	WATER AT SAT. 8A PCT.	EXCHANGBLE NA 5B4B MEQ.	KA 5D2 PCT.	SAR 5E	GYPSUM- AMT. 6F1B PERCENT	REQ. 6P3A	CLAY MINERALOGY LESS THAN 0.002 MM X RAY I 7A2E	II 7A2E	III 7A2E	DTA IV 7A3	PCT.	L.T. .0002 MM X RAY I 7A2E	II 7A2E	III 7A2E	MINERAL CODE	RELATIVE AMOUNT
9 302	0- 20		0.3	1				MI 5	MM 3							MI MICA	1 TRACE
9 303	58- 86		1.4	4				MT 4	MI 3							MM MT/MICA	2 SMALL
																MT MONTMORT	3 MODERATE
																	4 ABUNDANT
																	5 DOMINANT
																	6 INDETER.

SAMPLE PREPARATION CODES
A=STANDARD PREPARATION.

CROPLEY CLAY LOAM
S69Calif-36-26

Location: San Bernardino County, California; about 1/2 mile west of Highway 71, and about 45 feet north of Eucalyptus Avenue, SW¼, SW¼, SE¼ of sec. 16, T.2S., R.8W. Date of Sampling: June 11, 1969.
Description by: A. A. Knecht, W. Z. Brock. Collectors: A. A. Knecht, W. Z. Brock.
Classification: Chromic Pelloxerert, fine, montmorillonitic, thermic.
Vegetation: Small grain stubble and annual grasses. Climate: Average annual precipitation about 16 inches. Mean temperature about 61°F. Parent Material: Mixed alluvium. Topography: Alluvial fan, gently sloping, about 2 percent slopes. Elevation: 700 feet. Drainage: Moderately well drained; slow runoff, slow permeability. This area is above the flood plain and the water table is below five feet. Soil Moisture: This profile was dry down to 34 inches and moist below.
Remarks: Field reaction was determined by the Truog method. Two horizons (0-8" and 23-34") were analyzed in the Riverside Soil Survey Laboratory.

HORIZON	DESCRIPTION
	(0-20 cm.)
Ap	0 to 8 inches, dark gray (10YR 4/1) clay loam, black (10YR 2/1) moist; strong [illegible]
69302	very plastic; many very fine and fine roots; many very fine and fine vertical [illegible]
	(20-58 cm.)
AC1	8 to 23 inches, dark gray (10YR 4/1) clay loam, black (10YR 2/1) moist; moderate very coarse angular blocky structure; extremely hard, very firm, very sticky, very plastic; many very fine roots; many very fine and fine random interstitial pores; neutral (pH 7.0); gradual wavy boundary.
	(58-86 cm.)
AC2 RSL 69303	23 to 34 inches, dark grayish brown (10YR 4/2) clay loam, very dark grayish brown (10YR 3/2) moist; moderate very coarse prismatic structure; extremely hard, extremely firm, very sticky, very plastic; common very fine roots; common very fine and fine discontinuous tubular pores; intersecting slickensides; moderately alka-

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Chromic Pelloxerert, fine, montmorillonitic, thermic

SOIL Diablo clay SOIL Nos. S58Calif-1-1 LOCATION Alameda County, California

SOIL SURVEY LABORATORY Beltsville, Maryland LAB. Nos. 591000 - 591006

Depth (In.)	Horizon	1B1b Total Sand (2-0.05)	Total Silt (0.05-0.002)	Total Clay (< 0.002)	Sand: Very coarse (2-1)	Sand: Coarse (1-0.5)	Sand: Medium (0.5-0.25)	Sand: Fine (0.25-0.1)	Sand: Very fine (0.1-0.05)	Silt: 0.05-0.02	Silt: Int. III (0.02-0.002)	Silt: Int. II (0.2-0.02)	(2-0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	Coarse fragments 1A2c: 2A2 > 2 Pct.	2 - 19	19 - 76
		Pct. of < 2 mm															Pct. of < 76mm	
0-6	A11	12.1	42.0	45.9	0.0	0.1	0.2	3.3	8.5	14.1	27.9	25.2	3.6			<1		
6-15	A12	11.1	41.5	47.4	0.0	0.1	0.3	3.3	7.4	12.7	28.8	22.6	3.7			<1		
15-26	A13ca	9.3	43.7	47.0	0.1	0.1	0.2	2.6	6.3	11.5	32.2	19.8	3.0			<1		
26-32	A14ca	8.8	43.1	48.1	0.0	0.1	0.2	2.5	6.0	9.7	33.4	17.6	2.8			<1		
32-42	ACca	7.8	48.3	43.9	0.2	0.5	0.4	2.1	4.6	9.5	38.8	15.7	3.2			<1		
42-50	C2	6.8	56.0	37.2	0.4	0.6	0.4	1.1	4.3	16.3	39.7	21.5	2.5			<1		
50+	Dr	18.0	63.7	18.3	0.2	0.3	0.2	0.9	16.4	30.7	33.0	47.8	1.6					

Depth (In.)	6A1a Organic carbon Pct	6B2a Nitrogen Pct.	C/N	6C2b Ext. Iron as Fe Pct.	6E1e Carbonate as $CaCO_3$ Pct.		Bulk density: 4A1f 1/3 bar g/cc	Bulk density: 4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content: Pct.	Water content: 4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility LEf Pct	4D1 Extensibility LE Pct.	pH: 8C1a H_2O 1:1		
0-6	1.37	0.172	8													6.4		
6-15	0.92	0.135	7		1											7.4		
15-26	0.71	0.105	7		3											7.6		
26-32	0.64	0.102	6		4											7.7		
32-42	0.41	0.091	5		8											7.8		
42-50	0.25				8											7.9		
50+	0.10				4											8.0		

Depth (In.)	Extractable bases 5A1: 6N2b Ca	6O2b Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity: 5A1b NH_4OAc	5A3a Sum		6G1e KCl Ext. Al^{+++}	CEC 100 g clay					Base saturation: 5C3 Sum Pct.	Pct.
	meq/100 g																
0-6	22.9	9.0	0.1	1.0		3.1	32.6	36.2			71.0					91	
6-15	*		0.1	0.9			32.0				67.5						
15-26	*		0.2	0.9			30.5				64.9						
26-32	*		0.3	0.8			31.6				65.7						
32-42	*		0.3	0.8			26.5				60.4						
42-50	*		0.4	0.7			26.8				72.0						
50+	*		0.3	0.6			19.8				108.2						

Depth (In.)		Clay mineralogy (<0.002 mm.) 3A1: Mont.	Verm.	Mica 7A2	Kaolin.	
		X-ray				

* Calcareous

DIABLO CLAY
S58Calif-1-1

Location: 1325 feet east and 275 feet north of SW corner, Section 25, in SW1/4, SE1/4, SW1/4, Section 25, R. 1 E., T. 2 S., Bailey Ranch, Alameda County, California. Date of Sampling: November 10, 1958.

Description by: R. Ulrich. Collectors: R. E. Nelson, Larry Welch, Robert Mitchel, Leo Mayers, Lynn Whittig, R. Ulrich.

Classification: Chromic Pelloxerert, fine, montmorillonitic, thermic.

Vegetation: Annual brome grasses (soft chess and ripgut mostly), mouse barley, wild oats, poverty fescue, bur clover, star thistle, bull thistle, and mustard. Present use is annual grass range; may have been cultivated sometime in past. Climate: Rainfall is 14-15 inches (estimated); mean annual soil temperature about 60° F. Distinct summer-dry, winter-moist Mediterranean type climate. Parent Material: Interbedded calcareous fine grained sandstone and shale. Lime content is variable

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Chromic Pelloxerert, fine, montmorillonitic, thermic
SOIL Diablo clay SOIL Nos. S58Calif-1-2 LOCATION Alameda County, California

SOIL SURVEY LABORATORY Beltsville, Maryland LAB. Nos. 591007 - 591012

Depth (in.)	Horizon	1B1b Total Sand (2–0.05)	Total Silt (0.05–0.002)	Total Clay (< 0.002)	Sand: Very coarse (2–1)	Sand: Coarse (1–0.5)	Sand: Medium (0.5–0.25)	Sand: Fine (0.25–0.1)	Sand: Very fine (0.1–0.05)	Silt: 0.05–0.02	Silt: Int. III (0.02–0.002)	Int. II (0.2–0.02)	(2–0.1)	3A1a Carbonate clay	3A1a Non-carbonate clay	2A2 Coarse fragments > 2 Pct.	1A2a 2 - 19	19 - 76
		Pct. of < 2 mm															Pct. of < 76mm	
0-7	A11	18.4	42.7	38.9	0.0	0.2	0.6	6.0	11.6	14.3	28.4	30.3	6.8			<1		
7-17	A12	18.0	42.1	39.9	0.0	0.1	0.6	5.8	11.5	13.3	28.8	29.3	6.5			<1		
17-28	A13ca	11.4	48.5	40.1	0.1	0.2	0.5	3.7	6.9	10.6	37.9	20.3	4.5			<1		
28-37	Cca1	12.6	48.8	38.6	0.2	0.2	0.4	3.0	8.8	9.9	38.9	20.9	3.8			<1		
37-48	Cca2	24.7	50.7	24.6	0.0	0.1	0.2	3.7	20.7	20.3	30.4	44.1	4.0			<1		
48+	Dr	45.6	46.8	7.6	0.0	0.1	0.4	8.3	36.8	24.4	22.4	65.8	8.8			<1		

Depth (in.)	6A1a Organic carbon Pct.	6B2a Nitrogen Pct.	C/N	6C2b Ext. Iron as Fe Pct.	6E1e Carbonates $CaCO_3$ Pct.		Bulk density 4A1f 1/3 bar g/cc	4A1h Oven dry g/cc	g/cc	3B2 Cm	Water content Pct.	4B1c 1/3 bar Pct.	4B2 15 bar Pct.	4D1 Extensibility LEf Pct	4D1 Extensibility LE Pct.	pH 8C1a H_2O 1:1		
0-7	2.16	0.222	10													6.4		
7-17	1.04	0.145	7													6.9		
17-28	0.74	0.115	6		4											7.6		
28-37	0.54	0.091	6		6											7.7		
37-48	0.37	0.070	5		16											7.8		
48+	0.08				5											7.9		

Depth (in.)	Extractable bases 5A1 6N2b Ca	6O2b Mg	6P2a Na	6Q2a K	Sum of bases	6H2a Ext. Acidity	Cation Exch. Capacity 5A1b NH_4OAc	5A3a Sum		6G1e KCl Ext. Al^{+++}	CEC 100g clay					Base saturation 5C3 Sum	
	meq/100 g															Pct.	Pct.
0-7	28.6	5.8	0.1	1.1		3.3	34.6	38.9			88.9					92	
7-17	32.5	4.6	0.1	0.8		1.8	33.8	39.8			84.7					95	
17-28	*		0.1	0.7			32.4				80.8						
28-37	*		0.2	0.7			32.4				83.9						
37-48	*		0.1	0.5			22.2				90.2						
48+	*		0.1	0.5			19.3				253.9						

Depth (in.)	Clay mineralogy (<0.002 mm.) 3A1 Mont.	Verm.	Mica 7A2	Kaolin.	
	X-ray				

* Calcareous

DIABLO CLAY
S58Calif-1-2

Location: 1300 feet east and 1150 feet south of NW corner of Section 24, in SE1/4, NW1/4, NW1/4, Section 24, R. 1 E., T. 2 S., Beck Ranch, Alameda County, California.
Date of Sampling: November 11, 1958.
Description by: R. Ulrich. Collectors: R. E. Nelson, Larry Welch, Robert Mitchel, R. Ulrich.
Classification: Chromic Pelloxerert, fine, montmorillonitic, thermic.
Vegetation: Annual brome grasses (soft chess and ripgut mostly), annual rye, wild oats, bur clover, mustard, dock and other weeds. Present use is annual grass range; may have been cultivated sometime in the past. Climate: Rainfall is 14-15 inches (estimated); mean annual soil temperature about 60° F. Distinct summer-dry, winter-moist Mediterranean type climate. Parent Material: Interbedded calcareous fine grained sandstone and shale. Lime content is variable but it is usually weak or slightly calcareous although thin seams of limestone and other strongly calcareous sediments

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Diablo clay **LOCATION** Sonoma County, California

SOIL FAMILY: Chromic Pelloxerert, fine, montmorillonitic, thermic.

SOIL NOS. S61Calif-49-11 **LAB. NOS.** 61463 - 61469

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent)								3A1	2A2	TEXTURAL CLASS
		VERY COARSE SAND	COARSE SAND	MEDIUM SAND	FINE SAND	VERY FINE SAND	SILT	CLAY				

DIABLO CLAY
S61Calif-49-11

Location: Sonoma County, California. NW 1/4, NW1/4, section 14, T. 5 N., R. 7. W. projected; also starting from the center of the town of Petaluma find the point due northeast 3 1/4 miles where East Washington Street forms a "T" junction with Adobe Road. The site is 6000 feet northerly of this point on a line bearing 5° east of true north on a sloping hillside of aspect southeast. It is 1120 feet due southeast of the homestead (Beffa's milk barn). (Site spotted on photo No. CSH-1BB-165.) Date of Sampling: June 27, 1961.
Collectors: Reuben Nelson, Leo C. Klameth, V. C. Miller, G. Lambert, J. H. Kashiwagi, F. Graham.
Classification: **Chromic Pelloxerert, fine, montmorillonitic, thermic.**

Vegetation: Natural before cultivation - annual grasses and forbs, few scattered oaks. Use: Oat hay production. Climate: Subhumid mesothermal. Annual precipitation 24.7 inches. Mean January temperature 46.7° F., mean July temperature 66.3° F., mean annual temperature 57.5° F. Parent Material: Soft sandstone and shale of the Petaluma formation. Elevation: 340 feet. Relief: Smoothly rolling hills on 13 percent slope with southeast aspect. Drainage: Moderately well drained. Biological Activity: No worm or rodent activity apparent at this time of year.

HORIZON	DESCRIPTION
Ap RSL No. 61463	0 to 7 inches, gray (N 5/) clay, black (10YR 2/1) or very dark gray (10YR 3/1) when moist; granular; very hard when dry, extremely firm when moist; very sticky and very plastic when wet; abundant very fine roots; many very fine interstitial pores; im- [illegible]

SOIL SURVEY LABORATORY Riverside, California

SOIL TYPE Diablo clay **LOCATION** Sonoma County, California

SOIL FAMILY: Chromic Pellox rert, fine, montmorillonitic, thermic.

SOIL NOS. S61Calif-49-12 **LAB. NOS.** 61470 - 61477

DEPTH INCHES	HORIZON	1B1b PARTICLE SIZE DISTRIBUTION (in mm.) (per cent) 3A1									2A2	TEXTURAL CLASS
		VERY COARSE	COARSE	MEDIUM	FINE	VERY FINE						

DIABLO CLAY
S61Calif-49-12

Location: Sonoma County, California. Sartori Ranch - 1600 feet west and 1000 feet south of northerly most water tank on Manor Lane. SW1/4, NW1/4, section 13, T. 5 N., R. 7 W.
Date of Sampling: June 28, 1961.
Collectors: Reuben Nelson, Leo C. Klameth, V. C. Miller, G. Lambert, J. H. Kashiwagi, F. Graham.
Classification: **Chromic Pelloxerert, fine, montmorillonitic, thermic.**

Vegetation: Natural before cultivation - annual grasses and forbs and a few scattered oaks. Use: Oat hay production. Climate: Subhumid mesothermal. Annual precipitation 24.7 inches, mean January temperature 46.7° F., mean July temperature 66.3° F., mean annual temperature 57.5° F. Parent Material: Soft sandstone and shale of the Petaluma formation. Elevation: 400 feet. Relief: Smoothly rolling hills on 14 percent slope with southeast aspect. Drainage: Moderately well. Biological Activity: No worm or rodent activity apparent at this time of year.

HORIZON	DESCRIPTION
Ap RSL No. 61470	0 to 7 inches, dark gray (N 4/) clay, very dark gray (10YR 3/1) when moist, surface quarter inch is strong very fine subangular blocky, remainder of horizon weak coarse subangular blocky structure; very hard when dry, firm when moist, very sticky and very plastic when wet; abundant very fine and fine roots; many very fine and fine interstitial pores; slightly acid reaction, pH 6.5; abrupt, wavy boundary.
A11 RSL No. 61471	7 to 19 inches, very dark gray (N 3/) clay with common fine distinct mottles of light gray (10YR 7/1) when dry, black (10YR 2/1) with common fine distinct mottles of gray (10YR 5/1) when moist; massive; extremely hard when dry, extremely firm when moist; very sticky and very plastic when wet; abundant very fine and fine roots; common very fine tubular pores; few small scattered lime concretions and blotches; common slickensides; slightly acid reaction, pH 6.5; gradual, smooth boundary.
A12 RSL No.	19 to 30 inches, dark gray (5Y 4/1) clay, very dark grayish brown (2.5Y 3/2) when moist; massive breaking to weak fine subangular blocks; extremely hard when dry; very firm

U. S. DEPARTMENT OF AGRICULTURE
SOIL CONSERVATION SERVICE

SOIL FAMILY: Chromic Pelloxerert, fine, montmorillonitic, thermic

SOIL Diablo clay SOIL Nos. S65Calif-27-10 LOCATION Monterey County, California

SOIL SURVEY LABORATORY Riverside, California LAB. Nos. 65443 - 65448

		1B1b	Size class and particle diameter (mm) 3A1

DIABLO CLAY
S65Calif-27-10

Location: Monterey County, California. East 1/4 line of Sec. 25, T20S, R9E, about 6 miles east of San Lucas off Highway 198. (SE corner of SE$\frac{1}{4}$ of NE$\frac{1}{4}$.) Date of Sampling: May 19, 1965.
Description by: Lewis C. Leifer. Collectors: W. D. Nettleton and Benny R. Brasher; C. J. Weisel, R. D. Waggoner, F. J. Pecarich. Site selection by T. D. Cook.
Classification: **Chromic Pelloxerert, fine, montmorillonitic, thermic. (Grumusol).**
Vegetation: Range - soft chess, wild oats, ripgut, brome, filaree, annual fescue.
Climate: Precipitation 10 inches; mean annual temperature 62°F., mean January temperature 59°F., mean July temperature 68°F. Parent Material: Shale, Plio-Pleistocene, non-marine. Topography: Moderately steep slopes, locally slightly concave.

Drainage: Well drained, permeability slow, runoff medium. Soil Moisture: Very slightly moist to 18 inches and slightly moist to about 40 inches, very nearly dry below, mostly wilting point and below.
Remarks: Field determinations of pH by Truog Kit. Classification assumes intersecting


CALIFORNIA
10 0 10 20 30 40 50
OREGON
NEVADA
ARIZONA
MEXICO
PACIFIC
OCEAN
DEL NORTE
SISKIYOU
MODOC
HUMBOLDT
TRINITY
SHASTA
LASSEN
TEHAMA
PLUMAS
MENDOCINO
GLENN
BUTTE
SIERRA
NEVADA
YUBA
LAKE
COLUSA
SUTTER
LACER
YOLO
ELDORADO
SONOMA
NAPA
SACRAMENTO
ALPINE
AMADOR
SOLANO
CALAVERAS
MARIN
SAN JOAQUIN
TUOLUMNE
MONO
CONTRA COSTA
SAN FRANCISCO
OAKLAND
ALAMEDA
STANISLAUS
MARIPOSA
MERCED
SAN MATEO
SANTA CLARA
MADERA
SANTA CRUZ
FRESNO
SAN BENITO
MONTEREY
TULARE
KINGS
SAN LUIS OBISPO
KERN
SAN BERNADINO
SANTA BARBARA
VENTURA
LOS ANGELES
ORANGE
RIVERSIDE
SAN DIEGO
IMPERIAL
COLORADO RIVER
Crescent City
Yreka
Alturas
Eureka
Weaverville
Redding
Susanville
Red Bluff
Quincy
Willows
Oroville
Downieville
Nevada City
Colusa
Marysville
Yuba City
Lakeport
Ukiah
Auburn
Woodland
Placerville
Santa Rosa
Napa
Jackson
San Andreas
Bridgeport
Vallejo
Stockton
Sonora
San Rafael
Martinez
Mono Lake
Modesto
Mariposa
Redwood City
San Jose
Merced
Santa Cruz
Hollister
Madera
Fresno
Independence
Salinas
Visalia
Hanford
Bakersfield
San Luis Obispo
San Bernardino
Santa Barbara
Ventura
Riverside
Santa Ana
Sutton Sea
El Centro
SAN MIGUEL I.
SANTA ROSA I.
SANTA CRUZ I.
SANTA CATALINA I.
SAN NICHOLAS I.
SAN CLEMENTE
SAN DIEGO
Lake Almanor
Eagle Lake
Lake Tahoe
Mono Lake
Goose Lake
Upper Lake
Middle Alkali Lake
Lower Lake
CLEAR LAKE RESERVOIR
Klamath River
Trinity River
Eel River
Sacramento River
Sacramento
Joaquin
Tuolumne
Mariposa
Merced
Fresno
Kings
Colorado River
124°
123°
122°
121°
120°
119°
118°
117°
116°
115°
42°
41°
40°
39°
38°
37°
36°
35°
34°
33°